

Inflorescences of *Fargesia angustissima* T.P. Yi and *Yushania pauciramificans* T.P. Yi (Poaceae, Bambusoideae) shed light on the taxonomy of the Sino-Himalayan alpine bamboos

Xia-Ying Ye^{1,2}, Zu-Chang Xu², Yue-Hong Cheng³, Wei-Hua Wang¹, De-Zhu Li²

1 Agronomy and Life Science Department, Zhaotong University, Zhaotong, Yunnan 657000, China **2** Germplasm Bank of Wild Species, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming, Kunming, Yunnan 650201, China **3** Sichuan Wolong National Natural Reserve Administration, Aba, Sichuan, 623000, China

Corresponding author: De-Zhu Li (dzl@mail.kib.ac.cn)

Academic editor: Eduardo Ruiz-Sanchez | Received 25 August 2022 | Accepted 10 October 2022 | Published 9 December 2022

Citation: Ye X-Y, Xu Z-C, Cheng Y-H, Wang W-H, Li D-Z (2022) Inflorescences of *Fargesia angustissima* T.P. Yi and *Yushania pauciramificans* T.P. Yi (Poaceae, Bambusoideae) shed light on the taxonomy of the Sino-Himalayan alpine bamboos. *PhytoKeys* 215: 27–36. <https://doi.org/10.3897/phytokeys.215.94010>

Abstract

The taxonomy of the Sino-Himalayan alpine bamboos is controversial due to their complex evolutionary history and further complicated by the scarcity of inflorescence. Here, we supplement the description of the inflorescence of *Fargesia angustissima* T.P. Yi and *Yushania pauciramificans* T.P. Yi, which shed light on the taxonomy of *Fargesia* Franchet, *Borinda* Stapleton and *Yushania* Keng. *F. angustissima* has compressed inflorescence unilateral stretching out from reduced spathe, showing a transitional state between species with condensed inflorescence embraced by spathe-like bracts and species with open inflorescence without bracts. Considering that extensive gene flow existed between several clades of *Fargesia* found in recent studies, a broadly-defined *Fargesia* s. l. should be adopted. Meanwhile, the inflorescence of *Y. pauciramificans* has typical characteristics of *Yushania*, such as axilla with tuberculate glands, rachilla internodes ciliate and cylindrical florets, supporting the delimitation of *Yushania*.

Keywords

Borinda, *Fargesia*, inflorescence, reproductive characters, *Yushania*

Introduction

Although the taxonomy of bamboos has entered a new stage since the proliferation of molecular phylogenetic and phylogenomic studies (e.g., Bamboo Phylogeny Group 2012; Attigala et al. 2016; Zhang et al. 2020; Ye et al. 2021b), morphological characters play an important role in the naming and identification of the species. Reproductive characters are traditionally assumed to be critical in bamboo evolution and taxonomy, especially at the generic level (Li et al. 2006). However, due to variable blooming intervals, ranging from a couple of decades up to 120 years (Janzen 1976), many bamboo species were described without inflorescence information (Yi 1986, 1988a; Stapleton 1994; Keng and Wang 1996; Ohrnberger 1999; Li et al. 2006). This has caused the confusion in the definition of some genera, including some Sino-Himalayan alpine bamboos, e.g., *Fargesia* Franchet, *Yushania* Keng f. and *Borinda* Stapleton (Guo et al. 2002; Stapleton 2021).

Fargesia was delimited as having short-necked pachymorph rhizome with uni-caespitose culms and compressed inflorescence subtended by several small or large spathes, while *Yushania* has long-necked rhizome with diffuse culms and open inflorescence without bracts (Li et al. 2006; Yi et al. 2008; Shi et al. 2022). *Borinda* was described as clumping temperate bamboos, similar to *Yushania* for its inflorescence with reduced bracts, and *Fargesia* for its short-necked pachymorph rhizome (Stapleton 1994). Moreover, most species of *Borinda* were transferred from *Fargesia* (Stapleton 1998, 2006, 2021). Although morphological differentiation of rhizome and inflorescence has been used to distinguish these genera, the intermediate state of rhizome and inflorescence between them makes the genus delimitation very ambiguous. As a result, the bamboo accounts of the “Flora of China” recognize two genera, i.e., *Fargesia* and *Yushania* (Li et al. 2006).

In our recent molecular analyses of this taxonomically difficult group based on the double digest-restriction-site associated DNA sequencing (ddRAD) analyses (Ye et al. 2019), *Yushania* was resolved as a well-supported monophyletic lineage, demonstrating the phylogenetic importance of the rhizome type. If considering the rhizome type alone, *Fargesia yunnanensis* Hsueh & T.P. Yi needs to be transferred into *Yushania* for its generally long rhizome neck (12–35 cm). Actually, this species was nested in the ‘Fargesia1’ clade in the analysis of Ye et al. (2019) with high support. Therefore, more information on inflorescence knowledge should be provided for the delimitation of *Yushania*.

Fargesia was resolved as a polyphyly (Wang et al. 2017; Zhang et al. 2019; Zhou et al. 2019; Zhou et al. 2020; Ye et al. 2021a) and divided into several clades with high support in the recently ddRAD analyses (Ye et al. 2019). Stapleton (2021) transferred species in the ‘Fargesia3’ + ‘Fargesia4’ + *F. angustissima* clade of Ye et al. (2019) and several species sampled by Zhang et al. (2019) into *Borinda* based on the molecular phylogenetics and some floral and vegetative characteristics. Concurrently, Stapleton (2021) considered that *Fargesia* s. s. possesses tightly unilateral racemes and only distributed along the Qinling Mountains. In this case, most of the species

originally described in *Fargesia* could not be retained in this genus. Nevertheless, several species which were transferred into *Borinda* shared the floral characteristics of *Fargesia* s. s., with raceme enclosing by spathe-like sheaths and protruding from unilateral side, such as *F. edulis* Hsueh & T.P. Yi and *F. adpressa* T.P. Yi (Li et al. 2006; Shi et al. 2022). This indicates that limited reproductive characters cannot distinguish *Borinda* from *Fargesia* appropriately. Therefore, more knowledge of reproductive features should be provided to improve our understanding of the relationship of *Borinda*, *Fargesia* and *Yushania*.

In recent field surveys, we collected the floral and vegetative specimens for two bamboo species. A supplementary description of the inflorescence of these two species is presented here, providing new information on the delimitation of alpine bamboos.

Materials and methods

We collected two specimens with both floral and vegetative organs during our field work in Yunnan (*YXY2020023*) and Sichuan (*WL2021001*), China. Morphological studies were based on the living individuals in the field, specimens, and literature (Yi 1985, 1988a, b; Keng and Wang 1996; Li et al. 2006; Shi et al. 2022). Flowering and fruit materials were dissected under an OLYMPUS DP80 digital microscope at Germplasm Bank of Wild Species of the Kunming Institute of Botany. The morphological terminology follows McClure (McClure 1966).

Results

According to our observation and comparison of the type specimens and original literature (Yi 1985, 1988b) and bamboo accounts of “Flora Reipublicae Popularis Sini-cae” (Keng and Wang 1996), “Flora of China” (Li et al. 2006) and “Illustrated Flora of Bambusoideae in China” (Shi et al. 2022), we identified specimen *YXY2020023* to be *Yushania pauciramificans* T.P. Yi based on paniculate inflorescence on terminating leafy branches without spathes subtending, pachymorph rhizomes with long neck (20–50 cm in length), culms 2–3.2 m, internodes terete, branches 1–3 at lower nodes, 5–6 at upper, culm sheaths cartilaginous, with erect gray setae and absent auricles. *WL2021001* was identified to be *Fargesia angustissima* T.P. Yi according to panicles on terminating leafy branches subtended by slightly expanded bracts, pachymorph rhizomes with short neck (2–5 cm in length), unicaespitose culms with fine ridged internodes, culm node less prominent than sheath scar, culm sheaths persistent, which were longer than internodes, narrowly triangular, apically papery, linear, and narrowed for distal 1/3–1/2 of length, sparsely brown setulose, leaf blade abaxially proximally white-gray pubescent. All voucher specimens were deposited at Kunming Institute of Botany, Chinese Academy of Sciences (KUN), and epitype of these two species are also designated here (Turland et al. 2018).

Taxonomic treatment

Fargesia angustissima T.P. Yi

“油竹子” (You Zhu Zi)

Fig. 1

Fargesia angustissima T.P. Yi in J. Bamboo Res. 4(2): 21–22. pl. 4. 1985; Keng f. & Z. P. Wang, Fl. Reipubl. Poppularis. Sin. 9(1): 437. pl. 50, 1–8. 1996; D. Z. Li and Stapleton in Z. Y. Wu, P. H. Raven & D. Y. Hong, Fl. China 22: 85–86. 2006. L. B. Zhang in C. Y. Wu, P. H. Raven & H. Y. Hong, Fl. China Illustr. 22: 110. pl. 110: 1–7. 2007. ‘Type’: CHINA. Sichuan: Wenchuan County, Genda Township, 1550 m alt., live on limestone slope, 22 Sept. 1974, T.P. Yi 74450 (holotype, SCFI!); *ibid.*, 31°4.27’N, 103°19.64’E, 1434 m alt., 23 Dec. 2021, WL2021001 (epitype designated here, KUN, 1546903!).

≡ *Yushania ferax* subsp. *angustissima* (T.P. Yi) Demoly in Bambou. Bull. A. E. B., Sect. France. 46: 8. 2005.

≡ *Borinda angustissima* (T.P. Yi) Stapleton in Sida. 22(1): 332, 2006.

Description. Culms densely unicaespitose, 4–7 m tall, 1–2 cm in diameter; internodes terete, 28–35 cm long, glabrous, initially white powdery, longitudinal ribs very prominent; culm walls 1.5–2.5 mm thick; sheath scars prominent. Buds oblong. Branches 5–10 per node, slender. Culm sheaths persistent or gradually deciduous, much longer than internodes, distantly papery and narrowly banded, abaxially sparsely brown setulose, margins initially densely ciliate; auricles absent; oral setae 3–5, 5–7 mm long; ligules ca. 1 mm tall; blade reflexed, linear, narrower than apex of sheath, margins usually serrulate, readily deciduous. Foliage leaves 3–5 per branchlet; auricles absent; oral setae 5–8, 2–3 mm long; ligules convex, ca. 0.5 mm tall, external ligule pubescent; blades (1.7) 3.4–9.5 × 0.3–0.7 cm, narrowly lanceolate, abaxially proximally pubescent, second veins 2 (3) pairs, transverse veins distinct.

Flowering branches 18–60 cm long, with secondary branches; raceme composed of 1–3 spikelets, open to dense, flowering branchlet with terminal leaves 1–2, gradually deciduous, subtended by slightly inflated foliage-leaf-like sheaths, initially stretching out from one side of the sheaths; axes terete, glabrous, usually with a bract at the base of pedicels, bract lanceolate, 5 mm long. Spikelet purple-green to dark purple, 2–4 cm long, 7–10 mm wide, clustered closely; pedicels slender, curved or undulate, glabrous, 7–15 mm long; florets 2–8, 1.5–2 cm long; rachilla internode 2 mm long, expanded and pubescent at apex. Glumes 2, papery, apex acuminate, the first one narrowly lanceolate, 9–12 mm long, 2 mm wide, pubescent, apically awned, ca. 3 mm long; the second one ovoid-lanceolate, 11–15 mm long, ca. 2 mm wide, pubescent, apically awned, ca. 5 mm long. Lemma papery, ovoid-lanceolate, 15–19 mm long, 4–5 mm wide, abaxially scabrous, densely white setose, apically awned, 5–6 mm long; palea shorter than lemma, thinly papery, 5–10 mm long, 2-keeled, sulcate between




Figure 1. *Fargesia angustissima* T.P. Yi **A** habitat, showing flowering population **B** branches and internode **C** clum **D** branchlet, showing foliage leaves **E** culm sheath **F** flowering branches **G** inflorescence **H** anatomy of florets **I** fruit.

keels, upper part of keels ciliate, apex obtuse. Lodicules 3, membranous, transparent, elliptical-triangular, margins ciliate. Stamens 3, filaments free, anthers yellow, gradually dark purple, ovary long-ovoid, glabrous. Stigmas 3, plumose. Caryopses oblong, dark brown, ventrally grooved, 6–9 mm long, ca. 1–2 mm in diameter, glabrous, apex with persistent style.

Phenology. New shoots May to August. Flowering December to April; fruiting May to June.

Distribution and habitat. *Fargesia angustissima* is known from Dujiangyan, Wenchuan and Chongzhou of western Sichuan, and Beichuan and Pingwu of northwestern Sichuan, and mainly occurs on the steep limestone slope or along the stream at an elevation of 800–1800 m.

Additional specimens examined. CHINA. Sichuan: Beichuan County, Caijiaping, near Xiaozhaizigou Nature Reserve, 09 Nov. 2017, Y. X. Zhang 17142 (KUN!).

***Yushania pauciramificans* T.P. Yi**

“少枝玉山竹” (Shao Zhi Yu Shan Zhu)

Fig. 2

Yushania pauciramificans T.P. Yi in Bull. Bot. Res. 8(4): 71–73. pl. 5. 1988; Keng f. & Z. P. Wang, Fl. Reipubl. Poppularis. Sin. 9(1): 547. pl. 164, 6–8. 1996; D. Z. Li and Stapleton in Z. Y. Wu, P. H. Raven & D. Y. Hong, Fl. China 22: 72. 2006. L. B. Zhang in C. Y. Wu, P. H. Raven & H. Y. Hong, Fl. China Illustr. 22: 96. pl. 96: 6–8. 2007. ‘Type’: CHINA. Yunnan: Xinping County, Ailao Mountain, Liangshan, 2510 m alt., under forest, 1 Sept. 1986, *T.P. Yi* 86237 (holotype, SCFI!); *ibid.*, Gasa Town, 23°57'N, 101°33.90'E, 2257 m alt., 28 May 2020, *YXY2020023* (epitype designated here, KUN, 1546904!).

Description. Culms diffuse, 1.5–3.5 m tall, 0.6–1.2 cm in diameter; internodes terete, 15–27 cm long, initially with a white powdery ring below nodes, glabrous; culm walls 2.5–3.5 mm thick, cavity small; nodes weakly prominent; sheath scar obviously prominent, woody. Branches 1–3 at lower nodes, ca. 5 at upper. Culm sheaths persistent, triangularly oblong, 2/5–1/2 as long as internodes, cartilaginous, gray setose abaxially, margins densely setose; auricles absent; oral setae 2–4, erect, deciduous; ligule 1–1.5 mm tall, glabrous; blades linear lanceolate, glabrous, reflexed. Foliage leaves 2–5 per branchlet; sheath margins glabrous; auricles absent; oral setae 5–7, slightly curved; ligule 0.5–1 mm tall; blades 5.2–16 × 1.1–2.8 cm, lanceolate or elliptic-lanceolate, base broadly cuneate or rounded, glabrous, secondary veins 4–6 pairs, transverse veins distinct.

Flowering branches 6–22 cm long, lower nodes with secondary flowering branchlets; inflorescence open, paniculate, terminal on leafy branches, composed of 5–15 spikelets, axilla with tuberculate glands, subtended by a small bract; axes terete, 2–10 cm long, glabrous. Spikelet dark purple, 3–6 cm long; pedicels slender, 1.3–3 cm long, usually slightly sinuous, glabrous; florets 2–5, 1.2–3 cm long, cylindrical, apical floret sterile and tubulose; rachilla internodes slightly flattened, ca. 5 mm long, gray white pubescent, apex more densely, margins gray white ciliate. Glumes 2, apically awned, ca. 1 mm, the first one narrowly lanceolate, 4–7 mm long, ca. 1 mm wide, distally white pubescent; the second one ovoid-lanceolate, 6–10 mm long, 1.5–2 mm wide,




Figure 2. *Yushania pauciramificans* T.P. Yi **A** habitat **B** branches **C** bud **D** culm sheath **E** branchlet, showing foliage leaves **F** flowering branches **G** inflorescence **H** spikelet **I** anatomy of florets.

distally white pubescent. Lemma mucronate, papery, 7–10 mm long, 3–4 mm wide, densely white setose, apically awned, ca. 1 mm; palea slightly shorter than lemma, thinly papery, 6–9 mm long, 2-keeled, densely pubescent, apex obtuse, 2-cleft. Lodicules 3, membranous, transparent, elliptical-triangular, margins ciliate. Stamens 3, filaments free, anthers yellow, ovary long-ovoid, glabrous. Pistil short, stigmas 2, plumose. Fruits unknown.

Phenology. New shoots August. Flowering April to June.

Distribution and habitat. *Yushania pauciramificans* is known from Xinping, south-central Yunnan, mainly distributed in the evergreen broadleaved forest at an elevation of 2250–2500 m.

Discussion

Reproductive features play an important role in the delimitation of the genera of alpine bamboos, and can improve our understanding of the relationship of *Borinda*, *Fargesia* and *Yushania*. Although *Fargesia angustissima* was recombined into *Borinda* in the analysis of Stapleton (2006), its inflorescence characters provide new insight in its delimitation. According to the description, the leaf sheath underneath the inflorescence of *F. angustissima* is inflated but smaller than the spathe of some species of *Fargesia* s. s., such as *F. funiushanensis* T.P. Yi, *F. qinlingensis* T.P. Yi & J. X. Shao, while more similar to those with spikelets stretching out from one side of sheaths and arranging relatively loosely (Zhang and Ren 2016). These characteristics indicate that the inflorescence of *F. angustissima* is in a transitional state between compressed and open ones (Fig. 3). Moreover, *F. angustissima* possesses an independent position on the molecular phylogenetic trees and has a special habitat (Ye et al. 2019; Shi et al. 2022). The florets, phylogenetic position, and habitat of *F. angustissima* imply that it is different from other species of *Fargesia* and could be treated as a new genus, albeit following a narrower genus concept. Additionally, as currently circumscribed, *Borinda* is a paraphyletic group without distinct synapomorphies despite some hair-like vegetative morphology and gene flow occurring frequently between it and *Fargesia*1 clade (Ye et al. 2019; Stapleton 2021; Ye et al. 2021a). Considering the variable vegetative characters and insufficient reproductive features, any new combination of these alpine bamboos should be made cautiously, especially when extensive gene flow exists. Thus, we support the “Flora of China” in adopting a broadly-defined *Fargesia* s. l., rather than *Borinda* to minimize nomenclatural change.

The flowering branches and flowers of *Yushania pauciramificans* are similar to those of species of *Yushania*, both have similar paniculate inflorescence, axilla with tuberculate glands, ciliate rachilla internodes and florets. Combined with the reduced gene flow between *Yushania* and *Fargesia* revealed by D-statistic tests, the inflorescence state of *Y. pauciramificans* supports the monophyly of *Yushania* further (Ye et al. 2019).



Figure 3. Comparison of inflorescence **A** *Fargesia funiushanensis* T.P. Yi **B** *Fargesia angustissima* T.P. Yi **C** *Yushania pauciramificans* T.P. Yi.

Acknowledgements

We thank Peng-Fei Ma, Cen Guo, Yu-Jiao Wang, Ling Mao, Jia-Wei Sun of Kunming Institute of Botany, Chinese Academy of Science, and the guide of Gasa Town, Xiping, Yunnan, for their assistance with collecting specimens. The study was funded by the National Natural Science Foundation of China (No. 32260307), the Strategic Priority Research Program of Chinese Academy of Sciences (No. XDB31000000), and the Applied and Fundamental Research Foundation of Yunnan Province (No. 2019FD059).

References

- Attigala L, Kathriarachchi HS, Clark LG (2016) Taxonomic Revision of the Temperate Woody Bamboo Genus *Kuruna* (Poaceae: Bambusoideae: Arundinarieae). *Systematic Botany* 41(1): 174–196. <https://doi.org/10.1600/036364416X690570>
- Bamboo Phylogeny Group (2012) An updated tribal and subtribal classification of the bamboos (Poaceae: Bambusoideae). *The Journal of American Bamboo Society* 24(1): 1–10.
- Guo ZH, Chen YY, Li DZ (2002) Phylogenetic studies on the *Thamnocalamus* group and its allies (Gramineae: Bambusoideae) based on *ITS* sequence data. *Molecular Phylogenetics and Evolution* 22(1): 20–30. <https://doi.org/10.1006/mpev.2001.1039>
- Janzen DH (1976) Why bamboos wait so long to flower. *Annual Review of Ecology and Systematics* 7(1): 347–391. <https://doi.org/10.1146/annurev.es.07.110176.002023>
- Keng PC, Wang ZP (1996) *Flora Reipublicae Popularis Sinicae*, Vol. 9(1). Science Press, Beijing, 761 pp.
- Li DZ, Wang ZP, Zhu ZD, Xia NH, Jia LZ, Guo ZH, Yang GY, Stapleton CMA (2006) Tribe Bambuseae (Poaceae). In: Wu ZY, Raven PH, Hong DY (Eds) *Flora of China* Vol 22. Science Press, Beijing and Missouri Botanical Garden Press, St. Louis, 7–180.
- McClure FA (1966) *The Bamboos: A Fresh Perspective*. Harvard University Press, 110–112. <https://doi.org/10.4159/harvard.9780674428713>
- Ohrnberger D (1999) *The bamboos of the world: annotated nomenclature and literature of the species and the higher and lower taxa*. Access Online via Elsevier, 596 pp.
- Shi JY, Zhang YX, Zhou DQ, Ma LS, Yao J (2022) *Illustrated Flora of Bambusoideae in China* (vol. 2). Science Press, Beijing and Springer, Singapore, 488 pp. <https://doi.org/10.1007/978-981-16-2758-3>
- Stapleton CMA (1994) The bamboos of Nepal and Bhutan. Part II: *Arundinaria*, *Thamnocalamus*, *Borinda*, and *Yushania* (Gramineae: Poaceae, Bambusoideae). *Edinburgh Journal of Botany* 51(02): 275–295. <https://doi.org/10.1017/S0960428600000883>
- Stapleton CMA (1998) New combinations in *Borinda* (Gramineae: Bambusoideae). *Kew Bulletin* 53(2): 453–459. <https://doi.org/10.2307/4114510>
- Stapleton CMA (2006) New taxa and combinations in cultivated bamboos (Poaceae: Bambusoideae). *SIDA, Contributions to Botany* 22: 331–332. <http://www.jstor.org/stable/41968582>

- Stapleton CMA (2021) We need to talk about *Fargesia*: new combinations and a new genus in the temperate Sino-Himalayan bamboos (Poaceae: Bambusoideae). The Journal of American Bamboo Society 31: 1–16. https://bamboo.org/_uploads/pdfs/JABSv31_Stapleton_20210510.pdf
- Turland NJ, Wiersema JH, Barrie FR, Greuter W, Hawksworth DL, Herendeen PS, Knapp S, Kusber WH, Li DZ, Marhold K, May TW, McNeill J, Monro AM, Prado J, Price MJ, Smith GF (Eds) (2018) International Code of Nomenclature for algae, fungi, and plants (Shenzhen Code) adopted by the Nineteenth International Botanical Congress Shenzhen, China, July 2017. Regnum Vegetabile 159. Koeltz Botanical Books, Glashütten. <https://doi.org/10.12705/Code.2018>
- Wang XQ, Ye XY, Zhao L, Li DZ, Guo ZH, Zhuang HF (2017) Genome-wide RAD sequencing data provide unprecedented resolution of the phylogeny of temperate bamboos (Poaceae: Bambusoideae). Scientific Reports 7(1): e11546. <https://doi.org/10.1038/s41598-017-11367-x>
- Ye XY, Ma PF, Yang GQ, Guo C, Zhang YX, Chen YM, Guo ZH, Li DZ (2019) Rapid diversification of alpine bamboos associated with the uplift of the Hengduan Mountains. Journal of Biogeography 46(12): 2678–2689. <https://doi.org/10.1111/jbi.13723>
- Ye XY, Ma PF, Guo C, Li DZ (2021a) Phylogenomics of *Fargesia* and *Yushania* reveals a history of reticulate evolution. Journal of Systematics and Evolution 59(6): 1183–1197. <https://doi.org/10.1111/jse.12719>
- Ye XY, Zhang YX, Li DZ (2021b) Two new species of *Yushania* (Poaceae: Bambusoideae) from South China, with a taxonomic revision of related species. Plant Diversity 43(6): 492–501. <https://doi.org/10.1016/j.pld.2021.03.001>
- Yi TP (1985) Classification and distribution of the food bamboos of the giant panda. Journal of Bamboo Research 4(2): 20–45.
- Yi TP (1986) Studies on the genus *Yushania*. Journal of Bamboo Research 5(1): 8–66.
- Yi TP (1988a) A study of the genus *Fargesia* from china. Journal of Bamboo Research 7(2): 6–15.
- Yi TP (1988b) Six new species of bamboos from China. Bulletin of Botanical Research 8(4): 71–73.
- Yi TP, Shi JY, Ma LS, Wang HT, Yang L (2008) Iconographia Bambusoidearum Sinicarum. Science Press, Beijing, 766 pp.
- Zhang YQ, Ren Y (2016) Supplementary description of flowers and flowering branches of four *Fargesia* and one *Drepanostachyum* species (Bambusoideae, Poaceae), and notes on their taxonomy. Nordic Journal of Botany 34(5): 565–572. <https://doi.org/10.1111/njb.00975>
- Zhang YQ, Zhou Y, Hou XQ, Huang L, Kang JQ, Zhang JQ, Ren Y (2019) Phylogeny of *Fargesia* (Poaceae: Bambusoideae) and infrageneric adaptive divergence inferred from three cpDNA and nrITS sequence data. Plant Systematics and Evolution 305(1): 61–75. <https://doi.org/10.1007/s00606-018-1551-y>
- Zhang YX, Guo C, Li DZ (2020) A new subtribal classification of Arundinarieae (Poaceae, Bambusoideae) with the description of a new genus. Plant Diversity 42(3): 127–134. <https://doi.org/10.1016/j.pld.2020.03.004>
- Zhou Y, Zhang YQ, Xing XC, Zhang JQ, Ren Y (2019) Straight From the Plastome: Molecular Phylogeny and Morphological Evolution of *Fargesia* (Bambusoideae: Poaceae). Frontiers in Plant Science 10: 981. <https://doi.org/10.3389/fpls.2019.00981>
- Zhou Y, Li WW, Zhang YQ, Xing XC, Zhang JQ, Ren Y (2020) Extensive reticulate evolution within *Fargesia* (s.l.) (Bambusoideae: Poaceae) and its allies: Evidence from multiple nuclear markers. Molecular Phylogenetics and Evolution 149: e106842. <https://doi.org/10.1016/j.jmpev.2020.106842>