

Breynia hiemalis (Phyllanthaceae, Phyllanthaeae), a new species from Yunnan, south-west China

Feng Yang¹, Chao Chen^{2,3}, Jing-Yi Ye¹, Jian-Yong Wu⁴, Huan-Chong Wang^{5,6}

1 School of Life Sciences, Yunnan University, Kunming 650091, Yunnan, China **2** Key Laboratory of Tropical Forest Ecology, Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences, Mengla 666303, Yunnan, China **3** Yuanjiang Savanna Ecosystem Research Station, Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences, Yuanjiang 653300, Yunnan, China **4** Yuxi Forestry and Grassland Bureau, Yuxi 653100, Yunnan, China **5** School of Ecology and Environmental Science, Yunnan University, Kunming 650091, China **6** Herbarium of Yunnan University, Kunming 650091, Yunnan, China

Corresponding author: Huan-Chong Wang (hchwang@ynu.edu.cn)

Academic editor: Peter Bruyns | Received 12 April 2022 | Accepted 1 August 2022 | Published 26 August 2022

Citation: Yang F, Chen C, Ye J-Y, Wu J-Y, Wang H-C (2022) *Breynia hiemalis* (Phyllanthaceae, Phyllanthaeae), a new species from Yunnan, south-west China. *PhytoKeys* 206: 75–86. <https://doi.org/10.3897/phytokeys.206.85241>

Abstract

Breynia hiemalis Huan C. Wang & Feng Yang (Phyllanthaceae), of sect. *Cryptogynium* (Müll.Arg.) Welzen & Pruesapan in subg. *Breynia*, is described from Yunnan, south-west China. It is known from only a single locality in the valley of the Yuanjiang River, and usually occurs in the understory of the savanna vegetation. It is characterized by its broadly elliptic to orbicular leaf blades, shallowly plate-like calyces of the staminate flowers, ovaries with clearly erose rim and urceolate capsules. Morphological comparisons with similar species are also presented.

Keywords

Breynia sect. *Cryptogynium*, endemism, *Sauropus*, savanna, Yuanjiang River

Introduction

The family Phyllanthaceae, a segregate from Euphorbiaceae *sensu lato*, is a pantropical group of herbs, shrubs and treelets (Kathriarachchi et al. 2005; Hoffmann et al. 2006). It currently consists of about 2000 species, with more than 1200 placed in the largest tribe Phyllanthaeae (Govaerts et al. 2000; Hoffmann et al. 2006). Within the Phyllanthaeae, the generic classification is still contentious (Hoffmann et al. 2006;

Wagner and Lorence 2011; Pruesapan et al. 2012; Van Welzen et al. 2014; Jangid and Gupta 2016; Bouman et al. 2022). Recent molecular studies clearly demonstrated that *Phyllanthus* L. (in the traditional sense), the largest genus of Phyllanthaceae, with more than 800 species, is paraphyletic since *Glochidion* J. R. Forst. & G. Forst., *Breynia* J. R. Forst. & G. Forst. and *Synostemon* F. Muell. are nested within it (Bouman et al. 2021, 2022). Some authors suggested placing most species of the Phyllanthaceae in *Phyllanthus*. This would make *Phyllanthus* a large and morphologically heterogeneous group with more than 1200 species (e.g. Hoffmann et al. 2006; Wagner and Lorence 2011; Ralimanana et al. 2013; Kato and Kawakita 2017; Falcón-Hidalgo et al. 2020). Conversely, other authors (for example, Pruesapan et al. 2008; Pruesapan 2010; Van Welzen and Pruesapan 2010; Pruesapan et al. 2012; Bouman et al. 2018, 2021) prefer to divide *Phyllanthus s. l.* into several smaller, monophyletic genera that can be characterized morphologically. More recently, Bouman et al. (2022) split *Phyllanthus s. l.* into thirteen monophyletic and morphologically recognizable genera, where *Breynia* was kept as a separate genus.

Breynia and *Sauropus* Blume in the strict sense are closely related; both share bifid or emarginate styles, non-apiculate anthers, smooth seeds, and they generally possess sepal scales (Van Welzen et al. 2014). From the results of a well-sampled phylogenetic analysis, Van Welzen et al. (2014) expanded *Breynia* to include the south-east Asian species of *Sauropus* and reinstated the Australian genus *Synostemon*, previously relegated to a section of *Sauropus*, making each genus monophyletic and morphologically definable. The newly circumscribed *Breynia* is a moderately-sized genus of 85 to 90 species (Van Welzen et al. 2014; Bouman et al. 2022). It contains three infrageneric groups, including subg. *Breynia* and subg. *Sauropus* (Blume) Welzen & Pruesapan, with subg. *Breynia* subdivided into sect. *Breynia* and sect. *Cryptogynium* (Müll. Arg.) Welzen & Pruesapan. Members of *Breynia* are distributed mainly from Australia to tropical and subtropical Asia, with a center of diversity in south-east Asia (Van Welzen 2003; Bouman et al. 2022; POWO 2022). Twenty species of *Breynia* are found in China, including 15 species formerly placed in *Sauropus*, and most species occur in the southern and south-western regions (Li 1994; Li and Esser 2008; Li and Gilbert 2008). Recently, a new species of *Breynia* was described from Yunnan, south-west China by Yang et al. (2022). This brings the total number of Chinese species to 21, of which five species belong to sect. *Cryptogynium*, seven species belong to sect. *Breynia*, and nine species belong to subg. *Sauropus*.

During fieldwork in Yuanjiang National Natural Reserve, Yunnan Province (SW China), in December 2015 and January 2022, we came across a dwarf plant with discoid staminate flowers and 3 stigmas spreading horizontally from the top of obconical ovary, obviously belonging to *Breynia* according to the generic delimitation of Van Welzen et al. (2014) and Bouman et al. (2022). This could not be assigned to any previously known species from China (Li and Gilbert 2008), or from any of the adjacent south-east Asian countries, i.e. Vietnam, Laos, Thailand and Myanmar (Beille 1927; Pham 2003; Van Welzen 2003; Chakrabarty and Balakrishnan 2018; Van Welzen 2022; Van Welzen and Chayamarit 2022). Comparison with morphologically similar species supports recognizing this plant as an undescribed species.

Materials and methods

The new species was studied both in the field and in the herbarium. The collections of similar species housed at KUN, PE, XTBG, YUKU, and digital images available at the JSTOR Global Plants (<https://plants.jstor.org/>), the Chinese Virtual Herbarium (<http://www.cvh.ac.cn/>), and the Global Biodiversity Information Facility (<https://www.gbif.org/>) were reviewed. Surveys of pertinent taxonomic literature (for example, Beille 1927; Li 1994; Pham 2003; Van Welzen 2003; Ma et al. 2006; Li and Gilbert 2008; Van Welzen et al. 2014; Chakrabarty and Balakrishnan 2018; Van Welzen 2022; Van Welzen and Chayamarit 2022) were carried out. Measurements were made under a stereomicroscope (Olympus SZX2, Tokyo, Japan) using a ruler and a metric vernier caliper.

Taxonomy

Breynia hiemalis Huan C. Wang & Feng Yang, sp. nov.

urn:lsid:ipni.org:names:77303992-1

Figs 1–3

Type. CHINA. Yunnan Province: Yuanjiang County, Pupiao, 600–700 m, 23°28'37"N, 102°10'37"E, in savanna on a mountain slope, 12 Jan. 2022, *H. C. Wang et al. YJ16225* (holotype YUKU-02074690!; isotypes YUKU!, PE!, HITBC!).

Diagnosis. *Breynia hiemalis* can easily be distinguished from all morphologically similar species by plants glabrous throughout, by its broadly elliptic to orbicular and relatively small (4–21 × 4–17 mm) leaves, calyx of staminate flower shallowly plate-like, ovary rim conspicuously erose, and the urceolate capsule with a raised and lobed apical rim.

Description. Dwarf shrubs or subshrubs, 10–20 (–30) cm tall, monoecious, glabrous throughout, with phyllanthoid branching. Main stems more or less procumbent to ascending, brown, with 4 shallow ribs, sometimes rooting at the lower nodes; branches green, deciduous, ascending, 3–8 cm long. *Cataphylls* lanceolate, to 1 mm long, arranged spirally at the base of the plagiotropic branchlets. *Leaves* on ultimate branchlets distichous, simple; stipules triangular-lanceolate, usually auriculate basally, 1.5–2.0 mm long; *petiole* 1.2–1.6 × 0.3–0.5 mm; *blade* broadly elliptic to orbicular, rarely slightly ovate, papery, 4–21 × 4–17 mm, length/width ratio 1–1.5, base rounded to broadly cuneate, margin entire, flat, apex usually rounded, sometimes truncate, retuse, rarely mucronate, adaxially green, abaxially grey or slightly glaucous; venation pinnate, lateral veins 4 or 5 pairs, reticulate veins obscure. *Inflorescences* axillary, *peduncles* very short, ± 0.1 × 0.1 mm, with minute bracts, male or female flowers usually solitary, staminate flowers proximal, pistillate flowers usually distal. *Staminate flowers:* *pedicel* slender, ± 7 mm long; *calyx* shallowly plate-like, ± 4 mm in diam., red, 6-lobed; lobes biseriate, broadly obovate, slightly fleshy, 0.9–1.2 × 1.1–1.3 mm, apex obtuse or retuse, scales present; *stamens* 3, filaments connate, androphore ± 0.2 mm long, splitting horizontally, branches up to 0.5 mm long with anthers underneath, anthers ± 0.3 × 0.3 mm. *Pistillate flowers:* *pedicel* ± 3 mm

long, thickening upwards; calyx ± 6 mm in diam., greenish, whitish yellow, or pinkish, lobes biseriate, obovate, subcoriaceous, outer lobes $2.5\text{--}3.0 \times \pm 2.5$ mm, slightly longer and wider than the inner, inner lobes $\pm 2.5 \times 2.1\text{--}2.5$ mm, apex obtuse to truncate, shortly and abruptly acuminate; **ovary** obconical, 1.2–1.6 mm in diam., 3-locular, 2 ovules per locule, rim present at the apex, obviously erose; stigmas 3, spreading horizontally from top of ovary, apex split and recurved through $\pm 180^\circ$, sepals persistent and enlarged to $\pm 5 \times 4$ mm in fruit. **Capsules** urceolate, $\pm 4 \times 5\text{--}6$ mm, with a raised, lobed apical rim and persistent stigmas.




Figure 1. *Breyenia hiemalis*. (Drawn by Jing-Yi Ye from type specimen *H. C. Wang et al. YJ16225*) **A** habit **B** pistillate flower (apical view) **C** fruit **D** staminate flower (apical view) **E** staminate flower (lateral view).




Figure 2. *Breynia hiemalis* **A** habit **B** phyllanthoid branch showing pistillate flower and staminate flowers **C** stipule **D** staminate flower (apical view) **E** pistillate flower (apical view) **F** calyx in fruit (dorsal view) **G** fruit. Photographed by H. C. Wang from type locality in January 2022.



Figure 3. Holotype of *Breytia hiemalis* (YUKU-02074690).




Figure 4. South-western China, showing the known distribution (red star) of *Breynia hiemalis*. The blue lines represent the rivers.

Phenology. Flowering from December to January, fruiting from January to February.

Etymology. The epithet “*hiemalis*” is Latin for “belonging to winter”, referring to the flowering period of this new species.

Distribution and habitat. *Breynia hiemalis* appears to be rare and is endemic to Yunnan, south-west China. It is known from only a single locality in the valley of the Yuanjiang River, which flows from Yunnan (south-west China) through northern Vietnam to the Gulf of Tonkin (Fig. 4). The climate in Yuanjiang valley is characterized by a long dry season (the dry season can be further divided into a cool dry season (November to February) and a hot dry season (March to April)), with an annual average

temperature of 24 °C and a mean annual evaporation capacity of 2700–3800 mm, that is three to six times higher than the mean annual precipitation (600–800 mm), and with 80–90% of the precipitation concentrated in the wet season (from May to October) (Jin 2002; Shen et al. 2010; Zhou et al. 2017). *Breynia hiemalis* grows in savanna on a mountain slope (Fig. 5) at elevations of 500–700 m, together with *Lannea coromandelica* (Houtt.) Merr. (Anacardiaceae), *Adina cordifolia* (Roxb.) Brandis (Rubiaceae), *Bauhinia brachycarpa* Wall. ex Benth. (Fabaceae), *Tephrosia purpurea* (L.) Pers. (Fabaceae), *Woodfordia fruticosa* (L.) Kurz (Lythraceae), *Waltheria indica* L. (Malvaceae), *Jasminium mesnyi* Hance (Oleaceae), *Searsia paniculata* (Wall. ex G. Don) Moffett (Anacardiaceae), *Heteropogon contortus* (L.) P. Beauv. ex Roem. & Schult. (Poaceae), and others.

Additional specimens examined (Paratypes). China. Yunnan: Yuanjiang County, Pupiao, 17 Dec. 2015, *H. C. Wang et al.* YJ736 (YUKU).

Taxonomic notes. According to Van Welzen et al. (2014) and Bouman et al. (2022), *B. hiemalis* should be assigned to sect. *Cryptogynium* because of its horizontal anthers (the androphore splits apically into three horizontal arms with the anthers hanging underneath) and ovary with a rim. Prior to the present study, only five species of sect. *Cryptogynium* were recorded in China, namely *B. compressa* (Müll. Arg.) Chakrab. & N. P. Balakr. (a member of the *B. quadrangularis* (Willd.) Chakrab. & N. P. Balakr. complex which was recognized as a distinct species by Chakrabarty and Balakrishnan (2015)), *B. delavayi* (Croizat) Welzen et Pruesapan, *B. pierrei* (Beille) Welzen et Pruesapan, *B. similis* (Craib) Welzen et Pruesapan and *B. tsiangii* (P. T. Li) Welzen et Pruesapan (Li and Gilbert 2008; Van Welzen et al. 2014; Chakrabarty and Balakrishnan 2015). *Breynia hiemalis* shows some resemblance to *B. compressa* and *B. delavayi* in its dwarf habit and axillary inflorescences. However, *B. hiemalis* can be distinguished from *B. compressa* by several characters, namely stems more or less procumbent to ascending (vs. erect or arching in *B. compressa*), calyx of the staminate flower shallowly plate-like (vs. star-shaped), lobes broadly obovate (vs. suborbicular or squarish), apex obtuse or retuse (vs. emarginate-truncate to deeply bilobulate) (Fig. 6: D, E), ovary rim erose (vs. retuse) (Fig. 6: A, B), capsule urceolate (vs. ovoid), with raised and lobed apical rim (vs. with low and smooth apical rim). Additionally, *B. hiemalis* flowers in winter (from December to January), whereas *B. compressa* flowers from summer to autumn (from April to October). *Breynia hiemalis* differs strikingly from *B. delavayi* in its obscure reticulate veins (vs. reticulate veins elevated on both surfaces) and calyx of the staminate flower shallowly plate-like (vs. star-shaped) (Fig. 6: D, F).

Of the species of *Breynia* found in south-east Asia, *B. hiemalis* is also morphologically similar to *B. granulosa* (Airy Shaw) Welzen & Pruesapan, from eastern Thailand. Nevertheless, *B. granulosa* differs from *B. hiemalis* in having obovate leaves (vs. broadly elliptic to orbicular, rarely slightly ovate in *B. hiemalis*), calyx of staminate flower campanulate (vs. shallowly plate-like), androphores ± 0.8 mm (vs. ± 0.2 mm) long, smaller pistillate flowers, usually 4–5 mm (vs. ± 6 mm) in diam., stigmas ascending (vs. horizontally spreading) (Fig. 6: A, C) and ovoid (vs. urceolate) capsules. *Breynia hiemalis* is also similar to *B. poilanei* (Beille) Welzen et Pruesapan from Vietnam, but it clearly differs from the latter by its more or less procumbent to ascending stems (vs.

erect in *B. poilanei*), 0.1–0.2 (–0.3) m (vs. 1.5 m) tall, branches 3–8 cm (vs. 5–15 cm) long, stipules triangular-lanceolate (vs. triangular), 1.5–2.0 mm (vs. 0.5 mm) long, leaves broadly elliptic to orbicular, rarely slightly ovate (vs. ovate, rarely orbicular), capsules 4 mm (vs. up to 10 mm) wide. A key to distinguish the members of *Breynia* sect. *Cryptogynium* in China is given below.




Figure 5. Habitat of *Breynia hiemalis* **A** distant view of the type locality **B** nearby view.




Figure 6. *Breynia hiemalis* (**A**, **D**), *B. compressa* (**B**, **E**), *B. granulosa* (**C**), *B. delavayi* (**F**). **A–C** pistillate flowers **D–F** staminate flowers.

Key to Chinese species of *Breynia* sect. *Cryptogynium*

- 1 Shrubs 1–3 m tall 2
 – Dwarf shrubs or subshrubs 0.1–1.0 m tall 4
 2 Leaves ovate, 0.7–3.7 × 0.5–1.9 cm *Breynia similis*
 – Leaves ovate to lanceolate, 6–10 × 3–4 cm 3
 3 Leaves usually ovate, rarely ovate-lanceolate, papery, base rounded, apex caudate-acuminate *B. tsiangii*
 – Leaves ovate-lanceolate to lanceolate, coriaceous, base cuneate, apex acuminate *B. pierrei*
 4 Calyx of staminate flower shallowly plate-like *B. hiemalis*
 – Calyx of staminate flower star-shaped 5
 5 Branches pubescent, leaves orbicular, base shallowly cordate, truncate or rounded, lateral veins robust, reticulate nerves prominent *B. delavayi*
 – Branches glabrous; leaves ovate, elliptic or nearly rounded, base rounded or broadly cuneate, lateral veins slender, reticulate nerves obscure ... *B. compressa*

Acknowledgements

We thank the curators of the Yuanjiang Savanna Ecosystem Research Station who helped us with collecting. We express our gratitude to Pingping Li for the distribution map of the new species. We are grateful to Dr. Libing Zhang (Missouri Botanical Garden) for critically reading the manuscript and improving the English, and we give special thanks to two anonymous reviewers and the editor for their valuable comments and suggestions. This research was supported by the National Natural Science Foundation of China (grant no: 31960040) and the Second Tibetan Plateau Scientific Expedition and Research (STEP) programme (2019QZKK0502).

References

- Beille L (1927) *Breynia* J. R. Forster et G. Forster and *Sauropus* Blume. In: Lecomte MH (Ed.) Flore Générale de l'Indo-Chine, vol.5. Masson, Paris, 631–658.
- Bouman RW, Keßler PJA, Telford IRH, Bruhl JJ, Van Welzen PC (2018) Subgeneric delimitation of the plant genus *Phyllanthus* L. (Phyllanthaceae). *Blumea* 63: 167–198. <https://doi.org/10.3767/blumea.2018.63.02.14>
- Bouman RW, Keßler PJA, Telford IRH, Bruhl JJ, Strijk JS, Saunders RMK, Van Welzen PC (2021) Molecular phylogenetics of *Phyllanthus sensu lato* (Phyllanthaceae): Towards coherent monophyletic taxa. *Taxon* 70(1): 72–98. <https://doi.org/10.1002/tax.12424>
- Bouman RW, Keßler PJA, Telford IRH, Bruhl JJ, Strijk JS, Saunders RMK, Esser H-J, Falcón-Hidalgo B, Van Welzen PC (2022) A revised phylogenetic classification of tribe Phyllantheae (Phyllanthaceae). *Phytotaxa* 540(1): 001–100. <https://doi.org/10.11646/phytotaxa.540.1.1>

- Chakrabarty T, Balakrishnan NP (2015) Reinstatement of four taxa synonymized under *Breynia quadrangularis* with proposal for a new subgenus *Hemisauropus* (Phyllanthaceae). *Journal of Economic and Taxonomic Botany* 39(3–4): 415–419.
- Chakrabarty T, Balakrishnan NP (2018) Indo-Burmese Phyllanthaceae, A taxonomic Revision. Bishen Singh Mahendra Pal Singh, Dehra Dun, 1–437.
- Falcón-Hidalgo B, Fuentes SB, Berazáin RI, Borsch T (2020) Phylogenetic relationships and character evolution in Neotropical *Phyllanthus* (Phyllanthaceae), with a focus on the Cuban and Caribbean taxa. *International Journal of Plant Sciences* 181(3): 384–305. <https://doi.org/10.1086/706454>
- Govaerts R, Frodin DG, Radcliffe-Smith A (2000) World checklist and bibliography of Euphorbiaceae (with Pandaceae), vols 1–4. Royal botanic gardens, Kew.
- Hoffmann P, Kathriarachchi H, Wurdack KJ (2006) A phylogenetic classification of Phyllanthaceae (Malpighiales; Euphorbiaceae *sensu lato*). *Kew Bulletin* 61(1): 37–53.
- Jangid PP, Gupta S (2016) Systematic wood anatomy of the tribe Phyllanthae (Phyllanthaceae, Euphorbiaceae *s. l.*) from India: Implication in reinstatement of *Phyllanthus*, *Glochidion* and allies. *Nordic Journal of Botany* 34(4): 496–512. <https://doi.org/10.1111/njb.01072>
- Jin ZZ (2002) Floristic features of dry-hot and dry-warm valleys, Yunnan and Sichuan. Yunnan Science and Technology Press, Kunming, 1–255.
- Kathriarachchi H, Hoffmann P, Samuel R, Wurdack KJ, Chase MW (2005) Molecular phylogenetics of Phyllanthaceae inferred from five genes (plastid *atpB*, *matK*, *3'ndbF*, *rbcL*, and nuclear *PHYC*). *Molecular Phylogenetics and Evolution* 36(1): 112–134. <https://doi.org/10.1016/j.ympev.2004.12.002>
- Kato M, Kawakita A (2017) Obligate pollination mutualism. Springer, Tokyo, 310 pp. <https://doi.org/10.1007/978-4-431-56532-1>
- Li BT (1994) *Sauropus* Blume and *Breynia* J. R. Forster et G. Forster. *Flora Reipublicae Popularis Sinicae*, vol. 44 (1). Science Press, Beijing, 162–184.
- Li BT, Esser HJ (2008) *Breynia* J. R. Forster et G. Forster. *Flora of China*, vol. 11. Science Press, Beijing & Missouri Botanical Garden Press, St. Louis, Missouri, USA, 207–209.
- Li BT, Gilbert MG (2008) *Sauropus* Blume. *Flora of China*, vol. 11. Science Press, Beijing & Missouri Botanical Garden Press, St. Louis, Missouri, USA, 202–207.
- Ma HY, Peng H, Ying WQ (2006) *Sauropus* Blume and *Breynia* J. R. Forster et G. Forster. *Flora of Yunnan*, vol. 10. Science Press, Beijing, China, 136–146.
- Pham HH (2003) *An Illustrated Flora of Vietnam*, vol. 2. Young Publishing House, Ho Chi Minh City, 208–216.
- POWO (2022) Plants of the World Online. Royal Botanic Gardens, Kew. <http://www.plantsoftheworldonline.org/> [accessed 25. 1. 2022]
- Pruesapan K (2010) The rise and fall of *Sauropus* (Phyllanthaceae): a molecular phylogenetic analysis of *Sauropus* and allies. Doctoral thesis: Netherlands Centre for Biodiversity Naturalis (section NHN), PITA group, Leiden University Branch. <https://hdl.handle.net/1887/16170> [accessed 25. 1. 2022]
- Pruesapan K, Telford IRH, Bruhl JJ, Draisma SGA, Van Welzen PC (2008) Delimitation of *Sauropus* (Phyllanthaceae) based on plastid *matK* and nuclear ribosomal ITS DNA sequence data. *Annals of Botany (Oxford)* 102(6): 1007–1018. <https://doi.org/10.1093/aob/mcn193>

- Pruesapan K, Telford IRH, Bruhl JJ, Van Welzen PC (2012) Phylogeny and proposed circumscription of *Breynia*, *Sauropus* and *Synostemon* (Phyllanthaceae), based on chloroplast and nuclear DNA sequences. *Australian Systematic Botany* 25(5): 313–330. <https://doi.org/10.1071/SB11005>
- Ralimanana H, Hoffmann P, Rajeriarison C (2013) Taxonomic revision of *Phyllanthus* (Phyllanthaceae) in Madagascar and the Comoro Islands III: Subgenera Swartziani, Afroswartziani and Emblica. *Kew Bulletin* 68(4): 535–558. <https://doi.org/10.1007/s12225-013-9485-6>
- Shen R, Zhang JL, He B, Li F, Zhang ZM, Zhou R, Ou XK (2010) The structure characteristic and analysis on similarity of grassland community in dry-hot valley of Yuanjiang River. *Shengtai Huanjing Xuebao* 19(12): 2821–2825.
- Van Welzen PC (2003) Revision of the Malesian and Thai species of *Sauropus* (Euphorbiaceae: Phyllanthoideae). *Blumea* 448(2): 319–391. <https://doi.org/10.3767/000651903X675017>
- Van Welzen PC (2022) Flora Malesiana Euphorbiaceae. Naturalis Biodiversity Center, Leiden. <http://www.nationaalherbarium.nl/Euphorbs/specB/BreyniaT.htm> [accessed 25.6.2022]
- Van Welzen PC, Chayamarit K (2022) Flora of Thailand Euphorbiaceae. Naturalis Biodiversity Center, Leiden; Forest Herbarium, National Park, Wildlife and Plant Conservation Department, Bangkok. <http://nationaalherbarium.nl/ThaiEuph/ThBspecies/ThBreyniaT.htm> [accessed 25.6.2022]
- Van Welzen PC, Pruesapan K (2010) Four new species of *Breynia* (Phyllanthaceae / Euphorbiaceae *sensu lato*) and one new combination from Thailand and Malaysia. *Thai Forest Bulletin (Botany)* 38: 111–119.
- Van Welzen PC, Pruesapan K, Telford IRH, Esser HJ, Bruhl JJ (2014) Phylogenetic reconstruction prompts taxonomic changes in *Sauropus*, *Synostemon* and *Breynia* (Phyllanthaceae tribe Phyllanthaeae). *Blumea* 59(2): 77–94. <https://doi.org/10.3767/000651914X684484>
- Wagner WL, Lorence DH (2011) A nomenclator of Pacific oceanic island *Phyllanthus* (Phyllanthaceae), including *Glochidion*. *PhytoKeys* 4(0): 67–94. <https://doi.org/10.3897/phytokeys.4.1581>
- Yang F, Li PP, Liu JL, Wang QP, Wang HC (2022) *Breynia pseudorostrata* (Phyllanthaceae), a new species from Yunnan, southwest China. *Phytotaxa* 539(2): 210–216. <https://doi.org/10.11646/phytotaxa.539.2.8>
- Zhou Z, Gu BJ, Sun H, Zhu H, Tang YH (2017) Molecular phylogenetic analyses of Euphorbiaceae tribe Euphorbieae, with the description of a new genus, *Tsaiodendron* gen. nov., from south-western China. *Botanical Journal of the Linnean Society* 184(2): 167–184. <https://doi.org/10.1093/botlinnean/box023>