

Erythroxylum austroguangdongense (Erythroxylaceae), a new species from Guangdong, China

Chun-Mei He¹, Xin-Xin Zhou², Xue-He Ye^{2,3}, Weijun Chen⁴, Yi-Hua Tong^{2,5}

1 Guangdong Provincial Key Laboratory of Silviculture, Protection and Utilization, Guangdong Academy of Forestry, Guangzhou, 510520, China **2** Key Laboratory of Plant Resources Conservation and Sustainable Utilization & Key Laboratory of Digital Botanical Garden of Guangdong Province, South China Botanical Garden, Chinese Academy of Sciences, Guangzhou, 510650, China **3** Zhongkai University of Agriculture and Engineering, Guangzhou, 510225, China **4** Zhuhai Charmview International Travel Co. LTD, Zhuhai, 519000, China **5** Center of Conservation Biology, Core Botanical Gardens, Chinese Academy of Sciences, Guangzhou, 510650, China

Corresponding author: Yi-Hua Tong (yh-tong@scbg.ac.cn)

Academic editor: Hugo de Boer | Received 31 March 2022 | Accepted 1 July 2022 | Published 29 July 2022

Citation: He C-M, Zhou X-X, Ye X-H, Chen W, Tong Y-H (2022) *Erythroxylum austroguangdongense* (Erythroxylaceae), a new species from Guangdong, China. *PhytoKeys* 202: 133–138. <https://doi.org/10.3897/phytokeys.202.84688>

Abstract

Erythroxylum austroguangdongense (Erythroxylaceae), a new species from Guangdong Province, China, is described and illustrated. This new species is morphologically most similar to *E. calyptratum*, but is distinguished by the leathery leaf blade with fewer pairs of secondary veins and flowers borne on leafless nodes of the basal part of the current branch with much longer pedicels and sub-rectangular petal appendages. This is the second native species of *Erythroxylum* recorded from China.

Keywords

Coca family, morphology, taxonomy

Introduction

The genus *Erythroxylum* P. Browne (Erythroxylaceae), with about 264 species, is widely distributed in tropics and subtropics with the center of its diversity in Neotropics (POWO 2022). Many species of this genus contain the substance cocaine, and can be used medicinally as a narcotic. In China, only two species of *Erythroxylum* are recorded,

i.e., the introduced *E. novogranatense* (D. Morris) Hieron. and the widely distributed *E. sinense* Y. C. Wu (Liu and Bartholomew 2008).

During field surveys of medicinal plant resources in Guangdong Province, we encountered an unknown *Erythroxylum* species with white flowers, while the species previously recorded from China either has whitish-yellow (*E. novogranatense*) or pinkish (*E. sinense*) flowers. After a morphological comparison with specimens in the herbarium (IBSC) and consulting the relevant literature (e.g. Guillaumin 1907; Chung 1996; Hô 2003; Liu and Bartholomew 2008; Harwood and Chayamarit 2011; Komada et al. 2018), we concluded that this unknown species is morphologically distinct from all the other species previously known in China and its surrounding countries. Thus, we describe and illustrate it as a new species.

Materials and methods

Flowering and fruiting material was collected from Zhuhai and Taishan, Guangdong Province, China during several field trips from 2019 to 2022. Descriptions were based on both living and dried collections, which were deposited at the herbarium of South China Botanical Garden, Chinese Academy of Sciences (IBSC). Measurements were performed with a ruler, and small plant parts were observed and measured under a stereo microscope (Mshot-MZ101).

Taxonomic treatment

Erythroxylum austroguangdongense C. M. He, X. X. Zhou & Y. H. Tong, sp. nov.

urn:lsid:ipni.org:names:77302515-1

Fig. 1

Type. CHINA. Guangdong Province: Taishan, Chaliao Ao, 22°12'54.87"N, 112°57'34.84"E, 563 m a.s.l., 3 April 2021 (fl.), *Xin-Xin Zhou & Yue-Yao Liu LSX303* (holotype: IBSC, isotypes: IBSC).

Diagnosis. Most similar to *E. calytratum* Komada & Tagane in having reddish brown to grayish brown branches with dense lenticels, and white petals with appendages, but distinguished by the leathery (vs. thinly chartaceous) leaf blade with fewer pairs of secondary veins (6–8 pairs vs. 11–15 pairs), and flowers borne on leafless nodes of the basal part of current branch (vs. in leaf axils) with longer pedicels (1–1.5 cm vs. 5.2–7 mm) and sub-rectangular petal appendage (vs. bilobed appendage with each lobe consisting of a short anterior auricle and a large posterior auricle). A more detailed comparison of the two species is shown in Table 1.

Description. Shrubs deciduous, 1.5–2 m tall, dioecious. Young branches greenish, old branches reddish-brown to grayish-brown, lenticellate. Stipule triangular to narrowly triangular, 1.8–3 × 0.7–0.8 mm, margin entire when young, gradually fimbriate,

Table 1. Morphological comparison of *Erythroxylum austroguangdongense* and *E. calyptratum*.

Characters	<i>E. austroguangdongense</i>	<i>E. calyptratum</i>
Leaf blade texture	Leathery	Thinly chartaceous
Leaf blade shape	Elliptic or lanceolate	Elliptic, oblong, oblong-lanceolate or ovate
Leaf blade color	Dark green and shining adaxially	Pale green adaxially
Secondary veins	6–8 pairs	11–15 pairs
Flower	Solitary on leafless nodes of the basal part of current branch	Solitary to 3-fascicled in leaf axils
Pedicel length	1–1.5 cm	5.2–7 mm
Petal appendage	Sub-rectangular	Bilobed and each lobe consisting of a short anterior auricle and a large posterior auricle

or dissected, caducous when old. Leaves alternate, simple; petiole 3.5–6 mm long; blade elliptic or lanceolate, 4–7.7 × 1.6–2.5 cm, leathery, apex acuminate, acumen to 5 mm long, base cuneate or attenuate, margin entire, dark green and shining adaxially, pale green abaxially, midrib sunken adaxially, prominent and pale yellow to brownish-yellow abaxially, secondary veins 6–8 pairs, almost flat, faintly visible on both surfaces, tertiary veins reticulate, obscure adaxially, slightly visible abaxially. Flowers solitary on leafless nodes of the basal part of current branch, pedicel 1–1.5 cm long; bracteole triangular, ca. 0.8 mm long, margin fimbriate or serrately dissected; calyx 1.6–1.8 mm long, lobes 5, ovate-triangular, ca. 1 mm long, apex brownish, acute; petals 5, white, oblong, 3.7–4 mm long, apex obtuse or rounded, adaxially with one appendage attached ca. 1.5 mm above from base, appendage sub-rectangular, ca. 1.6 × 1 mm, papillate throughout, slightly concave on both upper and lower margin. Stamens or staminodes 10. Male flowers: stamens with different length of filaments arranged alternately, short filaments ca. 3.7 mm long, long ones ca. 4 mm long, all basally connate into a tube, tube ca. 2 mm long, densely covered with papillary trichomes; anthers ca. 1.1 mm long; sterile pistil ca. 4 mm long. Female flowers: staminodes ca. 1.2 mm long, staminodal tube ca. 1 mm long, anther absent; ovary ellipsoidal, 2–2.5 mm long, 1–1.3 mm in diam., 3-locular, with 1 fertile locule, styles 3, entirely free at base, ca. 1 mm long including stigma, stigma clavate, ca. 0.5 mm long, reflexed, with papillae. Young fruits green, turning to yellowish, ripening red, ovoid to reniform, apex oblique, 7–9 × 2.5–3 mm.

Etymology. The species epithet is named after the distribution area of this new species, South Guangdong.

Vernacular name. 南粤古柯 (Chinese pinyin: nán yuè gǔ kē).

Distribution and habitat. This species is currently known only from Guangdong Province, China. Three populations are found in Yangchun (E’huangzhang), Zhuhai (Fenghuang Mountain) and Taishan (Gudou Mountain) respectively. It grows in evergreen broadleaved forests at elevations of 170–800 m.

Conservation Status. According to the field observation, the number of mature individuals of *Erythroxylum austroguangdongense* is less than 100. Thus, it is assigned a status of ‘Endangered’ (EN) following the IUCN Red List categories and criteria (IUCN Standards and Petitions Committee 2022). Since most of its distribution area is under the protection of E’huangzhang Natural Reserve and Gudou Mountain Natural Reserve, and it is not economically valuable, the threat risk seems to be low.




Figure 1. *Erythroxylum austroguangdongense* **A** flowering branches **B** leafy branches **C** female flowers **D** male flower **E** stipule, showing fimbriate margin **F** flower bud, showing calyx **G** petals, adaxial (left), lateral (middle) and abaxial (right) view **H** androecium **I** staminoles and pistil **J** cross section of ovary, showing one fertile locule (low right) **K–L** fruits. Scale bars: 3 mm (**H**); 2 mm (**E–G, I**); 1 mm (**J**). Photographs **A–B** by Xin-Xin Zhou **D, K** by Wei-Jun Chen **H** by Xue-He Ye and others by Chun-Mei He.

Phenology. Flowering in March-April and fruiting in April-August.

Additional specimens examined (paratypes). CHINA. Guangdong, Taishan: Banyuzui, 448 m a.s.l., 28 March 2019 (fl.), *Lei Jiang, Jin-Fan Lin, Jin-Ye Feng, Fu-Jun Chen & Jie-Lin Chen GDS-00291* (IBSC); Chaliao Ao, 205 m a.s.l., 25 April 2019 (fr.), *Teng-Hui Guo, Jin-Fan Lin & Jia-Ming Chen GDS-00484* (IBSC); *ibid.*, 19 May 2019 (fr.), *Taishan Expedition 440781190519024LY* (IBSC). Yangchun: Bajia, Xianjiadong Reservoir, 800 m a.s.l., 1 August 2001 (fr.), *Hua-Gu Ye 6094* (IBSC); Bajia Reservoir, 750 m a.s.l., 5 August 2001 (fr.), *Hua-Gu Ye 6199* (IBSC). Zhuhai: Fenghuang Mountain, 177 m a.s.l., 2 April 2021 (young fr.), *Yi-Hua Tong, Wei-Jun Chen & Feng Ling TYH-2526* (IBSC); *ibid.* 20 March 2022 (fl.), *Wei-Jun Chen TYH-2561* (IBSC).

Discussion

This new species is the second native species reported from China, and it differs from previously recorded species, i.e., *Erythroxylum sinense*, in having leathery (vs. chartaceous) leaf blades with faintly visible (vs. prominent) veins and white (vs. pinkish) petals with sub-rectangular (vs. ligule-like) appendages.

Erythroxylum species, especially *E. coca* Lam. and *E. novogranatense*, are well known for their tropane alkaloids, such as cocaine (Aynilian et al. 1974). According to Lv et al. (2022), 383 compounds, including diterpenes, triterpenes, flavonoids, alkaloids, and other derivatives, have been found in 67 *Erythroxylum* species. Investigating this new taxon for its phytochemical constituents could lead to the discovery of novel sources of these compounds, as well as possibly new compounds unknown to science.

Acknowledgements

We are grateful to Mr. Feng Ling and Mr. Yue-Yao Liu for their help during field trips. Our thanks also go to the curators of IBSC for allowing us to examine their specimens. This study was funded by the Guangdong Provincial Special Fund for Natural Resource Affairs on Ecology and Forestry Construction with the project “Survey of Woody Plant Germplasm Resources in Guangdong”, the National Survey of Traditional Chinese Medicine Resources from National Administration of Traditional Chinese Medicine (grant no. GZY-KJS-2018-004) and Forestry Science and Technology Innovation Specific Project of Guangdong Province (grant no. 2018KJ CX002).

References

Aynilian GH, Duke JA, Gentner WA, Farnsworth NR (1974) Cocaine content of *Erythroxylum* species. *Journal of Pharmaceutical Sciences* 63(12): 1938–1939. <https://doi.org/10.1002/jps.2600631223>

- Chung RCK (1996) Erythroxylaceae. In: Soepadmo S, Wong KM, Saw LG (Eds) Tree Flora of Sabah and Sarawak. Vol. 2. Ampang Press, Kuala Lumpur, 168–174. <https://doi.org/10.26525/TFSS2010>
- Guillaumin A (1907) Érythroxylacées. In: Gagnepain F, Humbert H, Lecomte H (Eds) Flore Générale de l'Indo-Chine. Vol. 1. Masson, Paris, 589–592.
- Harwood B, Chayamarit K (2011) Erythroxylaceae. In: Santisuk T, Larsen K (Eds) Flora of Thailand. Vol. 11. The Forest Herbarium, Bangkok, 8–13.
- Hô PH (2003) An Illustrated Flora of Vietnam. Vol. 2. published by author, Ho Chi Minh, 951 pp.
- IUCN Standards and Petitions Committee (2022) Guidelines for using the IUCN Red List Categories and Criteria. Version 15. Prepared by the Standards and Petitions Committee. <http://www.iucnredlist.org/documents/RedListGuidelines.pdf>
- Komada N, Tagane S, Nguyen VN, Hoang TH, Hoang TS, Toyama H, Nagamasu H, Naiki A, Yahara T (2018) *Erythroxylum calyptratum* (Erythroxylaceae), a new species from Mt. Fansipan, northern Vietnam. *Phytotaxa* 347(4): 279–284. <https://doi.org/10.11646/phytotaxa.347.4.4>
- Liu QR, Bartholomew B (2008) Erythroxylaceae. In: Wu ZY, Raven PH, Hong DY (Eds) Flora of China. Vol 11. Science Press, Beijing & Missouri Botanical Garden Press, St. Louis, 39–40.
- Lv YL, Tian T, Wang YJ, Huang JP, Huang SX (2022) Advances in chemistry and bioactivity of the genus *Erythroxylum*. *Natural Products and Bioprospecting* 12(1): e15. <https://doi.org/10.1007/s13659-022-00338-z>
- POWO (2022) *Erythroxylum* P.Browne. <https://powo.science.kew.org/taxon/24933-1> [accessed 30 March 2022]