

Rubus dianchuanensis sp. nov. (Rosaceae) from Sichuan and Yunnan, southwest China

Qiu-Ping Wang¹, Yu-Ran Li¹, Qiang-Chun Huang², Huan-Chong Wang^{1,3}

1 School of Ecology and Environmental Science, Yunnan University, Kunming 650500, Yunnan, China

2 School of Life Sciences, Yunnan University, Kunming 650500, Yunnan, China **3** Herbarium of Yunnan University, Kunming 650091, Yunnan, China

Corresponding author: Huan-Chong Wang (hchwang@ynu.edu.cn)

Academic editor: A. Sennikov | Received 15 February 2022 | Accepted 22 March 2022 | Published 01 April 2022

Citation: Wang Q-P, Li Y-R, Huang Q-C, Wang H-C (2022) *Rubus dianchuanensis* sp. nov. (Rosaceae) from Sichuan and Yunnan, southwest China. *PhytoKeys* 193: 141–150. <https://doi.org/10.3897/phytokeys.193.82287>

Abstract

Rubus dianchuanensis, a new name for the species previously named as *R. vicarius* by W. O. Focke in 1911, is proposed. A detailed description, illustrations, and remarks on its phenology, ecology, and geographic distribution are provided. This raspberry (subg. *Idaeobatus*) is endemic to China and was only found in Sichuan and Yunnan, southwest China. Morphologically, it is most similar to *R. ovatisepalus* but clearly differs from the latter by having leaf densely white or grey tomentose abaxially, usually leaf-like bracts at the base of inflorescence, 1–6 cm long pedicels, and triangular-ovate sepals with acute to caudate apex.

Keywords

Endemism, later homonym, *Rubus subornatus*, synonymy, taxonomy

Introduction

The genus *Rubus* was established by Linnaeus (1753) in his *Species Plantarum* and ten species were described there. Currently, about 700 species of *Rubus* are recognized, making it one of the largest genera of Rosaceae (Robertson 1974; Lu and Boufford 2003). Plants of this genus are usually shrubs, rarely subshrubs or perennial herbs, and more or less prickly. Their leaves are compound or simple, flowers are pentamerous and mostly bisexual, fruits are fleshy aggregates of drupelets, and tori are usually convex, conical or cylindrical (Kalkman 1993; Lu and Boufford 2003; Wang and Wang 2019).

Members of *Rubus* are distributed nearly worldwide except for Antarctica and can be found in most types of land biomes from tropical to subarctic regions (Gustafsson 1942; Spies and Du Plessis 1984; Hummer 1996; Lu and Boufford 2003). There are more than 250 species of *Rubus* in East Asia, and this region is the center of diversity for the subgenera *Malachobatus* and *Idaeobatus* (Wang and Wang 2019). More than 200 species are recorded in China, and most of them occur in the southern and southwestern provinces (Lu and Boufford 2003). Recently, new species and nomenclatural changes of *Rubus* in China have been constantly reported (e.g., Huang and Hu 2009; Byalt 2011; Sun and Boufford 2012; Wang et al. 2013, 2017, 2019; Wang and Wang 2019).

During our fieldwork and the herbarium studies on a taxonomic revision for the Chinese species of *Rubus*, we encountered a raspberry difficult to assign to any species recognized by Yu and Lu (1985) and Lu and Boufford (2003). Further research showed that it should be identified as *R. vicarius* Focke, which had been synonymized with *R. subornatus* Focke previously (Yu and Lu 1985; Lu and Boufford 2003). This plant represents a separate species, therefore, should be resurrected. Nevertheless, Focke's name is a later homonym of *R. vicarius* Sudre (1902); consequently, a new name for this distinctive species is required.

Materials and methods

We studied the newly named species both in the field and the herbaria. Type specimens (or type photos) of accepted names and their synonyms in *Rubus* subg. *Idaeobatus* were extensively examined and compared, as well as herbarium materials from CDBI, IBSC, KUN, P, PE, PYU and YUKU (acronyms after Thiers 2022). Pertinent taxonomic literature (e.g., Focke 1877, 1910, 1911, 1914; Yu and Lu 1985; Lu and Boufford 2003) were extensively consulted. Morphological studies were carried out on dried material under a stereomicroscope, and measurements were conducted using a ruler or a metric vernier calliper.

Taxonomy

***Rubus dianchuanensis* Huan C. Wang & Q. P. Wang, sp. nov.**

urn:lsid:ipni.org:names:77296909-1

Figs 1, 2, 3A1–A5

Type. China. Sichuan Province: Liangshan Prefecture, Muli County, on the way from Wujiao to Yiji, 27°58'21.73"N, 100°41'51.20"E, 3300–3500 m a.s.l., 23 July 2021, Q. P. Wang et al. ML12992 (holotype YUKU!, isotypes YUKU!).

Rubus vicarius Focke in Sargent, Pl. Wils. 1: 56. 1911, *nom. illeg.*, non Sudre (1902: 12). Type: China. Sichuan Province, Leshan City, Wa Shan, in thickets, 1500–2100 m a.s.l., July to August 1908, E. H. Wilson 948 (BM!, NYBG!, US!).

Diagnosis. *Rubus dianchuanensis* is most similar to *R. ovatisepalus* Huan C. Wang, but clearly differs from the latter by its leaf abaxially densely white or grey tomentose, bracts in the inflorescence often leaf-like, pedicels 1–6 cm long, sepals triangular-ovate and with acute to caudate apex.

Description. Arching shrubs, 1 to 2 m high, deciduous or semi-evergreen. Stems with dense ferruginous glandular hairs and soft eglandular hairs. Branchlets cylindric, grey-green to brown, villous, with curved prickles and nearly straight, ferruginous, 1–2 mm long, glandular hairs. Leaves imparipinnate, usually 3-foliolate, rarely 5-foliolate. Stipules persistent, linear, 5–8 mm long, ca. 1.5 mm wide, pubescent, with glandular hairs, base shortly adnate to petioles. Petioles 0.5–4 cm long, petiolule of




Figure 1. *Rubus dianchuanensis* **A** habit **B** bract at the base of inflorescence **C** bract at the upper part of inflorescence **D** calyx **E** flower (side view) **F** aggregate fruit with persistent calyx.

terminal leaflets 1–3.5 cm long, lateral leaflets sessile or subsessile; petiolule and rachis with glandular hairs, intermixed pubescence and erect or recurved prickles. Leaf blades cordate or ovate-cordate in outline, papery, adaxially pubescent and with sparse glan-




Figure 2. *Rubus dianchuanensis* **A, B** habit **C** abaxial surfaces of leaves **D** portion of branchlet showing glandular hairs **E** flower **F** an immature aggregate fruit with calyx.

dular hairs, abaxially densely white or grey tomentose throughout, with sparse glandular hairs along veins. Terminal leaflets cordate, broadly ovate or ovate, 3–11 cm long, 2–7 cm wide, apex acute to acuminate, base rounded to subcordate; margin slightly lobed or not, double serrate; nervation pinnate, with 6–9 lateral veins on each side of the midrib. Lateral leaflets ovate or elliptic, apex acute, base cuneate to round, slightly oblique, 1.5–7 cm long, 1–5 cm wide, lateral veins 5–7 paired. Terminal inflorescences racemose-cymose, 4–10-flowered, 5–15 cm long; bracts at the base usually leaf-like, simple, ovate, ovate-lanceolate or lanceolate, with similar indumentum as the leaves,



Figure 3. *Rubus dianchuanensis* (A1–A5) and *R. subornatus* (B1–B5) A1 a flowering branch showing terminal inflorescence A2 portion of stem showing indumentum A3 bract at the base of inflorescence A4 flower (side view) A5 mature aggregate fruit B1 habit B2 portion of the stem B3 bracts B4 flower (side view) B5 mature aggregate fruit.

2.5–11 cm long, 1–5 cm wide, apex acute to acuminate, base rounded to subcordate; bracts at the upper portion linear, 4–12 mm long, ca. 1 mm wide, pubescent, with glandular hairs. Axillary flowers usually solitary, rarely 2–3-flowered. Pedicels 1–6 cm in length, densely pilose, with dense glandular hairs and curved prickles. Flowers 1–1.5 cm in diameter. Calyx grey-green or reddish, abaxially with soft hairs and glandular hairs; sepals triangular-ovate, erect or spreading after anthesis, 5–10 mm long, 2–4 mm wide, margin grey tomentose and entire, apex acute to caudate. Petals pink to white, obovate, 5–8 mm long, 4–5 mm broad, apex repand, base shortly clawed. Stamens numerous in 2 whorls; filaments linear, glabrous, ca. 5 mm long. Pistils numerous; ovaries sparsely pilose, styles glabrous. Aggregate fruit ovoid, orange-red to red.

Taxonomic notes. *Rubus dianchuanensis* was firstly collected by Ernest Henry Wilson in 1908 from Wa Shan (Leshan City) in western Sichuan, southwest China. Based on Wilson's collection, Focke (1911) published "*R. vicarius* n. form. (?)" with a Latin description in his monograph *Species Ruborum*. However, the name *R. vicarius* Focke was not validly published there under Article 36.1 of the Shenzhen Code (Turland et al. 2018). Shortly afterwards, in July 1911, the name *R. vicarius* Focke was definitely accepted by Focke (in Sargent 1911) and accompanied by a complete and direct reference, namely "Bibl. Bot. LXXII 211 (Spec. Rub.) (1911)", to his previous Latin description, it was therefore validated. Unfortunately, the name *R. vicarius* had been previously used by Sudre (1902) for a European species; thus, Focke's name as a later homonym was illegitimate (Article 53.1 of the Shenzhen Code).

Morphologically, *Rubus dianchuanensis* is most similar to *R. ovatisepalus* Huan C. Wang (Fig. 4), a species described recently from northwestern Yunnan and southeastern Xizang, southwest China (Wang and Wang 2019), in having dense glandular hairs throughout the plant and the racemose-cymose terminal inflorescences. However, *R. dianchuanensis* differs markedly from the latter by its leaf abaxially densely white or grey tomentose (*vs.* sparsely pubescent, with glandular hairs), bracts at base of the inflorescence usually leaf-like, ovate, ovate-lanceolate or lanceolate (*vs.* lanceolate to linear), 2.5–11×1–5 cm (*vs.* 0.7–1.2×0.1–0.2 cm), flower usually larger, 1–1.5 cm (*vs.* 0.8–1.2 cm) in diameter, pedicels 1–6 cm (*vs.* 0.7–1.5 cm) long, apex of sepals acute to caudate (*vs.* long acuminate to caudate).

Rubus dianchuanensis is also similar to *R. subornatus* Focke (including its variety *R. subornatus* var. *melanodenus* Focke) (Fig. 3B1–B5), with which *R. vicarius* Focke had been erroneously synonymized by Yu and Lu (1985) as well as Lu and Boufford (2003). Nevertheless, *R. dianchuanensis* differs from it by its not glaucous stems usually covered with dense ferruginous glandular hairs and soft eglandular hairs (*vs.* more or less glaucous, glabrous), terminal inflorescences racemose-cymose (*vs.* corymbose), bracts at base of the inflorescence often leaf-like, rarely trifoliolate, ovate, ovate-lanceolate or lanceolate (*vs.* linear) (Fig. 3: A3, B3), pedicels usually longer, 1–6 cm (*vs.* 1–2.5 cm) long, calyx abaxially with grey pubescent (*vs.* intermixed tomentose) and dense ferruginous glandular hairs (*vs.* sparse or not), without needle-like prickles (*vs.* with needle-like prickles), and slightly pink to white (*vs.* purplish-red) petals. Some specimens of *R. dianchuanensis* had been identified as *R. phoenicolasius* Maxim., but it



Figure 4. *Rubus ovatisepalus* **A** habit **B** abaxial surfaces of leaflets **C** mature aggregate fruit.

is well differentiated from the latter by stems and branches with short glandular hairs (*vs.* long glandular hairs) and without bristles (*vs.* dense bristles), terminal inflorescence racemose-cymose (*vs.* short racemes) 5–15 cm (*vs.* 1–6 cm) long, pedicels 1–6 cm (*vs.* 0.5–1.5 cm) long, flowers 1–1.5 cm (*vs.* 0.6–1.5 cm) in diameter, calyx without bristles (*vs.* with dense bristles), sepals triangular-ovate (*vs.* lanceolate). A detailed morphological comparison between these four species is summarized in Table 1.

Phenology. *Rubus dianchuanensis* flowering from June to August, fruiting from July to September.


Etymology. The specific epithet “dianchuanensis” refers to the Yunnan (called *dian* for short in Chinese) and Sichuan (called *chuan* for short in Chinese) provinces, where this species occurs.

Distribution and habitat. *Rubus dianchuanensis* is endemic to southwest China, where it has been collected from western Sichuan and northwestern Yunnan (Fig. 5). It usually occurs at elevations ranging from 2500–3600 meters and grows in open woods and thickets.

Additional specimens examined (Paratypes): CHINA. **Sichuan:** Muli County, Damadian, 3000 m a.s.l., 16 Aug 1937, *T. T. Yu* 7740 (PE); Shimian County, Liziping Country, 2700 m a.s.l., 28 Jul 1978, *Shimian Team* 78-0875 (SM); Shimian County, 1955, *C. C. Hsieh* 41198 (IBSC, PE); Kangding City, Zheduotang village, 3100 m a.s.l., 1 Aug 1963, *K. C. Kuan et all* 1218 (PE); same location, 3450 m a.s.l., 5 July 1953, *X. L. Jiang* 36185 (IBK, IBSC, PE); same location, 3600 m a.s.l., 16 Jul 1953, *W. P. Fang & X. L. Jiang* 36323 (IBK, ISBC, PE); Kangding County, 2750 m a.s.l., 24 Jun 1980, *Z. J. Zhao* 112962 (CDBI) and *Z. J. Zhao* 119262 (PE); Kangding

Table 1. A morphological comparison of *Rubus dianchuanensis* with its relatives.

	<i>R. dianchuanensis</i>	<i>R. ovatisepalus</i>	<i>R. subornatus</i>	<i>R. phoenicolasius</i>
Indumentum of stems	dense glandular hairs	dense glandular hairs	glabrous	dense glandular hairs and bristles
Abaxial indumentum of leaf blade	densely grey tomentose	sparsely pubescent, with glandular hairs	densely grey tomentose	densely grey tomentose
Terminal Inflorescence	racemose-cymose	racemiform cymes	corymbose	short racemes
Length of pedicel	1–6 cm	0.7–1.5 cm	1–2.5 cm	0.5–1.5 cm
Diameter of flower	1–1.5 cm	0.8–1.2 cm	2–3 cm	0.6–1.5 cm
Petal colour	white or slightly pink	white or slightly pink	purplish-red	white
Petal vs. sepal	petal slightly longer than sepals	petal shorter than sepals	petal shorter than sepals	petal much shorter than sepals

**Figure 5.** Geographical distribution of *Rubus dianchuanensis* (red dots).

City, Yajiageng, Laoyunshachang, 3318 m a.s.l., 101°58'17"E, 29°56'00"N, 28 Aug 2008, *Y. L. Peng & W. G. Tu Gaoxf-0856* (KUN); Xiaojin County, 3500 m a.s.l., 1 Jul 1959, *Z. G. Liu 0412* (CDBI, PE); Lixian County, 26 Aug 1957, *X. Li 74160* (IBSC, PE, NAS); Lixian County, Miyaluo village, 25 Jul 1958, *Z. L. Wu 33375* (PE; CAF);

Barkam City, Barkam County, Dalangjiao River, 2300–2900 m a.s.l., 12 Jul 1960, *Sichuan Medicine Source Survey Team 22297* (NAS, SM); Barkam City, 2800 m a.s.l., 11 Jul 1957, *H. F. Zhou & Z. Y. Zhang 22772* (IBSC, NAS, KUN, PE); Barkam City, Dalangzugou, 2700 m a.s.l., 27 Aug 1957, *X. Li 72288* (IBSC, NAS, PE); Heishui County, Naizigou, 2900 m a.s.l., 22 Jul 1957, *X. Li 73260* (IBSC, NAS, KUN, PE). **Yunnan:** Ninglang County, Lugu Lake, 27°39'21"N, 100°48'36"E, 2500–2600 m a.s.l., 6 Aug 2015, *H. C. Wang et al. LGH8164* (YUKU).

Acknowledgements

We would like to thank the curators and staff of the herbaria from which specimens have been used in this study and Dr. Piotr Kosiński for his comments on the manuscript. We are grateful to Dr. Zhang Libing (Missouri Botanical Garden) for his valuable discussion on the nomenclature of *Rubus vicarius* Focke. This research was supported by the National Natural Science Foundation of China (grant no: 31960040) and the Second Tibetan Plateau Scientific Expedition and Research (STEP) programme (2019QZKK0502).

References

- Byalt VV (2011) *Rubus laue* Byalt, a new name for *R. multisetosus* T.T. Yu et L.T. Lu (Rosaceae). *Novosti Sistematiki Vysshikh Rastenii* 42: 232–234.
- Focke WO (1877) Synopsis *Ruborum* Germaniae. die deutschen Brombeerarten ausführlich beschrieben und erläutert, Müller Verlagsbuchhandlung, Bremen, 434 pp.
- Focke WO (1910) Species *Ruborum*. Monographiae generis Rubi prodromus I. *Bibliotheca Botanica* 17 (Heft 72): 1–120. <https://doi.org/10.5962/bhl.title.15533>
- Focke WO (1911) Species *Ruborum*. Monographiae generis Rubi prodromus II. *Bibliotheca Botanica* 17 (Heft 72): 121–223. <https://doi.org/10.5962/bhl.title.15533>
- Focke WO (1914) Species *Ruborum*. Monographiae generis Rubi prodromus III. *Bibliotheca Botanica* 19 (Heft 83): 1–274.
- Gustafsson A (1942) The origin and properties of the European blackberry flora. *Hereditas* 28(3–4): 249–277. <https://doi.org/10.1111/j.1601-5223.1942.tb03279.x>
- Huang JY, Hu JM (2009) Revision of *Rubus* (Rosaceae) in Taiwan. *Taiwania* 54: 285–310.
- Hummer KE (1996) *Rubus* diversity. *HortScience* 31(2): 182–183. <https://doi.org/10.21273/HORTSCI.31.2.182>
- Kalkman C (1993) Rosaceae.—In: van Steenis CGGJ (Ed.) *Flora Malesiana*, ser. 1, spermatophyta, Vol. 11(2). Leiden University, Leiden, 227–351.
- Linnaeus C (1753) *Species Plantarum*. L. Salvius, Stockholm, 1200 pp.
- Lu LD, Boufford DE (2003) *Rubus* Linnaeus. In: Wu ZY, Raven PH (Eds) *Flora of China* 9. Science Press, Beijing & Missouri Botanical Garden Press, St. Louis, 195–285.
- Robertson KR (1974) The genera of Rosaceae in the Southeastern United States. *Journal of the Arnold Arboretum* 55(2): 303–332, 344–401, 611–662. <https://doi.org/10.5962/p.67288>

- Sargent CS (1911) *Plantae Wilsonianae: an enumeration of the woody plants collected in western China for the Arnold Arboretum of Harvard University Part 1*. Cambridge University Press, Cambridge, 144 pp.
- Spies JJ, Du Plessis H (1984) The genus *Rubus* in South Africa, I. Chromosome numbers and geographical distribution of species. *Bothalia* 15(3/4): 591–596. <https://doi.org/10.4102/abc.v15i3/4.1853>
- Sudre H (1902) Notes critiques sur les plantes distribuées: N°287 *Rubus vicarius* Sud. *Bulletin de l'Association Pyrénéenne pour l'Échange des Plantes* 10: e12.
- Sun Y, Boufford DE (2012) *Rubus naruhashii* (Rosaceae), a new name for *R. clivicola* E. Walker. *Shokubutsu Kenkyu Zasshi* 87: 135–136.
- Thiers B (2022) [continuously updated] Index Herbariorum: a global directory of public 22 herbaria and associated staff. New York Botanical Garden's Virtual Herbarium. <http://sweetgum.nybg.org/ih>
- Turland NJ, Wiersema JH, Barrie FR, Greuter W, Hawksworth DL, Herendeen PS, Knapp S, Kusber WH, Li DZ, Marhold K, May TW, McNeill J, Monro AM, Prad J, Price MJ, Smith GF (2018) International Code of Nomenclature for algae, fungi, and plants (Shenzhen Code) adopted by the Nineteenth International Botanical Congress Shenzhen, China, July 2017. *Regnum Vegetabile* 159. Koeltz Botanical Books, Glashütten. <https://doi.org/10.12705/Code.2018>
- Wang HC, Wang QP (2019) *Rubus ovatisepalus* (Rosaceae), a new species from Yunnan and Xizang, southwest China. *Annales Botanici Fennici* 56(4–6): 227–230. <https://doi.org/10.5735/085.056.0405>
- Wang HC, Wang YH, Sun H (2013) Nomenclatural changes in *Rubus* (Rosaceae) mostly from China. *Phytotaxa* 114(1): 58–60. <https://doi.org/10.11646/phytotaxa.114.1.6>
- Wang HC, Zhang RZ, Liang ZL, He ZR (2017) A new species and two new synonyms in Chinese *Rubus* (Rosaceae). *Annales Botanici Fennici* 54(1–3): 105–109. <https://doi.org/10.5735/085.054.0316>
- Wang QP, Yang F, Wang HC (2019) A new species and new synonym in *Rubus* subgenus *Cylactis* (Rosaceae). *Phytotaxa* 400(1): 043–047. <https://doi.org/10.11646/phytotaxa.400.1.6>
- Yu TT, Lu LT (1985) *Rubus* Linnaeus. In: Yu TT (Ed.) *Flora Reipublicae Popularis Sinicae* 37. Science Press, Beijing, 10–218.