

Impatiens liupanshuiensis (Balsaminaceae), a new species from Guizhou, China

Tao-Hua Yuan¹, Yi Chen¹, Shuang Yu¹, Liu-Yi Ren¹,
Rong-Xin Huang¹, Mei-Jun Li¹, Xin-Xiang Bai¹

¹ Forestry College, Guizhou University CN-5500252 Guiyang, China

Corresponding author: Xin-Xiang Bai (254715174@qq.com)

Academic editor: L. J. Dorr | Received 29 October 2021 | Accepted 19 February 2022 | Published 10 March 2022

Citation: Yuan T-H, Chen Y, Yu S, Ren L-Y, Huang R-X, Li M-J, Bai X-X (2022) *Impatiens liupanshuiensis* (Balsaminaceae), a new species from Guizhou, China. *PhytoKeys* 192: 37–44. <https://doi.org/10.3897/phytokeys.192.77269>

Abstract

Impatiens liupanshuiensis (Balsaminaceae), belonging to *I.* subgen. *Impatiens*, is recognised as a new species from Guizhou, China and it is described and illustrated. It is morphologically similar to *I. xanthocephala* W.W. Sm. in its yellow flowers, extremely small basal lobes on lateral united petals, broadly-dolabriform distal lobes and funnelliform lower sepal. However, it is distinctive in the number of lateral sepals, teeth on the margin of lateral sepals, the recurvature of the dorsal petal, the number of lateral veins, the shape and size of the lamina and the type of lamina margin. A detailed description of the new species and colour photographs are provided. Its geographical distribution and morphology are also compared to similar species.

Keywords

China, Flora of Guizhou, *Impatiens*, morphology, taxonomy

Introduction

The genus *Impatiens* Linnaeus (1753: 937–938) (Balsaminaceae) contains over 1000 species distributed primarily in the Old World tropics and subtropics (Grey-Wilson 1980; Yu 2012). A few temperate outliers have also been identified in the

Northern Hemisphere (Yu et al. 2016). Five conspicuous diversification centres in the Paleotropical Regions can be recognised: tropical Africa; Madagascar; southern India and Sri Lanka; the eastern Himalaya; and Southeast Asia area in the broad sense (Song et al. 2003; Yuan et al. 2004). In China, 349 species of *Impatiens* have been recorded (Yuan et al. in press), the overwhelming majority of them being distributed in the southwest region, including Yunnan, Sichuan, Tibet and Guizhou (Chen 2001; Yu 2012). Since 2021, according to our incomplete statistics, eight new species of *Impatiens* have been described from China (Gu et al. 2021; Liao et al. 2021; Peng et al. 2021a, b; Song et al. 2021a, b, c). In Guizhou, 61 species of *Impatiens* have been recorded (Peng et al. 2021a; Yu et al. 2021). Additionally, there is one other species of *Impatiens* from Guizhou currently in press (Ren et al. 2022).

It is well known that *Impatiens* species are notoriously difficult to classify (Hooker 1908; Grey-Wilson 1980) due to their semi-succulent stems and fleshy leaves, which make it difficult to dry and preserve herbarium specimens. As flowers are extremely fragile, they are difficult to separate and reconstruct from dried specimens, especially when critical floral parts are folded and coalesced. However, since the floral morphology is hyper-variable, the shape and size of petals and sepals are indispensable for identification. Thus, during field exploration, it is essential to make detailed records and separate the parts of the flowers on soft paper in situ.

In September 2016, during a botanical investigation in Zhongshan District, western Guizhou Province, China, we encountered an unusual species, which is only distributed in the *Fargesia spathacea* Franch. (Poaceae) community in Xiaojiucaiping Mountain, the highest mountain of Guizhou (up to 2900 m alt.). The population of this unusual species is large. It resembles *Impatiens xanthocephala* W.W. Sm. (1920: 207–208) (Smith 1920) and *I. undulata* Y.L.Chen & Y.Q.Lu (1990: 23–25) (Chen and Lu 1990) in its yellow flowers, but the highly recurved dorsal petal is distinctive, allowing the anthers to be fully exposed. In September 2020 and August 2021, we revisited Xiaojiucaiping Mountain for a further field investigation to record morphological characters of the species. After careful examination of the relevant specimens and literature (Xiong and Luo 1989; Chen 2001; Chen et al. 2007; Yu 2012; Kuang et al. 2014; Kuang 2015; Luo and Deng 2015; Peng et al. 2021a), the authors decided that the species was hitherto undescribed and close to *I. xanthocephala* W.W. Sm. Hence, it is described here as a new species with a detailed description and illustrations.

Materials and methods

The material for this study was mainly collected in the field at the type locality of the new species in 2020. Herbarium specimens were made carefully. The morphological characteristics of the new species were measured from fresh material and dried herbarium specimens by use of a ruler.

Taxonomic treatment

Impatiens liupanshuiensis X.X.Bai & T.H.Yuan, sp. nov.

urn:lsid:ipni.org:names:77295796-1

Figs 1, 2

Type. CHINA. Guizhou: Liupanshui City, Zhongshan District, Xiaojiucaiping Mountain, 2887 m alt., 26°51'8.76"N, 104°41'33.67"E, 24 Aug 2021, *Xin Xiang Bai et al.* BXX368 (holotype: GZAC!; isotype: PE!).

Diagnosis. This species is similar to *Impatiens xanthocephala* W.W. Sm. in its yellow flowers, funnellform lower sepal, broadly-dolabriform distal lobes of lateral united petals and extremely small basal lobes, but it can be distinguished by having 2 (vs. 4) lateral sepals with ciliate (vs. entire) margin, dorsal petal recurved (vs. patent), lateral veins 8–10 pairs [vs. 2–3(4) pairs], apex of the distal lobes of the united petals retuse (vs. rounded), margin serrulate (vs. remotely crenate or shallowly undulate), lamina ovate-lanceolate or lanceolate (vs. ovate or ovate-oblong), 5–10.5 × 1.7–2.8 cm (vs. 1–2 × 0.8–1 cm) and lamina margin serrulate (vs. remotely crenate or shallowly undulate).

Description. Perennial herb, 30–70 cm tall. Stem erect or procumbent basally, branched, upper part brown pubescent. Leaves alternate, petiolate, petioles 0.3–1 cm long. Lamina 5–10.5 × 1.7–2.8 cm, ovate-lanceolate or lanceolate, apex acuminate, base cuneate, with 1–3 pairs of basal glands, margin serrulate, teeth ending in setae, adaxial surface remotely puberulous on veins, abaxial surface glabrous, lateral veins 8–10 pairs. Inflorescences axillary, 2-flowered; peduncles 2–4 cm long, pedicels 1.5–2 cm long, bracteate above middle; bracts persistent, lanceolate, 2–4 mm long, margin denticulate. Flowers yellow, ca. 2 cm deep. Lateral sepals 2, obliquely ovate or ovate, inequilateral, 3–5 mm long, 2–3 mm wide, remotely denticulate on one or both sides, abaxial mid-vein slightly thickened, green, hyaline both sides, apex acuminate, aristate. Lower sepal funnellform, 0.5 cm deep excluding spur, mouth vertical, 0.5–0.9 cm wide, base abruptly contracted into an incurved spur, spur 1–1.5 cm long. Dorsal petal orbicular, 8–10 mm in diam., recurved at anthesis, abaxial mid-vein carinate, with a curved rostrum towards apex, base truncate, apex retuse. Lateral united petals 2-lobed, sessile, 13–18 mm long, basal lobes oblong, extremely small, 1–2 mm wide, distal lobes broadly-dolabriform, 1–1.3 cm long, 0.9–1 cm wide, apex retuse, base with reddish patches; auricle inflexed, small. Stamens 5, filaments linear, anthers obtuse. Ovary erect, fusiform, 5-carpellate. Capsule cylindrical, 1.5–2 cm long, seeds ellipsoid or ovoid, surface with warty protrusions.

Phenology. This species was observed flowering from July to October and fruiting from September to November.

Distribution. This species is currently known from only one population in Xiaojiucaiping Mountain, Zhongshan District, Liupanshui City, Guizhou, China. The population is large, with about 3000–3500 individuals.




Figure 1. *Impatiens liupanshuiensis* **A** habit **B** population **C, D** flower in face view **E, F** flower in lateral view **G** flower in dorsal view **H** flower dissected **I** lateral sepals (remotely denticulate on one or both sides) **J** leaves **K** stamens and pistils **L** different phases of the recurvature of the dorsal sepal **M** capsules **N** seeds. Photographs by X. X. Bai.




Figure 2. *Impatiens liupanshuiensis* **A** plant **B** flower in face view **C** flower in lateral view **D** lateral sepals **E** bract **F** dorsal sepal **G** lateral united petal **H** lower sepal **I** capsule **J** stamens and pistils. (Drawn by Y. Chen from holotype and corresponding colour photographs).

Ecology. This species was collected growing in *Fargesia spathacea* shrubs on the side of the plank road at an elevation of 2730–2887 m in Xiaojiucaiping Mountain, which is the highest mountain in Guizhou. The main associated species were *F. spathacea*, *Impatiens lecomtei*, *Sorbaria arborea* C.K. Schneid. (Rosaceae), *Pteridium revolutum* (Blume) Nakai (Dennstaedtiaceae), *Hypericum patulum* Thunb. (Hypericaceae), *Eupatorium lindleyanum* DC. (Asteraceae) and *Senecio scandens* Buch.-Ham. ex D. Don (Asteraceae).

Etymology. The specific epithet ‘liupanshuiensis’ refers to the type locality, Liupanshui City, Guizhou, China. The Chinese name is given as “六盘水凤仙花”.

Conservation status. *Impatiens liupanshuiensis* is currently known only from its type locality, Zhongshan District, Guizhou, China, where it is distributed in several

places with about 3000–3500 individuals known in the population. As Xiaojiucaiping Mountain is a part of the Axilixi Landscape and Famous Scenery, it is an area exposed to significant human disturbance. The conservation status can be evaluated as Vulnerable (VU) C2a(ii), based on the IUCN Red List Categories and Criteria (IUCN 2019).

Discussion

Yu et al. (2016) divided the genus *Impatiens* into two subgenera, subgen. *Clavicarpa* S.X. Yu ex S.X. Yu & Wei Wang and subgen. *Impatiens*, based on the number of carpels, the number of ovules per carpel, the shapes of the capsule, the number of pollen apertures and the pollen grain shape in polar view. In our study, *I. liupanshuiensis* is nested within subgen. *Impatiens* by its 5-carpellate ovary, cylindrical capsule and many ovules per locule. The new species is morphologically similar to *I. xanthocephala*, but *I. liupanshuiensis* has two lateral sepals with remote teeth on the margin, a more peculiar dorsal petal and more lateral veins and its flowers are shorter than its lamina. *I. liupanshuiensis* is also similar to *I. undulata*. To better distinguish the new species morphologically from *I. xanthocephala* and *I. undulata*, we list more details in Table 1.

The distributions of these three species are geographically isolated from each other: *Impatiens liupanshuiensis* is confined to western Guizhou Province, *I. xanthocephala* is recorded in southwest Sichuan Province and *I. undulata* is recorded in central Sichuan Province and southern Chongqing Municipality (Yuan et al. in press).

Table 1. Comparison of morphological characters in *Impatiens liupanshuiensis*, *I. xanthocephala* (data from Smith 1920 and Chen et al. 2007) and *I. undulata* (data from Chen and Lu 1990 and Chen et al. 2007).

Characters/Species	<i>I. liupanshuiensis</i>	<i>I. xanthocephala</i>	<i>I. undulata</i>
Plant height	30–70 cm	10–16 cm	40–100 cm
Stem indumentum	upper part brown pubescent	glabrous	glabrous
Leaf indumentum	adaxial surface puberulous on veins	glabrous	glabrous
Lamina	ovate-lanceolate or lanceolate, 5–10.5 × 1.7–2.8 cm	ovate or ovate-oblong, 1–2 × 0.8–1 cm	ovate or ovate-orbicular, 2–7 × 1.5–5 cm
Lamina margin	serrulate	remotely crenate or shallowly undulate	undulate or obtusely crenate
Lateral veins	8–10 pairs	2–3(4) pairs	6–7 pairs
Flower size	small, ca. 2 cm deep	large, ca. 3.5 cm deep	small, 2 cm deep
Dorsal petal	orbicular, 8–10 mm, recurved, abaxial mid-vein carinate, a curved rostrum towards apex	suborbicular, 5–6 mm, abaxial mid-vein carinate at middle	suborbicular, 3–5 mm, abaxial mid-vein carinate
Lateral sepal	2, inequilateral, remotely denticulate on one side or both sides	4, outer 2 orbicular, inner 2 ovate, equilateral, entire	2, equilateral, entire
United lateral petals	apex of distal lobes retuse	apex of distal lobes rounded	apex of distal lobes retuse
Lower sepal	funnelform	funnelform	salverform
Anthers	obtuse	obtuse	acute

Acknowledgements

The authors are grateful to Prof. Yuanxin Xiong for assistance on species identification and for providing valuable suggestions. We also thank Wenke Dong, Chenghua Yang and Chaoyi Deng for their helpful suggestions. Additionally, we would like to thank the anonymous reviewers and editors for their constructive comments and suggestions on our manuscript. This research was supported by the National Natural Science Foundation of China (31960328) and Investigation of Wild Ornamental Plant Resources in Guizhou (701256192201).

References

- Chen YL (2001) Balsaminaceae. In: Chen YL (Ed.) *Flora Reipublicae Popularis Sinica* (Vol. 47(2)), Sapindales. Science Press, Beijing, 1–243.
- Chen YL, Lu YQ (1990) Six new species of *Impatiens* from Sichuan. *Yunnan Zhi Wu Yan Jiu* 12(1): 22–30.
- Chen YL, Akiyama S, Ohba H (2007) Balsaminaceae. In: Wu ZY, Raven PH, Hong DY (Eds) *Flora of China* (Vol. 12). Science Press, Beijing & Missouri Botanical Garden Press, St. Louis, 43–113.
- Grey-Wilson C (1980) *Impatiens* of Africa. CRC Press, Rotterdam, 235 pp.
- Gu JZ, Yang N, Peng YC, Song YX, Guo SW, Cong YY (2021) *Impatiens shangjiangensis* (Balsaminaceae), a new species from Yunnan, China. *Annales Botanici Fennici* 58(4–6): 197–201. <https://doi.org/10.5735/085.058.0401>
- Hooker JD (1908) Les espèces du genre *Impatiens* dans l'herbier du muséum de Paris. *Nouvelles Archives du Muséum d'Histoire Naturelle, Paris, sér. 4, 10*: 233–272.
- IUCN (2019) Guidelines for using the IUCN Red List Categories and Criteria. Version 14. Prepared by the Standards and Petitions Subcommittee. <http://www.iucnredlist.org/documents/RedListGuidelines.pdf>
- Kuang RP, Duan LD, Gu JZ, Cai XZ, Cong YY, Liu KM (2014) *Impatiens liboensis* sp. nov. (Balsaminaceae) from Guizhou, China. *Nordic Journal of Botany* 32: 463–467. <https://doi.org/10.1111/j.1756-1051.2013.00194.x>
- Kuang RP (2015) Study on the flora in Guizhou/southern Sichuan and phylogenetic problems of *Impatiens* L. Ph.D. Thesis, Hunan Normal University, China.
- Liao RL, Cai L, Yu ZY, Wang YH, Sun WB (2021) *Impatiens wutaishanensis* (Balsaminaceae), a new species from Southeast Yunnan, China. *PhytoKeys* 176: 43–53. <https://doi.org/10.3897/phytokeys.176.58825>
- Linnaeus C (1753) *Species Plantarum* 2. Laurentius Salvius, Stockholm, 937–938.
- Luo Y, Deng LX [Eds] (2015) *Impatiens*. Guizhou Vascular Plant Catalogue. China Forestry Publishing House, Beijing, 342–344.
- Peng S, Cong YY, Tian J, Zhang CF, Hu GW, Wang QF (2021a) *Impatiens bullatisepala* (Balsaminaceae), a new species from Guizhou, China. *Phytotaxa* 500(3): 217–224. <https://doi.org/10.11646/phytotaxa.500.3.5>

- Peng S, Rono CP, Yang JX, Wang JJ, Hu GW, Wang QF (2021b) Description of a new species and lectotypification of two names in *Impatiens* Sect. *Racemosae* (Balsaminaceae) from China. *Plants* 10(9): e1812. <https://doi.org/10.3390/plants10091812>
- Ren LY, Chen Y, Yuan TH, Huang RX, Li MJ, Bai XX (2022) *Impatiens bijieensis* (Balsaminaceae), a new species from karst plateau in Guizhou, China. *PhytoKeys* 192: 1–10. <https://doi.org/10.3897/phytokeys.192.77517>
- Smith WW (1920) Diagnoses species novarum in herbario Horti Regi Botanici Edinburgensis cognitarum. CCCCLI-D. *Notes from the Royal Botanic Garden Edinburgh* 12: 206–207.
- Song Y, Yuan YM, Küpfer P (2003) Chromosomal evolution in Balsaminaceae, with cytological observations on 45 species from Southeast Asia. *Caryologia* 56(4): 463–481. <https://doi.org/10.1080/00087114.2003.10589359>
- Song YX, Peng S, Cong YY, Zheng YM (2021a) *Impatiens rapiformis*, a new species of *Impatiens* with root tuber from Yunnan, China. *Nordic Journal of Botany* 39(5): njb.03151. <https://doi.org/10.1111/njb.03151>
- Song YX, Cong YY, Kuang RP, Peng YC, Zheng YM (2021b) *Impatiens sterilis*, a rare new species of *Impatiens* with reduced sterile inflorescences from Yunnan, China. *Taiwania* 66(1): 48–52. <https://doi.org/10.6165/TAI.2021.66.48>
- Song YX, Xiao Y, Peng S, Cong YY, Hu GW (2021c) Two new species of *Impatiens* from China, and taxonomic insights into the *Longifilamenta* Group, which is endemic to China. *Plants* 10(8): e1697. <https://doi.org/10.3390/plants10081697>
- Xiong YX, Luo YC (1989). Balsaminaceae. In: Li YK (Ed.) *Flora Guizhouensis*, Tomus 7. Sichuan National Press, Chengdu, 626–641.
- Yu SX (2012) *Balsaminaceae of China*. Peking University Press, Beijing, 1–214.
- Yu SX, Janssens SB, Zhu XY, Lidén M, Gao TG, Wang W (2016) Phylogeny of *Impatiens* (Balsaminaceae): Molecular and morphological evidence into a new classification. *Cladistics* 32(2): 179–197. <https://doi.org/10.1111/cla.12119>
- Yu S, Li MJ, Yuan TH, Ren LY, Bai XX (2021) Species diversity and geographic distribution of wild *Impatiens* in Guizhou Province. *Xibei Zhiwu Xuebao* 41: 863–871.
- Yuan YM, Song Y, Geuten K, Rahelivololona E, Wohlhauser S, Fischer E, Smets E, Küpfer P (2004) Phylogeny and biogeography of Balsaminaceae inferred from ITS sequences. *Taxon* 53(2): 391–403. <https://doi.org/10.2307/4135617>
- Yuan TH, Li MJ, Ren LY, Huang RX, Chen Y, Bai XX (in press) A dataset on the diversity and geographical distributions of wild *Impatiens* in China. *Shengwu Duoyangxing*.