

Taxonomic note of *Parnassia* (Celastraceae) in China II: population surveys reveal that *P. guilinensis* is conspecific to *P. xinganensis*

Xiao-Song Dai^{1,2}, Dao-Zhang Min³, Bo Yang¹, Ding Wu¹, Bo Li³

1 Jiangxi Key Laboratory of Plant Resources and Biodiversity, Jingdezhen University, Jingdezhen 333000, China **2** School of Life Sciences, Nanchang University, Nanchang 330031, China **3** Research Centre of Ecological Sciences, College of Agronomy, Jiangxi Agricultural University, Nanchang 330045, China

Corresponding author: Ding Wu (parnassia@126.com); Bo Li (hanbolijx@163.com)

Academic editor: M.P. Simmons | Received 4 January 2021 | Accepted 30 January 2021 | Published 10 February 2021

Citation: Dai X-S, Min D-Z, Yang B, Wu D, Li B (2021) Taxonomic note of *Parnassia* (Celastraceae) in China II: population surveys reveal that *P. guilinensis* is conspecific to *P. xinganensis*. *PhytoKeys* 172: 67–73. <https://doi.org/10.3897/phytokeys.172.62749>

Abstract

Based on investigation of populations of *Parnassia guilinensis* and *P. xinganensis*, examination of herbarium specimens (including types), as well as consultation of protologues and distributions, *P. guilinensis* is hereby reduced to a synonym of *P. xinganensis*. *P. xinganensis* is endemic to northeastern Guangxi Province of China and characterized by having elliptic to ovate leaves and staminodes 3–5-branched with globose glands. Field photographs and an updated morphological description of *P. xinganensis* are provided.

Keywords

Endemic species, morphology, staminode, synonymy, taxonomy

Introduction

Parnassia L. consists of small, glabrous and rosulate perennial herbs, with several morphologically distinguishable traits, including a solitary, terminal, bisexual and pentamerous flower borne on an unbranched scape; five showy staminodes; and one-by-one stamen movement (Ku 1987; Ku and Hultgård 2001; Sandvik and Totland 2003; Ren 2010; Armbruster et al. 2014; Ren and Bu 2014). Species of *Parnassia* predominantly

occur in arctic and alpine regions of Europe, Asia, and North America with a center of diversification in mountainous areas in Pan-Himalaya and southwest China (Phillips 1982; Ku 1987; Ku and Hultgård 2001; Wu et al. 2003; Simmons 2004; Wu 2005). According to the most updated checklist, *Parnassia* contains 61 species, 2 subspecies, 11 varieties and 1 form (Shu et al. 2017). However, results from several taxonomic investigations after Shu et al. (2017) have changed the number of species in the genus. These changes include: *P. lanceolata* var. *oblongipetala* T.C. Ku was reduced to a synonym of *P. yunnanensis* Franch. (Shu and Zhang 2017), *P. venusta* Z.P. Jien, *P. degeensis* T.C. Ku and *P. kangdingensis* T.C. Ku were reduced to synonyms of *P. cacuminum* Hand.-Mazz. (Shu and Zhang 2017), *P. chengkouensis* T.C. Ku and *P. dilatata* Hand.-Mazz. were reduced to synonyms of *P. wightiana* Wall. ex Wight et Arn. (Wang et al. 2018), *P. brevistyla* (Brieg.) Hand.-Mazz. and *P. leptophylla* Hand.-Mazz. were reduced to synonyms of *P. delavayi* Franch. (Yu et al. 2018), *P. tibetana* Z.P. Jien ex T.C. Ku, *P. nubicola* subsp. *occidentalis* Schönbeck-Temesy and *P. nubicola* var. *nana* T.C. Ku were reduced to synonyms of *P. nubicola* Wall. ex Royle (Ma et al. 2020). Meanwhile, some new species were described, including *P. zhengyuana* M.X. Ren et J. Zhang and *P. simianshanensis* M.X. Ren, J. Zhang et Z.Y. Liu (Zhang et al. 2019).

In the taxonomy of *Parnassia*, the shape of basal leaves, the characteristics of petals (entire or flat, divided into lobes or filiform rays, respectively), and the shape of staminodes (i.e., number and depth of staminode branches, shape of staminode lobes, with globose glands at apex or not) were considered to be of great significance in species delimitation (Ku 1987; Ku and Hultgård 2001; Wu 2005). *P. guilinensis* G.Z. Li et S.C. Tang (Tang and Li 1999) and *P. xinganensis* C.Z. Gao et G.Z. Li (Gao and Li 1983) were described based on specimens collected from the same mountain: Mao'er Mountain, Xing'an County, Guilin City, Guangxi Province, China. The type specimen of *P. xinganensis* (11 December 1978, G.Z.Li 62923) was collected at an altitude of 1200 m on the mountain while that of *P. guilinensis* (23 August 1998, G.Z.Li and S.C. Tang M93) was collected at 580 m. Tang and Li (1999) noted that *P. guilinensis* was morphologically similar to *P. xinganensis* but could be distinguished from the latter by the staminodes being 5-lobed and the apex of lobes with globose glands, whereas the staminodes of *P. xinganensis* are 3- or rarely 4-lobed without globose glands at the lobe apex. In both *Flora Reipublicae Popularis Sinicae* (Ku 1995) and *Flora of China* (Ku and Hultgård 2001), the staminodes of *P. xinganensis* were described as eglandular. However, when examining the type specimen of *P. xinganensis* (Fig. 1), we found that the apex of the staminode lobes obviously bear globose glands and the number of lobes range from three to five, strongly indicating that *P. xinganensis* and *P. guilinensis* are the same species. In order to clarify the relationship of the two sympatric taxa, we visited the Mao'er Mountain in September and October of 2020 and conducted population surveys of *Parnassia* species from low to high altitudes. Population observations clearly revealed that there are no differences among individuals at 580 m and 1200 m (Fig. 2), confirming that the later described species *P. guilinensis* is conspecific with the former one, *P. xinganensis*. We thus reduce *P. guilinensis* to a synonym of *P. xinganensis*, and provide an updated description of *P. xinganensis* based on population observations.

Materials and methods

Type specimens of *P. guilinensis* and *P. xinganensis* deposited at IBK, as well as other *Parnassia* specimens collected from Mao'er Mountain, Xing'an County, Guangxi Province, China, preserved in GXMG, IBK, IBSC, KUN and PE (acronyms according to Thiers 2020+) were carefully examined under a stereo dissecting microscope (Stereo Zoom Leica S8 APO, Leica Microsystems 2017). Morphological traits were measured using a ruler and a micrometer based on both herbaria and fresh materials. Field investigations were carried out during September to October of 2020 in Mao'er Mountain and some individuals of *P. xinganensis* collected from different altitudes were transplanted to the greenhouse at Jingdezhen University for further observation and to be photographed. Voucher specimens were deposited at the herbarium of Jingdezhen University.

Taxonomic treatment

Parnassia xinganensis C.Z. Gao & G.Z. Li, 1983: 19.

Figs 1, 2

Type. CHINA. Guangxi: Guilin City, Xing'an County, Mao'er Mountain, streamsid es in valleys, alt. 1200 m, 11 December 1978, *G.Z.Li* 62923 (holotype: IBK00185227!; isotype: IBK00200466!).



Figure 1. Type specimens of *Parnassia xinganensis* (A) and *P. guilinensis* (B).




Figure 2. Morphological comparisons between *Parnassia xinganensis* (A1–A5) and *P. guilinensis* (B1–B5) collected from their type localities of Mao’er Mountain at altitudes of 1200 m and 580 m, respectively A1, B1 habitat A2, B2 flower A3, B3 calyces, staminodes and ovary A4, B4 petals A5, B5 variation of staminodes.

=*Parnassia guilinensis* G.Z. Li & S.C. Tang, syn. nov. Type: CHINA. Guangxi: Guilin City, Xing’an County, Mao’er Mountain, streamside, alt. 580 m, 23 August 1998, G.Z.Li & S.C.Tang M93 (holotype: IBK00200636!).

Description. Perennial herbs, glabrous. Rhizome sympodial, robust. Stems 1–8, 5–20 cm tall, usually with 1 cauline leaf near middle. Basal leaves (4–) 8–13 (–22); petiole (1–) 4.5–6.5 (–9.5) cm long; leaf blade elliptic, obovate-elliptic, oblong-ovate to ovate, abaxially gray-white, adaxially green, (1–) 3–5.5 (–7.5) × (0.8–) 1.5–2.5 (–3) cm, inconspicuously 5–7-veined on both surfaces, midvein prominent, base rounded, subtruncate to cuneate, apex obtuse to acute. Cauline leaf sessile, amplexicaul, ovate or ovate-triangular, 0.9–2.4 × 0.5–1.6 cm. Flowers 1.5–2.3 cm in diam.; hypanthium shortly campanulate or inconspicuous. Sepals green, elliptic to ovate, 3.7–5.2 × 2.3–3.7 mm, 5-veined, margin entire, apex obtuse. Petals spreading, white, elliptic to broadly obovate, 8.5–12 × 6.8–8.3 mm, 5-veined basally, base attenuate into a claw, 2–3 mm long, margin entire or slightly undulate, apex rounded, obtuse or emarginate. Anthers ellipsoid; filaments 2–7 mm long; staminodes flat, 3–3.5 mm long, 3–5-branched to middle, branchedes globose glandular at apex. Ovary superior, greenish, ovoid, slightly sunken into hypanthium; styles short, ca. 1–1.5 mm long; stigma 3-lobed, lobes oblong, spreading. Capsule ovoid, trigonous, 5–10 mm long, 3-valved. Seeds minute, oblong, ca. 1 mm long.

Phenology. Flowering – late June to November; fruiting – August to December.

Distribution and habitat. The species is endemic to northeastern Guangxi Province (recorded only in Xing’an County and Ziyuan County), China, and grows in clefts of the moist rocks along streams or under waterfalls, at an elevation of 400–1350 m.

Additional specimens examined. CHINA. Guangxi: Guilin City, Xing’an County, Mao’er Mountain, streamsides, moist rocks in valleys, alt. 580 m, 23 August 1998, *G.Z.Li & S.C.Tang M211* (IBK00200637!); under forests, alt. 611 m, 29 September 2014, *Xing’an Expedition 450325140929020LY* (GXMG0110865!); Ziyuan County, Mao’er Mountain, on rocks under forests, near streams, 6 December 1980, *G.Z.Li 10120* (IBK!).

Conservation status. At present, *P. xinganensis* has been reported only from two counties in northeastern Guangxi Province of China. Based on our field investigations, there are numerous mature individuals and young seedlings which could be easily discovered along streams and under waterfalls, indicating the population survives and regenerates well. Additionally, the Mao’er Mountain has been projected to a national nature reserve of China in 2003, and ranks among one of the earliest national nature reserves founded in Guangxi Province. It is apparent that the species will not be severely affected by human activities, thus we propose to list *P. xinganensis* as Least Concern (LC) according to the IUCN Red List categories and criteria (IUCN 2017).

Acknowledgements

We are grateful to the curators of GXMG, IBK, IBSC, KUN and PE for providing access to specimens; Dr. Li-Na Dong, Mr. Fang-Min Hu and Prof. Hui-Chun Du for their field assistance; Dr. Yu-Min Shu for sharing some *Parnassia* literature; and to Prof. Guang-Zhao Li for his comments on the manuscript. This work was supported by the National Natural Science Foundation of China (grants no. 41561014, 31900181).

References

- Armbruster WS, Corbet SA, Vey AJM, Liu SJ, Huang SQ (2014) In the right place at the right time: *Parnassia* resolves the herkogamy dilemma by accurate repositioning of stamens and stigmas. *Annals of Botany* 113(1): 97–103.
- Gao CZ, Li GZ (1983) New species of Saxifragaceae from Guangxi. *Guihaia* 3(1): 19–23.
- IUCN (2017) Guidelines for using the IUCN red list categories and criteria, ver. 13. IUCN Species Survival Commission.
- Ku TC (1987) A revision of the genus *Parnassia* (Saxifragaceae) in China. *Bulletin of Botanical Research* 7(1): 1–61.
- Ku TC (1995) *Parnassia*. In: Lu LT, Huang SM (Eds) *Flora Reipublicae Popularis Sinicae* (Vol. 35). Science Press, Beijing, 9–66.
- Ku TC, Hultgård UM (2001) *Parnassia*. In: Wu ZY, Raven PH (Eds) *Flora of China* (Vol. 8). Missouri Botanical Garden Press, St. Louis, Missouri, USA, and Science Press, Beijing, China, 95–118.
- Ma YH, Zhou HL, Shu YM (2020) Taxonomic note on *Parnassia* (Celastraceae): The identity of *P. nubicola*. *PhytoKeys* 154: 103–109. <https://doi.org/10.3897/phytokeys.154.54042>
- Phillips RB (1982) Systematics of *Parnassia* (Parnassiaceae): generic overview and revision of North American taxa. PhD Thesis. University of California, Berkeley.
- Ren MX (2010) Stamen movements in hermaphroditic flowers: Diversity and adaptive significance. *Journal of Plant Ecology* 34(7): 867–875.
- Ren MX, Bu ZJ (2014) Is there ‘anther–anther interference’ within a flower? Evidence from one-by-one stamen movement in an insect-pollinated plant. *PLoS ONE* 9(1): e86581. <https://doi.org/10.1371/journal.pone.0086581>
- Sandvik SM, Totland O (2003) Quantitative importance of staminodes for female reproductive success in *Parnassia palustris* under contrasting environmental conditions. *Canadian Journal of Botany* 81(1): 49–56. <https://doi.org/10.1139/b03-006>
- Shu YM, Zhang ZX (2017) Four new synonyms and a new combination in *Parnassia* (Celastraceae). *PhytoKeys* 77: 81–88. <https://doi.org/10.3897/phytokeys.77.11513>
- Shu YM, Yu HY, Mu XY, Zhang ZX (2017) Checklist and typification of *Parnassia* (Celastraceae). *Phytotaxa* 309(1): 1–22. <https://doi.org/10.11646/phytotaxa.309.1.1>
- Simmons MP (2004) Parnassiaceae. In: Kubitzki K (Ed.) *The families and genera of vascular plants* (Vol. 6). Springer Verlag, Berlin, 291–296. https://doi.org/10.1007/978-3-662-07257-8_33
- Tang SC, Li GZ (1999) Two new plant species and one new recorded genus from Guangxi. *Guihaia* 19(4): 341–343.
- Thiers B (2020+) [Continuously updated] Index Herbariorum: A global Directory of Public Herbaria and Associated Staff. New York Botanical Garden. <http://sweetgum.nybg.org/science/ih/>
- Wang YP, Chen S, Shu YM, Li B, Wu D (2018) Taxonomic note of *Parnassia* (Celastraceae) in China: The identity of *P. wightiana*. *Phytotaxa* 375(1): 130–134. <https://doi.org/10.11646/phytotaxa.375.1.10>

- Wu D (2005) Phylogeny and biogeography of the genus *Parnassia*. PhD Thesis. Kunming Institute of Botany, Chinese Academy of Science.
- Wu ZY, Lu AM, Tang YC, Cheng ZD, Li DZ (2003) The family and genera of angiosperms in China: a comprehensive analysis. Science Press, Beijing, 1209 pp.
- Yu HY, Guo FY, Shu YM, Zhang ZX (2018) Taxonomic note of *Parnassia* (Celastraceae) in China: A reassessment of Subsect. *Xiphosandra*. *PhytoKeys* 114: 43–54. <https://doi.org/10.3897/phytokeys.114.30551>
- Zhang YJ, Zhang J, Liu ZY, Ren MX (2019) *Parnassia zhengyuana* sp. nov. and *P. simianshanensis* sp. nov.: Two new species of *Parnassia* (Celastraceae) from karst caves and Danxia landform in southwest China. *Nordic Journal of Botany* 37(10): e02414. <https://doi.org/10.1111/njb.02414>