

Splitting the *Pisonia* birdcatcher trees: re-establishment of *Ceodes* and *Rockia* (Nyctaginaceae, Pisonieae)

Elson Felipe Sandoli Rossetto¹, Marcos A. Caraballo-Ortiz²

1 Departamento de Biologia Animal e Vegetal, Centro de Ciências Biológicas, Universidade Estadual de Londrina, Campus Universitário, Rodovia Celso Garcia Cid, PR 445, Km 380, 86057-970, Londrina, PR, Brazil **2** Smithsonian Institution, National Museum of Natural History, Department of Botany, MRC 166, Washington, DC 20013-7012, USA

Corresponding author: Elson Felipe Sandoli Rossetto (rossetto.felipe@gmail.com)

Academic editor: Alexander Sukhorukov | Received 29 January 2020 | Accepted 8 April 2020 | Published 7 July 2020

Citation: Rossetto EFS, Caraballo-Ortiz MA (2020) Splitting the *Pisonia* birdcatcher trees: re-establishment of *Ceodes* and *Rockia* (Nyctaginaceae, Pisonieae). *PhytoKeys* 152: 121–136. <https://doi.org/10.3897/phytokeys.152.50611>

Abstract

Several genera of Nyctaginaceae, currently merged under *Pisonia*, have been described for the Indo-Pacific region. Results from a recent molecular phylogenetic study of tribe Pisonieae showed that *Pisonia* is non-monophyletic and comprises three well-supported lineages: one including typical *Pisonia* and allies (*Pisonia* s.str.), a clade of species which corresponds to the original description of *Ceodes* and a third lineage whose single representative was formerly treated under the monotypic genus *Rockia*. Thus, as part of an effort to achieve a natural classification for tribe Pisonieae, this work proposes to re-establish *Ceodes* and *Rockia* to accommodate taxa with inconspicuous glands on anthocarps, recognising 21 species (20 for the former and one for the latter), of which 16 are new combinations: *Ceodes amplifolia* **comb. nov.**, *Ceodes artensis* **comb. nov.**, *Ceodes austro-orientalis* **comb. nov.**, *Ceodes brownii* **comb. nov.**, *Ceodes cauliflora* **comb. nov.**, *Ceodes coronata* **comb. nov.**, *Ceodes diandra* **comb. nov.**, *Ceodes gigantocarpa* **comb. nov.**, *Ceodes gracilescens* **comb. nov.**, *Ceodes lanceolata* **comb. nov.**, *Ceodes merytifolia* **comb. nov.**, *Ceodes muelleriana* **comb. nov.**, *Ceodes rapaensis* **comb. nov.**, *Ceodes sechellarum* **comb. nov.**, *Ceodes taitensis* **comb. nov.** and *Ceodes wagneriana* **comb. nov.** A general distribution of each species recognised in this work is also included, along with line drawings and colour pictures of representative species of *Ceodes*, *Pisonia* and *Rockia* and an updated dichotomous key based on reproductive characters for the nine genera (*Ceodes*, *Cephalotomandra*, *Grajalesia*, *Guapira*, *Neea*, *Necopsis*, *Pisonia*, *Pisoniella* and *Rockia*) comprising the tribe Pisonieae.

Résumé

Plusieurs genres de Nyctaginaceae actuellement fusionnés sous *Pisonia* ont été décrits pour la région Indo-Pacifique. Les résultats d'une récente étude phylogénétique moléculaire de la tribu Pisonieae ont montré que *Pisonia* est non monophylétique et comprend trois lignées bien supportées: une comprenant *Pisonia*

typique et ses alliés (*Pisonia* s.str.), un clade d'espèces qui correspond à la description originale de *Ceodes* et une troisième lignée dont l'unique représentant était auparavant traité sous le genre monotypique *Rockia*. Ainsi, dans le cadre d'un effort pour parvenir à une classification naturelle de la tribu Pisonieae, ce travail proposons de rétablir les *Ceodes* et *Rockia* pour accueillir des taxons avec des glandes discrètes sur les anthocarps, reconnaissant 21 espèces (20 pour les premières et une pour les dernières), dont 16 sont de nouvelles combinaisons: *Ceodes amplifolia* **comb. nov.**, *Ceodes artensis* **comb. nov.**, *Ceodes austro-orientalis* **comb. nov.**, *Ceodes brownii* **comb. nov.**, *Ceodes cauliflora* **comb. nov.**, *Ceodes coronata* **comb. nov.**, *Ceodes diandra* **comb. nov.**, *Ceodes gigantocarpa* **comb. nov.**, *Ceodes gracilescens* **comb. nov.**, *Ceodes lanceolata* **comb. nov.**, *Ceodes merytifolia* **comb. nov.**, *Ceodes muelleriana* **comb. nov.**, *Ceodes rapaensis* **comb. nov.**, *Ceodes sechellarum* **comb. nov.**, *Ceodes taitensis* **comb. nov.** et *Ceodes wagneriana* **comb. nov.** Une distribution générale de chaque espèce reconnue dans ce travail est également incluse, ainsi que des dessins au trait et des images en couleur des espèces représentatives de *Ceodes*, *Pisonia* et *Rockia*, et préparé une clé dichotomique mise à jour basée sur les caractères reproductifs des neuf genres (*Ceodes*, *Cephalotomandra*, *Grajalesia*, *Guapira*, *Neea*, *Neeopsis*, *Pisonia*, *Pisoniella* et *Rockia*) comprenant la tribu Pisonieae.

Keywords

Calpidia, Caryophyllales, flora of Hawaii, flora of the Indo-Pacific, flora of oceanic islands, *Heimerliodendron*, island endemics, *Timeroylea*

Introduction

The tribe Pisonieae Meisn. in Nyctaginaceae (Caryophyllales) contains the most diverse woody assemblage of the family, represented by over 200 species distributed mainly in the tropical and subtropical regions of the New World (Douglas and Spellenberg 2010). Although members of Pisonieae are present – and often common – in all types of Neotropical habitats and are important components for many ecosystems, taxonomic delimitations at the generic and species levels are still obscure and in urgent need of updated treatments. The tribe currently comprises seven accepted genera (*Cephalotomandra* H.Karst & Triana, *Grajalesia* Miranda, *Guapira* Aubl., *Neea* Ruiz & Pav., *Neeopsis* Lundell, *Pisonia* L. and *Pisoniella* (Heimerl) Standl.), all of them restricted to the New World except *Pisonia*, which has a pantropical distribution (Douglas and Spellenberg 2010). However, other genera have been erected in the last three centuries to include some of the Indo-Pacific taxa with dried anthocarps and inconspicuous glands along anthocarp ribs, which have been either recognised as accepted or treated as synonyms of *Pisonia* by different authors, resulting in a convoluted taxonomic history that we aim to clarify below and that is also summarised in Figure 1.

Ceodes J.R. Forst. & G.Forst., which was described by Forster and Forster (1776), is the oldest of the Indo-Pacific genera with its type species *Ce. umbellifera* J.R.Forst. & G.Forst. collected in Tanna Island at Vanuatu, characterised by the absence of stalked glands along the ribs of anthocarps (Seemann 1863). Later, Du Petit-Thouars (1804) established the genus *Calpidia* Thouars, whose detailed description is based on material collected in Mauritius and which, just as *Ceodes*, differed from *Pisonia*, mainly by the absence of glandular emergences on the surface of its anthocarps (Du Petit-Thouars 1806; see Figs 2, 3). The protologues of *Ceodes* and *Calpidia* describe essentially the same diagnostic characters and it is possible that Du Petit-Thouars was unaware that




Figure 1. Diagram depicting the history of classification for *Calpidia*, *Ceodes*, *Heimerliodendron*, *Pisonia* and *Rockia*. Major taxonomic treatments are shown within boxes. Authors are shown in bold and publication years of treatments and genera within parentheses.

Ceodes had been described 28 years earlier, as he did not mention this genus in either of his two publications on *Calpidia*.

In a global treatment of Nyctaginaceae, prepared by Choisy (1849), he merged *Calpidia* under *Pisonia* and, as with Du Petit-Thouars, he did not include *Ceodes* in this treatment. Over 30 years later, Bentham and Hooker (1880) maintained both *Calpidia* and *Ceodes* as synonyms of *Pisonia*. However, they recognised the monotypic genus *Timeroyea* Montrouz. (using the orthographic variant “*Timeroya*”) from New Caledonia, which besides *Ceodes* and *Calpidia*, represents a third genus with inconspicuous glands on its anthocarps, but was characterised by having many (25–30) stamens (Montrouzier 1860; Beauvisage 1901).

Nine years after Bentham and Hooker’s publication, Heimerl (1889) presented his first tribal treatment for Nyctaginaceae, in which he split *Pisonia* into six sections, resulting in an expanded delimitation of this genus. *Timeroyea* was reduced to a section of *Pisonia* (as *P. sect. Timeroya*), while *P. sect. Prismatocarpae* was established to accommodate taxa with up to 15 stamens, resulting in two sections of *Pisonia* with inconspicuously-glanded taxa. In this treatment, *Calpidia* was listed as a synonym of *Pisonia*, while *Ceodes* was not mentioned at all.

Subsequent work by Heimerl (1913a), based on the examination of additional material, led him to propose the split of *Pisonia* sensu lato with the reinstatement of *Calpidia* to embrace all taxa placed under *P. sect. Prismatocarpae* and *P. sect. Timeroya*. Here, *Calpidia* differed from other *Pisonia* sections by the absence of bracteoles, having a reduced perisperm that forms gelatinous traces and starch accumulation in the embryo and by its geographic distribution. In the following publication which included palynological analysis, Heimerl (1913b) proposed new combinations to *Calpidia*, including some of the newly described species by Warburg (1891) and Bargagli-Petrucci (1901) and described the monotypic genus *Rockia* Heimerl to accommodate the Hawaiian endemic *P. sandwicensis* Hillebr. Although the anthocarps of *Rockia* have inconspicuous glands located along the ribs, Heimerl (1913b) distinguished it from *Calpidia* by the presence of bracteoles and the number of pores in pollen grains, where *Rockia* has sessile flowers with one bract and two bracteoles at its base and tricolpate pollen,

while *Calpidia* has pedicellate flowers lacking any bracts or bracteoles at the upper portion of the pedicels and pollen with four or more colpi, resulting in many apertures.

In his last comprehensive synthesis of Nyctaginaceae, Heimerl (1934) maintained *Rockia* as an accepted genus and additionally recognised three sections in *Calpidia*, based on the presence of a rostrum – a beaked or filamentous extension at the apex of the anthocarps – and the number of stamens: *Ca.* sect. *Excelsae* (= *P.* sect. *Prismatocarpaceae*), *Ca.* sect. *Paucistaminatae* and *Ca.* sect. *Timeroyea*. *Calpidia* sect. *Timeroyea*, represented by the New Caledonian endemic *Ca. artensis* (Montrouz.) Heimerl, is characterised by flowers with a high number of stamens (≥ 30), while the Indo-Pacific widespread *Ca.* sect. *Excelsae*, which includes *Ca. excelsa* (Blume) Heimerl and *Ca. brunoniana* (Endl.) Heimerl, has flowers with 6–30 stamens. On the other hand, species with less than five stamens and a rostrum were placed into *Ca.* sect. *Paucistaminatae*, most of whose representatives occur in Papua New Guinea (e.g. *Ca. longirostris* (Teijsm. & Binn.) Heimerl).

Unlike with his former treatment, Heimerl (1934) mentioned *Ceodes*, but did not evaluate its status under the argument that he lacked enough information about this genus to reach a taxonomic decision, even when Skottsberg (1926), years before, had acknowledged the priority of *Ceodes* over *Calpidia*. Skottsberg re-established *Ceodes* under the argument that there were extant original specimens and that the scant description of the genus was similar to other names published during the late 18th century. However, Skottsberg's treatment was restricted to the plants from Hawai'i and included only two species in *Ceodes* (*Ce. brunoniana* (Endl.) Skottsbg. and *Ce. forsteriana* (Endl. ex Walp.) Skottsbg.) and listed a third one (*Ce. excelsa* (Blume) Skottsbg.) as a questionable species. As Heimerl (1934) still did not accept *Ceodes* and retained *Calpidia* in his new treatment of Nyctaginaceae, Skottsberg (1936) published a work focused on the nyctaginaceous trees from Hawai'i where he reinstated his views on the priority of *Ceodes*, this time recognising only one species (*Ce. umbellifera*, including *Ce. excelsa* and *Ce. forsteriana* as synonyms). In this same work, he also described the genus *Heimerlia* Skottsbg. to accommodate *Ce. brunoniana*, a species characterised by having hermaphroditic flowers. Following Skottsberg's views, Heimerl (1937) finally accepted the priority of *Ceodes* over *Calpidia* and described a new form for *Ce. umbellifera* (*Ce. umbellifera* f. *amplifolia* Heimerl), but did not effectuate any transfers from the sections of *Calpidia* he previously published. Two years later, in their first paper on a series of publications on the plants from Papua New Guinea, Merrill and Perry (1939) proposed two new combinations for species of *Ceodes*. Finally, Skottsberg (1941) corrected the name *Heimerlia* to *Heimerliodendron* Skottsbg. after noticing that the former had been already described for a fungus.

In a new and drastically different treatment for the group, Stemmerik (1964a) proposed a broad definition for *Pisonia* which re-incorporated all taxa with inconspicuous glands along anthocarps. In his revision, which was restricted to the Indo-Pacific taxa, he merged *Calpidia*, *Ceodes*, *Heimerliodendron* and *Rockia* within *Pisonia*. The three sections of *Calpidia*, recognised by Heimerl in 1934, were transferred to *Pisonia*, where *P.* sect. *Prismatocarpaceae* sensu Heimerl (1889) (same as *Ca.* sect. *Excelsae* sensu Heimerl 1934) was restored. *Rockia* was merged into *P.* sect. *Paucistaminatae* (sensu Heimerl 1934) along with the taxa with pedicellate flowers and having an anthocarp rostrum. Therefore, the delimitation of the three sections of *Pisonia*, proposed

by Stemmerik (1964a), was based on characters used by Heimerl (1934) to define his sectional ranks, such as type of glands on anthocarps, presence of a rostrum and number of stamens. However, he did not consider the absence of bracts and bracteoles at the upper portion of the pedicels, presence of starch in the embryo and number of apertures in pollen grains, as he argued that at least the pollen structure was not a constant character and, therefore, had no utility separating genera (Stemmerik 1964b).

A recent phylogenetic study of tribe Pisonieae, based on molecular data (Rossetto et al. 2019), indicated that *Pisonia*, as delimited by Stemmerik (1964a, b), is non-monophyletic. The current definition of *Pisonia* places taxa with inconspicuous glands along anthocarp ribs (Fig. 3A–C, F) into two distinct, well-supported lineages (i.e. clades A and C sensu Rossetto et al. 2019; Fig. 4), while typical *Pisonia* and its allies with glandular emergences (Fig. 3D–E) are restricted to clade B (*Pisonia* s.str.; Fig. 4). In clade A, formed by taxa carrying pedicellate flowers without bracteoles, members of the *P.* sect. *Prismatocarpae* and *P.* sect. *Timeroyea* (sensu Stemmerik 1964a) are included in the clade *Ceodes*, although these sections were not clustered in natural groups. Concurrently, *P. sandwicensis* from the *P.* sect. *Paucistaminatae* is placed in clade C as sister to the Neotropical genera *Guapira* and *Neea* (Rossetto et al. 2019; Fig. 4). Therefore, in order to simplify the classification of the tribe by designating monophyletic genera for the two independent lineages with inconspicuous glands within the tribe, it is necessary to resurrect *Ceodes* and *Rockia*. The objective of this work is to re-establish these two genera and to provide new combinations where necessary. To facilitate recognition in herbaria and in the field, we also provide colour pictures and line drawings of representative species of *Ceodes*, *Pisonia* and *Rockia* and a dichotomous key for the nine genera comprising the tribe Pisonieae.

Materials and methods

For the taxonomic treatment, we compiled accepted names following Stemmerik's revision (1964a), which is the most recent comprehensive treatment for Pisonieae in the Pacific region. We also consulted other more recent regional treatments and species descriptions and provided new generic combinations of the taxa that, according to our understanding, are currently considered as accepted (Friedmann 1986; Fosberg 1987; Philcox and Coode 1994; Florence 2004; Whistler 2004). Additional information on geographic distributions was obtained from Heimerl (1913b) and Stemmerik (1964b). Generic descriptions were based on Heimerl (1913b; 1934), while Skottsberg (1936) was used specifically for the description of pollen structure of *Ceodes*.

Results and conclusions

Here we re-established the genera *Ceodes* and *Rockia*, recognising 20 species for the former and one for the latter. Sixteen out of the 20 species, recognised for *Ceodes*, represent new combinations (see Taxonomic treatment section). The re-establishment of *Ceodes* and *Rockia* provides an important step to refine our knowledge of the taxonomy and evolution

of Pisonieae from the Indo-Pacific region. This work also has considerable implications for estimates of regional biodiversity, as many species of *Ceodes* are island endemics, while *Rockia* would be a genus restricted to the Hawaiian Archipelago (Wagner et al. 2005).

Some species of Pisonieae have been reported as dominant components of the vegetation from remote islands, in part because their sticky anthocarps can travel long distances attached to the feathers of seabirds (St. John 1951; Airy-Shaw 1952). Therefore, future studies on the taxonomy, ecology and biogeography of Pisonieae will help us understand how interactions with pollinators, seed dispersers (Walker et al. 1991; Murphy and Legge 2003; Burger 2005) and mycorrhiza (Hayward and Hynson 2014) have contributed to promote endemism in trees with a high dispersal capability.

Taxonomic treatment

***Ceodes* J.R.Forst. & G.Forst., Char. Gen. Pl., ed. 2: 141. 1776.**

- ≡ *Pisonia* sect. *Prismatocarphae* Heimerl, Nat. Pflanzenfam. 3(1b): 29. 1889. Type (designated by Stemmerik in *Blumea* 12: 277. 1964): *Pisonia umbellifera* (J.R.Forst. & G.Forst.) Seem. (≡ *Ceodes umbellifera* J.R.Forst. & G.Forst.), **syn. nov.**
- = *Calpidia* Thouars, Hist. Vég. Îles Austral. Afriq. 37, pl. 10. 1804. Type: *Calpidia oblonga* J.St.-Hil., **syn. nov.**
- = *Heimerlia* Skottsbo., Svensk Bot. Tidskr. 30: 738. 1936 (non Höhn. 1903). Type: *Heimerlia brunoniana* (Endl.) Skottsbo., **syn. nov.**
- = *Heimerliodendron* Skottsbo., Svensk Bot. Tidskr. 35: 364. 1941. Type: *Heimerliodendron brunonianum* (Endl.) Skottsbo., **syn. nov.**
- = *Pisonia* sect. *Paucistaminatae* (Heimerl) Stemm., *Blumea* 12: 277. 1964 ≡ *Calpidia* sect. *Paucistaminatae* Heimerl, Nat. Pflanzenfam. 16c: 125. 1934. Type (designated by Stemmerik in *Blumea* 12: 277. 1964): *Pisonia longirostris* Teijsm. & Binn. (≡ *Calpidia longirostris* (Teijsm. & Binn.) Heimerl), **syn. nov.**
- = *Pisonia* sect. *Timeroyea* (Montrouz.) Heimerl, Nat. Pflanzenfam. 3(1b): 29. 1889, ‘*Timeroya*’ ≡ *Timeroyea* Montrouz., Mém. Acad. Roy. Sci. Lyon, Sect. Sci. 10: 247. 1860. Type (designated by Stemmerik in *Blumea* 12: 277. 1964): *Pisonia artensis* (Montrouz.) Barg.-Petr. (≡ *Timeroyea artensis* Montrouz.), **syn. nov.**
- = *Vieillardia* Brong. & Gris, Bull. Soc. Bot. France 8: 375. 1861 (non Montrouz. 1860). Type: *Vieillardia austrocaledonica* Brong. & Gris., **syn. nov.**

Type. *C. umbellifera* J.R.Forst. & G.Forst.

Description. Habit and phyllotaxy. Dioecious or hermaphroditic trees or shrubs, leaves (sub)opposite or (sub)verticillate clustered at apex of branches.

Inflorescence. Axillary, terminal or occasionally cauliflorous, arranged in compound cymes.

Flowers. Unisexual (with vestiges of another sex) or rarely hermaphrodite, pedicellate, bracteoles absent at the upper portion of the pedicels, perianth campanulate

(Fig. 2A, D) to funnel-shaped, stamens 2 to many (> 30), long or shortly exerted (Fig. 2D) and stigma penicillate or less frequently fimbriate, exerted (Fig. 2A).

Anthocarp. Leathery or woody (but never fleshy), ellipsoid, prismatic or fusiform, sometimes with a rostrum at apex (Fig. 3B) and 4–5 longitudinal ribs with inconspicuous sticky glands (Figs 2B, C, 3A–C).

Pollen. Six and 12 colpi geometrically arranged.

Perisperm. Often scarce, gelatinous or mealy.

1. *Ceodes amplifolia* (Heimerl) E.F.S.Rossetto & Caraballo, comb. nov.

urn:lsid:ipni.org:names:77210103-1

≡ *Ceodes umbellifera* f. *amplifolia* Heimerl, Occas. Pap. Bernice Pauahi Bishop Mus. 13: 38. 1937. (Basionym).

Distribution. French Polynesia (Austral Islands) (Florence 2004).

2. *Ceodes artensis* (Montrouz.) E.F.S.Rossetto & Caraballo, comb. nov.

urn:lsid:ipni.org:names:77210104-1

≡ *Timeroyea artensis* Montrouz., Mém. Acad. Roy. Sci. Lyon, Sect. Sci. 10: 247. 1860. (Basionym).

Distribution. New Caledonia (Stemmerik 1964a).

3. *Ceodes austro-orientalis* (J.Florence) E.F.S.Rossetto & Caraballo, comb. nov.

urn:lsid:ipni.org:names:77210105-1

≡ *Pisonia austro-orientalis* J. Florence, Fl. Polynésie Franç. 2: 307. 2004. (Basionym).

Distribution. French Polynesia (Gambier Islands) (Florence 2004).

4. *Ceodes brownii* (J.Florence) E.F.S.Rossetto & Caraballo, comb. nov.

urn:lsid:ipni.org:names:77210106-1

≡ *Pisonia brownii* J.Florence, Fl. Polynésie Franç. 2: 308. 2004. (Basionym).

Distribution. French Polynesia: Nuku Hiva (Florence 2004).




Figure 2. Field images for representative species of *Ceodes*, *Pisonia* and *Rockia* (Nyctaginaceae) from the Pacific Islands **A** *Ceodes taitensis*. Branch with pistillate flowers **B** *Ceodes brunoniana*. Ripe anthocarps (fruits) exuding sticky secretions (arrow) **C, D** *Ceodes umbellifera*. Branch with ripe anthocarps (**C**) and staminate flowers at anthesis (**D**) **E, F** *Pisonia grandis* R.Br. Staminate flowers at anthesis (**E**) and ripe anthocarps (**F**) **G–I** *Rockia sandwicensis*. Staminate (**G**) and pistillate (**H**) flowers at anthesis and ripe anthocarps (**I**) Photo credits: **A, F** by J.-Y. Meyer **B** by L. Jensen **C** by C.-I Peng **D, E, I** by F. Starr and K. Starr **G, H** by K. Magnacca.

5. *Ceodes brunoniana* (Endl.) Skottsbo., Acta Horti Gothob. 2: 231. 1926.
Figs 2B, 3A

≡ *Pisonia brunoniana* Endl., Prodr. Fl. Norf. 43. 1833. (Basionym).

Distribution. Hawai‘i (Hawai‘i, Lana‘i, Maui, Moloka‘i, O‘ahu), Lord Howe and Norfolk Islands and New Zealand (Northern Island) (Heimerl 1913b; Wagner et al. 2005).

Note. *Pisonia brunoniana* Endl., which was considered by Stemmerik (1964a) a synonym of *P. umbellifera*, is treated here as an accepted name (as *Ce. brunoniana*), following Sykes (1987), who clarified the key characters to separate both species.

6. *Ceodes cauliflora* (Scheff.) E.F.S.Rossetto & Caraballo, comb. nov.

urn:lsid:ipni.org:names:77210107-1

≡ *Pisonia cauliflora* Scheff., Natuurk. Tijdschr. Ned.-Indië 32: 417. 1872. (Basionym).

Distribution. Mariana Islands, Solomon Islands and Lesser Sunda Islands, Moluccas and western Papua New Guinea (Stemmerik 1964b).

7. *Ceodes corniculata* (Barg.-Petr.) Merr. & L.M.Perry, J. Arnold Arbor. 20: 327. 1939.

≡ *Pisonia corniculata* Barg.-Petr., Nuov. Giorn. Bot. Ital. ser. 2, 8: 615. 1901. (Basionym).

Distribution. Bacan Islands (Moluccas) and western Papua New Guinea (Heimerl 1913b; Stemmerik 1964b).

8. *Ceodes coronata* (Heimerl) E.F.S.Rossetto & Caraballo, comb. nov.

urn:lsid:ipni.org:names:77210108-1

≡ *Ceodes umbellifera* var. *coronata* Heimerl, Occas. Pap. Bernice Pauahi Bishop Mus. 13: 41. 1937. (Basionym).

Distribution. Rapa Iti (French Polynesia) (Florence 2004).

9. *Ceodes diandra* (Pulle) E.F.S.Rossetto & Caraballo, comb. nov.

urn:lsid:ipni.org:names:77210110-1

≡ *Pisonia diandra* Pulle, Nova Guinea 8: 629. 1912. (Basionym).

Distribution. Papua New Guinea (Stemmerik 1964b).

10. *Ceodes gigantocarpa* (Heimerl) E.F.S.Rossetto & Caraballo, comb. nov.

urn:lsid:ipni.org:names:77210111-1

≡ *Calpidia gigantocarpa* Heimerl, Oesterr. Bot. Z. 63: 284. 1913. (Basionym).




Figure 3. Comparison of size and morphology of ripe fruits (anthocarps) amongst members of *Ceodes*, *Pisonia* and *Rockia* (Nyctaginaceae). The outline of infructescences are shown below each anthocarp **A** *Ceodes brunoniana* (based on *St. John* 11272 (US-01258187)) **B** *Ceodes longirostris* (based on *Brass* 2972 (HUH-00046918)). Note the extremely long rostrum at the tip of the anthocarp (indicated with an arrow) **C** *Ceodes umbellifera* (based on *Foxworthy* 593 (US-03661041)) **D** *Pisonia aculeata* L. (based on *Caraballo* 3464 (IJ)) **E** *Pisonia grandis* (based on *Fosberg* 24357 (US-00959523)) **F** *Rockia sandwichensis* (based on *Lorence* 6305 (US-00452890)). Illustration credit: Ramos Sepúlveda.

Distribution. New Caledonia (Stemmerik 1964a).

11. *Ceodes gracilescens* (Heimerl) E.F.S.Rossetto & Caraballo, comb. nov.
 urn:lsid:ipni.org:names:77210112-1

≡ *Calpidia gracilescens* Heimerl, Oesterr. Bot. Z. 63: 285. 1913. (Basionym).

Distribution. Tahiti (French Polynesia) (Florence 2004).

12. *Ceodes lanceolata* (Poir.) E.F.S.Rossetto & Caraballo, comb. nov.
 urn:lsid:ipni.org:names:77210113-1

≡ *Calpidia lanceolata* Poir., Encycl. Suppl. 2: 38. 1811. (Basionym).

Distribution. Mauritius and Réunion islands (Philcox and Coode 1994).

Note. We disagree with Stemmerik's (1964a) view on *Pisonia lanceolata* (Poir.) Choisy, which he considered a synonym of *P. umbellifera*. According to Friedmann (1986), *P. lanceolata* is a species with affinities to *P. brunoniana*, from which it can be distinguished by the pattern of incisions in the flower.




Figure 4. Generic relationships within tribe Pisonieae (Nyctaginaceae) showing the inferred positions of *Ceodes* (Clade A, in blue), *Pisonia* (Clade B, in green) and *Rockia* (Clade C, in yellow) (Rossetto et al. 2019).

13. *Ceodes longirostris* (Teijsm. & Binn.) Merr. & L.M.Perry, J. Arnold Arbor. 20: 328. 1939.

Fig. 3B

≡ *Pisonia longirostris* Teijsm. & Binn., Natuurk. Tijdschr. Ned.-Indië 25: 401. 1863. (Basionym).

Distribution. Solomon Islands, Lesser Sunda Islands, Sulu Archipelago (Philippines), Moluccas and Papua New Guinea (Stemmerik 1964b).

14. *Ceodes merytifolia* (Whistler) E.F.S.Rossetto & Caraballo, comb. nov.

urn:lsid:ipni.org:names:77210114-1

≡ *Pisonia merytifolia* Whistler, Rainforest Trees Samoa: 192. 2004, '*merytafolia*'. (Basionym).

Distribution. Samoa Archipelago (Whistler 2004).

15. *Ceodes muelleriana* (Warb.) E.F.S.Rossetto & Caraballo, comb. nov.

urn:lsid:ipni.org:names:77210115-1

≡ *Pisonia muelleriana* Warb., Bot. Jahrb. Syst. 13: 304. 1891. (Basionym).

Distribution. Solomon Islands and Papua New Guinea (Stemmerik 1964b).

16. *Ceodes rapaensis* (J.Florence) E.F.S.Rossetto & Caraballo, comb. nov.

urn:lsid:ipni.org:names:77210116-1

≡ *Pisonia rapaensis* J.Florence, Fl. Polynésie Franç. 2: 317. 2004. (Basionym).

Distribution. French Polynesia (Rapa Iti) (Florence 2004).

17. *Ceodes sechellarum* (F.Friedmann) E.F.S.Rossetto & Caraballo, comb. nov.

urn:lsid:ipni.org:names:77210117-1

≡ *Pisonia sechellarum* F.Friedmann, Bull. Mus. Natl. Hist. Nat., B, Adansonia, sér. 4, 8: 384. 1986 (publ. 1987). (Basionym).

Distribution. Seychelles (Silhouette Island) (Friedmann 1986).

18. *Ceodes taitensis* (Heimerl) E.F.S.Rossetto & Caraballo, comb. nov.

urn:lsid:ipni.org:names:77210118-1

Fig. 2A

≡ *Calpidia taitensis* Heimerl, Oesterr. Bot. Z. 63: 288. 1913. (Basionym).

Distribution. French Polynesia (Society Islands) (Florence 2004).

19. *Ceodes umbellifera* J.R.Forst & G.Forst., Char. Gen. Pl., ed. 2 142, t. 71. 1776.

Figs 2C, D, 3C

Distribution. Widespread across the Indo-Pacific islands (Pramanick et al. 2015).

20. *Ceodes wagneriana* (Fosberg) E.F.S.Rossetto & Caraballo, comb. nov.

urn:lsid:ipni.org:names:77210119-1

≡ *Pisonia wagneriana* Fosberg, Phytologia 62: 177. 1987. (Basionym).

Distribution. Hawai'i (Kaua'i) (Fosberg 1987).

***Rockia* Heimerl, Oesterr. Bot. Z. 63: 289. 1913.**

Type. *R. sandwicensis* (Hillebr.) Heimerl.

Description. *Habit and phyllotaxy.* Dioecious trees or shrubs, leaves (sub)opposite or (sub)verticillate clustered at apex of branches.

Inflorescence. Axillary, terminal, arranged in compound cymes.

Flowers. Unisexual (with vestiges of another sex), sessile, with one bract and two bracteoles present at the base, male perianth campanulate, stamens 10–26, exserted (Fig. 2G), female perianth tubular or fusiform (Fig. 2H), stigma fimbriate, exserted.

Anthocarp. Leathery, elongated fusiform, with 5 ribs covered by inconspicuous glands excreting sticky substances (Figs 2I, 3F).

Pollen. Tricolpate, with 3 apertures distant 120° from each other.

Perisperm. Abundant, gelatinous.

1. *Rockia sandwicensis* (Hillebr.) Heimerl, Oesterr. Bot. Z. 63: 290. 1913.

Figs 2G–I, 3F

≡ *Pisonia sandwicensis* Hillebr., Fl. Hawaiian Isl. 369. 1888. (Basionym).

Distribution. Hawai‘i (Hawai‘i, Kaua‘i, Lana‘i, Maui, Moloka‘i, O‘ahu) (Stemmerik 1964a; Wagner et al. 2005).

Key to genera from tribe Pisonieae

The following dichotomous key is compiled to separate the nine currently accepted genera within Pisonieae, based on reproductive features because vegetative (i.e. leaves and twigs) characters do not seem to provide enough resolution to help set apart these genera. Unfortunately, members of Pisonieae tend to have fugacious reproductive seasons and many collections in herbaria lack flowers and/or fruits. Thus, we support the recommendation made by Caraballo-Ortiz and Trejo-Torres (2017) on preparing multiple vouchers from a plant or population across seasons to document the full phenology of species and their range of morphological variation.

- 1 Staminate flowers with inserted stamens..... 2
- Staminate flowers with exserted stamens 4
- 2 Stamens about 30..... *Cephalotomandra*
- Stamens 5–13 3
- 3 Leaves drying blackish; inflorescences in corymbose cymes; flowers usually with urceolate corolla. Widespread across the Neotropics..... *Neea*
- Leaves greenish when dry; inflorescences in dichasium; flowers with campanulate corolla. Restricted to Guatemala *Neeopsis*
- 4 Flowers pedicellate, lacking bracts or bracteoles at the upper portion of the pedicels..... 5
- Flowers sessile, subtended by one bract and two bracteoles..... 6

- 5 Inflorescence in simple umbel; glandular emergences along anthocarp ribs. Neotropics..... ***Pisoniella***
- Inflorescence in compound cymes; inconspicuous glands along anthocarp ribs. Indo-Pacific..... ***Ceodes***
- 6 Anthocarps red- or violet coloured, more or less fleshy when ripe..... ***Guapira***
- Anthocarps dry when ripe7
- 7 Anthocarps winged, lacking sticky glands..... ***Grajalesia***
- Anthocarps not winged, sticky glands present **8**
- 8 Anthocarp ribs covered by glandular emergences. Pantropical ***Pisonia***
- Anthocarp ribs covered by inconspicuous glands. Endemic to Hawai‘i..... ***Rockia***

Acknowledgements

The authors thank Dr. Alexander Sukhorukov (Lomonosov Moscow State University) for translating texts from German and constructive editorial suggestions. We also appreciate the useful comments provided by Dr. Lucas Majure (Florida Museum of Natural History) to improve the quality of this manuscript. The artist Ramos Sepúlveda (Puerto Rico) prepared the elegant line drawings presented in Figure 3. We acknowledge the generous assistance of Carrie Craig (Smithsonian Institution) and Mike Skinner (Pensoft) proofreading the manuscript. The following botanists kindly provided permission to use their pictures in this publication: Ching-I Peng (Academia Sinica, Taiwan; deceased), Forest and Kim Starr (Starr Environmental, Hawai‘i), Jean-Yves Meyer (Délégation à la Recherche, Tahiti, French Polynesia & Tetiaroa Society), Karl Magnacca (Hawai‘i) and Lawrence Jensen (University of Auckland, New Zealand). We are grateful to the following online resources and institutions for providing virtual access to specimens and literature: Biodiversity Heritage Library (BHL), Harvard University Herbaria (HUH), International Plant Names Index (IPNI), JSTOR Global Plants, Missouri Botanical Garden (MO), New York Botanical Garden (NY), Smithsonian Institution (US) and the Smithsonian’s Flora of the Hawaiian Islands website.

References

- Airy-Shaw HK (1952) On the distribution of *Pisonia grandis* R. Br. (Nyctaginaceae), with special reference to Malaysia. *Kew Bulletin* 7(1): 87–97. <https://doi.org/10.2307/4117674>
- Bargagli-Petrucci G (1901) Le specie di *Pisonia* della regione di Monsoni. *Nuovo Giornale Botanico Italiano* 8: 603–625.
- Beauvisage G (1901) Genera Montrouzierana, plantarum Nova Caledoniae. *Annales de la Société Botanique de Lyon* 26(2): 49–96. <https://doi.org/10.3406/linly.1901.4713>
- Bentham G, Hooker JD (1880) Genera plantarum ad exemplaria imprimis in herbariis kewensibus servata definita, Vol. 3, pt. 1. Reeves & Co., London.

- Burger AE (2005) Dispersal and germination of seeds of *Pisonia grandis*, an Indo-Pacific tropical tree associated with insular seabird colonies. *Journal of Tropical Ecology* 21(3): 263–271. <https://doi.org/10.1017/S0266467404002159>
- Carballo-Ortiz MA, Trejo-Torres JC (2017) Two new endemic tree species from Puerto Rico: *Pisonia horneae* and *Pisonia roqueae* (Nyctaginaceae). *PhytoKeys* 86: 97–115. <https://doi.org/10.3897/phytokeys.86.11249>
- Choisy JD (1849) Nyctaginaceae. In: de Candolle A (Ed.) *Prodromus Systematis Naturalis Regni Vegetabilis*, Vol. 13, pt. 2. Masson, Paris, 425–458.
- Douglas N, Spellenberg R (2010) A new tribal classification of Nyctaginaceae. *Taxon* 59(3): 905–910. <https://doi.org/10.1002/tax.593018>
- Du Petit-Thouars LMA (1804) *Histoire des végétaux recueillis sur les îles de France, La Réunion (Bourbon) et Madagascar*. Huzard, Paris.
- Du Petit-Thouars LMA (1806) *Histoire des végétaux recueillis dans les Îles Australes d’Afrique*. Tourneisen, Paris. <https://doi.org/10.5962/bhl.title.163949>
- Florence J (2004) *Flore de la Polynésie Française*, Vol. 2. Collection Faune et Flore tropicales 41. IRD Éditions, Publications Scientifiques du Muséum National D’Histoire Naturelle, Paris.
- Forster JR, Forster G (1776) *Characteres Generum Plantarum*. B. White, T. Cadell, & P. Elmsly, London.
- Fosberg FR (1987) A new *Pisonia* (Nyctaginaceae) from the Hawaiian Islands. *Phytologia* 62(3): 177–178.
- Friedmann F (1986) Étude de la structure du périanthe chez des *Pisonia* paléotropicaux et description de *P. sechellarum* sp. nov. (Nyctaginaceae). *Bulletin du Muséum National d’Histoire Naturelle*. Section B. *Adansonia* 8(4): 383–392.
- Hayward J, Hynson NA (2014) New evidence of ectomycorrhizal fungi in the Hawaiian Islands associated with the endemic host *Pisonia sandwicensis* (Nyctaginaceae). *Fungal Ecology* 12: 62–69. <https://doi.org/10.1016/j.funeco.2014.09.001>
- Heimerl A (1889) Nyctaginaceae. In: Engler A, Prantl K (Eds) *Die Natürlichen Pflanzenfamilien*, Vol. 1b. Engelmann, Leipzig, 14–32.
- Heimerl A (1913a) Über die Nyctaginaceen-Gattung *Calpidia*. *Österreichische Botanische Zeitschrift* 63(1): 19–21. <https://doi.org/10.1007/BF01640846>
- Heimerl A (1913b) Die Nyctaginaceen-Gattungen *Calpidia* und *Rockia*. *Österreichische Botanische Zeitschrift* 63(7): 279–290. <https://doi.org/10.1007/BF01642830>
- Heimerl A (1934) Nyctaginaceae. In: Engler A, Harms H (Eds) *Die Natürlichen Pflanzenfamilien*, ed. 2, Vol. 16c. Engelmann, Leipzig, 86–134.
- Heimerl A (1937) Nyctaginaceae of southeastern Polynesia and other Pacific islands. *Occasional Papers of Bernice Pauahi Bishop Museum of Polynesian Ethology and Natural History* 13(4): 27–47.
- Merrill ED, Perry LM (1939) *Plantae Papuanae Archboldianae*, I. *Journal of the Arnold Arboretum* 20(3): 324–345.
- Montrouzier JXH (1860) *Flore de L’île Art (près de la Nouvelle Calédonie)*. *Mémoires de l’Académie Royale des Sciences, Belles Lettres et Arts de Lyon*. Section des Sciences 10: 173–254.

- Murphy S, Legge S (2003) A first record of a land-bird being trapped by *Pisonia umbellifera*. *The Sunbird* 33(3): 111–112.
- Philcox D, Coode MJE (1994) Nyctaginacées (family 141). In: Bosser J, Cadet T, Guého J, Marais W (Eds) *Flore des Mascareignes: La Réunion, Maurice, Rodrigues* (136. Myoporacées à 148. Hydnoracées). The Sugar Industry Research Institute, Mauritius, ORSTOM, Paris & The Royal Botanic Gardens, Kew, 1–12.
- Pramanick DD, Maiti GG, Mondal MS (2015) Taxonomic study of the genus *Pisonia* L. (Nyctaginaceae) in India. *Annals of Plant Sciences* 4(8): 1179–1185.
- Rossetto EFS, Faria AD, Ruas PM, Ruas CF, Douglas NA, Ribeiro JELS (2019) Clarifying generic delimitation in Nyctaginaceae tribe Pisonieae after more than a century of taxonomy confusion. *Botanical Journal of the Linnean Society* 189(4): 378–396. <https://doi.org/10.1093/botlinnean/boz001>
- Seemann B (1863) On the genus *Ceodes* of Forster. *Journal of Botany, British and Foreign* 1: 244–246.
- Skottsberg C (1926) Vascular plants from the Hawaiian Islands. I. *Acta Horti Gothoburgensis* 2: 185–284.
- Skottsberg C (1936) The arboreal Nyctaginaceae of Hawaii. *Svensk Botanisk Tidskrift* 30: 722–743.
- Skottsberg C (1941) *Heimerliodendron* nov. nom. *Svensk Botanisk Tidskrift* 35: 364.
- St. John H (1951) The distribution of *Pisonia grandis* (Nyctaginaceae). *Webbia* 8: 225–228. <https://doi.org/10.1080/00837792.1951.10669600>
- Stemmerik JF (1964a) Flora Malesianae Precursores XXXVIII: Notes on *Pisonia* L. in the Old World (Nyctaginaceae). *Blumea* 12: 275–284.
- Stemmerik JF (1964b) Nyctaginaceae. In: van Steenis CGGJ (Ed.) *Flora Malesiana*, Vol. 6. Wolters-Noordhoff Publishing, Groningen, 450–468.
- Sykes WR (1987) The parapara, *Pisonia brunoniana* (Nyctaginaceae). *New Zealand Journal of Botany* 25(3): 459–466. <https://doi.org/10.1080/0028825X.1987.10413362>
- von Höhnelt FXR (1903) Mykologische Fragmente. *Annales Mycologici* 1(5): 391–414.
- Wagner WL, Herbst DR, Lorence DH (2005) Flora of the Hawaiian Islands website. <http://botany.si.edu/pacificislandbiodiversity/hawaiianflora/index.htm> [accessed 03.15.2020]
- Walker TA, Chaloupka MY, King BR (1991) *Pisonia* islands of the Great Barrier Reef, part 1. *Atoll Research Bulletin* 350: 1–23. <https://doi.org/10.5479/si.00775630.350-1.1>
- Warburg O (1891) Beiträge zur Kenntnis der papuanischen Flora. *Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie* 13: 230–455.
- Whistler WA (2004) Rainforest trees of Samoa: A guide to the common lowland and foothill forest trees of the Samoan Archipelago. *Isle Botanica*, Honolulu.