

Linum aksehirense (sect. *Dasylinum*, Linaceae), a new species from Central Anatolia (Turkey)

Osman Tugay¹, Deniz Ulukuş²

1 Selçuk University, Department of Pharmaceutical Botany, Faculty of Pharmacy, Selçuklu, Konya, Turkey

2 Selçuk University, Department of Biotechnology, Faculty of Sciences, Selçuklu, Konya, Turkey

Corresponding author: Deniz Ulukuş (dulukus@selcuk.edu.tr)

Academic editor: R. Riina | Received 10 September 2019 | Accepted 9 November 2019 | Published 10 December 2019

Citation: Tugay O, Ulukuş D (2019) *Linum aksehirense* (sect. *Dasylinum*, Linaceae), a new species from Central Anatolia (Turkey). PhytoKeys 136: 23–34. <https://doi.org/10.3897/phytokeys.136.46477>

Abstract

Linum aksehirense (Linaceae) is described as a new species known from the slopes of the Sultan Mountains in the Akşehir district of Konya in Central Anatolia (Turkey). It is most similar to *L. pubescens* Banks & Sol. and *L. anisocalyx* P.H. Davis, from which it is easily distinguished by its stem leaf shape, sepal shape and petal colour. Seed and pollen surface ornamentations were photographed under SEM microscopy to explore micromorphological characters distinguishing the new species from close relatives. In addition, photographs of living material, a distribution map, ecological details, and an identification key are provided.

Keywords

Endemic, Konya, Linaceae, *Linum*, taxonomy

Introduction

The genus *Linum* Linnaeus (Linaceae) is comprised of about 200 species in the Linaceae family. *Linum* is distributed mainly in North America, the Balkan Peninsula, Anatolia, as well as in Eurasia and Africa (Robertson 1971), but it is also represented in South America and Australasia.

Linum was first described by Linnaeus (1753). The first comprehensive study on the genus was made by Planchon (1847, 1848). The most recent worldwide treatment of *Linum* was done by Winkler (1931). According to Winkler (1931) and Planchon (1847, 1848), *Linum* is divided into five sections. The genus has been the subject of several taxonomic studies and regional revisions, such as those focusing on American

and South African *Linum* species (Rogers 1963, 1981; Mildner and Rogers 1978), as well as Flora Europaea (Ockendon and Walters 1968), Flora of the U.S.S.R. (Yuzepchuk 1974), and Flora of Turkey and the Aegean Islands (Davis 1967). The genus plays an important role in the economic and social development of humans. For example, the seeds of *L. usitatissimum* are used nutritionally and medicinally. Linseed oil is also a significant source of inks, varnishes, and lubricants (McDill et al. 2009).

Linum is represented by four sections in Turkey: sect. *Syllinum* Grisebach (1843: 115), sect. *Linastrum* Planchon (1847: 597), sect. *Dasylinum* Planchon (1847: 598), and sect. *Linum* Planchon (1847: 598). With 54 taxa (Yılmaz 2018), Turkey is one of the most important centres of genetic diversity for the genus *Linum*. The new species described in this paper brings the number to 55 taxa in Turkey, 26 (47%) of which are endemic.

Linum sect. *Dasylinum* is characterised by having perennial or annual life cycles and distinct morphological characteristics. Leaves are alternate, often hairy. Petals usually have coherent claws and are blue, pink or white. Capsules are hairy or glabrous (Davis 1967). There have been numerous studies on the morphology of *Linum* in Turkey by several authors (Davis 1967; Yılmaz and Kaynak 2008, 2010; Yılmaz 2010). There have also been investigations on a worldwide scale concerning the palynology (Erdtman 1969; Rogers and Xavier 1971; Saad 1961, 1962; Xavier et al. 1980; Rogers 1985; Talebi et al. 2012), karyology (Ray 1944; Harris 1968; Rogers et al. 1972) and anatomy (Winkler 1931; Metcalfe and Chalk 1950) of the genus. There has been just one study on *Linum* seeds, which covers several species of the genus (Özcan and Zorlu 2009).

From a biogeographical and evolutionary point of view, McDill et al. (2009) reported that section *Linum* and *Dasylinum* were not monophyletic and that *Linum* appears to have arisen in Eurasia, from which it spread to Africa, North America, South America and Australasia.

We collected samples of what we suspected could be a new species of *Linum* while conducting field work around the Sultan Mountains between 2011 and 2017. The specimens were checked against the Flora of Turkey and the East Aegean Islands, as well as neighbouring floras (Flora Iranica, Flora Iraq, Flora of the U.S.S.R. and Flora Europaea). We subsequently decided that it was a species new to science.

The aim of this study was to describe the new species, named *Linum aksehirense*, occurring in the Sultan Mountains (Akşehir, Konya), in terms of its morphology, palynology and seed micromorphology.

Material and methods

Field collections were carried out in the Sultan Mountains from 2011 to 2017. Collected specimens were dried using traditional procedures and were deposited in the KNYA Herbarium. They were identified using the Flora of Turkey and the East Aegean Islands using stereo microscopy and checking them against other *Linum* accounts found in relevant literature. The taxonomical description of the species was made according to Davis (1967).

For palynological investigations, pollen slides were prepared according to Wodehouse's (1935) technique. The pollen micromorphology of *L. aksehirense* was scrutinized by scanning electron microscopy (SEM) techniques. For SEM, pollen grains were directly placed on double-sided carbon tape affixed to aluminium stubs, covered with gold with a Hummle VII sputter coater and photographed at magnifications of 2000× to 7000× with a JEOL-5600. SEM micrographs were used to determine exine sculpturing of the pollen. Pollen terminology was based on Punt et al. (2007). For seed investigations, SEM micrographs were used to determine seed coat sculpturing. The terminology of Stearn (1983) and Özcan and Zorlu (2009) was adopted to describe the SEM aspects of the seed coat.

Taxonomy

Linum aksehirense Tugay & Ulukoş, sp. nov.

urn:lsid:ipni.org:names:77203431-1

Figures 1–6

Diagnosis. *Linum aksehirense* belongs to *Linum* sect. *Dasylinum*. The new species is similar to *L. pubescens* but is distinguished by its spatulate lower stem leaves that are not evanescent (*vs.* oblong-spatulate evanescent), subequal lanceolate sepal shape (*vs.* subequal linear), petals 27–33 mm (*vs.* 18–27 mm) and petals that are blue-violet with a yellowish limb base (*vs.* pink with a bluish limb base).




Figure 1. Distribution map of *Linum aksehirense* (triangle), and closely related *Linum pubescens* (circle) and *Linum anisocalyx* (square) in Turkey.



Figure 2. Holotype specimen of *Linum aksehirensis* Tugay & Ulukuş.




Figure 3. General view of habit and flower **A–C** *Linum aksehirens*.




Figure 4. Scanning electron micrographs of seeds *Linum aksehirense* species **A** general view and **B** surface sculpturing pattern.




Figure 5. SEM micrographs of the pollen grains of *L. aksehirense* **A–C** polar, equatorial view and ornamentation (*O. Tugay* 14.542 & *D. Ulukuş*).

Type. TURKEY. B3 Konya; Akşehir, Sultan Mountains, slopes in *Pinus nigra* forest, 1150 m alt., 38°19.230'N, 31°23.181'E, 01 August 2017, *O. Tugay* 14.542 & *D. Ulukuş* (holotype KNYA, isotypes KNYA 28.229).

Description. The plant is a green annual. Flowering stems villous, erect, 22–33 cm; sterile shoots few. Lower stem leaves spatulate, not evanescent, 10–13 × 2–4 mm; median stem leaves lanceolate-elliptic, acute, 16–19 × 2–3.5 mm, 1–3 nerved.




Figure 6. Scanning electron micrographs of seeds sculpturing patterns *Linum* species **A** *L. aksehirense* **B** *L. anisocalyx* **C** *L. pubescens*.

Bract (similar to median stem leaves) stipitate glandular margined, lanceolate, 14–15 × 3–3.5 mm. Cymes divaricate, lax, few-flowered; flowers 3–7 per stem; pedicels 1–2 mm, not elongated in fruit. Sepals subequal, lanceolate, 12–14 × 2–3 mm, bearing long hairs and stipitate glandular margined. Petals blue-violet (blue when dry) with a yellowish base to the limb, 27–33 mm. Capsule 5 × 5 mm diam. with 1–1.5 mm beak. Seeds elliptic-oblong, 2–2.5 × 1–1.5 mm, brown, glossy. Sculpturing of seed coat is reticulate-rugolose-ruminata (Fig. 4A–B).

Paratypes. TURKEY. B3 Konya; Akşehir, slopes, 1170 m alt., 38°20'N, 31°24'E, 31 August 2011, *O. Tugay* 7.182 (KNYA); Akşehir, Hıdırlık, slopes, 1100 m alt., 38°19'N, 31°23'E, 01 August 2017, *O. Tugay* 14.520 & *D. Ulukuş* (KNYA).

Ecology. *Linum aksehirense*, which is endemic to Turkey, grows between 1100 and 1170 m elevation on slopes with *Pinus nigra* forest. The vegetation of this habitat is composed mainly of herbaceous and suffruticose plants including: *Agrimonia eupatoria* L. subsp. *asiatica*, *Campanula lyrata* Lam. subsp. *lyrata*, *Centaurea virgata* Lam., *Cistus laurifolius* L., *Cota tinctoria* (L.) J.Gay ex Guss. var. *tinctoria*, *Digitalis ferruginea* L. subsp. *ferruginea*, *Dianthus crinitus* Sm. var. *crinitus*, *Hedysarum varium* Willd. subsp. *varium*, *Phlomis armeniaca* Willd., *Pinus nigra* J.F.Arnold subsp. *nigra*, *Prunus divaricata* Ledeb. var. *divaricata*, *Scabiosa rotata* M.Bieb., *Securigera varia* (L.) Lassen, *Teucrium chamaedrys* L. subsp. *chamaedrys* and *Teucrium polium* L.

Phenology. The new species was observed flowering in July and collected fruiting from July to August.

Etyymology. The species epithet comes from ‘Akşehir’, where the new species is found.

Proposed Turkish name for the new species. Akşehir keteni.

Distribution and conservation status. *Linum aksehirense* is known from three localities in Konya province, in the Irano-Turanian phytogeographic region (Fig. 1). Its area of occupancy is estimated to be less than 500 km². The number of mature individual plants is estimated to be less than 250. Being an annual gives this new species a crucial advantage for survival against potential threats, which include the destruction of brush by locals, road construction and deterioration of habitat. Thus, according to criterion B and D, it can be included in the EN (Endangered) category (IUCN 2001; 2016).

Key to the species of *Linum* sect. *Dasylinum* in Turkey

- 1 Annuals..... 2
- Perennials..... 5
- 2 Petals c. 8 mm, free; sepals not or slightly longer than capsule..... *L. seljukorum*
- Petals 18 mm or more; coherent; sepals much longer than capsule..... 3
- 3 Petals 27–33 mm, blue-violet with a yellowish base to the limb *L. aksehirense*
- Petals 18–27 mm, pink with a bluish base to the limb 4
- 4 Sepals subequal *L. pubescens*
- Sepals very unequal, the outer two hiding the much shorter inner sepals.....
..... *L. anisocalyx*
- 5 Median cauline leaves margined by stalked glands..... 6
- Median cauline leaves not margined by glands 8
- 6 Inflorescence compact; petal claw 1/4 as long as limb..... *L. densiflorum*
- Inflorescence widely spreading, or rarely 1–3 flowered 7
- 7 Petal claw c. 1/2 as long as limb; median stem leaves not attenuate below
..... *L. hirsutum*
- Petal claw c. 1–1½ as long as limb; median stem leaves attenuate at both ends ...
..... *L. unquiculatum*
- 8 Plants very twiggy at base, with many sterile shoots; 1–7 flowers.... *L. olympicum*
- Plant herbaceous, with few or no sterile shoots; usually > 7 flowers..... 9
- 9 Cymes usually spreading, lax; leaves 1–3 nerved, oblong, linear or subspathulate;
petal claw c. 1/2 as long as limb *L. hirsutum*
- Cymes compact; leaves 3–7 nerved, broadly lanceolate; petal claw 1/4–1/3 as long
as limb *L. hypericifolium*

Pollen morphology

The pollen shape of the new species was subprolate (P/E: 1.13) with a polar axis of 53.85 ± 1.75 µm (mean \pm standard deviation) µm and an equatorial axis of

$47.70 \pm 4.70 \mu\text{m}$. The aperture was tricolpate. The colpus was long-acute ended with a colpus length of $34.82 \pm 3.26 \mu\text{m}$ and width of $12.91 \pm 2.23 \mu\text{m}$. Exine thickness was $0.5 \mu\text{m}$ and intine thickness was $0.4 \mu\text{m}$. Exine ornamentation was densely gemmate (Fig. 5 A–C).

Discussion

Linum aksehirense is similar to *L. pubescens*, *L. anisocalyx* and *L. viscosum* in morphology. However, it differs from these similar species in several vegetative and reproductive characters (Table 1).

According to Davis (1967), *L. anisocalyx* is closely related to *L. pubescens*, differing from *L. pubescens* primarily by its strongly dimorphic sepals (not dimorphic), which are rhomboid-lanceolate and glandular-margined.

Linum aksehirense differs from *L. pubescens* by its spatulate, not evanescent lower stem leaves (*vs.* oblong-spathulate evanescent), lanceolate-elliptic, acute median stem leaves, $16\text{--}19 \times 3\text{--}4 \text{ mm}$ (*vs.* oblong, subacute $12\text{--}23 \times 2\text{--}5 \text{ mm}$), subequal lanceolate sepals (*vs.* subequal linear), sepal size of $12\text{--}14 \times 2\text{--}3 \text{ mm}$ (*vs.* $9\text{--}12 \times 1\text{--}2 \text{ mm}$) and its petal colour, which is blue-violet with a yellowish limb base (*vs.* pink with a bluish limb base) (Table 1).

Linum aksehirense is similar to *L. anisocalyx*, differing in its lanceolate-elliptic, acute median stem leaves, $16\text{--}19 \times 3\text{--}4 \text{ mm}$ (*vs.* oblong, oblong, subacute $12\text{--}23 \times 2\text{--}5 \text{ mm}$), subequal lanceolate sepals (*vs.* very unequal elliptic-lanceolate), sepal size of $12\text{--}14 \times 2\text{--}3 \text{ mm}$ (*vs.* outer sepals $11\text{--}12 \times 3\text{--}3.5 \text{ mm}$, inner sepals $6\text{--}8 \times 2.5\text{--}3 \text{ mm}$), petal colour, which blue-violet with a yellowish limb base (*vs.* pink with a bluish limb base) and in its petal size, which is $27\text{--}33 \text{ mm}$ (*vs.* $18\text{--}26 \text{ mm}$) (Table 1).

Linum aksehirense can be distinguished from *L. viscosum*, by its spatulate not evanescent (*vs.* lanceolate, ovate-lanceolate, evanescent), sepal size of $12\text{--}14 \times 2\text{--}3 \text{ mm}$ (*vs.* $6\text{--}9 \times 1\text{--}1.5 \text{ mm}$), petal colour, which is blue-violet with a yellowish limb base (*vs.* pink), and in its petal size, which is $27\text{--}33 \text{ mm}$ (*vs.* $18\text{--}21 \text{ mm}$).

Xavier et al. (1980) described the basic pollen grain in *Linum* as subspheroidal, about $50 \mu\text{m}$ in diameter, isopolar, radially symmetric, tricolpate, colpi with pointed ends. However, in our study *L. aksehirense* had subprolate pollen shape. According to Talebi et al. (2012)'s palynologic study on four section of the Linaceae, including 15 taxa of *Linum*, in all examined taxa the pollen shape in polar view was circular (except in *L. densiflorum* where it was concave-triangular) and also the exine sculpturing pattern showed a clavate, pilate and gemmate to baculate form. Talebi et al. (2012) reported that pollen features of *L. densiflorum* consisted of an oblate-spheroidal pollen shape and small and large gemmate exine ornamentation. Our findings showed that *L. aksehirense* had subprolate pollen shape and the exine ornamentation was densely gemmate (Fig. 5 A–C). In terms of pollen micro-morphology, palynological results demonstrated that there are no clear differences among species of the same section. However, pollen shape can be used to distinguish species.

Özcan and Zorlu (2009) studied seed surface patterns *Linum* genus and found reticulate-ruminate patterns in sect. *Dasylinum*. Özcan and Zorlu (2009) showed that seed patterns provide characters to distinguish taxa at specific and infraspecific levels.

Table 1. Morphological comparison of *Linum aksehirense*, *L. pubescens*, *L. anisocalyx* and *L. viscosum*.

Characters	<i>L. aksehirense</i>	<i>L. pubescens</i>	<i>L. anisocalyx</i>	<i>L. viscosum</i>
Stem	unbranched or branched at the upper stem	branched at the median stem	branched at the base	unbranched or branched at the upper stem
Lower stem leaf shape	spathulate not evanescent	oblong-spathulate evanescent	oblong-spathulate evanescent	lanceolate, ovate-lanceolate evanescent
Median stem leaf shape	lanceolate-elliptic, acute, 16–19 × 3–4 mm	oblong, subacute 12–23 × 2–5 mm	oblong, subacute 12–23 × 2–5 mm	lanceolate, subacute 14–21 × 3–5 mm
Sepal shape	subequal lanceolate	subequal linear	very unequal elliptic-lanceolate	subequal lanceolate
Sepal size (mm)	12–14 × 2–3	9–12 × 1–2	outer sepals 11–12 × 3–3.5; inner sepals 6–8 × 2.5–3	6–9 × 1–1.5
Petal colour	blue-violet with a yellowish base to the limb	pink with a bluish base to the limb	pink with a bluish base to the limb	pink
Petal size (mm)	27–33	18–27	18–26	16–21

In this study, the micromorphological study of the seeds showed that there were clear differences among the studied species. According to our findings, *Linum aksehirense* had reticulate-rugulose-ruminate sculpturing while *L. pubescens* exhibited reticulate-rugulose-granulate and *L. anisocalyx* showed reticulate-rugulose (Fig. 6 A–C).

Conclusion

With the discovery of this new species, the number of species of *Linum* in Turkey has risen to 55. This study provides material and data to aid further research on this significant genus of the Linaceae.

Acknowledgements

The authors are grateful to the curators of AEF, ANK, E, G, GAZI, HUB, ISTE, ISTF, K, KNYA and LE who allowed us to study their *Linum* specimens. We would also like to thank Selçuk University, BAP (Project no: 09201154) for financial support. We wish to thank two anonymous reviewers for their careful reading of our manuscript and their many constructive comments and suggestions.

Reference

- Davis PH (1967) Flora of Turkey and the East Aegean Islands. University press, Edinburgh, 425–450.
- Erdtman G (1969) Handbook of Palynology. New York.
- Grisebach AHR (1843) Spicilegium Florae Rumelicae et Bithynicae Exhibens Synopsis Plantarum quas in aest(Vol. 1). Vieweg & Son, Brunsvigae, 407 pp.

- Harris BD (1968) Chromosome Numbers and Evolution in North American Species of *Linum*. *American Journal of Botany* 55(10): 1197–1204. <https://doi.org/10.1002/j.1537-2197.1968.tb07485.x>
- IUCN (2001) IUCN Red List Categories: Version 3.1. Prepared by the IUCN Species Survival Commission. IUCN, Gland and Cambridge.
- IUCN (2016) Standards and Petitions Subcommittee. Guidelines for Using the IUCN Red List Categories and Criteria. Version 12. Prepared by the Standards and Petitions Subcommittee. <http://www.iucnredlist.org/documents/RedListGuidelines.pdf>
- Linnaeus C (1753) *Species Plantarum*. Laurentius Salvius, Stockholm, 1200 pp.
- McDill J, Reppinger M, Simpson BB, Kadereit JW (2009) The phylogeny of *Linum* and Linaceae subfamily Linoideae, with implications for their systematics, biogeography, and evolution of heterostyly. *Systematic Botany* 34(2): 386–405. <https://doi.org/10.1600/036364409788606244>
- Metcalf CR, Chalk L (1950) *Anatomy of the Dicotyledons*. At The Clarendon Press, Oxford, 268–273.
- Mildner RA, Rogers CM (1978) Revision of the native South American species of *Linum* (Linaceae). *Phytologia* 39: 343–390.
- Ockendon DJ, Walters SM (1968) *Linum* L. In: Tutin TG, Heywood VH, Burges NA, Moore DM, Valentine DH, Walters SM, Webb DA (Eds) *Flora of Europea* (Vol. 2). Cambridge University Press, 206–211.
- Özcan T, Zorlu E (2009) A contribution to taxonomy of Turkish *Linum* based on seed surface patterns. *Biologia* 64(4): 723–730. <https://doi.org/10.2478/s11756-009-0123-6>
- Planchon JE (1847) Sur la Famille des Linées. *The London Journal of Botany* 6: 588–603.
- Planchon JE (1848) Sur la Famille des Linées. *The London Journal of Botany* 7: 165–186.
- Punt W, Hoen PP, Blackmore S, Nilsson S, Le Thomas A (2007) Glossary of pollen and spore terminology. *Review of Palaeobotany and Palynology* 143(1–2): 1–81. <https://doi.org/10.1016/j.revpalbo.2006.06.008>
- Ray C (1944) Cytological Studies on The Flax Genus, *Linum*. *American Journal of Botany* 31(4): 241–248. <https://doi.org/10.1002/j.1537-2197.1944.tb08027.x>
- Robertson KR (1971) The Linaceae in The Southeastern United States. *Journal of the Arnold Arboretum* 52: 649–665. <https://doi.org/10.5962/bhl.part.9119>
- Rogers CM (1963) Yellow flowered species of *Linum* in eastern North America. *Brittonia* 15(2): 97–122. <https://doi.org/10.2307/2805396>
- Rogers CM, Xavier KS (1971) Pollen Morphology as an Aid in Determining Relationships Among Some Widely Separated Old World Species of *Linum*. *Grana* 11(1): 55–57. <https://doi.org/10.1080/00173137109427412>
- Rogers CM, Mildner R, Harris BD (1972) Some Additional Chromosome Numbers in the Linaceae. *Brittonia* 24(3): 313–316. <https://doi.org/10.2307/2805668>
- Rogers CM (1981) A revision of the genus *Linum* in southern Africa. *Nordic Journal of Botany* 1(6): 711–722. <https://doi.org/10.1111/j.1756-1051.1981.tb01157.x>
- Rogers CM (1985) Pollen Morphology of The Monotypic Genus *Cliococca* (Linaceae). *Grana* 24(2): 121–123. <https://doi.org/10.1080/00173138509429923>

- Saad SI (1961) Pollen Morphology and Sporoderm Stratification in *Linum*. *Grana Palynology* 3: 109–129.
- Saad SI (1962) Palynological studies in the Linaceae. *Pollen et Spores* 4: 65–82.
- Stearn WT (1983) *Botanical Latin*. David & Charles. London, 566 pp.
- Talebi SM, Sheidai M, Atri M, Sharifnia F, Noormohammadi Z (2012) Palynological study of the genus *Linum* in Iran (a taxonomic review). *Phytologia Balcanica* 18(3): 293–303.
- Winkler H (1931) Linaceae, Tribe I. 3. Linoideae-Eulinae. In: Engler A, Prantl K (Eds) *Die Natürelchen Pflanzenfamilien Nebst Ihren Gattungen und Wichtigeren Arten, Insbesondere den Nutzpflanzen* (2nd edn.) (Vol. 19a.). W. Engelmann, Leipzig, 111–120.
- Wodehouse RR (1935) *Pollen grains*. McGraw-Hill, New York, 574 pp.
- Xavier KS, Mildner RA, Rogers CM (1980) Pollen morphology of *Linum*, sect. *Linastrum* (Linaceae). *Grana* 19(3): 183–188. <https://doi.org/10.1080/00173138009425002>
- Yılmaz Ö, Kaynak G (2008) A New Species of *Linum* (Linaceae) From West Anatolia, Turkey. *Botanical Journal of the Linnean Society* 156(3): 459–462. <https://doi.org/10.1111/j.1095-8339.2007.00763.x>
- Yılmaz Ö (2010) *Linum kaynakiae* Sp. Nov. (Sect. *Syllinum*, Linaceae) From Turkey. *Nordic Journal of Botany* 28(5): 605–612. <https://doi.org/10.1111/j.1756-1051.2010.00756.x>
- Yılmaz Ö, Kaynak G (2010) A New Taxon of *Linum* (Linaceae) From Southwest Anatolia, Turkey. *Novon* 20(4): 507–511. <https://doi.org/10.3417/2009003>
- Yılmaz Ö (2018) *Linum ayliniae* (Linaceae), a New Species from West Anatolia, Turkey. *Novon* 26(2): 174–179. <https://doi.org/10.3417/2018078>
- Yuzepchuk SV (1974) *Linum* L. In: Shishkin BK (Ed.) *Flora of U.S.S.R.*, 14: 67–112. *Acad. Sci. USSR* (Vol 14). Translation from Russian by Israel Programme of Scientific Translation, Jerusalem, 67–112.

Appendix I

Additional examined specimens

Linum pubescens; C5/6 Turkey, Hatay: near sea level, rocky limestone slopes, 07 May 1965, *Coodé 620 & Jones* (E!); C5/6 Syria/Turkey, Hatay: nr. Aleppo, or between there and Iskenderun, *Russell* (BM photo!); C6 Gaziantep: 30 km. S of Aintab, *Dinsmore 8.935!*; Urfa: Rum Kala'a, *Sint.* 1888: 268!; C8 Mardin: 11 km. W of Idil, 800 m, 05 May 1966, *Davis 42.427* (E!).

Linum anisocalyx; C5 İçel: plaine de Mersina, May 1855, *Balansa* (holo. K!); İçel: Mersin, 1896, *Siehe 199* (E!); İçel: Mersin to Kuzucubelen, 18 June 1950, 500 m, *Hub.-Mor.* 10.567!

Linum viscosum; Spain, Gerona: Roadside near Ripoll, 13 July 1979, *Rogers 13.567* (B photo!), Huesca, near Jaca, 8 July 1979, *Rogers 13.558* (B photo!); Austria: Ebersberg en Autriche. In campis aridis Sabulosis, 1812 (GDC photo!); Italy: Gênes, 08 July 1808, *Candolle*, (GDC photo!).