

Two new non-spiny *Solanum* species from the Bolivian Andes (Morelloid Clade)

Tiina Särkinen¹, Sandra Knapp², Michael Nee³

1 Royal Botanic Garden Edinburgh, 20A Inverleith Row, EH3 5LR Edinburgh, United Kingdom **2** Department of Life Sciences, Natural History Museum, Cromwell Rd, SW7 5BD London, United Kingdom **3** Missouri Botanical Garden, P.O. Box 299, St. Louis, MO 63166-0299, U.S.A.

Corresponding author: Tiina Särkinen (t.sarkinen@rbge.ac.uk)

Academic editor: Pavel Stoev | Received 28 December 2014 | Accepted 5 March 2015 | Published 20 March 2015

Citation: Särkinen T, Knapp S, Nee M (2015) Two new non-spiny *Solanum* species from the Bolivian Andes (Morelloid Clade). *PhytoKeys* 47: 97–109. doi: 10.3897/phytokeys.47.4423

Abstract

Two new Bolivian species are described from the Morelloid clade of *Solanum* (section *Solanum* in the traditional sense). *Solanum alliarifolium* M.Nee & Särkinen, **sp. nov.** is found in montane forests between 1,900 and 3,200 m and is morphologically most similar to *S. leptocaulon* Van Heurck & Müll.Arg., also from montane forests in southern Peru and Bolivia. *Solanum rhizomatum* Särkinen & M.Nee, **sp. nov.** is found in seasonally dry forests and matorral vegetation in lower elevations between 1,300 and 2,900 m and is most similar to *S. pygmaeum* Cav., a species native to sub-tropical Argentina but introduced in subtropical and temperate areas worldwide.

Resumen

Se describen dos nuevas especies bolivianas del clado Morelloid de *Solanum* (sección *Solanum* en sentido tradicional). *Solanum alliarifolium* M.Nee & Särkinen, **sp. nov.** se encuentra en bosques montanos entre 1,900 y 3,200 metros y es morfológicamente muy similar a *S. leptocaulon* Van Heurck & Müll.Arg., también presente en bosques montanos en el sur de Perú y Bolivia. *Solanum rhizomatum* Särkinen y M.Nee, **sp. nov.** se encuentra en bosques tropicales estacionalmente secos y en vegetación de matorral a elevaciones más bajas entre 1,300 y 2,900 m, y es más similar a *S. pygmaeum* Cav., una especie nativa para Argentina subtropical pero introducida en las zonas subtropicales y templadas del mundo.

Keywords

Bolivia, endemism, Morelloid Clade, tropical Andes, Solanaceae, *Solanum* section *Solanum*

Palabras clave

Andes tropicales, Bolivia, Clado Morelloid, endemismo, Solanaceae, *Solanum* sección *Solanum*

Introduction

Solanum is one of the most species-rich vascular plant genera in the tropical Andes (Jørgensen et al. 2011), where many new species continue to be described (e.g., Anderson et al. 2006; Stern and Bohs 2010; Knapp 2010a,b; Farrugia and Bohs 2010; Tepe et al. 2012; Särkinen et al. 2013a; Särkinen et al. 2015). Here we describe two new *Solanum* species from Bolivia that belong to the Morelloid clade, one of major clades of non-spiny solanums (Weese and Bohs 2007; Särkinen et al. 2013b).

The Morelloid clade is a group of ca. 75 species, most of which are endemic to the tropical Andes (Bohs 2005; Särkinen et al. in review). The clade includes five major groups (traditionally recognised as sections *Solanum*, *Campanulisolanum* Bitter, *Parasolanum* A.Child, *Chamasarachidium* Bitter, and *Episarcophyllum* Bitter), which are in the process of re-circumscription based on molecular results (Särkinen et al. in review). Section *Solanum* is the largest of these with ca. 52 species and ca. 580 published names and is the only group to occur outside of the Americas. The Morelloid clade is distinguished by its herbaceous or sub-shrubby habit, usually internodal inflorescences, small flowers and fruits, and the usual possession of stone cells in the fruits (Bitter 1911). These are sclerified structures that are usually white and spherical rather than flattened and brown or yellowish brown like the seeds. Although some studies have examined the taxonomy of the Old World and North American species of this group (Edmonds 1977, 1978; Schilling 1981), monographic treatment is needed to aid species identification and to clarify synonymy in South America, where most of the species diversity is found (Edmonds 1972; Barboza et al. 2013).

Recent taxonomic work focusing on producing a monographic treatment of the Morelloid clade has resulted in the description of various new species from Peru, Bolivia and Ecuador (Särkinen et al. 2013a; Särkinen et al. 2015). Two additional new species are described here from Bolivia. Descriptions are based on field work and examination of herbarium specimens from 20 herbaria (BH, BM, COL, CORD, CPUN, DUKE, E, F, GH, GOET, HUSA, HUT, K, LPB, MO, MOL, NY, S, UDBC, US, USM, USZ). Extent of Occurrence (EOO) and Area of Occupancy (AOO) were calculated using GeoCat (www.geocat.kew.org) with a 2 km² cell size for AOO calculation. Conservation status of each species was assessed using the IUCN (2014) criteria based on the GeoCat analyses (Bachman et al. 2011) combined with field knowledge. All specimens are cited in the text, and full data is provided in the Suppl. material 1 and on Solanaceae Source (www.solanaceaesource.org).

Taxonomic treatment

Solanum alliarifolium M.Nee & Särkinen, sp. nov.

urn:lsid:ipni.org:names:77145835-1

Figs 1–2

Diagnosis. Similar to *Solanum leptocaulon* Van Heurck & Müll.Arg., but differing in its slender creeping habit with stems rooting along nodes, broadly ovate to orbicular leaves with mostly undulate to shallowly lobed margins, and a stellate corollas lobed to the middle with lobes reflexed at anthesis.

Type. BOLIVIA. Santa Cruz. Prov. Vallegrande: 6.5 km by air SW of Guadalupe on rd to Pucará, at turnoff to Santa Ana, 18°36'S, 64°07'W, 2675 m, 15 Dec 1990 (fl, fr), M. Nee 40315 (holotype: LPB; isotypes: MO [MO-2537105], NY [NY00852828], USZ).

Description. Slender herb to 20–30 cm high, with multiple long, creeping stems arising from a central taproot. Stems rooting at nodes, 1–2 mm in diameter, up to 50 cm long, glabrous or sparsely pubescent with spreading translucent 4–6-celled simple uniseriate trichomes ca. 0.2 mm long. Sympodial units difoliate, not geminate. Leaves simple, 1.5–3.6 cm long, 0.9–2.3 cm wide, broadly ovate to orbicular; adaxial surface glabrous; abaxial surface glabrous or sparsely pubescent with appressed 1–3-celled simple uniseriate trichomes along veins and leaf margins; primary veins 3–4 pairs; base rounded to attenuate, occasionally decurrent; margins entire, undulate, or shallowly lobed; apex acute; petiole 0.7–1.5 cm long, sparsely pubescent with simple 1–3-celled uniseriate trichomes like those of the stems, especially on young leaves. Inflorescences 1.5–3.0 cm long, simple, lateral, leaf-opposing or internodal, with 2–6 flowers, sparsely pubescent with simple uniseriate 4–6-celled spreading trichomes; peduncle 1.0–3.0 cm long, 0.4–0.5 mm in diameter at the apex and 0.6 mm in diameter at the base; pedicels 0.6–0.9 cm long, ca. 0.4 mm in diameter at the base and ca. 0.5 mm in diameter at the apex, straight and spreading at anthesis, articulated at the base; pedicel scars spaced 0.2–1.5 mm apart. Buds globose, white or purple-tinged. Flowers 5-merous, all perfect, nodding; calyx tube ca. 1.4–1.5 mm long, the lobes 1.6–2.0 mm long, rectangular-deltate in outline with rounded to acute apices, somewhat spreading at anthesis, sparsely pubescent with simple 1–4-celled uniseriate trichomes; corolla 1.4–1.6 cm in diameter, white to pale or deep violet-blue, with a dark purple ring and yellow-green central star at the base, stellate, lobed to the middle, the lobes ca. 4.0–5.0 mm long, 2.0–2.5 mm wide, reflexed at anthesis, densely pubescent abaxially with 1–2-celled simple uniseriate trichomes, these usually shorter than the trichomes of stems and leaves, glabrous adaxially; filament tube 1.3–1.5 mm long; free portion of the filaments ca. 1.1–1.6 mm long, pubescent with 4–7-celled uniseriate trichomes at the base adaxially; anthers 3.5–4.0 mm long, 0.8–1.0 mm wide, ellipsoid to rectangular in outline, yellow, poricidal at the tips, the pores lengthening to slits with age; ovary globose, glabrous; style 5–6 mm long, exerted 1.0–1.7 mm beyond the anther cone, densely pubescent with 2–3-celled simple uniseriate trichomes in the basal 2/3;




Figure 1. Illustration of *Solanum alliarifolium*. **A** Habit **B** Inflorescence with details of indumentum of simple, multi-cellular hairs along the stem, and short ciliate hairs along leaf margins **C** Flower just before anthesis, with and without corolla lobes removed **D** Flower at anthesis **E** Stamens **F** Gynoecium **G** Fruit (**A–C, E–G** Nee 40315, **D** Vargas 787). Illustration by Bobbi Angell.

stigma clavate, minutely papillate. Fruit a globose berry, 4–5 mm in diameter, green when developing, the colour when mature unknown, with a few stone cell aggregates in each berry; fruiting pedicels 1.1–3.2 cm long, ca. 0.4 mm in diameter at the base, ca. 0.6 mm in diameter at the apex, spreading, becoming somewhat woody; fruiting calyx lobes 2.8–3.2 mm long, spreading. Seeds 15–20 per berry, ca. 1.5–1.7 mm long, ca. 1.2–1.3 mm wide, flattened, reniform, pale-brown, the sub-lateral hilum positioned close to the middle, the testal cells pentagonal in outline.

Distribution. Endemic to montane forests of the Eastern Bolivian Andes in the Departments of Chuquisaca, Cochabamba, and Santa Cruz, in open areas close to water sources, near rivers and moist depressions, and marshy meadows on sandy or rocky substrates, associated with *Podocarpus parlatorei* Pilg., *Alnus acuminata* Kunth, *Hesperomeles ferruginea* (Pers.) Benth., *Alchemilla pinnata* Ruiz & Pav., *Azorella biloba* (Schltdl.) Wedd., *Weinmannia fagaroides* Kunth, *Baccharis genistelloides* (Lam.) Pers., *Clethra scabra* Pers., *Myrsine coriacea* (Sw.) Roem. & Schult., *Symplocos nana* Brand, *Eleocharis* spp., *Chusquea* spp., *Morella pubescens* (Willd.) Wilbur, ferns, grasses and Apiaceae herbs; between 1,900 and 3,200 m elevation.

Ecology. Flowering and fruiting during the wet season, generally from October–March, with a single record known from August.

Etymology. The epithet refers to the leaf shape, which struck the collector of the type (MN) as like that of *Alliaria petiolata* (M.Bieb.) Cavara & Grande (Brassicaceae), a European species invasively adventive in the eastern USA and other temperate areas of the world.

Conservation status. We assign a preliminary IUCN threat status of Vulnerable (VU, B1) to *S. alliariifolium* based on the small extent of occurrence (EOO=16,136 km²). The area of occupancy is even smaller (AOO=40 km²) and would merit status as endangered (EN), but knowing that collection densities in the tropical Andes remain extremely low and that the collections are mainly along the sparse road network, we prefer basing our assessment on the extent rather than area of occurrence. No occurrences are known within the protected area network in Bolivia thus far, but collection data indicates that the species endures grazing pressures relatively well.

Specimens examined. BOLIVIA. CHUQUISACA: Prov. Belisario Boeto: ca. 5 km S of Nuevo Mundo at summit of rd to Villa Serrano, 2,300 m, 18 Oct 1997, *J.R.I. Wood 12710* (K). **Prov. Hernando Siles:** Primera Sección Monteagudo, Cantón Fernández, Comunidad Vallecito, 20°12'43"S, 64°18'03"W, 2,314 m, 11 Aug 2007, *M. Jiménez 603* (NY); Laguna Milagros, 20°17'16"S, 64°02'56"W, 1,993 m, 26 Dec 2005, *R. Lozano & M. Serrano 1787* (MO, NY); Primera Sección Monteagudo, Cantón Fernández, Comunidad Vallecito, 20°12'18"S, 64°17'58"W, 2,456 m, 9 Nov 2007, *J. Villalobos 927* (MO). **COCHABAMBA: Prov. Carrasco:** Siberia, 17°49'87"S 64°44'07"W, 2,940 m, 20 Feb 2005, *S. Altamirano & M. Alcázar 3075* (MO); Sehuenka–Totora, 2,800 m, Nov 1959, *M. Cárdenas 5716* (US); Jatun Pino, 3200 m, Jan 1961, *M. Cárdenas 5942* (K); near Lagunillas (Totora), 2,700 m, Jan 1951, *M. Cárdenas 4663* (US); narrow canyon of Rio Monte Puncu, 5 km NE of Monte Puncu, 10 km by air NW of Epizana, 17°33'S, 65°16'W, 2,700–2,750 m, 10 Mar 1988, *M. Nee 36631* (NY); de Episana a Sehuencas via Montepunca, 2,500 m, 25 Mar 1978, *C.M. Ochoa 12022* (US); 5 km above Sehuencas towards Monte Puncu, 2,500 m, 4 Feb 1995, *J.R.I. Wood 9317* (K). **SANTA CRUZ: Prov. Caballero:** 5.5 km (by rd) E of town of El Churro at pond along Pojo–Comarapa highway, 17°50'15"S 64°44'00"W, 2,940 m, 22 Jan 2006, *M. Nee & Jun Wen 53903* (MO, USZ). **Prov. Vallegrande:** Huasacañada, 5 km al S de la ciudad de Vallegrande, 18°31'30"S 64°05'42"W, 2,050 m, 3 Nov 1990, *I. Vargas C. 787* (MO, NY, USZ); carretera




Figure 2. Distribution map of *Solanum alliarifolium*.

entre Vallegrande y Pucará, 18°35'49"S 64°07'34"W, 2,673 m, 5 Mar 2005, *J.R.I. Wood 21774* (K, USZ).

Discussion. *Solanum alliarifolium* is distinct within the Morelloids in being a slender creeping herb rooting at the nodes, with broadly ovate to orbicular leaves with undulate to shallowly lobed margins. It is morphologically most similar to *S. leptocaulon* Van Heurck & Müll.Arg., which occurs in similar montane habitats in Bolivia and in southern Peru, but the latter species is a small scrambling shrublet with ovate-lanceolate leaves with entire margins. *Solanum leptocaulon* further differs from *S. alliarifolium* in having a campanulate corolla lobed only 1/3 of the way to the base, rather than a stellate corolla lobed to 2/3 to the base with the lobes clearly reflexed at anthesis.

***Solanum rhizomatum* Särkinen & M.Nee, sp. nov.**

urn:lsid:ipni.org:names:77145836-1

Figs 3–4

Diagnosis. Like *Solanum pygmaeum* Cav., but differing in having mostly 1-branched inflorescences with 6–15 flowers, anthers < 3.5 mm long, strongly recurving fruiting pedicels, and berries < 1 cm in diameter with fewer than 30 seeds.




Figure 3. Paratype of *Solanum rhizomatum* (Wood 11974, K).

Type. BOLIVIA. **Santa Cruz:** Prov. Vallegrande, 10 km (by air) NNW of Vallegrande, 18°23'S, 64°08'W, 1850 m, 1 Feb 1987, *M. Nee & G. Coimbra S. 33947* (holotype: LPB; isotypes: G, MO [MO-5894880], NY [NY00824501]).

Description. Rhizomatous herb with erect stems up to 15–50 cm tall arising from an underground rhizome. Stems 1.5–4.0 mm in diameter at base, slightly flexuose, terete to ridged, often slightly winged, often purple-coloured, glabrous to sparsely pubescent with appressed 1–4-celled simple uniseriate trichomes ca. 0.5 mm long. Sympodial units difoliate, not geminate. Leaves simple, 2.3–8.0 cm long, 1.2–4.3 cm wide, ovate-lanceolate; adaxial surface glabrous or sparsely pubescent with 1–2-celled spreading hairs along lamina and veins; abaxial surface pubescent only along veins; primary veins 4–6 pairs; base attenuate to decurrent; margins lobed to entire, often purple-tinged, pubescent with short, 1-celled simple uniseriate trichomes, if present lobes present throughout or most commonly only in the basal 1/3 of the blade; apex acute to acuminate; petiole 0.5–1.2 cm long, sparsely pubescent with spreading, simple uniseriate trichomes like those of the stems and leaves. Inflorescences 1.5–3.1 cm long, lateral and internodal, simple to 1-branched, with 6–15 flowers, sparsely pubescent with simple 1–4-celled uniseriate appressed trichomes; peduncle 1.0–2.4 cm long, and if branched, each branch with a rachis 3–4 mm long; pedicels 4–6 mm long, ca. 0.3 mm in diameter at the base and ca. 0.4 mm in diameter at the apex, straight and spreading at anthesis, articulated at the base; pedicel scars spaced 1–2 mm apart. Buds ovoid, white or purple-tinged. Flowers 5-merous, all perfect; calyx tube ca. 2.0–2.5 mm long, the lobes 1.0–1.5 mm long, triangular with acute apices, sparsely pubescent with simple 1–3-celled appressed uniseriate trichomes; corolla 1.2–1.5 cm in diameter, white or flushed with blue, with a yellow-green basal star, stellate, lobed 1/2 to 2/3 of the way to the base, the lobes 4.0–5.0 mm long, 2.5–3.0 mm wide, reflexed at anthesis, later spreading, densely pubescent abaxially with 1–2-celled simple uniseriate trichomes, these usually shorter than the trichomes of stems and leaves, glabrous adaxially; filament tube 1.2–1.5 mm long; free portion of the filaments 1.0–1.2 mm long, pubescent along internal side with spreading hairs like those of the stems and leaves; anthers 3.2–3.5 mm long, 0.9–1.0 mm wide, ellipsoid or rectangular in outline, yellow; ovary globose, glabrous; style 6–7 mm long, exerted 2.5–3.0 mm beyond the anther cone, densely pubescent with 4-celled simple uniseriate trichomes in the basal 2/3; stigma globose, minutely papillate. Fruit a globose berry, 6–7 mm in diameter, pale green (mature ?), with a few stone cell aggregates; fruiting pedicels 1.2–1.4 mm long, ca. 0.6 mm in diameter at the base, ca. 0.8 mm in diameter at the apex, strongly recurving; fruiting calyx lobes 2.5–3.5 mm long, appressed to the berry with the tips slightly reflexed. Seeds 15–25 per berry, 1.7–1.8 mm long, 1.4–1.5 mm wide, concave-reniform, pale brown, the hilum positioned towards the narrower end of the seed, the testal cells pentagonal in outline.

Distribution. Endemic to the arid interior valleys of the Bolivian Andes in the Departments of Cochabamba, Potosí, Santa Cruz, and probably Chuquisaca, growing in seasonally dry tropical forests and dry matorral vegetation, along slopes and on rocky and sandy soils, often found growing in moist depressions under the shade of




Figure 4. *Solanum rhizomatum*. **A** Habitat in seasonally dry forests, eastern Bolivian Andes, Vallegrande, Dept. Santa Cruz **B** Habit amongst low herbs in partial shade **C** Rhizome **D** Inflorescence (**A–D** Nee & Mendoza 57594). Photos by M. Nee, scale bars = 1 cm.

larger trees and thickets, associated with *Prosopis kuntzei* Harms ex Kuntze, *Jodina rhombifolia* (Hook. & Arn.) Reissek, legumes, grasses, columnar cacti, and Asteraceae herbs; between 1,300 and 2,900 m elevation.

Ecology. Flowering and fruiting during the wet season from Jan. to March.

Etymology. *Solanum rhizomatum* is named for its rhizomatous underground stem.

Conservation status. We assign a preliminary IUCN threat status of Least Concern (LC) to *S. rhizomatum* based on the known extent of the species occurrence (EOO=43,101 km²). The extremely small observed area of occupancy (AOO=48 km²) could merit endangered status (EN), but knowing that collection densities in the tropical Andes remain extremely low and considering that current collections are from >10 different localities, we prefer basing our threat status assessment on the extent rather than area of occurrence. It is not known whether *S. rhizomatum* is similar in its biology and vegetative spread to *S. pygmaeum*, and further studies may clarify this aspect of potential conservation assessments in the future. No populations are known thus far from the protected area network in Bolivia. The growth form that allows effective vegetative spreading would indicate that the species can withstand grazing pressures moderately well.




Figure 5. Distribution map of *Solanum rhizomatum*.

Specimens examined. BOLIVIA. COCHABAMBA: Prov. Campero: Mizque, 2,020 m, 20 Feb 1967, *R.F. Steinbach* 721 (US); ca. 5 km de Villa Granada hacia Peña Colorada, 18°12'10"S, 65°00'09"W, 2,201 m, 25 Feb 2004, *J.R.I. Wood* 20266 (K, LPB). **Prov. Cercado:** en la salida de Cochabamba hacia el Valle Alto pasando La Tranca y ca. 2 km antes de la Angostura, 17°30'24"S 66°05'35"W, 2,676 m, 10 Feb 2005, *J.R.I. Wood* 21590 (K). **POTOSÍ: Prov. Nor Chichas:** 4 km N of Cotagaita on road to Potosí, 20°50'S, 65°41'W, 2,900 m, 30 Mar 1997, *J.R.I. Wood* 11974 (K, LPB). **SANTA CRUZ: Prov. Caballero:** carretera de Pulquina a Saipina a 0.5 km al oeste de la cumbre, 18°05'58"S, 64°30'56"W, 1,788 m, 20 Feb 2003, *J.R.I. Wood* 19134 (BOLV, K). **Prov. Florida:** 3 km S of Mataral, 18°08'S, 64°13'W, 1,425 m, 6 Feb 1988, *M. Nee* 36252 (MO, NY, USZ); 4 km by road W of Mataral, 18°09'21"S, 64°15'17"W, 1,300 m, 22 Feb 1984, *G. Schmitt* 27A (MO). **Prov. Vallegrande:** Pueblo de Vallegrande, Cerro los Tres Pilares, 100–150 m antes de llegar a la cima sobre el sendero hacia el pueblo, 18°29'21"S, 64°07'13"W, 2,236 m, 19 Jan 2003, *M. Mendoza* 449 (K); Choroquetal, ca. 5 km de Vallegrande sobre la carretera a Mataral, entrando ca. 500 m, sobre la senda hacia Chacateal, 18°28'07"S, 64°07'25"W, 1,932 m, 17 Mar 2003, *M. Mendoza* 529 (K); 4 km SW El Trigal, 18°20'S, 64°10'W, 1,600 m, 8 Mar 1988, *M. Nee* 36536 (MO, NY, USZ); Las Cañas, 2500 m, 28 Feb 1984, *C.M. Ochoa* 15548 (US); 6.5 km (by air) NE of airport in Vallegrande, along bad dirt road down

into the Río San Blas valley, 18°26'33"S 64°03'12"W 1,795 m, 3 Jan 2011, *M. Nee & M. Mendoza 57594* (USZ).

Discussion. *Solanum rhizomatum* is most closely related to *S. pygmaeum* from central and coastal Argentina (see Barboza et al. 2013), another rhizomatous species of *Solanum* section *Solanum*. *Solanum rhizomatum* differs from *S. pygmaeum* in having mostly 1-branched inflorescences with 6–15 flowers, anthers 3.2–3.5 mm long, strongly recurving fruiting pedicels, and berries with 15–25 seeds, while *S. pygmaeum* always has simple (unbranched) inflorescences with 2–6 flowers, anthers usually >3.5 mm long, fruiting pedicels that are broadly spreading, and berries with >50 seeds. Although these sets of characters overlap to some extent, *S. pygmaeum* individuals are generally smaller than those of *S. rhizomatum* (10–20 cm tall), with smaller leaves 1–5 cm long and 0.5–2.2 cm wide, while *S. rhizomatum* grows 15–50 cm tall, with larger leaves 2.3–8.0 cm long and 1.2–4.3 cm wide. *Solanum pygmaeum* forms dense colonies in secondary habitats such as railroad sidings, and has been introduced and naturalised in Europe and North America, presumably in boats carrying wool from eastern coastal Argentina (Barboza et al. 2013).

As in many species of *Solanum*, variation in corolla colour occurs in *S. rhizomatum*, where corollas vary from white to pale lilac even within single individuals. Label information from *Nee & Mendoza 57594* notes changes in the corolla colour during development, where the corolla is white in bud, violet in anthesis, and darker after wilting.

Acknowledgements

We thank Pawel Ficinski, Samantha Murphy, Morvah George, and Raneer Prakash for help with herbarium loans and databasing of specimens, Bobbi Angell for preparing the illustration of *S. alliarifolium*, and the herbaria mentioned in the text for specimen loans. This work was supported by NSF grant DEB-0316614 “PBI *Solanum*: A worldwide treatment” to SK & MN, and by National Geographic Society Northern Europe Award GEFNE49-12 to TS.

References

- Anderson GJ, Prohens J, Nuez F, Martine C (2006) *Solanum perlongistylum* and *S. catilliflorum*, new endemic Peruvian species of *Solanum*, section *Basarthrum*, are close relatives of the domesticated pepino, *S. muricatum*. *Novon* 16(2): 161–167. doi: 10.3417/1055-3177(2006)16[161:SPASCN]2.0.CO;2
- Bachman S, Moat J, Hill A, de la Torre J, Scott B (2011) Supporting Red List threat assessments with GeoCAT: geospatial conservation assessment tool. *ZooKeys* 150: 117–126. doi: 10.3897/zookeys.150.2109
- Barboza GE, Knapp S, Särkinen T (2013) *Solanum* Grupo VII. Moreloide. In: Barboza GE (Ed.) *Flora Argentina: Flora Vasculare de la Republica Argentina, Dicotyledoneae, Solanaceae* Vol. 13. Instituto de Botanica Darwinion, San Isidro, Argentina, 231–264.

- Bitter G (1911) Steinzellkonkretionen im Fruchtfleisch beerentrager Solanaceen und deren systematische Bedeutung. *Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie* 45: 483–507.
- Bohs L (2005) Major clades in *Solanum* based on *ndhF* sequence data. In: Keating R, Hollowell VC, Croat TB (Eds) *A festschrift for William G. D'Arcy – The legacy of a taxonomist*. Missouri Botanical Garden Press, St. Louis, U.S.A., 27–49.
- Edmonds JM (1972) A synopsis of the taxonomy of *Solanum* sect. *Solanum* (Maurella) in South America. *Kew Bulletin* 27: 95–114. doi: 10.2307/4117874
- Edmonds JM (1977) Taxonomic studies on *Solanum* section *Solanum* (Maurella). *Botanical Journal of the Linnean Society* 75: 141–178. doi: 10.1111/j.1095-8339.1977.tb01482.x
- Edmonds JM (1978) Numerical taxonomic studies on *Solanum* L. section *Solanum* (Maurella). *Botanical Journal of the Linnean Society* 76: 27–51. doi: 10.1111/j.1095-8339.1978.tb01497.x
- Farrugia F, Bohs L (2010) Two new South American species of *Solanum* section *Crinitum* (Solanaceae). *PhytoKeys* 1: 67–77. doi: 10.3897/phytokeys.1.661
- IUCN (2014) Guidelines for using the IUCN Red List Categories and Criteria. Version 11. Prepared by the Standards and Petitions Subcommittee. <http://www.iucnredlist.org/documents/RedListGuidelines.pdf> [12 Dec 2014]
- Jørgensen PM, Ulloa Ulloa C, León B, León-Yáñez S, Beck SG, Nee M, Zarucchi JL, Celis M, Bernal R, Gradstein R (2011) Regional patterns of vascular plant diversity and endemism. In: Herzog SK, Herzog SK, Martínez R, Jørgensen PM, Tiessen H (Eds) *Climate Change and Biodiversity in the Tropical Andes*. Inter-American Institute for Global Change Research (IAI) and Scientific Committee on Problems of the Environment (SCOPE), 192–203.
- Knapp S (2010a) Four new vining species of *Solanum* (Dulcamaroid Clade) from montane habitats in tropical America. *PLoS ONE* 5(5): e10502. doi: 10.1371/journal.pone.0010502
- Knapp S (2010b) New species of *Solanum* (Solanaceae) from Peru and Ecuador. *PhytoKeys* 1: 33–51. doi: 10.3897/phytokeys.1.659
- Särkinen TS, Gonzáles P, Knapp S (2013a) Distribution models and species discovery: the story of a new *Solanum* species from the Peruvian Andes. *PhytoKeys* 16: 1–20. doi: 10.3897/phytokeys.16.6312
- Särkinen TS, Bohs L, Olmstead RG, Knapp S (2013b) A phylogenetic framework for evolutionary study of the nightshades (Solanaceae): a dated 1000-tip tree. *BMC Evolutionary Biology* 13: 214. doi: 10.1186/1471-2148-13-214
- Särkinen TS, Gonzáles P, Knapp S (2015) Four new non-spiny species of *Solanum* from South America (*Solanum* sect. *Solanum*). *PhytoKeys* 44: 39–64. doi: 10.3897/phytokeys.44.8693
- Särkinen TS, Barboza GE, Knapp S (in review) True Black nightshades: Phylogeny and delimitation of the Morelloid clade of *Solanum*. *Taxon*.
- Schilling EE (1981) Systematics of *Solanum* sect. *Solanum* (Solanaceae) in North America. *Systematic Botany* 6: 172–185. doi: 10.2307/2418547
- Stern S, Bohs L (2010) Two new species of *Solanum* (Solanaceae) from the Amotape-Huanca-bamba Zone of southern Ecuador and northern Peru. *PhytoKeys* 1: 53–65. doi: 10.3897/phytokeys.1.660

Tepé EJ, Ridley G, Bohs L (2012) A new species of *Solanum* named for Jeanne Baret, an overlooked contributor to the history of Botany. *PhytoKeys* 8: 37–47. doi: 10.3897/phytokeys.8.2101

Weese TL, Bohs L (2007) A three-gene phylogeny of the genus *Solanum* (Solanaceae). *Systematic Botany* 32: 445–463. doi: 10.1600/036364407781179671

Supplementary material I

Occurrence records of *Solanum alliariifolium* and *Solanum rhizomatum*

Authors: Tiina Särkinen, Sandra Knapp, Michael Nee

Data type: occurrence

Explanation note: Occurrence records of the two new *Solanum* species.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.