

Didymocarpus phuquocensis, a new species of Gesneriaceae from Phu Quoc Island, South-western Vietnam

Thi-Lien Tran¹, Ngoc-Sam Ly^{2,3}, Minh-Ngoc Tran¹,
Xuan-Truong Nguyen¹, Ngoc-Giang Cao¹, Hong-Dung Pham⁴

1 National Institute of Medicinal Materials, 3B Quang Trung, Hoang Kiem District, Ha Noi, Vietnam **2** Institute of Tropical Biology, VAST, 85 Tran Quoc Toan road, District 3, Ho Chi Minh City, Vietnam **3** Graduate University of Science and Technology, VAST, 18, Hoang Quoc Viet, Cau Giay District, Ha Noi, Vietnam **4** Phu Quoc National Park, 1 Nguyen Chi Thanh, Duong Dong Town, Phu Quoc District, Kien Giang Province, Vietnam

Corresponding author: Ngoc-Sam Ly (lysamitb@gmail.com)

Academic editor: Alan Paton | Received 13 December 2019 | Accepted 7 August 2020 | Published 4 September 2020

Citation: Tran T-L, Ly N-S, Tran M-N, Nguyen X-T, Cao N-G, Pham H-D (2020) *Didymocarpus phuquocensis*, a new species of Gesneriaceae from Phu Quoc Island, South-western Vietnam. *PhytoKeys* 159: 35–44. <https://doi.org/10.3897/phytokeys.159.47442>

Abstract

A new species of Gesneriaceae, *Didymocarpus phuquocensis*, is described and illustrated from Phu Quoc National Park, Kien Giang Province, South-western Vietnam. It is most similar to *D. pulcher*, *D. hookeri* and *D. punduanus* in having 3-verticillate petiolate leaves, morphologically similar calyx, corolla, stamens, pistil and fruit, but differs from all in the glandular-pubescent stems, petioles and leaf blades, 1(–2)-flowered cymes, longer corolla and fruit and longer and densely glandular-puberulent ovary. Data on distribution, ecology, phenology and provisional conservation assessment of the new species are given along with an illustration and a colour plate.

Keywords

Didymocarpus, Kien Giang, new species, taxonomy, Vietnam

Introduction

The genus *Didymocarpus* Wallich was established in 1819, based on the species *D. primulifolius* D. Don from Nepal. The genus has previously been considered to comprise about 180 species distributed in tropical Asia with a few scattered in Africa and Australia (Wallich 1819; Don 1825; Wang et al. 1998; Weber and Burt 1998; Weber et al. 2000). The taxonomic delimitation of the genus *Didymocarpus* has varied considerably over time (e.g. Burt 1998; Weber et al. 2000; Möller et al. 2011; Möller and Clark 2013; Li et al. 2015). By combining molecular phylogenetic data and morphological revision of *Didymocarpus*, recent studies have remodelled and reduced this number (Vitek et al. 2000; Weber et al. 2000, 2011a, b; Möller et al. 2011; Li et al. 2015; Möller et al. 2017), with some species being placed in other genera, particularly to *Henckelia* Spreng. (ca. 60 species (Weber et al. 2011b, Middleton et al. 2013)), *Petrocodon* Hance (more than 20 species (Weber et al. 2011a)) and *Tribounia* D.J. Middleton (2 species (Middleton and Möller 2012)). The genus, as currently recognised, has around 98 species (Möller et al. 2017). Species of *Didymocarpus* are distinguished from other genera of Gesneriaceae by having: lithophyte perennial habit, ovate to ovate-cordate lamina which is mostly glandular-hairy, tubular corolla with an oblique limb (rarely trumpet-shaped or bell-shaped), two fertile stamens, three staminodes, a capitate stigma, an orthocarpic ovary and a bivalve capsule which dehisces loculicidally (Weber et al. 2000). As currently circumscribed, *Didymocarpus* ranges from northwest India, eastwards through Nepal, Bhutan, northeast India, Burma (Myanmar), southern China, Vietnam, Laos, Cambodia, Thailand and the Malay Peninsula, with the highest species diversity being found in China and Thailand (Weber and Burt 1998; Weber et al. 2000; Nangngam and Maxwell 2013; Nangngam and Middleton 2014; Möller et al. 2016). In Vietnam, five species of *Didymocarpus* have been reported, namely *D. bonii* Pellegr., *D. kerrii* Craib, *D. poilanei* Pellegr., *D. pulcher* C.B. Clarke and *D. purpureobracteatus* W.W. Sm. (Pellegrin 1930; Pham 2000; Phuong and Xuyen 2012; Phuong et al. 2014), but *D. bonii* Pellegr. [= *Calcareoboaea bonii* (Pellegr.) Burt] has been recognised as *Petrocodon bonii* (Pellegr.) A. Weber & Mich. Möller (Weber et al. 2011a). Recently, *Didymocarpus puhoatensis* X. Hong & F. Wen was described from Central Vietnam (Hong et al. 2018).

During medicinal plant investigations in Phu Quoc National Park (NP), Kien Giang Province, south-western Vietnam, several interesting plants of a small species of Gesneriaceae were collected by the authors in 2018–2019. The flowers of these plants have a capitate stigma and other features characterising this plant as *Didymocarpus* (Wang et al. 1998, Weber et al. 2000). A critical examination of living flowers, herbarium specimens of these plants and comparison with type material and protologues of all closely-related species in Vietnam and neighbouring countries (e.g. Clarke 1874, 1883, 1884; Wallich 1829; Wang et al. 1998; Pham 2000; Nangngam and Maxwell 2013; Phuong et al. 2014; Nangngam and Middleton 2014; Sinha and Datta 2016; Roy 2017), suggested that these specimens were different from the other known *Didymocarpus* species. These plants with 3-verticillate, petiolate leaves, campanulate

calyx, funnellform corolla found in Phu Quoc NP show similarities in these characters with *D. pulcher*, *D. hookeri* C.B. Clarke and *D. punduanus* Wall. ex R.Br. However, it shows significant differences in its vegetative and floral structures (see Table 1) and we describe it here as a species new to science (see also Taxonomic Notes).

Materials and methods

The descriptions are mainly based on measurements from flowering material of living plants in the field, supplemented by measurements from herbarium specimens. Type specimens of the most closely-related species were examined from the herbaria material from the following herbaria: E, K, HN, IBK, P, VNM, VNMN and W (herbarium codes follow Thiers (2020)), as well as digitised specimen images of *Didymocarpus* species also being accessed from botanical websites (e.g. <https://science.mnhn.fr/>, <http://www.cvh.org.cn/>, <https://plants.jstor.org/>). All morphological characters were studied under a dissection microscope and are described using the general terminology and

Table 1. Morphological comparison of *C. puhoatensis* with its most closely-related taxa (based on Wallich 1829; Clarke 1874, 1883, 1484; Sinha and Datta 2016; Roy 2017).

Characters	<i>D. phuquocensis</i>	<i>D. pulcher</i>	<i>D. hookeri</i>	<i>D. punduanus</i>
Plant height	(8–)10–19 cm	ca. 30.5 cm	ca. 40 cm	20.3–25.4 cm
Stem	glandular-pubescent	puberulous, eglandular	villous, eglandular	pubescent, eglandular
Petiole	glandular-pubescent	puberulous, eglandular	villous, eglandular	pubescent, eglandular
Leaves	3-verticillate petiolate leaves, terminal whorl of smaller sessile leaves	3–4- verticillate petiolate leaves, terminal whorl of smaller sessile or subsessile leaves	usually 3–4- verticillate petiolate leaves, uppermost leaves sessile	3-verticillate petiolate leaves, 2 sessile leaves at the apex
Leaf blade				
Margins	serrate	crenulate to serrate	crenulate	shallowly crenulate
Base	attenuate or cuneate	auriculate cordate	cordate	cordate
Apex	attenuate to acute	short acute	subobtuse	subobtuse
Indumentum	glandular-pubescent	pubescent, eglandular	somewhat villous eglandular	puberulous, eglandular
Inflorescence	axillary or terminal 1(–2)-flowered cyme	axial and terminal many-flowered cyme	terminal many-flowered cyme	terminal many-flowered cyme
Bracts	oblong-lanceolate, abaxially densely multicellular glandular-pubescent	rounded/suborbic-ular, sparsely viscos pilose	rounded, glabrescent	ovate, nearly glabrous
Corolla	4.8–5.3 cm long, light purple, glabrous	2.5–3 cm, violet-purple, glabrous or outside sparsely multicellular-villous	ca. 1.4 cm long, pale yellow with rose marks or nearly white, outside pilose	ca. 2.2 cm long, purple, nearly glabrous
Ovary	ca. 3 cm long, densely multicellular glandular-puberulent	1.1–1.8 cm, glabrous	1.1–1.7 cm long, pubescent	0.7–1.3 cm long, glabrous
Style	sparsely glandular-puberulent	glabrous	glabrous	glabrous
Capsule	4.4–5.5 cm long	ca. 4 cm long	1.3–3.5 cm long	ca. 4 cm long

standard work of Wang et al. (1998) and Harris and Harris (2001). A distribution map was created using SimpleMapper (<http://www.simplemapper.net/>) (Shorthouse 2010). Conservation status was assessed using the IUCN Red List Categories and Criteria version 3.1 (IUCN 2019) and inferring from the GeoCAT website (<http://geocat.kew.org/editor>) (Bachman et al. 2011).

Taxonomic treatment

Didymocarpus phuquocensis N.S.Lý, T.L. Tran & N.G. Cao, sp. nov.

urn:lsid:ipni.org:names:77211380-1

Figures 1, 2

Diagnosis. *Didymocarpus phuquocensis* is most similar morphologically to *D. pulcher*, *D. hookeri* and *D. punduanus* in the 3-verticillate, petiolate leaves, the morphologically-similar calyx, corolla, stamens, pistil and fruit, but differs from all in the glandular-pubescent stems and petioles (vs. puberulous, villous, pubescent and eglandular of the latter three, respectively), glandular-pubescent leaf blades (vs. pubescent, somewhat villous, puberulous and eglandular, respectively), 1(–2)-flowered cymes (vs. many-flowered cymes of the latter three), longer corolla 4.8–5.3 cm long (vs. 2.5–3 cm in *D. pulcher*, ca. 1.4 cm in *D. hookeri* and ca. 2.2 cm in *D. punduanus*), longer and densely multicellular glandular-puberulent ovary ca. 3 cm long (vs. 1.1–1.8 cm and glabrous in *D. pulcher*, 1.1–1.7 cm and pubescent in *D. hookeri* and 0.7–1.3 cm and glabrous in *D. punduanus*) and longer fruits 4.4–5.5 cm long (vs. ca. 4 cm long in *D. pulcher*, 1.3–3.5 cm long of *D. hookeri* and ca. 4 cm long in *D. punduanus*).

Type. VIETNAM. Kien Giang Province: Phu Quoc District, Phu Quoc NP, Suoi Mo, 18 September 2018, 10°14'40.90"N, 104°2'14.15"E, 82 m elev., Lý Ngọc Sâm, Cao Ngọc Giang, Nguyễn Thị Liên, Ngô Minh Huyền, Hùng, Hà Văn Long, TNB-305 (Holotype: VNM, isotype: P, NIMM).

Description. Deciduous, perennial, epilithic herb, (8–)10–19 cm tall, stems 2.5–3.5 mm in diameter. **Dry season:** new vegetative buds produced from the rhizome which then develops during the rainy season. **Rainy season:** stem erect, (3–)4 nodes, pale greenish, densely white multicellular glandular-pubescent; the longest node separated from the base of stem 5.7–12.2 cm long, the medium nodes at the middle stem 1.5–4.5 cm long, the shortest nodes very shortly distanced (0.2–0.5 cm long) at the apex. **Leaves** 3-verticillate, petiolate in the 2nd and 3rd whorls, other whorls with smaller and sessile or subsessile leaves; blades coriaceous, adaxially light greenish, abaxially whitish-green, asymmetrically narrowly elliptic to elliptic-ovate, the largest ones 7.2–10 × 2.4–3.7, the smaller ones 1–4.6 × 0.5–2.1 cm, adaxially densely white multicellular glandular-pubescent, abaxially sparsely white multicellular glandular-pubescent, apex attenuate to acute, base lightly oblique, attenuate to cuneate, margin serrate; venation pinnate, with 5–6 of ascending secondary veins on each side of midrib, somewhat opposite, adaxially obscure, abaxially prominent, densely covered with indumentum as




Figure 1. *Didymocarpus phuquocensis* **A** flowering plants in natural habit **B** flowering plants showing 3-verticillate leaves (red arrow) **C** mature plant with dried fruit and new stems **D** flower (top view) **E** flower (side view) **F** flower (front view) **G** inflorescence with peduncle, bract and calyx **H** calyx (top view) **I** longitudinal section of calyx showing disc **J** longitudinal section of corolla showing anterior (lower) lip with fertile stamens **K** longitudinal section of corolla showing posterior (upper) lips and staminodes (red arrow) **L** close-up of fertile stamens **M** ovary and calyx **N** dried fruits. The coloured plate prepared by Ngoc-Sâm Lý.

the stem; petioles terete, unequal in length, 0.5–3.2 cm long [the longest ones 3–3.2 cm, the shortest ones 0.5–1 cm], 2–2.5 mm in diam., whitish-green, sometimes tinted greenish-purple above, with indumentum as the stem. **Inflorescences** terminal or sub-terminal, cyme 1(–2) flowered, pendent; **peduncle** slender, (1.3–)3.1–4.5 cm long, ca. 0.5 mm in diam., tinted reddish-green, covered with white multicellular glandular and glandular-pubescent; **pedicels** 5–11 mm long, ca. 0.7 mm in diam., pale green, with indumentum as the peduncle, but more sparse; **bracts** paired; lanceolate to oblong-lanceolate, 4–5 × 1–1.5 mm, apex round to acute, margin entire, green, adaxially sparsely multicellular glandular-pubescent, abaxially densely white multicellular glandular-pubescent. **Calyx** campanulate, 5–6.5 mm long, dull reddish, outside sparsely multicellular glandular-puberulent; tube 3–4 mm long, 2–2.5 mm in diam.; lobes triangular, (sub)equal, 5-lobed, symmetrical, 2–2.5 mm long, ca. 1.5–1.8 mm wide at base, apices acute. **Corolla** funnellform, 4.8–5.3 cm long, glabrous, light purple, paler at base; tube 3.8–4.1 cm long, base narrow, 2–3 mm in diameter, widening abruptly at 1.9–2.1 cm from the base, widest at throat, 1–1.1 cm in diam.; lobes (sub)orbicular; anterior (lower or abaxial) lip 3-lobed, unequal, the middle one 5–7 × 7–10 mm, the lateral ones 5–7 × 4.5–5 mm, apices rounded; posterior (upper or adaxial) lip 2-lobed, slightly equal, 5–6 × 7.5–8.5 mm, apices rounded. **Stamens** 2, inserted at 2.2–2.4 cm above the base of the corolla; filaments slender, white, glabrous, 7–8 mm long, glandular-puberulent on the connective; anthers brownish, oblong, 2–2.2 × ca. 1 mm, tips and bases rounded, white-bearded; **staminodes** 3, inserted ca. 3 mm below the stamens, reduced to filaments, equal in length, 2.5–3.5 mm long, glabrous, tips with few glandular-puberulent. **Disc** cupular, ca. 1 mm high, margin irregular sinuate. **Pistil** ca. 3.2 cm long; **ovary** cylindrical, greenish with white towards the base, ca. 3 cm long, 1–1.5 mm in diam., densely glandular-puberulent; **style** continuous with the top of the ovary, ca. 2 × 1 mm, whitish, sparsely glandular-puberulent; **stigma** irregular capitate, ca. 1 × 1 mm, concave, white, papillose. **Capsules** straight to slightly curved, linear, glabrous, 2-valved, loculicidal dehiscent, 4.4–5.5 cm long, 1–1.5 mm in diam., turning brown when ripe, calyx and style persistent. **Pollen** and **Seeds** not studied.

Distribution and habitat. This species grows in moist places and shaded areas in primary tropical evergreen forests, on granite bedrock along streams or on moist and mossy cliffs in Phu Quoc NP, at 80–530 m elev. (Fig. 2).

Phenology. Flowering and fruiting from May to October.

Etymology. The specific epithet “*phuquocensis*” was named after the type locality.

Provisional conservation status. At present, four relatively-large subpopulations consisting of about 2000 mature individuals/mature clumps have been observed in Phu Quoc NP. The extent of occurrence (EOO) and the area of occupancy (AOO) were estimated using the web Geospatial Conservation Assessment Tool or GeoCAT (Bachman et al. 2011) and the auto-value cells width 2 km were calculated to be 53.7 km² and 16 km², respectively. These two values meet the criteria B1 (EOO < 100 km²) for Critically Endangered and B2 (AOO < 500 km²) for Endangered, following the IUCN Red List Categories and Criteria Version 3.1 (IUCN 2019). Although the known habitat of the new species is protected as part of the National Park, its habitat is fragmented and still faces some risk due to loss of the habitat within some parts




Figure 2. Distribution map of *Didymocarpus phuquocensis* in Phu Quoc National Park, Phu Quoc Island, Kien Giang Province.

of its range (in particular the clearing of forest land for agricultural fields and building of numerous roads and tourism areas). Based on the IUCN Red List Criteria (IUCN 2019), we therefore provisionally assess this species as Endangered (EN B2ab(iii), C).

Other specimens examined (*Paratypes*). VIETNAM. Kien Giang Province: Phu Quoc Island, Phu Quoc NP, K7 peak, 10°21'53.42"N, 104°0'31.22"E, 356 m elev., 21 May 2019, *Lý Ngọc Sâm, Hà Văn Long, TNB-430* (VNM); the same locality, Nui Chua peak, 10°22'40.09"N, 104°2'6.24"E, 532 m elev., 22 July 2019, *Cao Ngọc Giang, Ngô Minh Huyền, Hà Văn Long, TNB-502* (VNM); the same locality, Ham Rong Mount, 10°24'6.34"N, 103°58'6.47"E, 351 m elev., 10 July 2019, *Cao Ngọc Giang, Ngô Minh Huyền, Hà Văn Long, TNB-508* (VNM).

Vernacular name. Vietnamese language: Song bé phú quốc.

Taxonomic notes. Morphologically, the 3-verticillate petiolate leaves of *D. phuquocensis* are shared with several species of *Didymocarpus*, such as *D. insulsus* Craib (north-eastern, Thailand), *D. tristis* Craib (Chanthaburi Province, south-eastern Thailand), *D. dongrakensis* B.L.Burtt (northeast Thailand) *D. newmanii* B.L.Burtt (Chanthaburi Province, south-eastern Thailand), *D. pulcher* (from India, Buhtan, Nepal, China to Vietnam), *D. hookeri* (Assam, Arunachal Pradesh, Meghalaya and Sikkim, India), *D. punduanus* (Assam, Meghalaya and Nagaland, India) (Wang et al. 1998; Nangngam and Maxwell 2013; Phuong et al. 2014; Sinha and Datta 2016; Roy 2017). Of these, the campanulate calyx of the new species is similar to *D. pulcher*, *D. hookeri* and *D. punduanus*, but distinguished from *D. insulsus*, *D. newmanii* and *D. tristis* which have the calyx 5-lobed to the base. *Didymocarpus phuquocensis* is most similar to *D. pulcher*, *D. hookeri*

and *D. punduanus* in the morphological characters of 3-verticillate petiole leaves, the same shape of calyx, corolla, stamens, pistil and fruit. The major differences between the new species and the three latter are outlined above in the diagnosis. Moreover, the shorter plant height ((8–)10–19 cm), the narrowly elliptic to elliptic-ovate leaf blades that have serrate leaf margins, attenuate to cuneate leaf base and attenuate to acute leaf apex, the oblong-lanceolate bracts being abaxially densely multicellular glandular-pubescent, the light purple corolla and sparsely glandular-puberulent styles of *D. phuquocensis* also distinguish it from *D. pulcher*, *D. hookeri* and *D. punduanus*. A detailed morphological comparison between *D. phuquocensis*, *D. pulcher*, *D. hookeri* and *D. punduanus* is provided in Table 1.

Acknowledgements

This study was financial supporting by the grant of the project *TNB.ĐT/14-19/C16* belonging to the Program of Tây Nam Bộ (Vietnam) enabling our expedition in Phu Quoc Island of Kien Giang Province where the new species was discovered. The authors are thankful to Director of the Board of the NIMM and Phu Quoc National Park and their staff for the field trip permit and facilities and to Mr Ha Van Long, Nguyen Minh Hung, Le Duc Thanh and Mrs Ngo Thi Minh Huyen for field assistance. Special thanks to Mr Stephen Maciejewski, Gesneriad Society and Prof Dr Michael LoFurno, Temple University, Philadelphia, the USA for editorial assistance.

References

- Bachman S, Moat J, Hill AW, de Torre J, Scott B (2011) Supporting Red List threat assessments with GeoCAT: Geospatial conservation assessment tool. *ZooKeys* 150: 117–126. <https://doi.org/10.3897/zookeys.150.2109>
- Burtt BL (1998) Taxonomic history of *Didymocarpus* and *Henckelia*. *Beitr Biol Pflanzen* 70: 365–375.
- Clarke CB (1874) *Commelynaceae et Cyrtandraceae Bengalenses*. Thacker, Spink & Co., 120 pp.
- Clarke CB (1883) *Cyrtandraceae*. In: Alphonso, de Candolle (Eds) *Monographiae Phanerogamarum* 5: 1–137. [G. Masson; Masson & amp., Parisiis.]
- Clarke CB (1884) *Gesneriaceae*. In: Hooker JD (Ed.) *Flora of British India* 4. L. Reeve & Co., London, 336–375.
- Don D (1825) *Gesneriaceae*. *Prodromus Florae Nepalensis*. Linnean Society. Gale, J., London, 89–90. [121–124.]
- Harris JG, Harris MW (2001) *Plant Identification Terminology: an Illustrated Glossary* (2nd). Spring Lake Pub., Spring Lake, Utah, 216 pp.
- Hong X, Li ZL, Maciejewski S, Wen F, Do TV (2018) *Didymocarpus puhoatensis* (Gesneriaceae), a new species from Vietnam. *PhytoKeys* 94: 87–93. <https://doi.org/10.3897/phytokeys.94.21650>

- IUCN (2019) Guidelines for Using the IUCN Red List Categories and Criteria. Version 14. Prepared by the Standards and Petitions Subcommittee. <http://www.iucnredlist.org/documents/RedListGuidelines.pdf> [accessed 15 August 2019]
- Li JM, Sun WJ, Chang Y, Yang WG (2015) Systematic position of *Gyrocheilos* and some odd species of *Didymocarpus* (Gesneriaceae) inferred from molecular data, with reference to pollen and other morphological characters. *Journal of Systematics and Evolution* 54(2): 113–122. <https://doi.org/10.1111/jse.12169>
- Middleton DJ, Möller M (2012) *Tribounia*, a new genus of Gesneriaceae from Thailand. *Taxon* 61(6): 1286–1295. <https://doi.org/10.1002/tax.616009>
- Middleton MJ, Weber A, Yao TL, Sontag S, Möller M (2013) The current status of the species hitherto assigned to *Henckelia* (Gesneriaceae). *Edinburgh Journal of Botany* 70(3): 385–404. <https://doi.org/10.1017/S0960428613000127>
- Möller M, Clark JL (2013) The state of molecular studies in the family Gesneriaceae: a review. In: Clark JR (Ed.) *Proceedings of the World Gesneriad Research Conference 2010*. Selbyana 31: 65–253.
- Möller M, Forrest A, Wei YG, Weber A (2011) A molecular phylogenetic assessment of the advanced Asiatic and Malesian didymocarpoid Gesneriaceae with focus on non-monophyletic and monotypic genera. *Plant Systematics and Evolution* 292(3–4): 223–248. <https://doi.org/10.1007/s00606-010-0413-z>
- Möller M, Wei YG, Wen F, Clark JL, Weber A (2016) You win some you lose some: Updated generic delineations and classification of Gesneriaceae – implications for the family in China. *Guihaia* 36: 44–60.
- Möller M, Nampy S, Janeesha AP, Weber A (2017) The Gesneriaceae of India: Consequences of updated generic concepts and new family classification. *Rheedea* 27(1): 23–41. <https://doi.org/10.22244/rheedea.2017.27.1.5>
- Nangngam P, Maxwell JF (2013) *Didymocarpus* (Gesneriaceae) in Thailand. *Gardens' Bulletin* (Singapore) 65(2): 185–225.
- Nangngam P, Middleton DJ (2014) Five new species of *Didymocarpus* (Gesneriaceae) from Thailand. *Thai Forest Bulletin. Botany* 42: 35–42.
- Pellegrin F (1930) Gesnéracées (Cyrtrandrées). In: Lecomte H (Ed.) *Flore Générale de L'Indochine* (Vol. 4(5)). Masson & Cie, Paris, 456–608.
- Pham HH (2000) Gesneriaceae. In: Pham HH (Ed.) *An Illustrated Flora of Vietnam*, Vol. 3. Young Publishing House, Tp. Ho Chi Minh, 12–29. [In Vietnamese with English summary]
- Puong VX, Xuyen DT (2012) A new record of species *Didymocarpus kerrii* Craib (Gesneriaceae Dumort) for the flora of Vietnam. *Journal of Biology* 34(1): 72–74. <https://doi.org/10.15625/0866-7160/v34n1.671>
- Puong VX, Dang QV, Xuyen DT (2014) Genus *Didymocarpus* Wall. and a new record of species *Didymocarpus purpureobracteatus* Smith for the flora of Vietnam from Xuan Lien natural reserve, Thanh Hoa province. *Academia Journal of Biology* 36(1): 45–49. <https://doi.org/10.15625/0866-7160/v36n1.4516>
- Roy S (2017) Taxonomic studies of the family Gesneriaceae of Indian part of eastern Himalaya. PhD Thesis, University of Kalyani, West Bengal, India.

- Shorthouse DP (2010) SimpleMappr, an online tool to produce publication-quality point maps. <http://www.simplemappr.net> [accessed 11 October 2019]
- Sinha BK, Datta S (2016) Taxonomic account on the family Gesneriaceae in Northeast India. *Nelumbo* 58(0): 1–43. <https://doi.org/10.20324/nelumbo/v58/2016/105932>
- Thiers B (2020) (continuously updated) Index Herbariorum: A global directory of public herbaria and associated staff. New York Botanical Garden's Virtual Herbarium. <http://sweetgum.nybg.org/science/ih/> [accessed 20 February 2019]
- Vitek E, Weber A, Burtt BL (2000) Names, types and current placement of the species hitherto referred to *Didymocarpus*, *Loxocarpus*, *Codonoboea*, *Platyadenia* and *Henckelia* (Gesneriaceae). *Annalen des Naturhistorischen Museums in Wien* 102: 477–530.
- Wallich N (1819) Notice of the progress of botanical science in Bengal. *Edinburgh Philosophical Journal* 1: 376–381. [letter to F. Hamilton]
- Wallich N (1829) A Numerical List of Dried Specimens of Plants in the East India Company's Museum: Collected Under the Superintendence of Dr. Wallich of the Company's botanic Garden at Calcutta. East India Company, London, 306 pp. <https://doi.org/10.5962/bhl.title.1917>
- Wang WT, Pan K, Li Z, Weitzman AL, Skog LE (1998) Gesneriaceae. In: Wu ZY, Raven PH (Eds) *Flora of China* (Vol. 18). Science Press, Beijing & Missouri Botanical Garden, St. Louis, 349–358.
- Weber A, Burtt BL (1998) Remodelling of *Didymocarpus* and associated genera (Gesneriaceae). *Beiträge zur Biologie der Pflanzen* Breslau 70: 293–363.
- Weber A, Burtt VL, Vitek E (2000) Materials for a revision of *Didymocarpus* (Gesneriaceae). *Annalen des Naturhistorischen Museums in Wien* 102: 441–475.
- Weber A, Wei YG, Puglisi C, Wen F, Mayer V, Möller M (2011a) A new definition of the genus *Petrocodon* (Gesneriaceae). *Phytotaxa* 23(1): 49–67. <https://doi.org/10.11646/phytotaxa.23.1.3>
- Weber A, Middleton DJ, Forrest A, Kiew R, Lim CL, Rafidah AR, Sontag S, Triboun P, Wei YG, Yao TL, Möller M (2011b) Molecular systematics and remodelling of *Chirita* and associated genera (Gesneriaceae). *Taxon* 60(3): 767–790. <https://doi.org/10.1002/tax.603012>