

The resurrection of *Neohattoria* Kamim. (Jubulaceae, Marchantiophyta): a six decade systematic conflict resolved through a molecular perspective

Juan Larraín¹, Benjamin Carter², Blanka Shaw², Jörn Hentschel³,
Lynika S. Strozier¹, Tatsuwo Furuki⁴, Jochen Heinrichs⁵,
Barbara Crandall-Stotler⁶, John Engel¹, Matt von Konrat¹

1 *Science & Education, The Field Museum, 1400 South Lake Shore Drive, Chicago, IL 60605-2496, U.S.A.* **2** *Department of Biology, Duke University, Durham, NC 27708-0338, U.S.A.* **3** *Department of Systematic Botany with Herbarium Haussknecht and Botanical Garden, Friedrich Schiller University, Fürstengraben 1, 07743 Jena, Germany* **4** *Natural History Museum & Institute, 955-2 Aoba-cho, Chuo-ku, Chiba-shi, Chiba 260-8682, Japan* **5** *Ludwig-Maximilians-Universität München, Department für Biologie I, Systematische Botanik und Mykologie, GeoBio-Center, Menzinger Straße 67, 80638 München, Germany* **6** *Southern Illinois University, Department of Plant Biology, Mail Code 6509, Carbondale IL 62901-6509, U.S.A.*

Corresponding author: Juan Larraín (musgoschiloe@gmail.com)

Academic editor: E. Cooper | Received 20 March 2015 | Accepted 16 April 2015 | Published 16 June 2015

Citation: Larraín J, Carter B, Shaw B, Hentschel J, Strozier LS, Furuki T, Heinrichs J, Crandall-Stotler B, Engel J, von Konrat M (2015) The resurrection of *Neohattoria* Kamim. (Jubulaceae, Marchantiophyta): a six decade systematic conflict resolved through a molecular perspective. *PhytoKeys* 50: 101–122. doi: 10.3897/phytokeys.50.4940

Abstract

The systematic placement of *Frullania herzogii* has been contentious since its description six decades ago. Over the years it has been interpreted as either a member of the genus *Frullania* or segregated into its own genus, *Neohattoria*, due to morphological similarities with both *Frullania* and *Jubula*. Here we provide molecular evidence that supports the recognition of the genus *Neohattoria* and its inclusion within the Jubulaceae, together with *Jubula* and *Nipponolejeunea*. Jubulaceae are placed sister to Lejeuneaceae rather than to the monogeneric Frullaniaceae.

Keywords

DNA sequence data, *Frullania*, Frullaniaceae, Japan, *Jubula*, Jubulaceae, Lejeuneaceae, liverwort, *Nipponolejeunea*

Introduction

The liverwort *Frullania herzogii* S.Hatt. was originally described by Hattori (1955) from a poor, sterile specimen collected on Mt. Hayachine in Iwate Prefecture, northern Honshu, Japan. Since that time the generic and even familial placement of the species has remained controversial. The species also has remained poorly known partially because of its seemingly limited distribution in the subalpine coniferous forest zones of Honshu and Hokkaido, Japan, and the Kuril Islands (Inoue et al. 1981, Stotler and Crandall-Stotler 1987). Hattori (1955) remarked that the leaf morphology, with acute teeth along the margin, differed from all the other Japanese *Frullania* Raddi species known by him. A few years later, in his monograph of Japanese Frullaniaceae, Kamimura (1961) erected the new genus *Hattoria* Kamim. to separate this taxon from other *Frullania* species. He stated that although his new genus superficially resembled species of *Cololejeunea* (Spruce) Schiffn. or *Frullania*, there was an important similarity between the branching patterns of *Hattoria herzogii* (S.Hatt) Kamim. and species in the genus *Jubula* Dumort. In both *Jubula* and *Hattoria*, the branches replace the lobule of the leaf at the point of insertion, and the leaf lobes are attached to both the main stem and to the branch. Although Kamimura (1961) noted the similarity of cell shape between *Hattoria* and *Frullania*, he considered the combination of branching architecture and leaf denticulation sufficient to recognize *Hattoria* as a distinct genus. A year later he had to give a new name, *Neohattoria* Kamim., to his recently described genus (Kamimura 1962), because of the almost simultaneous although earlier description of *Hattoria* by Schuster for a liverwort in the Lophoziaceae (Schuster 1961).

Later Schuster (1963), in a key for the Southern Hemisphere genera of liverworts, expanded the circumscription of *Neohattoria* to include two more species, *Frullania microscopica* Pearson from New Caledonia, and *F. parhamii* (R.M.Schust.) R.M.Schust. ex von Konrat, L.Söderstr. & A.Hagborg from Fiji. He based his taxonomic decision on the morphology of the reduced leaves on branch bases, the subfloral innovations, and the sharply delimited bracts and bracteoles of *F. microscopica*, and on the toothed leaf lobes of this species. Schuster (1963) did not provide any argument for placement of the Fijian *F. parhamii* in *Neohattoria*, other than the hyaline margins of the leaves that can be seen in this species and in *F. microscopica* (as inferred from the key). However, his key is restricted to the Southern Hemisphere and did not include the type of the genus, which completely lacks a hyaline border in leaf lobes. Schuster (1970) later expanded this generic concept even further, including the Australasian *F. rostrata* (Hook.f. & Taylor) Hook.f. & Taylor ex Gottsche, Lindenb. & Nees (as *Neohattoria australis* R.M.Schust.) and *F. hodgsoniae* von Konrat, Braggins, Hentschel & Heinrichs (as *Neohattoria rostrata* R.M.Schust.), the SE Asian *F. junghuhniana* Gottsche var. *tenella* (Sande Lac.) Grolle & S.Hatt. [as *Neohattoria perversa* (Steph.) R.M.Schust.], the New Caledonian *F. chevalieri* (R.M.Schust.) R.M.Schust. and *F. neocaledonica* J.J.Engel

(as *Neohattoria caledonica* R.M.Schust.). Of these, *F. hodgsoniae* is now considered a member of *F.* subg. *Diastaloba* Spruce sect. *Inconditum* von Konrat, Hentschel & Heinrichs (von Konrat et al. 2010), while the rest of the taxa are currently included in *Frullania* subg. *Microfrullania* (R.M.Schust.) R.M.Schust. The current taxonomic placement of these taxa is based on both morphological (Hattori and Mizutani 1982, Schuster 1992) and molecular evidence (Hentschel et al. 2009, von Konrat et al. 2012).

Asakawa et al. (1979) demonstrated, based on chemical compound differences, that Jubulaceae *sensu lato* should be divided into three families, i.e. Jubulaceae, Frullaniaceae and Lejeuneaceae. This view has been confirmed by most molecular phylogenies published to date (e.g., Forrest et al. 2006, Heinrichs et al. 2005, 2007). Asakawa et al. (1979) listed 11 morphological characters that support the separation of Frullaniaceae and Jubulaceae, and placed *Neohattoria* together with *Jubula* in the Jubulaceae. Hattori (1982, 1984, 1986) and Hattori and Mizutani (1982) also accepted the separation between Jubulaceae and Frullaniaceae and argued that *Amphijubula* R.M.Schust., a genus formerly considered by Schuster (1970, 1980) as intermediate between *Jubula* and *Frullania*, should be placed within *Frullania*. This view was first held by Engel (1978), who had earlier reduced *Amphijubula* to a synonym of *Frullania*.

In 1987, Stotler and Crandall-Stotler published a thorough treatise of the taxonomic history of *Neohattoria herzogii* (S.Hatt.) Kamim. in the context of a detailed re-evaluation of its morphology, including the discovery of immature female inflorescences. In that contribution they came to the conclusion that this taxon should be considered within the circumscription of *Frullania*, although in its own subgenus, *F.* subg. *Dentatilobi* Stotler & Crand.-Stot. Their conclusion was based on both vegetative and reproductive characters, including the morphology of the bracts surrounding the female gametangia, lobule anatomy, leaf cell pattern, and the morphology of regenerants. Although they recognized that leaf-lobe insertion, branch morphology, and morphology of stylus are more similar to *Jubula* than to *Frullania*, they concluded that on the basis of the *Frullania*-like inflorescences and regenerants, *Neohattoria* should be synonymized with *Frullania*. This synonymy was adopted by Grolle and Meister (2004) who described a morphologically similar plant from Oligocene amber from Bitterfeld (Germany) as *Frullania* (subg. *Dentatilobi*) *hamatosetacea* Grolle. However, this fossil species appears morphologically closer to *F.* subg. *Microfrullania* than to *Neohattoria*, and this issue will be explored in detail in a forthcoming monograph of the latter subgenus.

Lack of useable specimens has previously precluded inclusion of *Neohattoria* in molecular phylogenetic studies. As a result of recent collecting activities, fresh material became available that allowed for successful DNA extraction and amplification. In the present study, we use molecular sequence data to investigate the phylogenetic position of *Neohattoria*. We investigate whether the genus should be placed in the Frullaniaceae or the Jubulaceae and evaluate whether molecular evidence supports the recognition of *Neohattoria* as a distinct genus.

Methods

Microscopy

For the production of microscopic images an Olympus BX51 microscope was used, equipped with both a QICAM Fast1394 camera from QIMAGING (Surrey, Canada), and a slide scanner (moving platform stage attached between the objectives and the condenser) from Objective Imaging Ltd. (Cambridge, UK). The software “Surveyor” from the latter company was used for the digitally rendered images.

DNA extraction, PCR amplification and sequencing

We worked with two independent datasets to address two different questions, (1) what is the position of *Neohattoria* relative to the Frullaniaceae, Jubulaceae and Lejeuneaceae, and once we obtained results from these analyses, we asked (2) what is the position of *Neohattoria* within the Jubulaceae. For dataset 1 sequences were generated for two mitochondrial (*nad1*, *rps3*), and two chloroplast loci (*psbA*, *rbcL*), following DNA extraction, amplification and sequencing methods described by Shaw et al. (2003), and using primer sequences provided in Cooper et al. (2011). For dataset 2 we used the aforementioned plastid regions (*psbA* and *rbcL*) together with the nuclear ITS region following the methods described by Shaw et al. (2003), and the chloroplast *trnL-trnF* region, amplified and sequenced as described in von Konrat et al. (2012). All sequences were edited and manually aligned in PhyDE v0.9971 (www.phyde.de) following the alignment rules and hotspot definitions presented in Kelchner (2000), Olsson et al. (2009), and Borsch and Quandt (2009).

Taxon sampling and outgroup selection

For dataset 1 seven species of *Radula* were selected as outgroup taxa following the results already published in recent liverwort phylogenies (Davis 2004, Forrest et al. 2006, Feldberg et al. 2014, Heinrichs et al. 2005, 2007). The same criteria were undertaken for dataset 2, including all taxa with sequences available in GenBank for *Jubula* and *Nipponolejeunea* S.Hatt. (Ahonen 2006, Ahonen et al. 2003, Konstantinova and Vilnet 2011, Päscht et al. 2010, Wilson et al. 2004, 2007), using selected taxa of the Lejeuneaceae and species of *Frullania* as outgroup based on results from dataset 1. GenBank accession numbers for both newly generated sequences and for already published sequences are provided in Appendices 1 and 2 for datasets 1 and 2 respectively.

Phylogenetic inferences

Both datasets were analysed with PartitionFinder v1.1.0 (Lanfear et al. 2012, 2014) to develop best-fit partitioning schemes and models of molecular evolution. Dataset 1

was partitioned setting one separate data block for each of the four genes used, each of them divided in three according to each codon position; introns and/or spacers were coded as extra partitions. Dataset 2 was partitioned in four parts, corresponding to the regions included only, without inner codon partition for the coding regions analysed. For dataset 1, phylogenetic reconstructions under maximum likelihood (ML) were performed in GARLI v2.01 (Zwickl 2006), setting up seven different models for the eleven partitions determined by PartitionFinder. Two independent searches each with 100 bootstrap replicates were made, and the 50% majority-rule consensus tree from all obtained trees was obtained with SumTrees v3.3.1 included in the package DendroPy v3.12.2 (Sukumaran and Holder 2010). Bayesian Posterior Probabilities analyses (PP) were executed in MrBayes v3.2.2 (Huelsenbeck and Ronquist 2001, Ronquist and Huelsenbeck 2003) also with the partitioned data set as given by PartitionFinder, and setting a different model for the individual partitions from the available options in MrBayes, with all characters given equal weight and gaps treated as missing data. The default settings of the program for a priori probabilities were used. Four runs, each with four MCMC chains (one million generations each) were run simultaneously, with the temperature of the heated chain set to 0.2 (default setting). Chains were sampled every 100 generations. Calculation of the consensus tree and posterior probabilities of clades was based on the set of trees sampled after the chains had converged, as observed graphically using Tracer v1.5 (Rambaut and Drummond 2007). For dataset 2, phylogenetic reconstructions under ML were performed in GARLI v2.01 and Bayesian analyses were executed with MrBayes v3.2.2 following the protocols as described above. For this dataset only three different partitions were suggested by PartitionFinder, and the models given by this software for each partition were incorporated into the settings of both the ML and the Bayesian analysis. Trees were edited and support values added using TreeGraph v2.0.54-364 beta (Stöver and Müller 2010).

Results

The complete alignment for dataset 1 including all four regions mentioned above, with flanking areas pruned to avoid ambiguous readings, comprised 4818 characters for 54 accessions, of which 694 were parsimony informative. A total of 101 new sequences were generated for this study (Appendix 1). In the analysis of the *Neohattoria* sequences with accessions of the Frullaniaceae, Jubulaceae and Lejeuneaceae (dataset 1), *Neohattoria* is strongly supported (as defined by Pedersen et al. 2007) as one of three clades belonging to the Jubulaceae in both ML and Bayesian analyses, with accessions of *Nipponolejeunea*, resolved in a second clade and those of *Jubula*, in a third clade (Fig. 1), although the latter with low support (ML = 52, PP = 0.6). The Jubulaceae is resolved as sister to the Lejeuneaceae with strong support in both types of analysis. The position of the Frullaniaceae as sister to this latter clade (Jubulaceae + Lejeuneaceae) was strongly supported by the Bayesian analyses (PP = 1.0), but it was not recovered by the ML analyses. The Bayesian analyses also resolved *Neohattoria* as sister to the rest of the Jubulaceae (*Nipponolejeunea* + *Jubula*) with strong support (PP = 1.0).




Figure 1. Maximum likelihood (ML) tree showing the systematic position of *Neohattoria* relative to the Jubulaceae, Frullaniaceae and Lejeuneaceae. Wide black branches indicate ML bootstrap support > 90 % and PP > 0.95.




Figure 2. Maximum likelihood (ML) tree showing the systematic position of *Neohattoria herzogii* within the Jubulaceae. Only 1/2 of the length of the branch between the Frullaniaceae and the Lejeuneaceae/Jubulaceae clade is depicted. Wide black branches indicate ML bootstrap support > 90% and PP > 0.95.




Figure 3. *Neobattoria herzogii*. **A** Habit, dorsal view **B** Habit, ventral view with distal lobules detached **C** Regenerant shoot originating from a detached lobule **D** Lobule **E** Underleaf **F–K** Leaves. All from Furuki 22673 (F). Scale bar: 350 μm (**A**, **B**), 200 μm (**C**), 180 μm (**D**), 300 μm (**E**), 150 μm (**F–K**).

The complete alignment for dataset 2 including all four regions included, and after pruning the flanking areas to avoid ambiguous readings and deleting unalignable areas of the ITS region, comprised 3737 characters for 55 accessions, of which 548 were

parsimony informative. The four different regions were not equally represented in the matrix, as shown in Appendix 2. The results of the analyses (Fig. 2) confirm with strong support the placement of *Neohattoria* within the Jubulaceae (ML = 100, PP = 1.0), and forming a sister clade to *Nipponolejeunea*, although recovered with strong support only by the Bayesian analysis (ML = 64, PP = 0.97). *Jubula* was resolved as the sister clade to the *Neohattoria-Nipponolejeunea* clade, although with low support (ML = 65, PP = 0.5).

The voucher of *Neohattoria herzogii* used for DNA extraction is illustrated in Figure 3.

Discussion

Our molecular analyses support recognition of the genus *Neohattoria* as distinct from the genus *Frullania*, as first proposed by Kamimura (1961) almost 55 years ago. Moreover, our molecular analysis strongly supports its inclusion within the Jubulaceae, together with *Jubula* and *Nipponolejeunea*. A close relationship with *Jubula*, based on similarities in branch morphologies, was first suggested by Kamimura (1961, p. 94), and also accepted by Hattori et al. (1972). Inoue et al. (1981) provided new karyological, chemical and ecological data on *N. herzogii* and concluded that the biosystematic evidence collected suggested distance between *Jubula* and *Neohattoria*, but, nonetheless, retained *Neohattoria* in the Jubulaceae. While morphologically closer to *Jubula* than *Nipponolejeunea* to which it is sister, it is clearly not nested in the *Jubula* clade. This combination of molecular and morphological evidence, in fact, supports its recognition as a distinct genus in the Jubulaceae.

Circumscription and relationships of the Jubulaceae

Our results strongly support the position of the Jubulaceae (containing *Jubula*, *Nipponolejeunea* and *Neohattoria*) sister to the Lejeuneaceae, and the Frullaniaceae as sister of the latter clade, although without significant support (Fig. 1). These results agree with several molecular phylogenies (e.g. Ahonen 2004, Forrest et al. 2006, Heinrichs et al. 2005, 2007). Thus the traditional view of a widely circumscribed Jubulaceae including *Frullania* is further rejected in this study.

These three families (Frullaniaceae, Jubulaceae and Lejeuneaceae) share several morphological characters, including the leaves divided into two (or three) parts [lobe, lobule (and stylus)], the beaked perianths, the sporophyte enclosed in a stalked true calyptra, the bistratose capsule wall, and the vertically aligned elaters that are attached to the valve apices (Crandall-Stotler et al. 2009, Gradstein et al. 2001, Schuster 1992). However, these characters need to be carefully evaluated to understand their evolution and their role in demonstrating the history of these lineages. In the past, *Jubula*, *Frullania* and members of the Lejeuneaceae were placed in a single taxonomic group (the subtribe Jubuleae), based largely on the similarities among their sporophytes (e.g. Müller 1915). Verdoorn (1930) argued that based on most characters (e.g., number of archeogonia, seta form, and

lobule ontogeny) *Jubula* belongs nearest to *Frullania*, which later lead Schuster (1992, p. 6) to describe *Jubula* as a “*bona-fide* genus of Jubulaceae [= Frullaniaceae]”. Mizutani (1961) was the first to propose that, except for the lobule structure, *Jubula* had no alignment with *Frullania*, and subsequently placed *Jubula* into the Lejeuneaceae. However, Asakawa et al. (1979) concluded that chemically, both *Jubula* and *Frullania* are quite different from *Lejeunea* Lib. species. Interestingly, the phylogenetic analysis by Crandall-Stotler and Stotler (2000) of 40 gametophyte and 21 sporophyte characters distributed among 34 liverwort families, resolved *F. asagrayana* Mont. as sister to a clade containing *J. hutchinsiae* (Hook.) Dumort. *subsp. pennsylvanica* (Steph.) Verd. and *L. cavifolia* (Ehrh.) Lindb. However, in the systematic treatment of the same work (Crandall-Stotler and Stotler 2000) Jubulaceae is presented as including both *Jubula* and *Frullania*, whereas the Lejeuneaceae is presented as a separate family, following accepted classifications of the time. The revised version of that classification, incorporating some recent molecular data, presents the Frullaniaceae, Jubulaceae and Lejeuneaceae as three separate families within the suborder Jubulineae (Crandall-Stotler et al. 2008, 2009), which is accepted here but with the transfer of *Neohattoria* from the Frullaniaceae to the Jubulaceae.

Assessing the importance of different morphological characters in circumscribing Frullaniaceae, Jubulaceae and Lejeuneaceae has been a difficult problem, but there are several characters that are consistent with the molecular phylogenetic results presented here. In most Lejeuneaceae a true stylus does not develop, but instead a single, unstalked slime papilla is formed at the junction of the lobule base and the stem, while in *Jubula* and *Neohattoria* there is a one- or two-celled filament terminated by a slime papilla in this position (Crandall-Stotler and Guerke 1980, Stotler and Crandall-Stotler 1987). Both types of structures are clearly different from those of the Frullaniaceae, where the stylus is always formed by more than two cells and is usually very conspicuous. The Jubulaceae and Frullaniaceae can be clearly differentiated from the Lejeuneaceae by the lobule, which is almost free from the larger dorsal lobe, and typically modified into an inflated, balloon-like to helmet-shaped sac whose aperture is directed either toward the shoot base or toward the stem, with the exception of *Nipponolejeunea* which has Lejeuneaceae-like lobules. Guerke (1978) hypothesised that *Jubula* was more advanced than *Frullania* on the basis that *Jubula* has many specialized characteristics e.g., a highly reduced stylus, seta, and foot, and features associated with the sporeling. In contrast, Schuster (1992, p. 9) stated that taxa such as *Amphijubula microcaulis* (Gola) R.M.Schust. (\equiv *F. microcaulis* Gola), with a 16 + 4 seriate seta and monogynous gynoecia, diminish the distinctions between the two groups such that he prefers not to attempt a “subfamilial separation” at all. However, revision of the chemical, morphological, and ecological data provided support for the recognition of two subfamilies in the Jubulaceae (Guerke 1978, von Konrat 2004). Alternatively, Asakawa et al. (1979), on the basis of biochemical and morphological evidence, proposed two families: Jubulaceae (*Jubula*, *Neohattoria*) and Frullaniaceae (*Frullania*, *Stearia* S.Hatt., *Amphijubula*, and *Schusterella* S.Hatt.). Hattori (1982, 1984, 1986) and Hattori and Mizutani (1982) also accepted two families. This approach has been adopted in most recent hepatic floras and classifications (Paton 1999, Damsholt 2002, Casas et al. 2009, Crandall-Stotler et al. 2009, Frey and Stech 2009).

Schuster (1980, 1992) questioned the division into two families and argued that only the single family Jubulaceae should be recognized, but commented that this area of classification remains replete with ambiguities and contradictions. Interestingly, he also suggested that there was a possibility that *Neohattoria* might share a closer affinity to Jubulopsidaceae (= Lepidolaenaceae) than to Jubulaceae (Schuster 1996), a view first expressed when Grolle (1966) transferred *Jubula novae-zelandiae* E.A.Hodgs. & S.W.Arnell, which is the generic type of *Jubulopsis* R.M.Schust., to *Neohattoria*. However, recent molecular analyses (e.g., Heinrichs et al. 2005, Forrest et al. 2006) have demonstrated that *Jubulopsis* (= *Lepidolaena*) is far removed from the Jubulaceae.

Morphologically, the monogeneric Frullaniaceae can be differentiated from the Jubulaceae by: (1) plants usually with conspicuous secondary pigmentation, often reddish; (2) initial leaves of branches either trifid or bifid; and (3) spores with rosette-like protrusions. Conversely, in the Jubulaceae the plants are: (1) soft and without secondary pigmentation (thus usually dull green to pale brown); (2) the initial leaves of branches are small, subtriangular, and never tri- or bifid; and (3) the spores without rosette-like protrusions. The first two of these characters support the placement of *Neohattoria* within Jubulaceae rather than Frullaniaceae (spores remain unknown in *Neohattoria*).

Chemically, *Frullania* species in general, produce significant amounts of sesquiterpene lactones, diterpenoids, and bibenzyl derivatives, which are considered important chemosystematic markers of the group (Asakawa et al. 1981, 1983, 1987, Kraut et al. 1994). On the other hand, cyclocolorone and maalioxide have been isolated as major components of *Jubula hutchinsiae* (Hook.) Dumort. subsp. *japonica* (Steph.) Horik. & Ando (Asakawa et al. 1979); interestingly cyclocolorone is also widely distributed in the Porellaceae. In contrast, no members of *Jubula* or *Frullania* produce paraffinic hydrocarbons which are characteristic for *Neohattoria* (Inoue et al. 1981).

Interestingly, Schuster (1996) suggested that there was a possibility that *Neohattoria* might share a closer affinity to Jubulopsidaceae (= Lepidolaenaceae) than to Jubulaceae. This view was first expressed when Grolle (1966) transferred *Jubula novae-zelandiae* E.A.Hodgs. & S.W.Arnell, which is the type species of *Jubulopsis* R.M.Schust., to *Neohattoria*. However, preliminary unrooted trees made for this contribution including *Ascidiota* C.Massal., *Gackstroemia* Trevis., *Goebeliella* Steph., *Lepidogyna* R.M.Schust., *Lepidolaena* Dumort. (= *Jubulopsis*) and *Porella* L. together with representatives outside the Porellales, showed *Neohattoria* far away from Lepidolaenaceae but within Jubulaceae (results not depicted). These results are basically the same as the ones observed in recent molecular phylogenies (e.g. Heinrichs et al. 2005, Forrest et al. 2006), demonstrating that these groups are only distantly related to either the Jubulaceae or the Frullaniaceae.

Circumscription and relationships of *Neohattoria*

Our results place *Neohattoria* in the Jubulaceae with strong support, together with *Nipponolejeunea* and *Jubula*. Within the Jubulaceae, *Neohattoria* is resolved as sister to *Nipponolejeunea*, and this latter clade sister to *Jubula*, although this relationship is sen-

sitive to taxon sampling (cf. Figs. 1 and 2), and not strongly supported in the analyses. When describing the genus *Hattoria* (later renamed *Neohattoria*), Kamimura (1961) conceived it as a monotypic genus containing only the Japanese endemic *N. herzogii*. The singularity of this taxon was well described and illustrated, highlighting its closer affinities to *Jubula* instead of *Frullania*, mostly because of its branching pattern and leaf insertion: “[...] *the branch replaces the lobule of leaf in origin and the lobe is inserted partly to the stem and partly to the branch. The first leaf and underleaf of branches are much deformed, being the “Vorblätter” of Verdoorn (1930).*” (Kamimura 1961, p. 94). The characteristic combination of traits that led Kamimura to describe this new genus vanished when Schuster (1963, 1970) added more species in the circumscription of *Neohattoria* as explained above. Schuster (1970) still recognized the taxonomic singularity of *N. herzogii* when placing it in its own subgenus within *Neohattoria*, but failed to see the relationships of this taxon with other *Jubula* species, precisely because of his wide concept of *Neohattoria* that includes members of *F.* subg. *Microfrullania* and *F.* subg. *Diastaloba*.

Oil-bodies in *Neohattoria* are homogenous, usually more than ten per cell, and similar in size to chloroplasts (Hattori et al. 1972, Inoue et al. 1981). Hattori et al. (1972) reported 10–20 oil-bodies per leaf lobe median cell for *N. herzogii* and later Inoue et al. (1981, p. 25) reported a similar number “usually 7–15 per leaf-lobe cell (rarely up to 22)”. Hattori et al. (1972) stated that oil-bodies of *Neohattoria* are hyaline and homogenous, and Inoue et al. (1981) recorded in their specimen of *Neohattoria* that the oil-bodies were completely colourless and homogenous. However, they noted that sometimes they were faintly papillose with a few distinct granules; Inoue et al. (1981) were uncertain if this was due to degeneration of the oil-bodies. Reports of oil-body numbers for *Jubula* are ambiguous: although Guerke (1979) and Paton (1999) suggested they range between 3–7 in all *Jubula* taxa, Schuster (1992) stated that the oil-bodies are numerous in the North American material of *J. pennsylvanica* (\equiv *J. hutchinsiae* subsp. *pennsylvanica*), ranging from 6–16 per cell, and Mizutani (1961) reported 2–10 for Japanese *Jubula*. All authors agree that the oil-bodies in *Jubula* are faintly granular or homogeneous. In *Nipponolejeunea*, on the other hand, the oil-bodies range between 3–5(7) per cell, are hyaline to somewhat grayish, and are formed by 15–20 internal oil-globules (Mizutani 1961). In *Frullania* the oil-bodies are usually larger, finely to coarsely papillose rather than smooth, and few per cell, with their number generally increasing from the leaf-lobe marginal cells to the basal cells, except in the species that have basal ocelli; however, this number rarely reaches the number of oil-bodies seen in *Neohattoria* or *Jubula*. The average number of oil-bodies from the 22 species studied by von Konrat (2004) is 4.3 per median lobe cell. One remarkable exception is the North American species *F. stylifera* (R.M.Schust.) R.M.Schust., which has up to 16 oil-bodies per median cell (von Konrat 2004). A survey of over sixty species (including literature data) suggests that this is a rare condition in the genus (von Konrat 2004). Schuster (1992) described the oil-bodies of *Frullania* as formed of numerous oil-globules and usually appearing coarsely to finely papillose, the only exception being the oil-bodies of *F.* subg. *Microfrullania*, which are smooth and frequently

appear as almost homogeneous oil-droplets (von Konrat 2004). The oil-bodies of *Neohattoria* then appear closer to the other Jubulaceae genera in appearance (although smooth, homogeneous oil-bodies are also seen in *Frullania* subg. *Microfrullania*) and number, notwithstanding the number reported for *Nipponolejeunea* and some reports of *Jubula* taxa with fewer oil-body numbers.

Nomenclatural novelties

Neohattoria Kamim., Journal of Japanese Botany 37: 218. 1962.

≡ *Frullania* subg. *Dentatilobi* Stotler & Crand.-Stotl., Memoirs of The New York Botanical Garden 45: 542. 1987 (“*Dentatiloba*”). **syn.nov.** – Type: *Frullania herzogii* S.Hatt.

Acknowledgements

We thank Anders Hagborg (The Field Museum) and Lars Söderström (Norwegian University of Science and Technology) and the Early Land Plants Today (ELPT) project for access to nomenclatural data. Support from the Biodiversity Synthesis Center of the Encyclopedia of Life provided important funding to help foster international initiatives. The Biodiversity Heritage Library is acknowledged for the facility they provide that has greatly accelerated our effort. The generous support by the National Science Foundation (Awards No. 1145898, 1146168, and 0531730) is gratefully acknowledged. We also recognize the support of the Museum Collection Spending Fund, administered by The Field Museum, as well as curatorial support provided by Yarency Rodriguez, Lucia Kawasaki and Anna Balla (The Field Museum). Lauren Smith is acknowledged for providing the digital images used to compile the plates. Finally, we thank Matt Renner and an anonymous reviewer for their help in improving the manuscript.

References

- Ahonen I (2004) Molecular phylogeny of liverwort order Porellales (Marchantiophyta, Jungermanniopsida). In: Goffinet B, Hollowell VC, Magill R (Eds) Molecular Systematics of Bryophytes. Monographs in Systematic Botany from the Missouri Botanical Garden 98: 169–188.
- Ahonen I (2006) The taxonomic position of the genus *Nipponolejeunea* Hatt. Journal of the Hattori Botanical Laboratory 99: 319–342.
- Ahonen I, Muona J, Piippo S (2003) Inferring the phylogeny of the Lejeuneaceae (Jungermanniopsida): a first appraisal of molecular data. The Bryologist 106: 297–308. doi: 10.1639/0007-2745(2003)106[0297:ITPOTL]2.0.CO;2

- Asakawa Y, Tokunaga N, Toyota M, Takemoto T, Hattori S, Mizutani M, Suire C (1979) Chemosystematics of bryophytes II. The distribution of terpenoids in Hepaticae and Anthocerotae. *Journal of the Hattori Botanical Laboratory* 46: 67–76.
- Asakawa Y, Matsuda R, Toyota M, Hattori S, Ourisson G (1981) Terpenoids and bibenzyls of 25 liverwort *Frullania* species. *Phytochemistry* 20: 2187–2194. doi: 10.1016/0031-9422(81)80111-2
- Asakawa Y, Matsuda R, Toyota M, Takemoto T, Connolly JD, Phillips WR (1983) Sesquiterpenoids from *Chiloscyphus* sp., *Clasmatocolea* sp., and *Frullania* sp. *Phytochemistry* 22: 961–964. doi: 10.1016/0031-9422(83)85030-4
- Asakawa Y, Matsuda R, Cheminat A (1987) Bibenzyl derivatives from *Frullania* species. *Phytochemistry* 26: 1117–1122. doi: 10.1016/S0031-9422(00)82361-4
- Borsch T, Quandt D (2009) Mutational dynamics and phylogenetic utility of noncoding chloroplast DNA. *Plant Systematics and Evolution* 282: 169–199. doi: 10.1007/s00606-009-0210-8
- Casas C, Brugués M, Cros RM, Sérgio C, Infante M (2009) Handbook of liverworts and hornworts of the Iberian Peninsula and the Balearic Islands. Illustrated keys to genera and species. Institut d'Estudis Catalans, Secció de Ciències Biològiques, Barcelona, 177 pp.
- Cooper ED, Shaw AJ, Shaw B, Henwood MJ, Heslewood MM, Brown EA (2011) A multi-locus molecular phylogeny of the Lepidoziaceae: Laying the foundations for a stable classification. *Molecular Phylogenetics and Evolution* 59: 489–509. doi: 10.1016/j.ympev.2011.02.006
- Crandall-Stotler B, Guerke WR (1980) Developmental anatomy of *Jubula* Dum. (Hepaticae). *Bryologist* 83: 179–201. doi: 10.2307/3242131
- Crandall-Stotler B, Stotler RE (2000) Morphology and classification of the Marchantiophyta. In: Shaw AJ, Goffinet B (Eds) *Bryophyte biology*. Cambridge University Press, Cambridge, 21–70. doi: 10.1017/CBO9781139171304.003
- Crandall-Stotler B, Stotler RE, Long DG (2008) Morphology and classification of the Marchantiophyta. In: Goffinet B, Shaw AJ (Eds) *Bryophyte biology*, second edition. Cambridge University Press, Cambridge, 1–54.
- Crandall-Stotler B, Stotler RE, Long DG (2009) Phylogeny and classification of the Marchantiophyta. *Edinburgh Journal of Botany* 66: 155–198. doi: 10.1017/S0960428609005393
- Damsholt K (2002) *Illustrated flora of Nordic liverworts and hornworts*. Nordic Bryological Society, Lund, 837 pp.
- Davis EC (2004) A molecular phylogeny of leafy liverworts (Jungermanniidae: Marchantiophyta). *Monographs in Systematic Botany from the Missouri Botanical Garden* 98: 87–118.
- Engel JJ (1978) A taxonomic and phytogeographic study of Brunswick Peninsula (Strait of Magellan) Hepaticae and Anthocerotae. *Fieldiana Botany* 41: 1–319. doi: 10.5962/bhl.title.2426
- Feldberg K, Schneider H, Stadler T, Schäfer-Verwimp A, Schmidt A, Heinrichs J (2014) Epiphytic leafy liverworts diversified in angiosperm-dominated forests. *Scientific Reports* 4: 5974. doi: 10.1038/srep05974
- Forrest LL, Davis EC, Long DG, Crandall-Stotler BJ, Clark A, Hollingsworth ML (2006) Unraveling the evolutionary history of the liverworts (Marchantiophyta): multiple taxa, genomes

- and analyses. *The Bryologist* 109: 303–334. doi: 10.1639/0007-2745(2006)109[303:UTEHOT]2.0.CO;2
- Frey W, Stech M (2009) Marchantiophyta. In: Frey W (Ed.) *Syllabus of plant families – A. Engler’s Syllabus der Pflanzenfamilien. part 3: Bryophytes and seedless vascular plants. 13th edition, viii, Borntraeger, Berlin/Stuttgart, 13–115.*
- Gradstein SR, Churchill SP, Salazar-Allen N (2001) *Guide to the bryophytes of tropical America. Memoirs of the New York Botanical Garden* 86: 1–577.
- Grolle R (1966) *Miscellanea hepaticologica* (51–60). *Journal of Japanese Botany* 41: 141–147.
- Grolle R, Meister K (2004) *The liverworts in Baltic and Bitterfeld amber. Weissdorn, Jena, 1–91.*
- Guerke WR (1978) *A monograph of the genus Jubula Dumortier. Bryophytorum Bibliotheca* 17: 1–118.
- Hattori S (1955) *A remarkable Frullania species from northern Japan. Feddes Repertorium Specierum Novarum Regni Vegetabilis* 58: 53–54.
- Hattori S (1982) *A synopsis of New Guinean Frullania, Hepaticae. Journal of the Hattori Botanical Laboratory* 51: 203–271.
- Hattori S (1984) *New Caledonian Frullaniaceae. Journal of the Hattori Botanical Laboratory* 57: 405–426.
- Hattori S (1986) *A synopsis of New Caledonian Frullaniaceae. Journal of the Hattori Botanical Laboratory* 60: 203–237.
- Hattori S, Mizutani M (1982) *A status of Amphijubula (Hepaticae) with special reference to the seta anatomy. Journal of the Hattori Botanical Laboratory* 52: 441–448.
- Hattori S, Sharp AJ, Mizutani M (1972) *Schusterella, a new genus of Jubulaceae (Hepaticae). Journal of Japanese Botany* 20: 329–338.
- Heinrichs J, Gradstein SR, Wilson R, Schneider H (2005) *Towards a natural classification of liverworts (Marchantiophyta) based on the chloroplast gene rbcL. Cryptogamie, Bryologie* 26: 215–233.
- Heinrichs J, Hentschel J, Wilson R, Feldberg K, Schneider H (2007) *Evolution of leafy liverworts (Jungermanniidae, Marchantiophyta): estimating divergence times from chloroplast DNA sequences using penalized likelihood with integrated fossil evidence. Taxon* 56: 31–44.
- Hentschel J, von Konrat MJ, Pócs T, Schäfer-Verwimp A, Shaw AJ, Schneider H, Heinrichs J (2009) *Molecular insights into the phylogeny and subgeneric classification of Frullania Raddi (Frullaniaceae, Porellales). Molecular Phylogenetics and Evolution* 52: 142–156. doi: 10.1016/j.ympev.2008.12.021
- Holmgren PK, Holmgren NH, Barnett LC (1990) *Index herbariorum. Part I: The herbaria of the world (Regnum Veg. 120), eighth edition. New York Botanical Garden, New York, 693 pp.*
- Huelsenbeck JP, Ronquist F (2001) *MrBayes: Bayesian inference of phylogeny. Bioinformatics* 17: 754–755. doi: 10.1093/bioinformatics/17.8.754
- Inoue H, Asakawa Y, Gradstein SR (1981) *A biosystematic study of Neohattoria herzogii (Hatt.) Kamim. Bulletin of the National Science Museum, Tokyo, Series B, 7: 23–30.*
- Kamimura M (1961) *A monograph of Japanese Frullaniaceae. Journal of the Hattori Botanical Laboratory* 24: 1–109.

- Kamimura M (1962) On the genus *Neohattoria* Kamim. nom. nov. (Hepaticae). *Journal of Japanese Botany* 37: 26.
- Kelchner SA (2000) The evolution of non-coding chloroplast DNA and its application in plant systematics. *Annals of the Missouri Botanical Garden* 87: 482–498. doi: 10.2307/2666142
- Konstantinova NA, Vilnet AA (2011) *Jubula hutchinsiae* subsp. *caucasica* subsp. nov. (Jubulaceae, Marchantiophyta) – a new taxon from the western Caucasus. *Arctoa* 20: 227–238. doi: 10.15298/arctoa.20.18
- Kraut L, Mues R, Sim-Sim M (1994) Sesquiterpene lactones and 3-benzylphthalides from *Fruilania muscicola*. *Phytochemistry* 37: 1337–1346. doi: 10.1016/S0031-9422(00)90409-6
- Lanfear R, Calcott B, Ho SYW, Guindon S (2012) PartitionFinder: combined selection of partitioning schemes and substitution models for phylogenetic analyses. *Molecular Biology and Evolution* 2: 1695–1701. doi: 10.1093/molbev/mss020
- Lanfear R, Calcott B, Kainer D, Mayer C, Stamatakis A (2014) Selecting optimal partitioning schemes for phylogenomic datasets. *BMC Evolutionary Biology* 14: 82. doi: 10.1186/1471-2148-14-82
- Mizutani M (1961) A revision of Japanese Lejeuneaceae. *Journal of the Hattori Botanical Laboratory* 24: 115–302.
- Müller K (1915) Die Lebermoose Deutschlands, Oesterreichs u. d. Schweiz mit Berücksichtigung der übrigen Länder Europas, II. Abteilung. In: Dr. L. Rabenhorst's Kryptogamen-Flora von Deutschland, Oesterreich und der Schweiz. 2. Auflage, 6. Band, 24. Lieferung. Eduard Kummer, Leipzig, 593–656.
- Olsson S, Buchbender V, Enroth J, Hedenäs L, Huttunen S, Quandt D (2009) Phylogenetic analyses reveal high levels of polyphyly among pleurocarpous lineages as well as novel clades. *The Bryologist* 112: 447–466. doi: 10.1639/0007-2745-112.3.447
- Paton JA (1999) *The liverwort flora of the British Isles*. Harley Books, Colchester, 626 pp.
- Pätsch R, Hentschel J, Linares-Palomino R, Zhu R-L, Heinrichs J (2010) Diversification and taxonomy of the liverwort *Jubula* Dumort. (Jungermanniopsida: Porellales) inferred from nuclear and chloroplast DNA sequences. *Systematic Botany* 35: 6–12. doi: 10.1600/036364410790862515
- Pedersen N, Holyoak DT, Newton AE (2007) Systematics and morphological evolution within the moss family Bryaceae: A comparison between parsimony and Bayesian methods for reconstruction of ancestral character states. *Molecular Phylogenetics and Evolution* 43: 891–907. doi: 10.1016/j.ympev.2006.10.018
- Rambaut A, Drummond AJ (2007) Tracer, version 1.4. <http://beast.bio.ed.ac.uk/Tracer>
- Ronquist F, Huelsenbeck JP (2003) MRBAYES 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19: 1572–1574. doi: 10.1093/bioinformatics/btg180
- Schuster RM (1961) Studies in Lophoziaaceae. I. The genera *Anastrophyllum* and *Sphenolobus* and their segregates. *Revue Bryologique et Lichénologique* 30: 55–73.
- Schuster RM (1963) Studies on antipodal Hepaticae. I. Annotated key to the genera of antipodal Hepaticae with special reference to New Zealand and Tasmania. *Journal of the Hattori Botanical Laboratory* 26: 185–309.
- Schuster RM (1970) Studies on antipodal hepaticae, III. *Jubulopsis* Schuster, *Neohattoria* Kamimura and *Amphijubula* Schuster. *Journal of the Hattori Botanical Laboratory* 33: 266–304.

- Schuster RM (1980) New combinations and taxa of Hepaticae, I. *Phytologia* 45: 415–437.
- Schuster RM (1992) The Hepaticae and Anthocerotae of North America east of the hundredth meridian. Volume V. Field Museum of Natural History, Chicago, 854 pp.
- Schuster RM (1996) On *Jubulopsis* Schust. (Jungermanniales: Jubulopsidaceae fam. nov.) and its relationships. *Journal of Bryology* 19: 297–310.
- Shaw AJ, Cox CJ, Boles SB (2003) Polarity of peatmoss evolution: who says mosses have no roots? *American Journal of Botany* 90: 1777–1787. doi: 10.3732/ajb.90.12.1777
- Stotler RE, Crandall-Stotler B (1987) A re-evaluation of the genus *Neohattoria* (Jubulaceae). *Memoirs of the New York Botanical Garden* 45: 535–543.
- Stöver BC, Müller KF (2010) TreeGraph 2: Combining and visualizing evidence from different phylogenetic analyses. *BMC Bioinformatics* 11: 7. doi: 10.1186/1471-2105-11-7
- Sukumaran J, Holder MT (2010) DendroPy: A Python library for phylogenetic computing. *Bioinformatics* 26: 1569–1571. doi: 10.1093/bioinformatics/btq228
- Verdoorn F (1930) Die Frullaniaceae der Indomalaischen Inseln. (De Frullaniaceis VII). *Annales Bryologici Supplementary Volume* 1: 1–188. doi: 10.1007/978-94-015-5385-8
- von Konrat M (2004) A systematic study of the liverwort genus *Frullania* Raddi: encompassing a worldwide monograph of subg. *Microfrullania* (Schust.) Schust.; a revision of the New Zealand species & study of subsidiary species. PhD dissertation, The University of Auckland, NZ, xxviii + 408 pp.
- von Konrat M, Hentschel J, Heinrichs J, Braggins JE, Pócs T (2010) Forty-one degrees below and sixty years in the dark: *Frullania* sect. *Inconditum*, a new section of Australasian *Frullania* species including *F. colliculosa*, sp. nov., *F. hodgsoniae*, nom. and stat. nov., *F. aterima*, and *F. hattorii* (Frullaniaceae, Marchantiophyta). *Nova Hedwigia* 91: 471–500. doi: 10.1127/0029-5035/2010/0091-0471
- von Konrat M, de Lange P, Greif M, Strozier L, Hentschel J, Heinrichs J (2012) *Frullania knightbridgei*, a new liverwort (Frullaniaceae, Marchantiophyta) species from the deep south of Aotearoa-New Zealand based on an integrated evidence-based approach. *PhytoKeys* 8: 13–36. doi: 10.3897/phytokeys.8.2496
- Wilson R, Gradstein SR, Heinrichs J, Groth H, Ilkiu-Borges AL, Hartmann FA (2004) Phylogeny of Lejeuneaceae: a cladistic analysis of chloroplast *rbcl* sequences and morphology with preliminary comments on the mitochondrial *nad4–2* spacer region. In: Goffinet B, Hollowell VC, Magill RE (Eds) *Molecular Systematics of Bryophytes*. Monographs in Systematic Botany from the Missouri Botanical Garden 98: 189–202.
- Wilson R, Gradstein SR, Schneider H, Heinrichs J (2007) Unravelling the phylogeny of Lejeuneaceae (Jungermanniopsida): Evidence for four main lineages. *Molecular Phylogenetics and Evolution* 43: 270–282. doi: 10.1016/j.ympev.2006.10.017
- Zwickl DJ (2006) GARLI (Genetic Algorithm for Rapid Likelihood Inference). Version 2.01. <http://www.bio.utexas.edu/faculty/antisense/garli/garli.html>

Appendix I

Voucher information for data set 1. Information is presented in the following order: taxon name, collector followed by collection number (herbarium acronyms follow Holmgren et al. 1990), country: region (if known), GenBank accession numbers (*psbA/rbcL/rps3/nad1*). Lacking sequences are indicated by a dash (—). New sequences generated for this study are marked by an asterisk (*).

Acanthocoleus madagascariensis (Steph.) Kruijt, Pócs 97145/AA (GOET), Uganda, EF011843/DQ983649/—/—; *Blepharolejeunea chimantaensis* van Slageren & Kruijt, Pócs & Rico 00234/A (F), Venezuela, KF851876/—/—/KF852465; *Bryopteris filicina* (Sw.) Nees, Churchill, Magombo & Price 19855 (NY), Bolivia, AY607930/DQ439681/KF851576/KF852481; *Caudalejeunea reniloba* (Gottsche) Steph., Pócs et al. 01090/AB (F), Australia, KF851845/KF852294/KF851541/KF852441; *Ceratolejeunea coarina* (Gottsche) Schiffl., Zartman 1235.1 (DUKE), Brazil, AY607934/AY608026/—/KF852489; *Cololejeunea microscopica* (Taylor) Schiffl., Long & Rothenro 37789 (E), Scotland: Wester Ross, KF851954/KF852386/KF851651/KF852552; *Colura conica* (Sande Lac.) K.I.Goebel, Pócs & Streimann 9986/W (F), Australia: Queensland, KM817490*/KM817513*/KM817536*/KM817462*; *Colura imperfecta* Steph., Pócs & Pócs 07019/A (F), Thailand, KF851881/KF852327/—/KF852469; *Drepanolejeunea erecta* (Steph.) Mizut., Long 28691 (E), Bhutan, JF513393/JF513452/KF851515/JF513342; *Frullania albertii* Steph., Davis 295 (DUKE), Ecuador, AY607942/DQ439685/KM817549*/KM817477*; *Frullania atrata* (Sw.) Nees ex Mont., Dauphin 3306 (F), Costa Rica, KM817491*/—/KM817540*/KM817466*; *Frullania caulisequa* (Nees) Mont., Karst, Shaw & Gibbs 022 (DUKE), USA: North Carolina, KM817500*/KM817526*/KM817553*/KM817481*; *Frullania dilatata* (L.) Dumort., Stotler 4666 (SIU), Portugal, KM817502*/KM817528*/KM817555*/KM817482*; *Frullania eboracensis* Lehm., Stotler 80-4354 (ABSH), USA: Illinois, AY688827/AY688779/KM817547*/KM817475*; *Frullania ecklonii* (Spreng.) Spreng. ex Gottsche, Lindenb. & Nees, Pócs 02030/W (F), Kenya, KM817488*/KM817510*/KM817533*/KM817459*; *Frullania ericoides* (Nees) Mont., Long 35167 (E), China: Yunnan, KM817486*/KM817507*/KM817531*/KM817456*; *Frullania falciloba* Taylor ex Lehm., Engel, von Konrat & Braggins 26837 (F), New Zealand, KM817489*/KM817511*/KM817534*/KM817460*; *Frullania moniliata* (Reinw., Blume & Nees) Mont., Mizutani s.n. (ABSH), Japan, AY507484/AY507401/KM817548*/KM817476*; *Frullania nodulosa* (Reinw., Blume & Nees) Nees, Pócs & Pócs 03261/A (F) Fiji, KM817492*/KM817517*/KM817541*/KM817467*; *Frullania parhamii* (R.M.Schust.) R.M.Schust. ex von Konrat, L.Söderstr. & A.Hagborg, von Konrat, Braggins & Naikatini 6/16-5 (F), Fiji, —/KM817516*/KM817539*/KM817465*; *Frullania pycnantha* (Hook.f. & Taylor) Taylor ex Gottsche, Lindenb. & Nees, von Konrat 99/409 (F), New Zealand, KM817499*/KM817525*/—/KM817480*; *Frullania rostrata* (Hook.f. & Taylor) Hook.f. & Taylor ex Gottsche, Lindenb. & Nees, Engel, von Konrat & Braggins 27770 (F), New Zealand, —/KM817512*/KM817535*/KM817461*; *Frullania tamarisci* (L.) Dumort. 1, Stot-

ler 4661 (SIU), Portugal: Sintra, KM817501*/KM817527*/KM817554*/—; *Frullania tamarisci* 2, Long 35371 (E), France, KM817487*/KM817508*/KM817532*/KM817457*; *Frullanooides densifolia* Raddi, Gradstein 10171 (GOET), Ecuador, KF851930/KF852371/KF851634/KF852530; *Fulfordianthus pterobryoides* (Spruce) Gradst., Gradstein & Varon 11069 (GOET), Colombia, KF851931/KF852372/KF851635/KF852531; *Jubula hutchinsiae* (Hook.) Dumort. 1, Long 29077 (E), UK: England, —/KM817509*/—/KM817458*; *Jubula hutchinsiae* 2, Drehwald 3007 (GOET), Portugal, EF011746/AY548101/—/—; *Jubula hutchinsiae* subsp. *bogotensis* (Steph.) Verd., Gradstein s.n. (GOET), Mexico, EF011758/AY548100/—/—; *Jubula hutchinsiae* subsp. *javanica* (Steph.) Verd. 1, Konstantinova & Savchenko K479/1-07 (F), Russia, —/KM817506*/KM817542*/KM817468*; *Jubula hutchinsiae* subsp. *javanica* 2, Kodama s.n. (ABSH), Japan: Wakayama Pref., AY507492/AY507408/KF851585/JF513366; *Jubula hutchinsiae* subsp. *pennsylvanica* (Steph.) Verd., Risk 11005 (DUKE), USA, AY607954/KM817523*/KM817550*/—; *Lejeunea deplanata* Nees var. *deplanata*, Shaw F533 (DUKE), USA: North Carolina, KM817498*/KM817524*/KM817552*/KM817479*; *Lejeunea japonica* Mitt., Bakalin s.n. (F), Russia, —/KM817518*/KM817543*/KM817469*; *Lejeunea tuberculosa* Steph., Long 28596 (E), Bhutan, JF513394/JF513453/KF851518/JF513344; *Leptolejeunea elliptica* (Lehm.) Besch., Yamaguchi s.n. (F), Japan, KM817485*/KM817515*/KM817538*/KM817464*; *Lopholejeunea eulopha* (Taylor) Schiffn., Pócs et al. 08036/U (F), Fiji, KF851868/KF852314/—/—; *Marchesinia mackaii* (Hook.) Gray, Buryova 2181 (DUKE), UK: Wales, —/KF852356/KF851619/KF852515; *Mastigolejeunea auriculata* (Wilson) Steph., Shaw 6222 (DUKE), USA: Alabama, KF851917/KF852359/KF851622/KF852518; *Neohattoria herzogii* (S.Hatt.) Kamim., Furuki 22673 (F), Japan: Honshu, KM817504*/KM817530*/KM817557*/KM817484*; *Nipponolejeunea pilifera* (Steph.) S.Hatt., Ohnishi 5975 (HIRO), Japan, AM396291/AM392293/—/—; *Nipponolejeunea subalpina* (Horik.) S.Hatt., Ohnishi 5611 (GOET), Japan, AM396290/AM392292/—/—; *Odontolejeunea lunulata* (F.Weber) Schiffn., Picon et al. 00227/CE (F), Venezuela, —/KM817514*/KM817537*/KM817463*; *Ptychanthus striatus* (Lehm.) Nees, Pócs & Pócs 03288/O (F), Fiji, KF851872/KF852318/KF851558/KF852460; *Radula buccinifera* (Hook.f. & Taylor) Taylor ex Gottsche, Lindenb. & Nees, Engel, von Konrat & Braggins 23569 (F), New Zealand, KM817495*/KM817521*/KM817545*/KM817472*; *Radula cuspidata* Steph., Engel & von Konrat 23517 (F), New Zealand, KM817496*/—/KM817546*/KM817473*; *Radula grandis* Steph., Engel, von Konrat & Braggins 24847 (F), New Zealand, KM817494*/KM817520*/KM817544*/KM817471*; *Radula lindenbergiana* Gottsche ex C.Hartm., Stotler 4656 (SIU), Portugal, KM817503*/KM817529*/KM817556*/KM817483*; *Radula perrottetii* Gottsche ex Steph., Mizutani 15030 (F), Japan, —/DQ439700/KM817551*/KM817478*; *Radula ratkowskiana* K.Yamada, Engel, von Konrat & Braggins 24365 (F), New Zealand, KM817497*/KM817522*/—/KM817474*; *Radula tasmanica* Steph., Engel, von Konrat & Braggins 24874 (F), New Zealand, KM817493*/KM817519*/—/KM817470*; *Spruceanthus thozetianus* (Gottsche & F.Muell.) B.M.Thiers & Gradst., Pócs 01107/M (GOET), Australia,

AM396273/AM384877/—/—; *Stictolejeunea squamata* (Willd. ex F. Weber) Schiffn., Dauphin & Gonzalez 2134 (GOET), Costa Rica: Alajeula, KF851951/—/—/ KF852549; *Thysananthus convolutus* Lindenb., Gradstein 10205 (GOET), Indonesia: Java, KF851953/DQ983737/KF851650/KF852551.

Appendix 2

Voucher information for data set 2. Information is presented in the following order: taxon name, collector followed by collection number (herbarium acronyms follow Holmgren et al. 1990), country: region (if known), GenBank accession numbers (ITS region/rbcL/trnL-F/psbA). Lacking sequences are indicated by a dash (—). New sequences generated for this study are marked by an asterisk (*).

Frullania kunzei (Lehm.) Lehm. & Lindenb., Costa & Gradstein 3769 (GOET), Brazil, FJ380536/FJ380863/FJ380387/FJ380697; *Frullania nisquallensis* Sull., Doyle 11001 (GOET), USA, FJ380503/FJ380826/FJ380349/FJ380661; *Frullania peruviana* Gottsche, Schaefer-Verwimp & al. 24356 (GOET), Ecuador, FJ380543/FJ380870/FJ380394/FJ380704; *Frullanooides mexicana* van Slageren, Burghardt 4421a, Mexico, DQ987366/DQ983682/DQ987464/EF011851; *Fulfordianthus pterobryoides* (Spruce) Gradst., Dauphin 2518, Costa Rica, AM237145/DQ983684/AM237198/EF011832; *Jubula hutchinsiae* (Hook.) Dumort., Ahonen, Huttunen et Virtanen 3190 (H), Taiwan, AY125350/AY125946/AY144477/—; *Jubula hutchinsiae* subsp. *bogotensis* (Steph.) Verd. 1, Gradstein s.n. (GOET), Mexico: Veracruz, FN396818/—/FN398013/—; *Jubula hutchinsiae* subsp. *bogotensis* 2, Gradstein s.n. (GOET?), Mexico, DQ987273/AY548100/DQ987388/AM396281; *Jubula hutchinsiae* subsp. *bogotensis* 3, Gradstein 9449 (GOET), Costa Rica, FN396817/—/FN398012/—; *Jubula hutchinsiae* subsp. *bogotensis* 4, Frahm et al. 1313 (GOET), Peru, FN396816/—/—/—; *Jubula hutchinsiae* subsp. *caucasica* Konstant. & Vilnet 1, Konstantinova K456-5-07 (KPABG), Russia: Caucasus, JN836964/—/JN836974/—; *Jubula hutchinsiae* subsp. *caucasica* 2, Konstantinova K429-3-08 (KPABG), Russia: Caucasus, JN836961/—/JN836971/—; *Jubula hutchinsiae* subsp. *caucasica* 3, Konstantinova K462-1-08 (KPABG), Russia: Caucasus, JN836960/—/JN836970/—; *Jubula hutchinsiae* subsp. *caucasica* 4, Konstantinova K463-1-07 (KPABG), Russia: Caucasus, JN836962/—/JN836972/—; *Jubula hutchinsiae* subsp. *caucasica* 5, Konstantinova K371-1-08 (KPABG), Russia: Caucasus, JN836958/—/JN836968/—; *Jubula hutchinsiae* subsp. *caucasica* 6, Konstantinova K446-7-08 (KPABG), Russia: Caucasus, JN836959/—/JN836969/—; *Jubula hutchinsiae* subsp. *caucasica* 7, Konstantinova K443-14-08 (KPABG), Russia: Caucasus, JN836963/—/JN836973/—; *Jubula hutchinsiae* subsp. *hutchinsiae* 1, Long 29077 (GOET), UK: Devon, FN396813/—/FN398010/—; *Jubula hutchinsiae* subsp. *hutchinsiae* 2, Long 35296 (GOET), UK: Wales, FN396814/—/FN398011/—; *Jubula hutchinsiae* subsp. *hutchinsiae* 3, Schaefer-Verwimp & Verwimp 25675 (GOET), Portugal: Madeira, FN396811/—/FN397099/—; *Jubula hutchinsiae* subsp. *hutchinsiae* 4, Schaefer-

Verwimp & Verwimp 25796 (GOET), Portugal: Boaventura, FN396812/—/FN398009/—; *Jubula hutchinsiae* subsp. *hutchinsiae* 5, Drehwald & Reiner-Drehwald 3007 (GOET), Portugal, DQ987260/AY548101/DQ987380/AM396282; *Jubula hutchinsiae* subsp. *japonica* (Steph.) Horik. & Ando 1, Koponen et al. 54308 (H), China, AY125342/AY125938/AY144479/—; *Jubula hutchinsiae* subsp. *japonica* 2, Inoue BSE755 (GOET), Japan: Kochi, FN396809/—/—/—; *Jubula hutchinsiae* subsp. *japonica* 3, Gradstein & Mizutani 2958 (GOET), Japan: Miyazaki, FN396810/—/FN397098/—; *Jubula hutchinsiae* subsp. *japonica* 4, Bakalin P-68-10-08 (KPABG), Russia: Primorsky Kray, JN836967/—/JN836977/—; *Jubula hutchinsiae* subsp. *javanica* (Steph.) Verd. 1, Zhu et al. 3361 (HSNU), China: Hainan, FN396800/—/—/—; *Jubula hutchinsiae* subsp. *javanica* 2, Zhu et al. 20050903-7a (HSNU), China: Hainan, FN396801/—/—/—; *Jubula hutchinsiae* subsp. *javanica* 3, Long 34765 (GOET), China: Yunnan, FN396805/—/FN397095/—; *Jubula hutchinsiae* subsp. *javanica* 4, Pocs 98105/C (GOET), Viet Nam: Vin-Phuc, FN396807/—/—/—; *Jubula hutchinsiae* subsp. *javanica* 5, Pocs & Tran Ninh 98103/A2 (GOET), Viet Nam: Vin-Phuc, FN396808/—/FN397097/—; *Jubula hutchinsiae* subsp. *javanica* 6, Schaefer-Verwimp & Verwimp 18870/A (GOET), Malaysia: Pahang, FN396802/—/FN397094/—; *Jubula hutchinsiae* subsp. *javanica* 7, Zhu 555 (HSNU), China: Fujian, FN396806/—/FN397096/—; *Jubula hutchinsiae* subsp. *javanica* 8, Bakalin Kor-12-6-08 (KPABG), South Korea, JN836966/—/JN836976/—; *Jubula hutchinsiae* subsp. *javanica* 9, Schaefer-Verwimp & Verwimp 18935 (GOET), Malaysia: Pahang, FN396803/—/—/—; *Jubula hutchinsiae* subsp. *javanica* 10, Wang 685B (HSNU), China: Yunnan, FN396804/—/—/—; *Jubula hutchinsiae* subsp. *pennsylvanica* (Steph.) Verd. 1, Buck 39060 (H?), USA: West Virginia, AY776308/AY776303/AY776309/—; *Jubula hutchinsiae* subsp. *pennsylvanica* 2, Davison 5045 (UNAF), USA: Alabama, FN396819/—/—/—; *Jubula hutchinsiae* subsp. *pennsylvanica* 3, Davison 5201 (UNAF), USA: West Virginia, FN396821/—/FN398015/—; *Jubula hutchinsiae* subsp. *pennsylvanica* 4, Davison 4707 (UNAF), USA: Alabama, FN396822/—/FN398016/—; *Jubula hutchinsiae* subsp. *pennsylvanica* 5, Davison 3775a (UNAF), USA: Alabama, FN396823/—/FN398017/—; *Jubula hutchinsiae* subsp. *pennsylvanica* 6, Davison & Risk 2537 (UNAF), USA: Kentucky, FN396820/—/FN398014/—; *Jubula hutchinsiae* subsp. *pennsylvanica* 7, Konstantinova ACH-3-92 (KPABG), USA, JN836965/—/JN836975/—; *Jubula hutchinsiae* subsp. *pennsylvanica* 8, Davison 4690 (UNAF), USA, Alabama, FN396824/—/FN398018/—; *Jubula hutchinsiae* subsp. *pennsylvanica* 9, Hyatt 8212 (UNAF), USA: North Carolina, FN396825/—/FN398019/—; *Jubula hutchinsiae* subsp. *pennsylvanica* 10, Davison s.n. (UNAF), USA: North Carolina, FN396826/—/FN398020/—; *Neohattoria herzogii* (S.Hatt.) Kamim., Furuki 22673 (F), Japan: Honshu, KM817455*/KM817530*/KM817505*/KM817504*; *Nipponolejeunea pilifera* (Steph.) S.Hatt. 1, Ohnishi 5975 (HIRO), Japan, —/AM392293/FJ380228/AM396291; *Nipponolejeunea pilifera* 2, Higuchi 41359 (H?), Japan, AY776307/AY776304/AY776310/—; *Nipponolejeunea pilifera* 3, Masuzaki 510 (HIRO), Japan: Yakushima Is., —/AB476588/—/—; *Nipponolejeunea pilifera* 4, Ohnishi s.n. (H), Japan, AY125341/AY125937/AY144478/—; *Nippon-*

olejeunea subalpina (Horik.) S.Hatt. 1, Ohnishi 5611 (HIRO), Japan, DQ987289/AM392292/FJ380227/AM396290; *Nipponolejeunea subalpina* 2, Higuchi 41358 (H?), Japan, AY776306/AY776305/AY776311/—; *Ptychanthus striatus* (Lehm.) Nees, Gradstein 10217, Indonesia: Java, DQ987297/DQ983723/DQ987403/EF011777; *Schiffneriolejeunea nymannii* (Steph.) Gradst. & Terken, Gradstein et al. 10321, Malaysia, DQ987320/DQ983725/DQ987424/EF011801.