

# The taxonomic identity of *Didymostigma trichanthera* (Gesneriaceae)

Lihua Yang<sup>1</sup>, Chen Feng<sup>1</sup>, Ming Kang<sup>1</sup>, Fang Wen<sup>2,3</sup>

**1** Key Laboratory of Plant Resources Conservation and Sustainable Utilization, South China Botanical Garden, Chinese Academy of Sciences, Guangzhou 510650, China **2** Guangxi Key Laboratory of Plant Conservation and Restoration Ecology in Karst Terrain, Guangxi Institute of Botany, Guangxi Zhuang Autonomous Region and Chinese Academy of Sciences, Guilin 541006, China **3** Gesneriad Conservation Center of China, Guangxi Zhuang Autonomous Region, Guilin 541006, China

Corresponding author: Fang Wen ([wenfang760608@139.com](mailto:wenfang760608@139.com))

---

Academic editor: Y.-G. Wei | Received 20 December 2018 | Accepted 27 February 2019 | Published 26 August 2020

**Citation:** Yang L, Feng C, Kang M, Wen F (2020) The taxonomic identity of *Didymostigma trichanthera* (Gesneriaceae). In: In: Shui Y-M, Chen W-H, Ren M-X, Wen F, Hong X, Qiu Z-J, Wei Y-G, Kang M (Eds) Taxonomy of Gesneriaceae in China and Vietnam. *PhytoKeys* 157: 191–197. <https://doi.org/10.3897/phytokeys.157.32577>

---

## Abstract

Based on consulting original literature, the examination of specimens, and field investigations, *Didymostigma trichanthera* is shown to be conspecific with *Henckelia anachoreta*. Therefore, *Didymostigma trichanthera* is formally treated as a synonym of *Henckelia anachoreta* here.

## Keywords

China, Guangdong, *Didymostigma trichanthera*, *Henckelia anachoreta*, taxonomy

## Introduction

The genus *Didymostigma* W.T. Wang (1984) was once considered to be a monotypic genus, having only one species, *D. obtusum* (C.B. Clarke) W.T. Wang (1984). Subsequently, two new taxa, *D. leiophyllum* D. Fang & X.H. Lu (Fang et al. 1994) and *D. trichanthera* C.X. Ye & X.G. Shi (2005) were discovered and described. The type species of this genus, *D. obtusum*, is widely distributed from eastern Guangdong to southern Fujian, China (Wang et al. 1998; Li and Wang 2004; Wei et al. 2010). The

other two species, however, are regarded as narrowly endemic species and have only been found at their type localities (Fang et al. 1994; Ye and Shi 2005).


*Didymostigma trichanthera* was simply described based on a single collection (*Chuang-Xing Ye 5960*) from Nankunshan, Guangdong Province, China. Ye and Shi (2005) thought that *D. trichanthera* mainly differs from *D. obtusum* by its lanose fertile anthers, hairy filaments, and the unhidden pistil to the corolla tube. The species status of *D. trichanthera* was once doubted by Wei et al. (2010). They find the calyx of *D. trichanthera* is completely different from the type species *D. obtusum*, but they did not carry out further study on this questionable species. After consulting original literature about this genus, type species, and this so-called new species (Wang 1984; Ye and Shi 2005), carrying out field investigations in Nankunshan, and conducting examinations of type materials of *D. trichanthera*, we are convinced that *D. trichanthera* is not a species belonging to the genus *Didymostigma*. Our detailed morphological comparisons find that this so-called *Didymostigma* species is actually conspecific with *Henckelia anachoreta* (H.F. Hance) D.J. Middleton & Mich. Möller (Weber et al. 2011). Consequently, it is essential to reduce *Didymostigma trichanthera* to a synonym of *Henckelia anachoreta*.

## Material and methods

We performed detailed comparisons of type materials between *Didymostigma trichanthera* and the type species of *Didymostigma* (*D. obtusum*), and also between *D. trichanthera* and *Henckelia anachoreta*. The study of specimens was conducted in IBK, IBSC and SYS. We also checked high-resolution digital images of the specimens in A (<https://huh.harvard.edu/>), BM (<http://data.nhm.ac.uk/>), E (<http://www.rbge.org.uk/>), K (<https://www.kew.org/>), P (<https://science.mnhn.fr/institution/mnhn/search>), TI (<http://umdb.um.u-tokyo.ac.jp/Dshokubu/Tshokubu.htm>) and WU (<http://herbarium.univie.ac.at/index.htm>) by their web service, as well as via online databases, such as the Chinese Virtual Herbarium (<http://www.cvh.ac.cn/>), JSTOR Global Plants (<http://plants.jstor.org/>) and Specimens Database of Native Plants in Taiwan (<http://www.hast.biodiv.tw/Specimens/specimenQueryC.aspx>). Moreover, detailed morphological studies of both *H. anachoreta* and *Didymostigma obtusum* were undertaken based on plants from natural populations at Nankunshan. Some other field observations of these two species were carried out in Guangxi, Guangdong, Fujian of China.

## Results and discussion

We did not find any other *Didymostigma* species in Nankunshan in our field work, except for *D. obtusum* (Fig. 1B). However, *Henckelia anachoreta*, a species which is similar to *Didymostigma obtusum* in its vegetative characteristics to some extent, can



**Figure 1.** Living plants of *Henckelia anachoreta* (**A, C, D**) and *Didymostigma obtusum* (**B, E**) at Nankunshan. **A, B** habitat **C** side view of flower **D, E** opened corolla showing pistil and stamens. Scale bar: 1 cm. Arrow indicates calyx and stigma.

be easily found here (Fig. 1A). In fact, Nankunshan has been intensively botanized in recent years, because of its high biodiversity (Chen et al. 2017; Xu et al. 2017), and these studies also did not find *D. trichanthera*. As we know, the natural environment of Nankunshan has been well protected since it was listed as a national forestry park in 1993. Nevertheless, *D. trichanthera* has not been found or collected again since it was described in 2005. It is difficult to believe that *D. trichanthera*, if indeed a well characterized species, is represented only by its type collection.

Unfortunately, the holotype of *D. trichanthera* cannot be found in SYS. According to the isotype (Fig. 2B) preserved in A and the description made by Ye and Shi (2005), we can clearly find that the calyx of *D. trichanthera* is 5-lobed to near middle, with an obvious calyx tube, and the lobes are triangular (Fig. 2B). In contrast, the representative calyx of *Didymostigma* is 5-parted to near base, and the



**Figure 2.** **A** Holotype of *Henckelia anachoreta* (E.H. Hance 11377, BM-000041739!) and **B** isotype of *Didymostigma trichanthera* (Chuang-Xing Ye 5960, A-00135544!). Arrow indicates calyx.

lobes are lanceolate-linear (Fig. 1E; Wang 1984). Additionally, the typical stigma of *Didymostigma* is only ca. 1 mm long (Fig. 1E; Wang 1984). Therefore, although the detailed characteristics of the stigma of *D. trichanthera* have not been clearly observed in the isotype, the description (3–4 mm long) of it in the protologue (Ye and Shi 2005) indicates that it is not a typical character of *Didymostigma*. However, it seems that these two important characters of *D. trichanthera* are well matched with *Henckelia* Spreng. Our detailed comparisons find that there is no obvious difference between the isotype of *Didymostigma trichanthera* (Fig. 2B) and the holotype of *Henckelia anachoreta* (Fig. 2A), and between the images of *Didymostigma trichanthera* given by Ye and Shi (2005) and *Henckelia anachoreta* photo by us at Nankunshan (Fig. 1A, C). In fact, most of the descriptions of *Didymostigma trichanthera* in the protologue, especially the diagnosis characters (lanose fertile anthers, hairy filaments and the unhidden pistil to the corolla tube), are exactly matched with *Henckelia anachoreta*.

All the reasons mentioned above have prompted us to carefully consider that *Didymostigma trichanthera* is conspecific with *Henckelia anachoreta*. As a result, the taxonomic treatment of *Didymostigma trichanthera* needs to be made here.

## Taxonomic treatment

***Henckelia anachoreta* (H.F. Hance) D.J. Middleton & Mich. Möller in Weber et al. (2011: 774)**

- ≡ *Chirita anachoreta* H.F. Hance (1866: 231). *Roettlera anachoreta* (H.F. Hance) O. Kuntze (1891: 476). *Didymocarpus anachoretus* (H.F. Hance) H. Lév. (1906: 427).  
 = *Chirita minutiserrulata* B. Hayata (1915: 133). *Didymocarpus minutiserrulatus* (B. Hayata) Y. Yamamoto (1936: 72). Type: China. Taiwan: Boho, July 1911, *Inaba s.n.* (TI, not seen).  
 = *Didymostigma trichanthera* C.X. Ye & X.G. Shi (2005: 447), syn. nov. Type: China. Guangdong: Longmen County, Nankunshan National Forest Park, 17 August 2003, *Chuang-Xing Ye 5960* (Holotype SYS; Isotype A-00135544!).

**Type.** China. Guangdong: Qingyuan City, North River, 27 July 1864, *F.H. Hance 11377* (Holotype BM-000041739!; Isotype K-000858355!).

**Distribution and habitat.** *Henckelia anachoreta* is a common species with a wide distribution in China (Guangdong, Guangxi, Hunan, Taiwan, Xizang, Yunnan), India (Sikkim), Laos, Myanmar, northern Thailand and northern Vietnam (Weber et al. 2011). Plants often grow on moist rocks or ground surfaces in forest or near valley stream sides.

**Additional specimens examined. China.** Guangdong: Haifeng County, 15 August 1935, *W.T. Tsang 25492* (IBSC); Longmen County, 27 October 1981, *G.C. Zhang 280* (HGAS); Maomin County, 2 August 1956, *L. Deng 1751* (HITBC); Qingyuan County, 15 September 1936, *K.Z. Hou 74155* (IBK); Qujiang County, 13 August 1956, *Z. Huang 41855* (IBSC); Ruyuan County, 8 July 2014, *J.M. Li 7840, 7763* (HEAC); Wengyuan County, 16 August 1933, *X.Q. Liu 2053* (IBSC); Yangshan County, 5 July 1956, *L. Deng 1691* (IBSC). Guangxi: Fangchenggang County, 14 July 1908, *Anonymous s.n.* (PE); *ibid.* 7 July 1912, *K.K. Chung* (IBSC); *ibid.* 7 August 1933, *J.L. Zuo 23588* (IBSC); *ibid.* 25 August 1936, *W.T. Tsang 26748* (IBSC); *ibid.* 10 September 1936, *W.T. Tsang 26826* (IBSC); *ibid.* 9 July 2010, *Shiwandashan team 2619, 2656* (IBK); Gongcheng County, 14 August 1957, *Gongcheng team 195* (IBK); Hengxian County, 15 October 2007, *Y.Q. Su 15915* (GXMG); *ibid.* 7 September 2008, *Ching-I Peng 21784* (HAST); Jinxiu County, 8 September 1981, *Dayaoshan team 10146* (IBSC); *ibid.* 19 September 1981, *Dayaoshan team 10317* (IBSC); *ibid.* 12 September 1981, *Dayaoshan team 10488* (IBK); *ibid.* 1 November 1981, *Dayaoshan team 12255* (IBK); Jingxi County, 17 September 2010, *Y.S. Huang & L. Wu LYJX0509* (IBK); Lingle County, 15 August 1928, *R.C. Ching 6928* (IBSC); Luocheng County, 15 July 1931, *S.S. Sin 22411* (IBSC); Ningming County, 19 August 2010, *W.B. Xu & W.H. Wu NM396* (IBK); Pingxiang County, 27 August 1986, *Beijing team 0973* (PE); Shanglin County, 6 August 1973, *Y. Wang et al. 67046* (PE); *ibid.* 19 October 2011, *L. Wu & J.C. Yang D3372* (IBK); Wuming County, 5 August 2010,

*L. Wu* & *R.H. Jiang* D0235 (IBK); Xing'an County, 24 September 2014, *Xing'an team* 450325140924027LY (GXMG); Yongfu County, 21 July 1956, *H.F. Qin* 700342 (IBK); Zhaoping County, 11 August 1957, *C.Z. Jiang* & *M.S. Xia* 4069 (IBK). Hunan: Guidong County, 19 September 1977, *B.G. Li* 5533 (IBSC); Jiangyong County, 8 July 1959, *P.X. Tan* 62211 (IBK); 12 July 1959, *P.X. Tan* 63671 (FJSI). Taiwan: Kaohsiung hsien, 19 September 1991, *C.C. Wang* 588 (HAST); *ibid.* 13 September 1997, *W.L. Chiou* and *K.C. Yang* s.n. (WU); *ibid.* 16 September 2000, *C.I. Peng* 18073 (HAST); *ibid.* 12 August 2008, *C.I. Huang* 3463 (HAST); *ibid.* 6 November 1991, *C.I. Peng* 14739, 14770 (HAST); Pingtung hsien, 20 September 1990, *W.P. Leu* 551 (HAST); *ibid.* 23 August 2006, *C.I. Huang* 2831 (HAST); *ibid.* 8 October 2011, *P.F. Lu* 22985 (HAST). Xizang: Jilong County, 14 September 2008, *L.M. Gao et al.* GLM-081579 (KUN). Yunnan: Cangyuan County, 27 August 2013, *J.M. Li* 9485 (HEAC); Hekou County, 27 November 1992, *Y.Z. Wang* 92065 (PE); *ibid.* 18 August 1993, *Y.M. Shui* 003411 (PE); *ibid.* 2 October 2003, *J.M. Li* 1022 (PE); *ibid.* 9 October 2011, *M.T. Liu* LMT2011025 (PE); *ibid.* 17 August 2013, *Z.J. Qiu et al.* QZJ-0957 (PE); Jinping County, 12 August 1951, *P.Y. Mao* 314 (PE); *ibid.* 21 September 2006, *L.M. Gao* GLM-06283, GLM-06287 (KUN); *ibid.* 8 September 2012, *Jinping team* 5325300650 (IMDY); Luchun County, 30 September 1973, *D.D. Tao* 635 (KUN); *ibid.* 18 October 2000, *Y.M. Shui* & *W.H. Chen* 13123 (KUN); *ibid.* 23 October 2000, *Y.M. Shui* & *W.H. Chen* 13747 (KUN); *ibid.* 25 October 2000, *Y.M. Shui* & *W.H. Chen* 13905 (KUN); Maguan County, 19 August 2013, *Z.J. Qiu et al.* QZJ-0962 (PE); *ibid.* 18 September 2013, *P.W. Li* LPW2013144, LPW2013143 (PE); Malipo County, 10 August 2004, *J.M. Li* LJM-2004-54 (PE); *ibid.* 28 August 2012, *P.W. Li* LPW2012016 (PE); Menghai County, 24 August 2011, *J.M. Li* 82412 (HEAC); Menglian County, 6 August 1973, *Menglian team* 9967 (KUN); *ibid.* 14 August 1973, *Menglian Team* 10172 (KUN); Mengla County, 23 October 1959, *X.W. Li* 13540 (KUN); Pingbian County, 9 July 1934, *H.T. Tsai* 62481 (PE); *ibid.* 18 September 1939, *Q.W. Wang* 81896 (KUN); *ibid.* 20 September 1939, *Q.W. Wang* 81981 (KUN, PE); *ibid.* 28 September 1954, *K.M. Feng* 4697 (PE); *ibid.* 18 September 2012, *Pingbian team* (IMDY); Wenshan County, 14 August 1947 *K.M. Feng* 11242 (PE, IBSC); *ibid.* 20 August 1947, *K.M. Feng* 11376 (PE); Xichou County, 29 August 1947, *K.M. Feng* 11450 (PE, IBSC); Yanshan County, 19 October 1939, *Q.W. Wang* 84483 (PE).

**Vietnam.** Hà Tây: Mont-Bavi, 22 July 1886, *Anonymous* s.n. (P); *ibid.* 4 September 1886, *Anonymous* s.n. (P). Ha Giang: Vi Xuyen Dist., 7 September 2000, *Harder, D.K., Hieu, N.Q., Du, N.V.* 5302 (E). Thanh Hoa: Ba Thuoc Dist., 9 October 2003, *Averyanov, L.; Loc, P.K.; Doan, D.T.; Vinh, N.T.* HAL4197 (E). Tonkin: Sai Wong Mo Shan, 18 July-9 September 1940, *W.T. Tsang* 30389 (E). **Laos.** Khammouan: Kaeng Meaung landing on Nakai Nam Theun, 21 October 2005, *Newman, M F; Thomas, P I; Armstrong, K E; Sengdala, Khamphone* & *Lamxay, Vichith* LAO 385 (E). **Myanmar.** Haungry: 15 August 1919, *Kingdon-Ward, F.* 3536 (E). **Thailand.** Nakhon Ratchasima: Khao Yai Nat. Park, 22 October 1969, *C.F. van Beusekom, C. Charoenpol* 1833 (P); Siam: September 1910, *Q.J.G. Kew* 1417 (P).

## Acknowledgments

This work was supported by grants from the NSFC-Guangdong Natural Science Foundation Joint Project (U1501211), Science Research Foundation of Guangxi Academy of Sciences (no. 2017YJJ23022), Guangxi Natural Science Foundation Program (2015GXNSFB139004 & 2017GXNSFAA198006) and the National Natural Science Foundation of China (31860047). We thank Winston J. Goretsky (Calgary, Alberta, Canada) for reviewing and editing this work.

## References

- Chen HF, Cui XD, Zhang YY (2017) Plants of Nankunshan. China Forestry Publishing House, Beijing, 629 pp.
- Fang D, Qin HD, Lu XH (1994) New plants of Gesneriaceae from Guangxi of China. *Acta Phytotaxonomica Sinica* 32: 563–570.
- Hance HF (1866) Adversaria in stirpes imprimis Asiae orientalis critical minusve notas interjectis novarum plurimarum diagnosibus. *Annales des Sciences Naturelles, Botanique, sér.* 55: 202–261.
- Hayata B (1915) *Icones plantarum formosanarum nec non et contributiones ad floram formosanam* vol. 5. Government of Formosa, Taihoku, 358 pp.
- Kuntze O (1891) *Revisio Generum Plantarum* vol. 2. Arthur Felix, Leipzig, 375–1011.
- Léveillé H (1906) [“1905”] Les Gesneracées de la Chine. *Compte Rendu de l’Association Francaise Pour l’Avancement des Sciences* 34: 422–429.
- Li ZY, Wang YZ (2004) Plants of Gesneriaceae in China. Henan Science and Technology Publishing House, Zhengzhou, 721 pp.
- Wang WT (1984) Duo genera nova Gesneriacearum e sina australi. *Acta Phytotaxonomica Sinica* 22: 185–190.
- Wang WT, Pan KY, Li ZY, Weitzman AL, Skog LE (1998) Gesneriaceae. In: Wu ZY, Raven PH (Eds) *Flora of China*. Vol.18. Science Press, Beijing, & Missouri Botanical Garden Press, St. Louis, 244–401.
- Weber A, Middleton DJ, Forrest A, Kiew R, Lim CL, Rafidah AR, Sontag S, Triboun P, Wei YG, Yao TL, Möller M (2011) Molecular systematics and remodelling of *Chirita* and associated genera (Gesneriaceae). *Taxon* 60(3): 767–790. <https://doi.org/10.1002/tax.603012>
- Wei YG, Wen F, Möller M, Monro A, Zhang Q, Gao Q, Mou HF, Zhong SH, Cui C (2010) Gesneriaceae of South China. Guangxi Science and Technology Publishing House, Guilin, 777 pp.
- Xu YC, Cui XD, Zhang YY (2017) Wild ornamental plants of Nankunshan. China Forestry Publishing House, Beijing, 320 pp.
- Yamamoto Y (1936) *Observationes ad floram formosanam* XII. (Contributions from the herbaria of America and Europe. III). *Journal of the Society of Tropical Agriculture* 8: 63–77.
- Ye CX, Shi XG (2005) A new species of *Didymostima* (Gesneriaceae) from southern China. *Harvard Papers in Botany* 9: 447–448.