

# A new species of *Amentotaxus* (Taxaceae) from China, Vietnam, and Laos

Lian-Ming Gao<sup>1</sup>, Shao-Lin Tan<sup>1,2</sup>, Gui-Liang Zhang<sup>3</sup>, Philip Thomas<sup>4</sup>

**1** CAS Key Laboratory for Plant Diversity and Biogeography of East Asia, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming 650201, China **2** Kunming College of Life Science, University of Chinese Academy of Sciences, Kunming, Yunnan 650201, China **3** Hekou Branch of Administration Bureau of Daweishan National Nature Reserve, Hekou, Yunnan 661399, China **4** Royal Botanic Garden Edinburgh, 20A Inverleith Row, Edinburgh EH3 5LR, Scotland, UK

Corresponding author: Lian-Ming Gao ([gaolm@mail.kib.ac.cn](mailto:gaolm@mail.kib.ac.cn))

Academic editor: De-Zhu Li | Received 19 February 2019 | Accepted 29 April 2019 | Published 29 August 2019

**Citation:** Gao L-M, Tan S-L, Zhang G-L, Thomas P (2019) A new species of *Amentotaxus* (Taxaceae) from China, Vietnam, and Laos. In: Cai J, Yu W-B, Zhang T, Li D-Z (Eds) Revealing of the plant diversity in China's biodiversity hotspots. *PhytoKeys* 130: 25–32. <https://doi.org/10.3897/phytokeys.130.33956>

## Abstract

A new species *Amentotaxus hekouensis* L.M. Gao is described as new to science from Hekou, Yunnan of China, Lao Cai of Vietnam and Xiang Khouang of Laos. The new species is similar to *A. argotaenia* (Hance) Pilg. in linear or linear-lanceolate leaves, stomatal bands white and microsporophylls 6–8, each with 4–6 pollen sacs, but differs from the latter by its larger leaf size with 8–12.5 cm × 0.9–1.4 cm (vs. 2–11 cm × 0.5–1.1 cm in *A. argotaenia*), long acuminate leaf apex (vs. rounded to sharply triangular in *A. argotaenia*), stomatal bands with 25–30 rows (vs. 15–25 rows in *A. argotaenia*), stomatal bands equal to or slightly narrower than marginal bands (vs. narrower than marginal bands in *A. argotaenia*); pollen-cone racemes borne 1–2 (vs. 2–4 (10) in *A. argotaenia*), cones in 12–16 pairs (vs. ca. 12 pairs in *A. argotaenia*). Its distinctive nature has also been confirmed through DNA barcoding analysis of this genus. The new species is provisionally assessed as endangered (EN) due to its restricted distribution, small population size and the prevalence of habitat destruction within its range.

## Keywords

*Amentotaxus hekouensis*, New species, Endangered species, DNA barcoding, China


## Introduction

*Amentotaxus* Pilg. (Taxaceae), comprising of five or six species, is confined to southern China (including Taiwan), small areas of the eastern Himalayas and parts of Indo-China (Farjon 2010). All species of *Amentotaxus* are currently assessed as threatened at either national or global level (Wang and Xie 2004; IUCN 2016a). They usually occur in the understorey of moist submontane and montane semi-deciduous or evergreen forests (Gao et al. 2017). *Amentotaxus argotaenia* Pilg. is the most widespread species in this genus, recorded from most provinces of southern China as well as several areas of Vietnam and Laos (Fu et al. 1999; Farjon 2010; Nguyen et al. 2004; Averyanov et al. 2014). *Amentotaxus yunnanensis* H.L. Li is generally confined to the evergreen forests in the karst formation of southeast Yunnan, southwest Guizhou and adjacent areas of Vietnam and Laos (Fu et al. 1999; Farjon 2010; Averyanov et al. 2014). In 1996, *A. hatuyenensis* T.H. Nguyen was described from a single locality in Hagiang Province, northern Vietnam (Nguyen and Vidal 1996), but it is now regarded as a synonym of *A. yunnanensis*, based on morphological and molecular data (Phan et al. 2014; Gao et al. 2017). The remaining accepted species are restricted to Taiwan of China, eastern Himalayas and central Vietnam.

In September, 2006, specimens of an *Amentotaxus* tree were collected from limestone montane forest near Longyinzhong village, Nanxi town, Hekou county, Yunnan province, China. Its morphology differed markedly from *A. yunnanensis* which commonly occurs in this region. In 2012, a second collection with the same morphology was found in the same region. In our DNA barcoding study (Gao et al. 2017), specimens from Hekou formed a distinct clade with specimens with similar morphology from Lao Cai in Vietnam and Xiang Khoang in Laos. The putative new species differs from all other recognised species in the length and width of the leaves, the shape of the leaf apex, curvature of the leaf margin, stomatal band width and its ratio to the leaf marginal bands. In 2016, several further surveys collected specimens with male cones that showed differences in the number of pollen sacs and the length of the racemes. Data analysis of *trnL-F* and ITS sequences for the newly collected specimen GLM-164271 had an identical sequence with the specimen GLM-06209 (Am 20) and CKF250 (Am 24), used in our DNA barcoding study (Gao et al. 2017), lending further support to the recognition of a new species (Figure 1). Although female cones have not yet been collected, other species in the genus do not show significant differences in the female cones (Fu et al. 1999; Farjon 2010).

## Materials and methods

All measurements of the new *Amentotaxus* species were taken from dried herbarium specimens and field notes. In the description, length and width are represented as length × width. In total, eleven dried specimens of the new species were examined. All measurements of *A. argotaenia* (Hance) Pilg. and *A. yunnanensis* H.L. Li were based on literature (Fu et al. 1999; Farjon 2010) and our observations.



**Figure 1.** Neighbour-joining (NJ) tree of ITS and *trnL-F* sequences of *Amentotaxus* based on P-distance with bootstrap values above 50%.


## Taxonomic treatment

### *Amentotaxus hekouensis* L.M. Gao, sp. nov.

urn:lsid:ipni.org:names:60479341-2

Figs 2, 3, Table 1

**Diagnosis.** *Amentotaxus hekouensis* L.M. Gao resembles *A. argotaenia* (Hance) Pilg., but differs in its larger leaf size (8–12.5 cm × 0.9–1.4 cm), long acuminate leaf apex, leaf margin flat with slightly wavy, stomatal bands white or greenish-white with 25–30 rows, stomatal bands equal or slight narrower than marginal bands; pollen-cone racemes borne 1–2, cones in 12–16 pairs, microsporophylls 6–8, each with 4–6 pollen sacs.



**Figure 2.** *Amentotaxus hekouensis* L.M. Gao (from the holotype, drawn by Ling Wang). **A** Branchlets with male cone **B** pollen sacs.

**Table 1.** Morphological character comparison amongst the three regional *Amentotaxus* species.


Characters	<i>A. bekouensis</i>	<i>A. argotaenia</i>	<i>A. yunnanensis</i>
Length of leaf	8–12.5 cm	2–11 cm	3.5–10 cm
Width of leaf	9–14 mm	5–11 mm	8–12 mm
Texture of leaves	thin, leathery	thick, leathery	thick, leathery
Apex of the leaves	long acuminate	rounded to sharply triangular	obtuse or tapered
Width of stomatal bands	2.1–3.0 mm	1–2 mm	3–4 mm
No. rows of each stomatal band	25–30	15–25	ca. 40
Ratio of stomatal band/marginal band	0.75–1	< 1	> 2
Colour of marginal band	bright green in fresh	Yellowish-green	Yellowish-green in fresh
No. raceme per male cone	1–2	2–4 (10)	4–6

**Type.** CHINA. Yunnan: Honghe, Hekou County, Nanxi Town, Longyinzhong village, 22°41'8.9"N, 104°01'11.5"E, 926 m alt., 16 April 2016 (with male cone), G. L. Zhang, *GLM-164271* (holotype: KUN!; isotypes: KUN!)


**Morphological description.** Small tree to 4–5 m tall; branch cylindrical or sub-triangular, grey; leafy branchlets ascending or sub-erect, broadly rectangular to oblong-elliptic in outline, 6–12 × 11–20 cm, axis green in 1<sup>st</sup> year, greenish-yellow in 2<sup>nd</sup> and 3<sup>rd</sup> years, quadrangular or subterete in cross section. Leaves borne at 50–80° to branchlet axis, distichous, twisted at the short petiolate or nearly sessile base, petiole 2–4 mm long, almost opposite, 4–6 leaf pairs on each branchlet; leaves leathery, thin, linear or linear-lanceolate, 8–12.5 × 0.9–1.4 cm, straight, slightly falcate at the apex, cuneate at base, asymmetric, apex long acuminate, leaf margin flat or slightly down-curved, usually slightly wavy; leaf marginal band dark green in fresh, yellowish-green in dry, 2.5–3.2 mm wide; stomatal bands white or greenish-white, 2.1–3.0 mm wide, equal or slight narrower (> 3/4) to marginal bands, 25–30 rows, densely arranged; midvein slight sunken or flat adaxially, raised abaxially, 1.2–2.0 mm wide, narrower than the stomatal bands and marginal bands, green, same colour as the branchlet. Male-cone racemes borne 1–2, ca. 8.0 cm long; cones in 12–16 pairs, ovoid, cones at base maturing earlier than those at apex; microsporophylls 6–8, peltate, each with 4–8 pollen sacs. Ovulate cone and mature seed unknown. Male cones maturing March to April.

**Additional specimens examined.** CHINA. Yunnan: Hekou County, Nanxi Town, 20 Sept 2006, L. M. Gao et al. *GLM-06209* (KUN!); *ibid.*, 20 Oct 2012, L. M. Gao *GLM-123943* (KUN!), *ibid.*, 31 Jan 2016, L. M. Gao et al. *GLM-164258*, *GLM-164259* (KUN!), *ibid.*, 21 Oct. 2001, Y. M. Shui et al. *15078* (KUN); Funing County, Long-may, 24 April 1940, C. W. Wang *88812* (KUN); Malipo County, 4 Jan. 1940, C. W. Wang *86175* (KUN). Vietnam. Lao Cai: Muong Khuong, Nam Chay commune, Moi village, 21 Nov. 2009, CKF team *CKF250* (KUN, CPC, CMBG); Hoa Binh: Mai Chau, Pa Co commune, 3 Dec. 2003, S. G. Wu et al. *WP413* (KUN). Laos. Xiang Khouang: Phak Leung Mt., Tanh village, Dec. 2008, X. Gong *GX-5* (KUN).

**Distribution and ecology.** *Amentotaxus bekouensis* L.M. Gao has only been found in small disjunct areas in China (Funing, Malipo and Hekou County, Yunnan Province), Vietnam (Muong Khuong, Lao Cai Province; Mai Chau, Hoa Binh Province) and Laos (Xiang Khouang) (Figure 4). It is confined to the montane evergreen forest on lime-



**Figure 3.** *Amentotaxus bekouensis* L.M. Gao. **A** Habitat **B** habit **C** branchlet with adaxial leaves **D** branchlet with abaxial leaves **E** young shoot **F** branchlet and male cone racemes with microsporophylls **G** male cone racemes **H** microsporophylls and pollen sacs.



**Figure 4.** Geographical distribution of *Amentotaxus hekouensis* L. M. Gao.

stone mountains at elevations between 850 and 1200 (-1750) m. In part of its distribution along the border between China and Vietnam, it is sympatric with *A. yunnanensis*.

**Phenology.** Male cones of *Amentotaxus hekouensis* mature in March to April; seeds have not been seen so far.

**Etymology.** The specific epithet refers to the type location in Hekou County.

**Conservation status.** Several surveys have been carried out in this region since 2006 and only a few small trees of the new species are known. *Amentotaxus hekouensis* occurs in small and isolated patches of forests at very low densities. All the known localities are subject to degradation from surrounding agriculture and forest clearance. Given its montane habitat, the new species may also be susceptible to the impacts of climate change. The combination of a restricted, fragmented distribution with very limited Area of Occupancy, small population size and habitat conversion and destruction throughout its range indicate that its IUCN status should be Endangered under Criterion B2ab(ii,iii) (IUCN 2016b).

## Discussion

*Amentotaxus hekouensis* has distinct morphological features compared to the closely related *A. argotaenia* and the sympatric *A. yunnanensis* (Table 1). Its distinctive nature has also been supported by DNA barcoding analysis of this genus (Figure 1). Therefore,

the description of a new species is supported by both morphological and molecular evidence. Due to its restricted distribution, small population size and the prevalence of habitat destruction within its range, *A. hekouensis* is assessed as endangered (EN).

## Acknowledgements

We are grateful to Ms. Ling Wang from KUN for drawing the illustration of the holotype. We also thank Profs. Xun Gong, Yumin Shui from Kunming Institute of Botany, Chinese Academy of Sciences for providing the specimens from Laos and Vietnam of the new species. This work was supported by the Large-scale Scientific Facilities of the Chinese Academy of Sciences (Grant No: 2017-LSFGBOWS-02), the Strategic Priority Research Program of the Chinese Academy of Sciences (XDB31010000), and the Program of Science and Technology Talents Training of Yunnan Province, China (Grant No: 2017HA014). The Royal Botanic Garden Edinburgh is funded by the Rural and Environment Science and Analytical Services division of the Scottish Government.

## References

- Averyanov LV, Nguyen TH, Sinh KN, Pham TV, Lamxay V, Bounphanmy S, Lorphengsy S, Phan LK, Lanorsavanh S, Chantthavongsa K (2014) Gymnosperms of Laos. *Nordic Journal of Botany* 32(6): 765–805. <https://doi.org/10.1111/njb.00498>
- Farjon A (2010) *A Handbook of the World's Conifers*, 2 Vols. Brill, Leiden and Boston. <https://doi.org/10.1163/9789047430629>
- Fu LK, Li N, Mill RR (1999) Taxaceae. In: Wu ZY, Raven PH, Hong DY (Eds) *Flora of China*. Science Press, Beijing, Missouri Botanical Garden Press, St. Louis, Vol. 4, 89–96.
- Gao LM, Li Y, Phan LK, Yan LJ, Thomas P, Long KP, Möller M, Li DZ (2017) DNA barcoding of East Asian *Amentotaxus* (Taxaceae): Potential new species and implications for conservation. *Journal of Systematics and Evolution* 55(1): 16–24. <https://doi.org/10.1111/jse.12207>
- IUCN (2016a) The IUCN Red List of Threatened Species. Version 2015-4 <http://www.iucn-redlist.org> [accessed 12 February 2016]
- IUCN (2016b) Guidelines for Using the IUCN Red List Categories and Criteria, version 12 (February 2016). <http://www.iucnredlist.org/documents/RedListGuidelines.pdf> [downloaded 6 May 2016]
- Nguyen TH, Phan KL, Nguyen NDTL, Thomas PI, Farjon A, Averyanov L, Regalado J (2004) Vietnam Conifers: Conservation Status Review. Hanoi: Fauna and Flora International, Vietnam Program, 102–109.
- Nguyen TH, Vidal JE (1996) *Flore Cambodge, Laos and Vietnam*, Volume 28. Paris: Muséum National d'Histoire Naturelle, Laboratoire de Phanérogamie, 122–128.
- Phan KL, Phan KL, Le TML, Nguyen GS (2014) Molecular characterization of *Amentotaxus hatuyenensis* N.T. Hiep (Taxaceae) based on ITS1 partial sequence. *VNU Journal of Science: Natural Sciences and Technology* 30(3S): 46–52.
- Wang S, Xie Y (2004) *China Species Red List*. Higher Education Press, Beijing, 468 pp.