

A new species and two new records of *Goniothalamus* (Annonaceae) from Lao PDR

Bin Yang^{1,2}, Ren-Bin Zhu³, Hong-Bo Ding¹,
Somsanith Bouamanivong⁴, Yun-Hong Tan^{1,2}

1 Southeast Asia Biodiversity Research Institute, Chinese Academy of Sciences & Center for Integrative Conservation, Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences, Menglun, Mengla, Yunnan 666303, China **2** Center of Conservation Biology, Core Botanical Gardens, Chinese Academy of Sciences, Menglun, Mengla, Yunnan 666303, China **3** Chinese Union of Botanical Gardens Secretariat, Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences, Menglun, Mengla, Yunnan 666303, China **4** Ecology Division, Biotechnology and Ecology Institute, Ministry of Science and Technology, P.O. Box: 2279, Vientiane Capital, Lao PDR

Corresponding author: Yun-Hong Tan (tyh@xtbg.org.cn)

Academic editor: Xiao-Hua Jin | Received 12 August 2019 | Accepted 11 October 2019 | Published 10 January 2020

Citation: Yang B, Zhu R-B, Ding H-B, Bouamanivong S, Tan Y-H (2020) A new species and two new records of *Goniothalamus* (Annonaceae) from Lao PDR. In: Jin X-H, Xia N-H, Tan Y-H (Eds) Plant diversity of Southeast Asia-II. PhytoKeys 138: 17–25. <https://doi.org/10.3897/phytokeys.138.38995>

Abstract

Goniothalamus saccopetaloides Y.H. Tan & Bin Yang, a new species is described and illustrated from Laos. This species shows morphological similarities to *G. yunnanensis* W.T. Wang, but it differs from the latter by having almost fleshy, involute and saccate outer petals, subglobose monocarps, and single seeded monocarps. *Goniothalamus cheliensis*, and *G. calvicarpus* are new records for the Flora of Lao PDR. A key to *Goniothalamus* species indigenous to Laos is provided here.

Keywords

Goniothalamus, Laos, field survey, new species, new record

Introduction

Goniothalamus (Blume) Hook.f. & Thomson, comprising over 130 species of trees and shrubs, are one of the largest palaeotropical genera in the Annonaceae, mainly distributed from India and Sri Lanka to tropical Australia and the South Pacific Islands (Saun-

ders and Chalermglin 2008; Nakkuntod et al. 2009; Turner 2014; Thomas et al. 2017). The genus is characterized by flowers with two whorls of petals of which the inner petals are smaller than the outer ones; three inner petals form a distinctive mitriform dome over the reproductive organs acting as a pollination chamber (Saunders and Chalermglin 2008; Tang et al. 2015), and stamens with apical connectives. Although the genus *Mitrephora* J. D. Hooker & Thomson also shares these similar features, *Goniothalamus* can be distinguished easily by its linear-oblong stamens, inner petals each with a short claw or stipe (Li et al. 2011). The genus shows considerable diversity in flowers of different size and shape, hairy indumentum, and color usually cream, yellow or red at maturity, and fruit morphology, with fruit of different size and shape (Saunders and Chalermglin 2008; Tang 2014; Tang et al. 2015). Species of the genus are widely distributed in tropical South-East Asian lowland and submontane forests (Tang 2014). Several *Goniothalamus* species have been described in recent years, including four new species from Thailand (Saunders and Chalermglin 2008), four new species from Borneo (Turner and Saunders 2008), and a new species from Palawan, the Philippines (Tang et al. 2013). In Laos, *Goniothalamus* are represented by four species (Newman et al. 2007; Lee 2016), i.e., *Goniothalamus laoticus* (Finet & Gagnep.) Bân, *Goniothalamus repevensis* Pierre ex Finet & Gagnep., *Goniothalamus marcanii* Craib and *Goniothalamus saigonensis* Pierre ex Finet & Gagnep., whereas the latter two were respectively treated as synonyms of *Goniothalamus tamirensis* Pierre ex Finet & Gagnep. (Saunders and Chalermglin 2008) and *Goniothalamus gabriacianus* (Baill.) Ast (Li et al. 2011).

In recent years, the authors examined the flowering material of Annonaceae cultivated in the living collections of Xishuangbanna Tropical Botanical Garden (XTBG), Chinese Academy of Sciences (CAS) and found an unknown species, which is very distinctive and belongs to the genus *Goniothalamus*; the records showed that this accession was originally collected in 2002, from Lao PDR (although the exact location remains unknown). Based on the morphological characters, we compared it to all other currently known species and concluded that it is new to science. As part of the botanical inventory of China-Laos transboundary biodiversity conservation, we carried out floristic surveys in the Nam Ha National Biodiversity Conservation Area in Luang Namtha Province and Phou Hin Phee National Biodiversity Conservation Area in Oudomxay Province of the northern Laos. During the fieldwork in March and October of 2018, we encountered two species of *Goniothalamus* representing new records for the flora of Laos. Therefore, the new and noteworthy species of *Goniothalamus* from Laos are provided and updated in this study.

Material and methods

Our study of the new taxon was predominantly based on plant material newly collected in XTBG. The records showed that this accession was originally collected in 2002, from Laos, although the exact location remains unknown. Ten individuals cultivated in

two living collections were observed. Morphological characterizations were measured in the field. We compared our samples with type specimens of similar species deposited in herbaria. Specimens of *Goniothalamus* from Laos and neighboring regions were examined from the following herbaria: HITBC, IBSC, SING, A, K, BK, high-resolution digital images of specimens (especially types) from JSTOR Global Plants (<https://plants.jstor.org/>) and other virtual herbarium websites (<http://www.cvh.ac.cn/>), as well as the taxonomic literature for species identification.

Taxonomic treatments

Goniothalamus saccopetaloides Y.H.Tan & Bin Yang, sp. nov.

urn:lsid:ipni.org:names:77204192-1

Fig. 1

Diagnosis. *Goniothalamus saccopetaloides* is morphologically similar to *G. yunnanensis* W.T. Wang with elliptic-oblong to oblong leaf blades and broadly lanceolate, pinkish orange to reddish brown outer petals, but easily distinguished by having almost fleshy, involute, saccate outer petals, subglobose and single seeded monocarps.

Type. Lao PDR. Specific location unknown. Voucher from a cultivated plant at the Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences, 2 May 2019, B. Yang, XTBG-0054 (holotype, HITBC!).

Description. Shrub to small tree, to 3 m tall. Young branches glabrous. Leaf laminas 15.5–27.8 cm long, 4.5–8.2 cm wide, length/width ratio ca. 3.4, elliptic-oblong to oblong, apex acuminate, base cuneate to attenuate, papyraceous, glabrous abaxially and adaxially; midrib glabrous and (strong) prominent abaxially, glabrous and impressed adaxially; secondary veins 13–18 pairs, (slightly) impressed adaxially; tertiary veins percurrent, distinct; petioles 5–12 mm long, 2–3 mm in diameter, glabrous. Flowers 1–7, often on the main trunk (cauliflory) and on older branches (ramiflory), pendants; flowering pedicels 2–5 mm long, sparsely hairy; pedicel bracts 3–6, 1–2 mm long. Sepals 5–6 mm long, 5–7 mm wide, basally connate, apex acuminate, broad ovate, puberulous abaxially, glabrous adaxially, greenish-yellow, venation longitudinal slightly conspicuous adaxially, indistinct abaxially. Outer petals 1.4–2.7 cm long, 0.7–1.2 mm wide, length/width ratio 1.6–3.0, almost fleshy, involute, saccate, elliptic lanceolate to broadly lanceolate, sparsely puberulous abaxially, subglabrous adaxially, yellowish green in young stage, pinkish orange to reddish brown in mature stage, venation slightly distinct adaxially, occasionally slightly distinct abaxially. Inner petals 6.5–12 mm long, 5–10 mm wide, length/width ratio 1.1–2.2, broadly ovate to obovate, sparsely puberulous abaxially, pubescent adaxially, yellowish green in young stage, pinkish orange to reddish brown in mature stage, base attenuate to a 2–3 mm claw. Stamens ca. 80 per flower, 1.2–1.5 mm long, 0.6–0.7 mm wide; connectives truncate. Carpels 18–22 per flower, ovary 1.5–2 mm long, light green, with




Figure 1. *Goniothalamus saccopetaloides* Y.H.Tan & Bin Yang, sp. nov. **A** habit **B** leafy branch **C** flowers on main trunk **D** fruits on main trunk **E** leaves (adaxial and abaxial surface) **F** flowers **G–H** dissected flowers **I** carpels **J** longitudinally dissected monocarp **K** seeds. Photographed by R.B. Zhu and B. Yang.

white hairs; stigma and style 2–3 mm long, glabrous. Immature fruits green, with small white dots, mature fruits orange to red; fruiting pedicels 3–5 mm long, 1.5–2 mm in diameter, subglabrous. Monocarps one seeded, 12–13 mm long, 9–11 mm wide, length/width ratio 1.2–1.3, subglobose to ellipsoid, base and apex rounded, smooth, subglabrous, glossy, pericarp medium-thick, ca. 1.5 mm thick, stipes subsessile to 1.5 mm long, ca. 2 mm in diameter, glabrous. Seeds 9–10 mm long, ca. 7 mm wide, length/width ratio 1.3–1.4, ellipsoids, testa slightly rugose, dark brown to black brown, aril orange.

Etymology. The new species is named after its saccate and almost fleshy outer petals.

Phenology. *Goniothalamus saccopetaloides* has been observed in flowers from the end of March to May and in fruits from July to September.

Distribution and habitat. *G. saccopetaloides* was originally distributed in Laos; the exact location and its wild habitat remains unknown. Additional collections in the future may help to clarify its full distribution.

Conservation status. Due to insufficient field surveys so far, very limited details about its natural distribution and population status are currently known. Further investigation is required to determine more distribution sites and conservation status of the new species; at this moment we consider it as data deficient (DD) according to the IUCN Red List Categories (IUCN 2012).

Additional specimen examined (paratype). Lao PDR. Specific location unknown. Voucher from the cultivated plants at the Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences, 2 May 2019, B. Yang, XTBG-0055 (HITBC!).

Notes. *Goniothalamus saccopetaloides* is morphologically similar to *G. yunnanensis*; however, in addition to the description in the diagnosis, the two species can be easily distinguished by other characters. *Goniothalamus saccopetaloides* has 1 to 7 flowers, often on the main trunk (cauliflory) and on older branches (ramiflory), scarcely on young growth, whereas *G. yunnanensis* has 1 to 2 flowers, axillary, often on young growth, sometimes from leafless nodes (Li et al. 2011). Moreover, *G. saccopetaloides* has one ovule per carpel, subglobose monocarps and seeds with rounded apices, and *G. yunnanensis* has 2 ovules per carpel, ellipsoid monocarps and seeds with acute apices (Li et al. 2011).

New records for Laos

Goniothalamus calvicarpus Craib

Fig. 2 (A–D)

Goniothalamus calvicarpus Craib, Bull. Misc. Inform. Kew. 1922: 227 (1922). – TYPE: Thailand: Sukotai, Kao Luang, Sukhothai Province, Northern Thailand, 4 May 1922, A.F.G. Kerr 5946 (holotype: K!; isotypes: BK!, BM!).




Figure 2. *Goniothalamus calvicarpus* Craib **A** leafy branch **B** leaves (abaxial) **C** fruit **D** flower (epibiotic sepals); *Goniothalamus cheliensis* Hu **E** leafy branch **F** leaves (abaxial) **G** flower **H** flower (picked petals, showing stamens and carpels) **I** dissected flowers. Photographed by H.B. Ding.

Distribution and habitat. Thailand (Saunders and Chalermglin 2008), China (Li et al. 2011) and Northern Laos (Luang Namtha Province), mountain slopes, 800–1500 m.

Specimens examined. Laos: Tha Se Village, Nam Ha National Biodiversity Conservation Area, Luang Namtha Province, Northern Laos, tropical lower montane forest, 938 m, 20°49'59.42"N, 101°15'48.05"E, 21 Oct 2018, *Y.H. Tan et al.*, L0757 (HITBC); idem, 21 Oct 2018, *Y.H. Tan et al.*, L0764 (HITBC); idem, 26 Mar 2018, *Y.H. Tan et al.*, L0183 (HITBC); Nam Sing Village, Nam Ha National Biodiversity Conservation Area, Luang Namtha Province, Northern Laos, stone forest, 900 m, 20°45'09"N, 101°12'08"E, 25 Mar 2018, *Y.H. Tan et al.*, L0153 (HITBC).

Notes. There has been considerable taxonomic confusion between *G. calvicarpus* and *G. griffithii* Hook. f. & Thomson (Saunders and Chalermglin 2008), the Chinese material identified as *G. griffithii* to be more correctly placed in *G. calvicarpus* (Saunders and Chalermglin 2008; Li et al. 2011). *Goniothalamus calvicarpus* is relatively widely distributed in the lower mountain evergreen forest in Xishuangbanna, southern Yunnan, compared with our materials collected in Northern Laos, the characters of flowers and fruits are exactly in line with *G. calvicarpus*; it is also not very hard to encounter around the adjacent areas in Northern Laos bordering southern Yunnan.

Goniothalamus cheliensis Hu

Fig. 2 (E–I)

Goniothalamus cheliensis Hu, Bull. Fan Mem. Inst. Biol. 10: 122 (1940). – TYPE: China: Che-Li Hsien, Maan-Shang, Yunnan, Sep. 1936, *C.W. Wang* 78573 (holotype: TAI!, isotypes: A!, IBSC!).

Distribution and habitat. Northern Thailand (Saunders and Chalermglin 2008), China (Li et al. 2011) and Northern Laos (Oudomxay Province), Montane forests; 1000–1500 m.

Specimens examined. Laos: Maung Xai, Phou Hin Phee National Biodiversity Conservation Area, Oudomxay province, 1357 m, 20°43'19.12"N, 102°08'46.61"E, 30 March 2018, *Y.H. Tan et al.*, L0350 (HITBC); ibdem, 1372 m, 20°43'18.24"N, 102°08'47.31"E, 30 March 2018, *Y.H. Tan et al.*, L0351 (HITBC).

Notes. *Goniothalamus cheliensis* is a very distinctive species, with leaves, flowers and fruits very large and densely rust-colored hirsute (Saunders and Chalermglin 2008), according to our personal observation, the monocarps of *G. cheliensis* can be up to 15 cm long.

With the addition of *Goniothalamus saccopetaloides*, *Goniothalamus cheliensis*, and *Goniothalamus calvicarpus*, seven species are currently recognized in Laos (Newman et al. 2007; Saunders and Chalermglin 2008; Li et al. 2011; Lee 2016). A key is provided below to further elucidate the morphological differences among the species occurring in Laos.

Key to the *Goniotalamus* species in Laos and along with *G. yunnanensis*

- 1 Young branches densely hairy to velutinous, leaf midrib and petiole (very densely hairy abaxially) **2**
- Young branches glabrous, leaf midrib and petiole glabrous to sparsely hairy abaxially **3**
- 2 Leaf blade 50–76 × 13–22 cm, stamen connectives apiculate, monocarps oblong, 3–15 cm long and densely hairy *G. cheliensis*
- Leaf blade 12–17 × 3.7–4.7 cm, stamen connectives truncate, monocarps ovoid, less than 3 cm long and glabrous *G. tamirensis*
- 3 Adaxial surface of leaves with very prominent secondary and tertiary veins **4**
- Adaxial surface of leaves with impressed or only slightly prominent secondary and tertiary veins **6**
- 4 Stamen connectives apiculate, sepals 11–18.5 mm long *G. calvicarpus*
- Stamen connectives truncate, sepals 5–8 mm long **5**
- 5 Flowers on older branches and young growth, outer petals not fleshy, never involute, monocarps ellipsoid, 1.8–2.5 cm long, apex acute *G. yunnanensis*
- Flowers on main trunk and older branches, outer petals almost fleshy, involute, monocarps subglobose to ellipsoid, 1.2–1.3 cm long, apex rounded *G. saccopetaloides*
- 6 Ovules 8–10 per carpel, monocarps oblate, with its longitudinal ridge *G. laoticus*
- Ovules 1–2 per carpel, monocarps oblong-ellipsoid to ellipsoid, or elliptic-ovoid **7**
- 7 Leaf blades with short to long acuminate apices, outer petal venation distinct, sparsely puberulous to glabrous abaxially *G. repevensis*
- Leaf blades with acute to attenuate apices, outer petal venation indistinct, hispidulous abaxially *G. gabriacianus*

Acknowledgements

The authors cordially thank staff of the Biotechnology and Ecology Institute (BEI), Nam Ha National Biodiversity Conservation Area in Laung Namtha Province and Phou Hin Phee National Biodiversity Conservation Area, Oudomxay province of Lao PDR for their support of field works in the protected area. We are grateful to Dr. Liang Song, Dr. Hua-Zheng Lu, Dr. Su Li, Mr. Ren Li, Mr. Xiao-Dong Zeng, Ms. Zhong-Li Gan and Ms. Dong-Li Quan for their kind help in the field. This research was funded by Lancang-Mekong Cooperation (LMC) Special Fund (Biodiversity Monitoring and Network Construction along Lancang-Mekong River Basin project) and the CAS 135 program (No. 2017XTBG-F03), and the project of the Southeast Asia biodiversity research institute, Chinese Academy of Sciences (Y4ZK111B01).

References

- IUCN (2012) IUCN Red List Categories and Criteria, Version 3.1 (2nd edn). Gland and Cambridge, 32 pp.
- Lee YM (2016) A checklist of plants in Lao PDR. Korea National Arboretum of the Forest Service, 402 pp.
- Li BT, Gilbert MG, Schatz G (2011) *Goniothalamus*. In: Wu ZH, Raven PH, Hong DY (Eds) Flora of China (Vol. 19). Science Press, Beijing, and Missouri Botanical Garden Press, St. Louis, 684–687.
- Nakkuntod M, Su YCF, Seelanan T, Saunders RMK (2009) Molecular phylogenetic and morphological evidence for the congeneric status of *Goniothalamus* and *Richella* (Annonaceae). *Taxon* 58(1): 127–132. <https://doi.org/10.1002/tax.581013>
- Newman M, Ketphanh S, Svengsuksa B, Thomas P, Sengdala K, Lamxay V, Armstrong K (2007) A Checklist of the Vascular Plants of Lao PDR. Royal Botanic Garden, Edinburgh, 394 pp.
- Saunders RMK, Chalermglin P (2008) A synopsis of *Goniothalamus* species (Annonaceae) in Thailand, with description of three new species. *Botanical Journal of the Linnean Society* 156(3): 355–384. <https://doi.org/10.1111/j.1095-8339.2007.00762.x>
- Tang CC (2014) Systematics and phylogenetics of *Goniothalamus* (Annonaceae): a study of putative rapid, recent evolutionary radiation in a species-rich palaeotropical genus. PhD thesis, University of Hong Kong.
- Tang CC, Xue B, Saunders R (2013) A new species of *Goniothalamus* (Annonaceae) from Palawan, and a new nomenclatural combination in the genus from Fiji. *PhytoKeys* 32: 27–35. <https://doi.org/10.3897/phytokeys.32.6663>
- Tang CC, Thomas DC, Saunders RMK (2015) Molecular phylogenetics of the species-rich angiosperm genus *Goniothalamus* (Annonaceae) inferred from nine chloroplast DNA regions: Synapomorphies and putative correlated evolutionary changes in fruit and seed morphology. *Molecular Phylogenetics and Evolution* 92: 124–139. <https://doi.org/10.1016/j.ympev.2015.06.016>
- Thomas DC, Tang CC, Saunders RMK (2017) Historical biogeography of *Goniothalamus* and Annonaceae tribe Annoneae: Dispersal-vicariance patterns in tropical Asia and intercontinental tropical disjunctions revisited. *Journal of Biogeography* 44(12): 2862–2876. <https://doi.org/10.1111/jbi.13086>
- Turner IM (2014) *Goniothalamus*. In: Soepadmo E, Saw LG, Chung RCK, Kiew R (Eds) Tree flora of Sabah and Sarawak (Vol. 8). Forest Research Institute Malaysia Press, 42–70.
- Turner IM, Saunders RMK (2008) Four new species of *Goniothalamus* (Annonaceae) from Borneo. *Nordic Journal of Botany* 26(5–6): 329–337. <https://doi.org/10.1111/j.1756-1051.2008.00359.x>