

Synopsis of *Martinella* Baill. (Bignoniaceae), with the description of a new species from the Atlantic Forest of Brazil

Alexandre R. Zuntini¹, Lucia G. Lohmann¹

¹ Departamento de Botânica, Instituto de Biociências, Universidade de São Paulo, Rua do Matão, 277, 05508-090, São Paulo, SP, Brazil

Corresponding authors: Alexandre R. Zuntini (zuntini@gmail.com); Lucia G. Lohmann (llohmann@usp.br)

Academic editor: A. Paton | Received 1 January 2014 | Accepted 1 April 2014 | Published 9 May 2014

Citation: Zuntini AR, Lohmann LG (2014) Synopsis of *Martinella* Baill. (Bignoniaceae, Bignoniaceae), with the description of a new species from the Atlantic Forest of Brazil. *PhytoKeys* 37: 15–24. doi: 10.3897/phytokeys.37.6940

Abstract

Martinella has traditionally included two species, *Martinella iquitoensis* and *Martinella obovata*, that are characterized by the presence of interpetiolar ridges surrounding the stems and minute prophylls of the axillary buds. A third species, *Martinella insignis*, is here described as new, illustrated and compared to other species in the genus. *Martinella insignis* is the first record of the genus in the Atlantic Forest of Brazil, and differs from other species of *Martinella* by the yellow corolla (vs. red to dark purple) and 5-lobed calices (vs. 2–4-lobed).

Keywords

Martinella, Bignoniaceae, Bignoniaceae, Neotropics, Brazilian Atlantic Forest

Introduction

Martinella Baill. (1888) is strongly supported as monophyletic by molecular data (Lohmann 2006). Species of *Martinella* are well distinguished by the combination of minute triangular prophylls of the axillary buds, an interpetiolar ridge surrounding the stem, and bilobed or (2–)4-parted calyces (Lohmann and Taylor 2014); the latter two being considered putative synapomorphies of the genus (Lohmann 2006; Lohmann

and Taylor 2014). In addition to these features, the basal tubular portion of the corolla is slightly longer and much narrower than the calyx, leaving it loose within the calyx, while the upper portion of the corolla tube is abruptly inflated and campanulate, up to four times wider than the tubular portion (Fig. 1D). Gentry (1974) described this corolla morphology plus the red to purple color as a *Martinella*-type flower, and hypothesized that this flower was associated with pollination by hummingbirds. The only exception to Gentry's floral description is the position of the anthers (exserted or subserted), which are always included in *Martinella*.

The genus as currently circumscribed includes two species of neotropical lianas (Lohmann and Ulloa Ulloa 2013; Lohmann and Taylor 2014): *Martinella iquitoensis* A.Samp., restricted to the Amazon basin (Brazil, Colombia, Ecuador, Peru and Venezuela) and *Martinella obovata* (Kunth) Bureau & K.Schum., ranging from Central America to Northern South America and the Amazon basin (Lohmann et al. 2013). A third morphologically distinct species of *Martinella* was discovered during fieldwork in the Atlantic Forest of Brazil; its description here extends the known range of this genus. Here we present an overview of the genus and the separation of its species.

Materials and methods

This work is based on the study of herbarium collections of *Martinella* deposited in CEPEC, MBM, MBML, MO, NY, RB, SPF and VIES (herbarium acronyms follow Thiers 2013). Morphological descriptions are based on dried specimens, and follow the terminology of Radford et al. (1974). The parentheses in the descriptions indicate rare conditions. Micrographs of selected structures were made using a stereomicroscope and digitally processed through focus stacking. For the species distribution map, the geographical data of *M. insignis* were combined with those used in Lohmann et al. (2013) and plotted over a digital elevation model (GTOPO 30, available from the U.S. Geological Survey).

Taxonomic treatment

***Martinella* Baill., Hist. Pl. 10: 30. 1888.**

<http://species-id.net/wiki/Martinella>

Type. *Martinella martinii* (DC.) Baill. ex K. Schum. (= *Martinella obovata* (Kunth) Bureau & K. Schum.)

Lianas. *Roots* with tuberous portions. *Branches* terete, glabrous or puberulous, with trichomes simple or stipitate-glandular, with continuous interpetiolar ridges, without interpetiolar glands; prophylls minute, triangular, glabrous or puberulous. *Leaves* 3-foliate or 2-with the terminal leaflet modified into a simple or trifid tendril; leaflets membranous to coriaceous, margins entire (sinuate), with or without mite-domatia,

glabrous to puberulous, with glands on adaxial surface. *Inflorescences* axillary, arranged in racemes, panicles, thyrses or compound dichasia. *Flowers* with calyx tubular (campanulate), bilobed, irregularly 2–4-lobed, or 5-lobed, with lobes rounded or aristate, membranous, with few scattered glands; corolla deep purple, red or yellow, tubular in the basal portion and campanulate in the upper part, straight to weakly curved, membranous, glabrous outside, glabrous inside except with few glandular trichomes at stamen insertion; stamens included, glabrous, pollen in monads; ovary terete, smooth, glabrous or lepidote, with a single series of ovules on each placenta, style glabrous, stigma rhombic, glabrous. *Capsules* drying dark brown, linear, flattened, smooth, glabrous or puberulous, with calyx caducous; seeds oblong, winged, with wings opaque.

Martinella comprises three species, distributed from Mexico to eastern Brazil. The main features that distinguish the species are summarized in Table 1 and outlined in the key below.

Key to species of *Martinella*

- 1 Calyx 5-lobed; corolla yellow; eastern Brazil **1. *M. insignis***
- 1' Calyx 2–4-lobed; corolla deep wine to red; Antilles, Central America and Northern South America through Amazon basin **2**
- 2 Inflorescence arranged in thyrse or panicle; leaflet with cuneate base
..... **2. *M. iquitoensis***
- 2' Inflorescence arranged in raceme; leaflet with cordate to truncate or rarely cuneate base **3. *M. obovata***

1. *Martinella insignis* A.H. Gentry ex Zuntini & L.G. Lohmann, sp. nov.

urn:lsid:ipni.org:names:77138471-1

http://species-id.net/wiki/Martinella_insignis

Figs 1–2

Type. BRAZIL. Bahia: Itamaraju, Rodovia Itamarajú-Teixeira de Freitas, 3km de Itamaraju (BR-101). Fazenda Chapadão, 3 November 1983, R. Callejas, A. M de Carvalho & L. M. Silva 1629 (holotype MBM-94960!; isotypes CEPEC not seen, MO-3600686!, NY-00483568!, RB-232556!).

Diagnosis. *Martinella insignis* differs from *M. iquitoensis* and *M. obovata* by its 5-lobed calyces and yellow corollas, in contrast to 2–4-lobed calyces and dark purple to red corollas in these other species (Table 1).

Description. *Lianas*. Branches green, drying brownish, striated, densely covered with stipitate-glandular trichomes when young; prophylls 0.7–1.5 × 1.0 mm, densely covered with stipitate glandular trichomes, without patelliform glands (with few patelliform glands). *Leaves* 2-foliolate with the terminal leaflet generally modified into a trifold tendril; petioles terete, 34–64 mm long, covered with stipitate glandular trichomes; petiolules terete, 14–42 mm long, covered with stipitate glandular trichomes;



Figure 1. Morphology of *Martinella insignis*: **A** Habit **B** Node with interpetiolar ridge **C** Glandular stipitate trichomes **D** Flower (lateral view) **E** Calyx (opened) and gynoecium **F** Fruit **G** Root system with tuberous portions. From Zuntini 151 (A–E), Sucre 5519 (F) and Zuntini 321 (G).

leaflets weakly discolorous, membranous, ovate, with a long acuminate to caudate apex and a cordate base, margins entire (sinuate), 7.6–11.8 × 3.4–6.4 cm, glabrous except on margins and main veins of the abaxial surface where stipitate glandular trichomes are found, with pocket domatia on the axils of primary and secondary veins, with

Table 1. Morphological and geographical summary of *Martinella* species, based on Gentry (1977, 1982, 2009) and pers. obs.

Character	<i>M. insignis</i>	<i>M. iquitoensis</i>	<i>M. obovata</i>
Leaflet texture	membranous	coriaceous	membranous to coriaceous
Leaflet shape	ovate	elliptic	ovate (elliptic)
Leaflet base	cordate	cuneate	cordate to truncate (cuneate)
Leaf domatia	pocket	absent	absent
Tendril	trifid	simple (trifid)	trifid (simple)
Inflorescence	compound dichasium	thyrses or panicle	raceme
Calyx lobes	5; aristate	2–4; rounded	2–4; rounded
Corolla color	yellow	dark purple	dark purple to red (lilac)
Distribution	Eastern coast of Brazil	Amazon basin	Antilles, Central America and Northern South America through Amazon basin
Soil preference	sandy	sandy	generalist

few glands concentrated near base and scattered along the mid-vein on the adaxial surface. *Inflorescences* compound dichasia, with up to 7 branching orders, 9.2–12.3 cm long, sparsely to densely covered with stipitate glandular trichomes; bracts linear to narrowly elliptic, 8.4–24.7 × 0.8–3.4 mm, densely covered with stipitate glandular trichomes; pedicels terete, 5.8–19.4 mm, sparsely to densely covered with stipitate glandular trichomes. *Flowers* with calyx pale green, tubular (campanulate), 8.8–15.4 × 6.5–12 mm, sparsely covered with stipitate glandular trichomes except densely covered at the base, with few glands near the apex; lobes 5, very shallowly triangular, aristate, aristae 2.9–8.6 mm long, densely covered with stipitate glandular trichomes; corolla yellow, weakly curved, 29.5–48.3 mm long, tubular basal portion 12.7–18.9 long × 2.4–4.5 mm wide, upper campanulate portion 15.0–23.4 long × 9.2–17.2 mm wide, lobes subcircular, 3.8–9.8 × 6.1–10.9 mm, with ciliate margins; stamens in two lengths, longer ones 11.4–17.3 mm, shorter ones 12.1–15.8 mm, thecae 2.4–3.0 mm, glabrous; staminode 1.1–3.6 mm, glabrous; gynoeceum 29.2–39.6 mm long; ovary glabrous; style glabrous; stigma rhomboid, glabrous; nectariferous disk 2.3–3.0 × 1.0–1.8 mm. *Capsules* 33.6–73.0 × 1.1–1.2 cm, pubescent when immature, glabrous when developed. *Seeds* ca. 1.0 × 4.6 cm.

Distribution and habitat. *Martinella insignis* is restricted to the northern portion of the Brazilian Atlantic Forest, occurring predominantly in areas with sandy soils (Fig. 3).

Etymology. The species epithet means remarkable or clearly distinguishable. This epithet was probably selected by Alwyn Gentry as reference to the contrasting floral color among species of *Martinella*.

Phenology. Flowering specimens were collected between September and February and fruiting collections in January, September and November.

Conservation status. This species is considered Data Deficient [DD] according to IUCN Standards and Petitions Subcommittee (2014) since this taxon is only known from very few specimens, with little information about its distribution and abundance. Further field studies are needed so that its conservation status can be properly documented.




Figure 2. Details of *Martinella insignis*: **A** Flower **B** Leaflet base showing glands (abaxial face) **C** Mite-domatia between primary and secondary veins (abaxial side) **D** Root system with tuberous portions **E** Calyx detailing the aristae **F** Arista detail with glandular trichomes **G** Simple, tector trichomes on immature fruit. From Zuntini 151 (A–C, E, F), Zuntini 321 (D) and Demuner 4481 (G). Micrographs were obtained using focus stacking.

Discussion. *Martinella insignis* is the first species of *Martinella* found in the Atlantic Forest of Brazil. This new species clearly belongs to *Martinella* based on its prophylls, the continuous interpetiolar ridges and the corolla shape. However, *M. insignis* can be distinguished by the membranous leaflets, 5-lobed aristate calyces, and yellow corollas. In addition, *M. insignis* also has pocket-shaped leaf domatia (Fig. 2C) and a puberulous indument of glandular stipitate trichomes that covers almost all organs, with variable density (Fig. 2E–F). These trichomes may also be found in *M. obovata* and a few other species in Bignoniaceae, and are typically formed by a multicellular secretory head, supported by a uniseriate stalk (Nogueira et al 2013). Only corollas and fruits lack these; the corollas are glabrous, and fruits have simple, deciduous trichomes (Fig. 2G). Similar to the other species, the root system of *M. insignis* has unusual tuberous portions (Fig. 2D) that might represent an adaptation to the sandy soils, by accumulating water. However, the anatomical structure and function of these are yet unknown. Alwyn Gentry had already noted this new taxon, and had proposed the epithet “*insignis*” in sched.; his earlier findings are here accredited.

Additional collections examined. Brazil. Bahia: Guaratinga, Fazenda Vitória, 16°43'S, 39°46'W, 29 October 1979, L.A. Mattos Silva & H.S. Brito 634 (CEPEC, MO). Itabuna, Alcobaca para (ramal) S. Antonio, 24 January 1972, R.S. Pinheiro 1759 (CEPEC, MO). Itamaraju, S. Mori, L.A. Mattos Silva & T.S. Santos 10743 (CEPEC, MO), Itamaraju, Fazenda Riacho das Pedras, prop. Sr. Gersino Antônio Bronzon, 17°08'48"S, 39°21'53"W, 12 February 2007, R.A.X. Borges, A. Amorim, W.W. Thomas, L.C. Gomes, S. Sant'Ana & O. Cruz 825 (CEPEC, SPF). Espírito Santo: Linhares, Reserva Natural da Companhia Vale do Rio Doce (“Reserva de Linhares”), MME, 19°07'57.5"S, 40°04'06.3"W, 65m, 14 December 2007, A.R. Zuntini, W.A.A. Pires & G.S. Siqueira




Figure 3. Distribution of *Martinella* species: *M. insignis* (solid blue circle), *M. iquitoensis* (solid green square) and *M. obovata* (open red triangles). Elevation in meters, according to the scale presented on the right.

151 (CVRD, RB, SPF), A.R. Zuntini, E. Françoso, J. Lopes & V. Augusto 321 (SPF). Governador Lindenberg, Pedra de Santa Luzia, 420–590 m, 7 November 2007, V. Demuner, T.A. Cruz & R.R. Vervloet 4481 (MBML, SPF). Sooretama, Mata de tabuleiro situada ao Noroeste da sede da Reserva da Sooretama, 14 July 1969, D. Sucre 5519 (RB–photo).

2. *Martinella iquitoensis* A. Samp., *Ann. Acad. Bras. Sci.* 7: 122. 1935.

http://species-id.net/wiki/Martinella_iquitoensis

Martinella manaosiana A.Samp., *Bol. Mus. Nac. Rio de Janeiro* 12(3, 4): 84. 1936.

TYPE: BRAZIL. Amazonas: Manaus, 25 July 1931, A. Ducke *sn* (holotype: RB-24095!; isotype MO-2193049!, RB-24095 [second sheet]!).

Type. PERU. Loreto: Iquitos, 23 February 1924, J.G. Kuhlmann 1492 (holotype RB-22027!; isotypes, MO-2192060!, RB-22027 [second sheet]!).

Distribution and habitat. This species is distributed widely in the Amazon basin (Brazil, Colombia, Ecuador, Peru and Venezuela), typically in sandy soils (Lohmann and Taylor 2014; Fig. 3).

Conservation status. *Martinella iquitoensis* is distributed geographically through an area that is < 2000 km², with seven Rapoport (1982) sub-populations known to date and ≥ 20% of its known individuals occurring outside Protected Areas, making it susceptible to the current reduction and degradation of its habitat. Therefore, this species is here considered as Vulnerable [VU B2ab(ii,iii)] according to the IUCN criteria (IUCN 2012; IUCN Standards and Petitions Subcommittee 2014).

Discussion. This species was distinguished by Sampaio from *M. obovata* based on the corolla color and size, leaflet texture and size, tendril type, and calyx indument. However these characters have proven to be fairly variable, especially in *M. obovata*, leading to morphological overlap between those taxa. The lack of a clear morphological discontinuity combined with the sympatric distributions, make these species hard to separate. Moreover, the difficulty in delimitating these two species can be observed in the few treatments that dealt with those species, which is particularly evident in the contrasting species keys presented (Sampaio 1935; MacBride 1961; Gentry 1982; Gentry 2009).

A character that might help telling these species apart is the inflorescence structure: a thyrses or panicle in *M. iquitoensis* versus a raceme in *M. obovata*. This character, combined with leaflet base, is here proposed as diagnostic for each species; however, the examination of additional material is necessary to validate its usefulness and consistency.

3. *Martinella obovata* (Kunth) Bureau & K.Schum., in Mart., Fl. Bras. 8(2): 161, tab. 84. 1896.

http://species-id.net/wiki/Martinella_obovata

Spathodea obovata Kunth, Nov. Gen. Sp. (quarto ed.) 3: 147. 1818. [1819].

Bignonia obovata (Kunth) Spreng., Syst. Veg. 2: 830. 1825.

Macfadyena obovata (Kunth) Miers, Proc. Roy. Hort. Soc. London 3: 200. 1863.

Type. COLOMBIA. Turbaco, s.d., *Humboldt and Bonpland 1391* (holotype, P-Bonpl. [P00670823]!)

Distribution and habitat. This species is found from Central America and the Antilles through northern South America to the southern Amazon basin, in the Antilles, Belize, Bolivia, Brazil, Colombia, Costa Rica, Ecuador, French Guiana, Guatemala, Guyana, Honduras, Mexico, Panama, Peru, Surinam, and Venezuela, in different soil types and habitats (Lohmann and Taylor 2014; Fig 3).

Conservation status. *Martinella obovata* is distributed geographically through an area that is ≥ 2000 km², with < 20% of its known individuals occurring outside Protected Areas, and seventy-two Rapoport (1982) sub-populations known to date. Therefore, this species is here considered as Least Concern [LC] according to the IUCN criteria (IUCN 2012; IUCN Standards and Petitions Subcommittee 2014).

Discussion. This species is the most variable and has the largest distribution in the genus (Central America and Caribbean throughout Amazon basin). Such variation is

product of its phenotypic plasticity and wide ecological range, and is responsible for making this species hard to be distinguished from *M. iquitoensis* (as discussed above). Nonetheless, this high variation may also be seen as an evidence of a species complex, in which case, additional studies would be needed. For a complete list of synonyms see Gentry (1977).

Acknowledgments

We thank CNPq for an undergraduate fellowship to ARZ (113717/2006-7) and a Pq-1 grant to LGL; FAPESP for a research grant to LGL (2011/50859-2), two FAPESP/BIOTA-NSF/Dimensions of Biodiversity Grants (2012/50260-6 and 2013/50297-0), and Vale S.A. for financial and logistic support. We also thank Dr. Charlotte Taylor for her review and significant comments on this manuscript; Iván Jiménez, Seema Sheth and Trisha Consiglio for assistance with the establishment of the conservation status of taxa; the editor and anonymous reviewers for their constructive remarks; Dr. Rafaela Forzza (RB) and Dr. Adriana Lobão (UFF) for providing images of specimens; Klei Sousa for the illustration and Geovane Siqueira for assistance during field work.

References

- Baillon HE (1888) Bignoniaceae. In: Histoire des plantes 10(53). Librairie Hachette & Co., Paris, 1–58.
- Gentry AH (1974) Coevolutionary patterns in Central American Bignoniaceae. *Annals of the Missouri Botanical Garden* 61(3): 728–759. <http://www.jstor.org/stable/2395026>, doi: 10.2307/2395026
- Gentry AH (1977) Bignoniaceae. In: Flora of Ecuador (7). University of Gothenburg, Stockholm, 1–173.
- Gentry AH (1982) Bignoniaceae. In: Flora de Venezuela 8(4). Fondo Editorial Acta Científica Venezolana, Caracas, 1–433.
- Gentry AH (2009) Bignoniaceae. In: Flora de Colombia 25. Universidad Nacional de Colombia, Bogotá, 1–462.
- IUCN (2012) IUCN Red List Categories and Criteria: Version 3.1. Second edition. IUCN, Gland, Switzerland and Cambridge, UK.
- IUCN Standards and Petitions Subcommittee (2014) Guidelines for using the IUCN red list categories and criteria. Version 11. Prepared by the Standards and Petitions Subcommittee. <http://www.iucnredlist.org/documents/RedListGuidelines.pdf>
- Lohmann LG (2006) Untangling the phylogeny of neotropical lianas (Bignoniaceae, Bignoniaceae). *American Journal of Botany* 93(2): 304–318. doi: 10.3732/ajb.93.2.304
- Lohmann LG, Bell CD, Calió MF, Winkworth, RC (2013) Pattern and timing of biogeographical history in the Neotropical tribe Bignoniaceae (Bignoniaceae). *Botanical Journal of the Linnean Society* 171(1): 154–170. doi: 10.1111/j.1095-8339.2012.01311.x

- Lohmann LG, Ulloa Ulloa C (2013) Bignoniaceae. In: iPlants prototype checklist. <http://www.iplants.com> [accessed 17.08.2013]
- Lohmann LG, Taylor CM (2014) A new generic classification of Bignoniaceae (Bignoniaceae) based on molecular phylogenetic data and morphological synapomorphies. *Annals of the Missouri Botanical Garden* 99(3): 348–489.
- MacBride JF (1961) Bignoniaceae. In: *Flora of Peru*. Publications of the Field Museum of Natural History, Botany Series 13 (5C / 1): 3–101.
- Nogueira A, El Ottra JHL, Guimarães E, Machado SR, Lohmann LG (2013) Trichome structure and evolution in Neotropical lianas. *Annals of Botany* 112: 1331–1350. doi: 10.1093/aob/mct201
- Radford AE, Dickison WC, Massey JR, Bell CR (1974) *Vascular Plant Systematics*. Harper-Collins, New York, USA.
- Rapoport EH (1982) *Aerography: Geographical Strategies of Species*. Pergamon, New York, USA.
- Sampaio AJ (1935) Novas especies de Bignoniaceas. *Annaes da Academia Brasileira de Ciencias* 7(2): 111–127.
- Thiers B (2013) *Index Herbariorum: A global directory of public herbaria and associated staff*. New York Botanical Garden's Virtual Herbarium. <http://sweetgum.nybg.org/ih/>