

# *Caloneis nanyiensis* sp. nov. (Bacillariophyta) from Nanyi Lake of Anhui Province, China

Yong Fang<sup>1</sup>, Pan Yu<sup>2,3</sup>, Qing-Min You<sup>1</sup>, John Patrick Kociolek<sup>4</sup>, Yue Cao<sup>5</sup>, Quan-Xi Wang<sup>1</sup>, Zhi-Ping Qian<sup>1</sup>

<sup>1</sup> College of Life Sciences, Shanghai Normal University, Shanghai 200234, China

<sup>2</sup> School of Environment and Geographical Sciences, Shanghai Normal University, Shanghai 200234, China

<sup>3</sup> Yangtze River Delta Urban Wetland Ecosystem National Field Scientific Observation and Research Station, Shanghai 200234, China

<sup>4</sup> Museum of Natural History and Department of Ecology and Evolutionary Biology, University of Colorado, UCB 218, Boulder CO80309, USA

<sup>5</sup> Institute of Science and Technology, Shanghai Institute of Technology, Shanghai 201418, China

Corresponding author: Zhi-Ping Qian (qzp0313@shnu.edu.cn)

## Abstract

A new species, *Caloneis nanyiensis* sp. nov., is described from Nanyi Lake, the largest lake in southern Anhui Province, China. Observations were made using a light and scanning electron microscope documenting the size, shape and ultrastructure of the new species. *Caloneis nanyiensis* sp. nov. has rhombic valves with acutely rounded apices. The raphe is narrow and arched. An internal axial plate covers the alveoli, leaving small marginal openings bordered by costa, thickened and raised from the valve face. The striae are slightly radiate to parallel in the central area, becoming radiate towards the apices. By comparing the new species with similar *Caloneis* species, *C. nanyiensis* sp. nov. was confirmed to be sufficiently different with respect to valve size and striae density to be recognised as new to science. The new species lives in freshwater habitats and epiphytic on *Cladophora*. The discovery enhances our knowledge of the diversity of freshwater diatoms in China.

**Key words:** Axial area, *Caloneis*, girdle, raphe, rhombic valve



Academic editor: Bing Liu

Received: 31 December 2024

Accepted: 10 March 2025

Published: 17 April 2025

**Citation:** Fang Y, Yu P, You Q-M, Kociolek JP, Cao Y, Wang Q-X, Qian Z-P (2025) *Caloneis nanyiensis* sp. nov. (Bacillariophyta) from Nanyi Lake of Anhui Province, China. *PhytoKeys* 255: 113–122. <https://doi.org/10.3897/phytokeys.255.145664>

Copyright: © Yong Fang et al.

This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0).

## Introduction

Cleve and Grove initially described the genus *Caloneis* at the subgroup level, originally within the genus *Navicula* (Cleve and Grove 1891). Unfortunately, they did not provide a valid description; however, a few years later, Cleve provided a valid generic description in the *Synopsis of the Naviculoid Diatoms* (Cleve 1894). The type species of the genus *Caloneis* was indicated by Boyer (1927) to be *Caloneis amphisbaena* (Bory) Cleve (1894). The primary diagnostic features of this genus include valves that are linear-lanceolate to elliptical with capitate or rostrate ends. The shapes of axial and central areas are variable. Distal raphe fissures are usually distinct. The chambered striae give the appearance of one to two longitudinal lines. The striae of *Caloneis* are composed of fine alveoli. The outer wall of the alveoli is perforated by many rows of small, round poroids occluded by hymens and the inner wall of each alveolus typically opens to the inside of the valve by one fairly large, transapically elongate areola (Round et

al. 1990). *Caloneis* is considered a widely distributed genus with a very broad ecological range, including freshwater, brackish and marine environments (Levkov and Williams 2014). To date, 392 taxonomically accepted species of *Caloneis* are listed in AlgaeBase (Guiry and Guiry 2025). After 2000, several new species have been reported from various localities from around the world (Lange-Bertalot et al. 2004; Metzeltin et al. 2005; Metzeltin and Lange-Bertalot 2007; Levkov and Williams 2014).

Traditionally, *Caloneis* and *Pinnularia* Ehrenberg have been considered distinct genera, the diagnostic features between the two genera mainly including the genus *Caloneis* having alveolate striae that are usually thinner and denser than those of *Pinnularia* (Hustedt 1935; Round et al. 1990; Mann 2001; Levkov and Williams 2014; Kulikovskiy et al. 2023). Molecular studies conducted thus far indicate that both the genus *Pinnularia* and the genus *Caloneis* are not monophyletic (Bruder et al. 2008; Souffreau et al. 2011; Kulikovskiy et al. 2023). These findings suggest that the genus *Pinnularia* may require further subdivision, which remains unresolved. With the in-depth study of molecular systematics, we believe that the taxonomic status of the genera *Pinnularia* and *Caloneis* will be more accurately and clearly divided.

In China, the vast majority of new *Caloneis* species were published before 2000, including *Caloneis bacillum* f. *latilanceolatum* Zhu and Chen (1995), *Caloneis chansiensis* Skvortzov (1935), *C. elongata* var. *constricta* Cheng and Chin (1980), *Caloneis fasciata* var. *pekinensis* Skvortzov (1928), *Caloneis holsstii* var. *tibetica* Jao (1964), *Caloneis hunanensis* Chen and Zhu (Zhu and Chen 1989), *Caloneis lepidula* var. *angustata* Skvortzov (1976), *Caloneis patagonica* var. *sinica* Skvortzov (1938a), *C. platycephala* Cheng and Chin (1980), *Caloneis schroderi* var. *densestriata* Skvortzov (1976), *Caloneis schumanniana* var. *biconstricta* f. *minor* Zhu and Chen (1995), *Caloneis silicula* var. *hankensis* Skvortzov (1929), *Caloneis schumanniana* f. *gracilis* Skvortzov (1935), *Caloneis silicula* var. *hinganica* Skvortzov (1976) and *Caloneis sphagnicola* Skvortzov (1938b). After 2000, only one taxon has been reported in this genus: *Caloneis clevei* var. *parallela* Skvortzov ex Gololobova and Kulikovskiy (Skvortzov 2012).

During the investigation of freshwater diatom diversity in Nanyi Lake, we discovered a new species, which is described here as *Caloneis nanyiensis* sp. nov. The purpose of this study was to document and formally describe the species, based on both light microscope (LM) and scanning electron microscope (SEM) observations and to compare it with other morphologically similar species of the genus.

## Material and methods

Diatom samples were collected from Nanyi Lake (31°01'–31°10'N, 118°50'–119°3'E), Anhui Province, China, in August 2018. The lake area is 210 km<sup>2</sup>. Nanyi Lake was formed by the differentiation of the ancient Danyang Lake. The ancient Danyang Lake system was a stagnant lake formed by long-term siltation and water accumulation in a newly-constructed fault depression. The climate of the Nanyi Lake area, as reported by the Langxi County meteorological station, belongs to the North subtropical monsoon humid climate zone. The main features are: mild climate, four distinct seasons, hot and rainy season, abundant rainfall and ample sunshine (Jia et al. 2021).

In the field, several water chemistry characteristics were recorded, including: pH, temperature, salinity, total dissolved solids (TDS) and conductivity. These parameters were measured using a YSIPro Plus multiparameter meter (YSI, Ohio, USA). In the field, collections of attached algae were scraped from the surfaces of stones using (sterilised?) toothbrushes and/or a knife and the samples were placed in a bottle, preserved with formalin (4% final concentration) and sealed.

In the laboratory, the diatom valves were cleaned of organic matter using the Microwave Accelerated Reaction System (Model MARS, CEM Corporation, USA) (Parr et al. 2004). The digestion followed a pre-programmed digestion scheme (temperature: 180 °C, ramp: 15 min, hold: 15 min) (Yu et al. 2019). After digestion, samples were alternatively centrifuged (5 minutes at 3500 rpm) and washed with distilled water (approximately five times) until the pH of the sample was approximately neutral. The cleaned material was kept in 95% ethanol. Cleaned diatom specimens were mounted on glass slides in Naphrax for light microscopy (LM) or air-dried on to coverslips and mounted on to Cu stubs for observation with a scanning electron microscope (SEM). LM studies were made with a ZEISS AXIO Imager A2 microscope fitted with DIC optics and a 1.4 numerical aperture, 100× oil immersion objective. SEM examination was conducted using a Hitachi SU-8010 (2 kV, working distance less than 6 mm). Images were compiled with Adobe Photoshop CS6. Morphological terminology follows Round et al. (1990). Both unprepared (field) samples and prepared slides of mounted material are housed in the Laboratory of Algae and Environment, Department of Biology, Shanghai Normal University (SHTU).

## Results

### *Caloneis nanyiensis* Pan Yu & Qing-Min You, sp. nov.

Figs 1–3


**Type material. Holotype.** Specimen circled on slide NYH-20180801 (= Fig. 1A), deposited in the Herbarium of Shanghai Normal University (SHTU), China.

**Isotype.** Specimen circled on slide 652047 (= Fig. 1D), deposited in the Herbarium of University of Colorado, Boulder, USA.

**Type locality.** China. Anhui Province: Nanyi Lake, sampling site with the coordinates 31°01'N, 118°50'E. Diatom samples collected by Pan Yu, 19 August 2018.

**Description. LM** (Fig. 1A–M): Valve length 14.5–28.5 µm, width 8.0–10.2 µm. Valves rhombical in shape with acutely rounded apices. Raphe narrow and arched, with slightly unilaterally bent central pores and slightly curved terminal fissures. Axial area linear-lanceolate, becoming wider at the central area. Striae are slightly radiate to parallel in the central area, becoming more radiate towards the apices, 16–18 in 10 µm.

**SEM valve exterior** (Fig. 2A–E): Valve face usually uneven, with slightly raised ends and a slightly concave middle (Fig. 2A, B). Central area nearly rhombic, with a width accounting for 1/4–1/3 of the valve, the striae pattern in the central area is obviously shorter. (Fig. 2A, B, E). Raphe branches arched with proximal raphe ends weakly unilaterally deflected and dilated, drop-like in shape (Fig. 2A, B, E). Distal raphe fissures hooked and continuing on to the mantle (Fig. 2C, D). Striae multiseriate, composed of 4–5 rows of small, rounded areolae, which




**Figure 1.** *Caloneis nanyiensis* sp. nov. LM  $\times 1000$  **A–M** thirteen valves showing a size diminution series; note that the valves rhombic and raphe narrow and arched. **A** Illustration of holotype specimen **D** illustration of isotype specimen. Scale bars: 10  $\mu\text{m}$  (**M**).

are occluded by heavy silicified, perforated hymenes (Fig. 2D, E). Girdle with a single row of linear poroids (Fig. 2A).

**SEM valve interior** (Fig. 3A–E): The raphe is filiform and slightly arched (Fig. 3A). The proximal raphe endings are deflected to the same sides (Fig. 3B, C) and curve towards a central nodule that is positioned to one side of the central area. The raphe terminates distally as an elevated helictoglossa (Fig. 3D, E). An axial plate covers the alveoli, leaving small marginal openings bordered by costae thickened and raised from the valve face (Fig. 3A, B). Striae composed of 2–4 areolae are present on the mantle (Fig. 3D, E).


**Etymology.** The new species is named after the type locality, Nanyi Lake.

**Distribution and ecology.** So far, the new species has only been collected at the type locality in Nanyi Lake. The habitat of the new species is characterised by pH 8.1, water temperature 30.2  $^{\circ}\text{C}$ , TDS 0.204  $\text{g}\cdot\text{l}^{-1}$ , conductivity 175.7  $\mu\text{s}\cdot\text{cm}^{-1}$ , collected in one sample (NYH–20180801) on *Cladophora*. In the type sample, this new species occurred at less than 2% relative abundance, established from a total count of 400 valves. Amongst the species co-occurring with *Caloneis nanyiensis* sp. nov., only four species accounted for more than 5% of



**Figure 2.** *Caloneis nanyiensis* sp. nov. SEM views of the external valve **A, B** external view of an entire valve note the raphe narrow and arched **C** details of the apices on the external valve; note the distal raphe fissures hooked and continuing on to the mantle **D** details of the striae on the external valve; note the Striae multiseriate, composed of small rounded areolae and areolae are occluded by heavy silicified, perforated hymenes **E** showing the central area. Scale bars: 5 µm (**A, B**); 2 µm (**C, E**); 1 µm (**D**).

the assemblage: *Fragilaria pararumpens* Lange-Bertalot, Hofmann & Werum (Hofmann et al. 2013) (25.75%), *Achnantheidium jackii* Rabenhorst (Rabenhorst 1861) (18.75%), *Aulacoseira granulata* (Ehrenberg) Simonsen (Simonsen 1979) (11.5%) and *Encyonopsis microcephala* (Grunow) Krammer (Krammer 1997) (%).



**Figure 3.** *Caloneis nanyiensis* sp. nov. SEM views of valve interior. **A** Internal view of an entire valve; note the raphe is filiform and slightly arched **B, C** details of the proximal raphe endings which are deflected to the same side **D, E** details of the distal raphe ends terminating as helictoglossae and the striae are composed of 2–4 areolae on the mantle. Scale bars: 5  $\mu\text{m}$  (**A**); 2  $\mu\text{m}$  (**B, C**); 1  $\mu\text{m}$  (**D, E**).

### Discussion

The new species described in this study possesses all the ultrastructural features of the genus *Caloneis* (Round et al. 1990), including having a narrow, slightly arched raphe with slightly unilaterally bent central pores and slightly

curved terminal fissures, an axial plate covering the alveoli and striae parallel in the central area, becoming radiate towards the apices.

*Caloneis nanyiensis* sp. nov. can be compared to several species in the same genus, based on similarities in the outline and structure of the valve, including *C. caribeana* Metzeltin & Lange-Bertalot (Metzeltin and Lange-Bertalot 2007), *C. permagna* (Bailey) Cleve (Cleve 1894) and *Caloneis schumanniana* var. *lancettula* Hustedt (Hustedt 1930). The morphological characteristics of *C. nanyiensis* and these similar species are summarised in Table 1 to facilitate their comparison. The outline of the valves of *C. nanyiensis* is rhombical with acutely rounded ends, while that of *C. caribeana* is rhombical to rhombic-lanceolate with acutely rounded ends, *C. permagna* is rhombic-lanceolate with acutely rounded ends and *C. schumanniana* var. *lancettula* is narrow-lanceolate with broadly rounded ends. Additionally, the valves of *C. nanyiensis* are shorter (14.5–28.5 µm) than *C. permagna* (85–220 µm), *C. caribeana* (32–57 µm) and *C. schumanniana* var. *lancettula* (35–40 µm), as well as narrower (8.0–10.2 µm in *C. nanyiensis*) versus the breadth of valves in *C. permagna* (35–55 µm) and *C. caribeana* (14–16 µm). Furthermore, no central area was observed in the new species, but *C. caribeana* has a narrow fascia central area, *C. permagna* has an irregularly lanceolate central area and *C. schumanniana* var. *lancettula* has a transapically rectangular central area. The stria density of *C. nanyiensis* is higher (16–18/10 µm) than that of *C. permagna* (9–12/10 µm), while being lower than those of the other similar species. Individuals in the populations of *C. nanyiensis* examined here have a valve outline that is rhombical in shape, which, together with the slightly curved raphe, help to differentiate it from other species in the genus.

We also compared our new species with the smaller valves of *C. distinguenda* Levkov & Williams (Levkov and Williams 2014). However, the latter taxon is characterised by elliptic-lanceolate valves being 36 µm long and 15 µm wide. Additionally, the central area in smaller specimens of *C. distinguenda* is distinctly separated from axial area and transversally elliptic in shape. The density of the striae in smaller specimens of *C. distinguenda* is lower (14–16/10 µm) than that of *C. nanyiensis*.

*Caloneis* are common in alkaline, brackish and marine habitats, species of the genus *Caloneis* having a broad ecological niche (Levkov and Williams 2014).

**Table 1.** Comparison of morphological characteristics of *Caloneis nanyiensis* sp. nov. and closely-related taxa.

| Species/Feature           | <i>C. nanyiensis</i> sp. nov. | <i>C. caribeana</i> Metzeltin & Lange-Bertalot | <i>C. permagna</i> (Bailey) Cleve               | <i>C. schumanniana</i> var. <i>lancettula</i> Hustedt |
|---------------------------|-------------------------------|--|---|---|
| Valve length (µm)         | 14.5–28.5                     | 32–57  | 85–220  | 35–40   |
| Valve width (µm)          | 8.0–10.2                      | 14–16  | 35–55   | 8   |
| Valve outline             | Rhombical                     | Rhombical to rhombic-lanceolate                | Rhombical-lanceolate                            | Narrow-lanceolate                                     |
| Valve apices              | Acutely round                 | Acutely round                                  | Acutely round                                   | Broadly rounded                                       |
| Axial area                | Linear to lanceolate          | Narrow, linear                                 | Irregularly lanceolate                          | Narrow-lanceolate                                     |
| Central area              | Absent                        | Very narrow fascia                             | Irregularly lanceolate                          | Transapically rectangular                             |
| Raphe                     | Narrow and arched             | Distinctly curved                              | Straight  | Straight  |
| Density of striae (10 µm) | 16–18                         | 17–20  | 9–12  | 17–20   |
| References                | Current study                 | Metzeltin and Lange-Bertalot (2007)            | Cleve (1894), Krammer and Lange-Bertalot (1986) | Hustedt (1930), Krammer and Lange-Bertalot (1986)     |

For example, *Caloneis* is diverse in cold-water oligotrophic habitats, *C. limosa* prefers to inhabit alkaliphilic, oligotrophic and up to  $\beta$ -mesosaprobic water bodies, and *C. schumanniana* is classified as a planktonic-benthic, oligo-xenosaprobic and alkaliphilic species (Denys 1991; Stancheva et al. 2009; Levkov and Williams 2014). In the present study, *Caloneis nanyiensis* has only been found on *Cladophora* in Nanyi Lake. The water quality of Nanyi Lake reflects a moderate nutrient level. This investigation did not detect the presence of this novel species in any other location. It is postulated that this species may be a good indicator of water bodies with moderate nutrient levels. However, further ecological studies are needed to confirm this hypothesis.

## Acknowledgements

We would like to thank Lixuan Zhang, Linxin Lu, Bingwei Xing and Yang Yu for help in the field and in the preparation of samples for microscopy. We extend our heartfelt gratitude to the reviewer for the invaluable guidance and support provided. The insightful recommendations have been instrumental in enhancing the quality of this manuscript and we are deeply appreciative of your professional expertise and dedication.

## Additional information

### Conflict of interest

The authors have declared that no competing interests exist.

### Ethical statement

No ethical statement was reported.

### Funding

This research was funded and supported by National Natural Science Foundation of China (No. 32100165, 32170205).

### Author contributions

Formal analysis: YF, PY. Investigation: YC, QMY. Project administration: QXW, Writing – original draft: YF. Writing – review and editing: ZPQ, JPK, PY.

### Author ORCIDs

Pan Yu  <https://orcid.org/0000-0001-7937-2381>

John Patrick Kociolek  <https://orcid.org/0000-0001-9824-7164>

Zhi-Ping Qian  <https://orcid.org/0009-0008-7514-8073>

### Data availability

All of the data that support the findings of this study are available in the main text.

## References

Boyer CS (1927) Synopsis of North American Diatomaceae, Supplement, Part 2. Naviculatae, Surirellatae. Proceedings. Academy of Natural Sciences of Philadelphia 79: 229–583.



- Bruder K, Sato S, Medlin LK (2008) Morphological and molecular investigations of naviculoid diatoms IV. *Pinnularia* vs *Caloneis*. *Diatom* 24: 8–24.
- Cheng Z, Chin TG (1980) Notes on some new species and new records of diatoms from the Strait of Taiwan (Fujian Coast), China. V. *Coscinodiscus*, *Caloneis*. *Universitatis Amoiensis, Acta Scientiarum Naturalium* 19(3): 109–116, 126.
- Cleve PT (1894) Synopsis of the naviculoid diatoms. Part I. *Kongliga Svenska Vetenskapsakademiens Handlingar Series* 4 26(2): 1–194.
- Cleve PT, Grove E (1891) Sur quelques Diatomées nouvelles ou peu connues. *Le Diatomiste* 1: 64–68.
- Denys L (1991) A check-list of the diatoms in the Holocene deposits of the western Belgian coastal plain with a survey of their apparent ecological requirements. *Professional Paper Belgische Dienst* 246: 1–41.
- Guiry MD, Guiry GM (2025) *AlgaeBase*. World-wide electronic publication, National University of Ireland, Galway. <https://www.algaebase.org> [accessed 04 March 2025]
- Hofmann G, Werum M, Lange-Bertalot H (2013) *Diatomeen im Süßwasser-Benthos von Mitteleuropa. Bestimmungsflora Kieselalgen für die ökologische Praxis. Über 700 der häufigsten Arten und ihre Ökologie*. Koeltz Scientific Books, Königstein, 908 pp.
- Hustedt F (1930) *Bacillariophyta (Diatomeae) Zweite Auflage*. In: Pascher A (Ed.) *Die Süßwasser-Flora Mitteleuropas*. Heft 10. Verlag von Gustav Fischer, Jena, 466 pp.
- Hustedt F (1935) Untersuchungen über den Bau der Diatomeen, X und XI. *Berichte der Deutschen Botanischen Gesellschaft* 53(1): 3–41. <https://doi.org/10.1111/j.1438-8677.1935.tb01863.x>
- Jao C (1964) Some fresh-water algae from southern Tibet. *Oceanologia et Limnologia Sinica* 6(2): 169–192.
- Jia RY, Wang H, Li XX, Yang YY, Zou ZH, Wu LC, Liang MD, Yi R, Zhou Z, Xia QP (2021) Plant Investigation of Nanyi Lake Wetland in Xuancheng District, Anhui Province. *Bulletin of Biology* 56(5): 10–13.
- Krammer K (1997) *Die cymbelloiden Diatomeen. Eine Monographie der weltweit bekannten Taxa. Teil 2. Encyonema Part., Encyonopsis und Cymbellopsis*. *Bibliotheca Diatomologica* 37: 1–469.
- Krammer K, Lange-Bertalot H (1986). *Bacillariophyceae. 1. Teil: Naviculaceae* In: Ettl H, Gerloff J, Heynig H, Mollenhauer D (Eds) *Süßwasserflora von Mitteleuropa*, Band 2/1. Gustav Fisher Verlag, Jena, 876 pp.
- Kulikovskiy M, Glushchenko A, Kezlya E, Kuznetsova I, Kocielek JP, Maltsev Y (2023) The genus *Pinnularia* Ehrenberg (Bacillariophyta) from the Transbaikal Area (Russia, Siberia): Description of seven new species on the basis of morphology and molecular data with discussion of the phylogenetic position of *Caloneis*. *Plants* 12(20): 3552. <https://doi.org/10.3390/plants12203552>
- Lange-Bertalot H, Genkal SI, Vekhov NV (2004) New freshwater species of Bacillariophyta. *Biologia Vnutrennikh* 4: 12–17.
- Levkov Z, Williams DM (2014) Observations on *Caloneis* Cleve (Bacillariophyceae) species from the ancient lakes Ohrid and Prespa. *Nova Hedwigia. Beiheft* 143: 141–158. <https://doi.org/10.1127/1438-9134/2014/008>
- Mann DG (2001) A discussion of *Caloneis* and related genera. *Diatom* 17: 29–36. [https://doi.org/10.11464/diatom1985.17.0\\_29](https://doi.org/10.11464/diatom1985.17.0_29)
- Metzeltin D, Lange-Bertalot H (2007) Tropical diatoms of South America II. Special remarks on biogeography disjunction. *Iconographia Diatomologica* 18: 1–877.

- Metzeltin D, Lange-Bertalot H, García-Rodríguez F (2005) Diatoms of Uruguay. Compared with other taxa from South America and elsewhere. *Iconographia Diatomologica* 15: 1–736.
- Parr JF, Taffs KH, Lane CM (2004) A microwave digestion technique for the extraction of fossil diatoms from coastal lake and swamp sediments. *Journal of Paleolimnology* 31(3): 383–390. <https://doi.org/10.1023/B:JOPL.0000021857.32734.c6>
- Rabenhorst L (1861) *Die Algen Europas, Fortsetzung der Algen Sachsens, resp. Mittel-Europas*. Dresden Decades I-CIX.
- Round FE, Crawford RM, Mann DG (1990) *The Diatoms: Biology and Morphology of the Genera*. Cambridge University Press, Cambridge, 747 pp.
- Simonsen R (1979) The diatom system: Ideas on phylogeny. *Bacillaria* 2: 9–71.
- Skvortzov BV (1935) Diatomées récoltées par le Père E. Licent au cours de ses voyages dans le Nord de la Chine au bas Tibet, en Mongolie et en Mandjourie. *Publications du Musée Hoangho Paiho de Tien Tsin* 36: 1–43.
- Skvortzov BV (2012) New and little known fresh-and brackish water diatoms chiefly from the Eastern part of Asia and their geographical distribution. In: Kulikovskiy MS, Lange-Bertalot H, Metzeltin D, Witkowski A (Eds) *Lake Baikal: hotspot of endemic diatoms I*. *Iconographia Diatomologica* vol. 23, A.R.G. Gantner Verlag K.G., Ruggell, 749–861.
- Skvortzov BV (1928) Diatoms from ponds of Peking. *The Peking Society of Natural History Bulletin* 3: 43–48.
- Skvortzov BV (1929) A Contribution to the Algae, Primorsk District of Far East, U.S.S.R. Diatoms of Hanka Lake. *Memoirs of the Southern Ussuri Branch of the State Russian Geographical Society, Vladivostok*, 66 pp.
- Skvortzov BV (1938a) Diatoms from Chengtu, Szechwan, Western China. *Philippine Journal of Science* 66(4): 479–496.
- Skvortzov BV (1976) Moss diatom flora from River Gan in the northern part of Great Khingan Mountains, China, with description of a new genera *Porosularia* gen. nov. from Northern and Southern China. (First Part). *Quarterly Journal of the Taiwan Museum* 29(1–2): 111–152.
- Skvortzov BV (1938b) Diatoms from a mountain bog, Kaolingtze, Pinchiang-Sheng Province, Manchoukuo. *Philippine Journal of Science* 66(3): 343–362.
- Souffreau C, Verbruggen H, Wolfe AP, Vanormelingen P, Siver PA, Cox EJ, Mann DG, Van de Vijver B, Sabbe K, Vyverman W (2011) A time-calibrated multi-gene phylogeny of the diatom genus *Pinnularia*. *Molecular Phylogenetics and Evolution* 61(3): 866–879. <https://doi.org/10.1016/j.ympev.2011.08.031>
- Stancheva R, Manoylov K, Gillett N (2009) Morphological variation of the *Caloneis schumanniana* species complex (Bacillariophyceae) from different environmental conditions in North American streams. *Hydrobiologia* 635(1): 157–170. <https://doi.org/10.1007/s10750-009-9908-4>
- Yu P, You QM, Kociolek JP, Wang QX (2019) Three new freshwater species of the genus *Achnantheidium* (Bacillariophyta, Achnanthidiaceae) from Taiping Lake, China. *Fottea* 19(1): 33–49. <https://doi.org/10.5507/fot.2018.015>
- Zhu HZ, Chen JY (1989) *New species and varieties of diatom from Suoxiyu*. Science Press, Beijing, 33–37.
- Zhu HZ, Chen JY (1995) New taxa of diatom (Bacillariophyta) from Xizang (Tibet) (I). *Zhiwu Fenlei Xuebao* 33: 516–519.