

A new species of *Piper* (Piperaceae) with peltate leaves from Serranía de las Quinchas, Colombia

M. Alejandra Jaramillo¹, Dayro Rodríguez-Duque², Magda Escobar-Alba²

¹ Grupo Diversitas, Facultad de Ciencias Básicas y Aplicadas, Universidad Militar Nueva Granada, km 2 Vía Cajicá-Zipacquirá, Cajicá, Colombia

² Grupo de Investigación Biodiversidad y Conservación, Museo de Historia Natural “Luis Gonzalo Andrade”, Facultad de Ciencias, Universidad Pedagógica y Tecnológica de Colombia-Universidad Nacional de Colombia, Av. Central del Norte 39-115, Tunja-Boyacá, Tunja, Colombia

Corresponding author: M. Alejandra Jaramillo (maria.jaramillo@unimilitar.edu.co)

Abstract

Piper quinchasense is described and illustrated as a new species occurring in the understory of wet montane forest of the middle Magdalena Valley in Colombia, the easternmost portion of the Chocó Region. Its relationships are discussed with related taxa from the *Macrostachys* clade. An identification key for 35 Neotropical *Piper* species with peltate leaves is provided.

Key words: Boyacá, Chocó Region, *Macrostachys* clade, Piperales, tropical montane forests

Resumen

Piper quinchasense se describe e ilustra como una nueva especie que ocurre en el sotobosque de bosques húmedos montaños del valle medio del Magdalena en Colombia, la porción más oriental del Chocó Biogeográfico. Se discuten sus relaciones con otras especies del clado *Macrostachys*. Se presenta una clave de identificación para 35 especies de *Piper* Neotropical con hojas peltadas.

Palabras claves: bosque montano tropical, Boyacá, Chocó Biogeográfico, clado *Macrostachys*, Piperales



Academic editor: Elton John de Lirio

Received: 3 February 2023

Accepted: 5 May 2023

Published: 29 May 2023

Citation: Jaramillo MA, Rodríguez-Duque D, Escobar-Alba M (2023) A new species of *Piper* (Piperaceae) with peltate leaves from Serranía de las Quinchas, Colombia. *PhytoKeys* 227: 9–24, <https://doi.org/10.3897/phytokeys.227.101405>

Copyright: © M. Alejandra Jaramillo et al. This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0).

Introduction

Piper, with more than 2000 species (Callejas-Posada 2020), is one of the most species-rich genera among flowering plants (Frodin 2004). *Piper* is also a common element in the understory of Neotropical forests (Gentry 1990; Draper et al. 2021). Species of *Piper* are a key resource for bats of the genus *Carollia* (Fleming 2004; Bohlender et al. 2018; Yohe et al. 2021), and they are a critical element that supports diverse trophic networks that involve moths and parasitoid wasps (Wilson et al. 2012; Slinn et al. 2018). *Piper*'s enormous diversity of secondary metabolites is critical for their coexistence (Salazar et al. 2016a, 2016b) and diversification (Massad et al. 2022). And it is also of immense interest in the pharmaceutical industry (Perez Gutierrez et al. 2013; Salehi et al. 2019). This genus is, without doubt, an essential structural and trophic element

of the understory and lower strata of Neotropical forests (Salazar et al. 2016b; Draper et al. 2021).

Molecular phylogenetic studies have been instrumental in reviving the infrageneric classification of the genus, but also identifying convergence in morphological traits (Jaramillo and Callejas-Posada 2004; Trujillo et al. 2022). Molecular phylogenetics (Jaramillo and Manos 2001; Jaramillo et al. 2008) have validated the monophyly of groupings proposed in the mid-1800s (Kunth 1839; Miquel 1843). The infrageneric classification was not used during the 1900s, probably as the expeditions in the early 1900s produced too many species to classify. Today we know that *Piper* clades and subclades are easy to recognize with a combination of key morphological characters (Jaramillo et al. 2008). A formal infrageneric classification of *Piper* based on phylogeny is under preparation (Callejas pers. comm.). Molecular phylogenetics have been instrumental in clarifying the relationships of *Piper* species with “axillary” inflorescences, a polyphyletic set of taxa from the Chocó Region that were wrongly merged in the genus *Trianaeopiper* Trel. (Trelease 1928). Molecular phylogenetics demonstrated that *Trianaeopiper* is not monophyletic, revealing that the diagnostic character axillary inflorescence – which are shortened sympodial branches, is convergent (Jaramillo and Callejas-Posada 2004). Similarly, molecular phylogenetics have further supported that *Piper* species with peltate leaves are part of at least four Neotropical *Piper* clades (Trujillo et al. 2022). *Piper* classification, evolution and ecological studies have been greatly enriched with molecular phylogenetics.

Identifying *Piper* species continues to be difficult for the untrained eye, and many taxa are submerged in a few large, broadly distributed, but artificial taxa (Tebbs 1990, 1993). To the trained specialist, new *Piper* species are a common finding in the tropical forests, or even in the undetermined piles in the most important herbaria around the world. Additionally, molecular phylogenetics have confirmed the clade (subgeneric) affiliation of new species (Bornstein et al. 2014; Tepe et al. 2014; Trujillo et al. 2022). During our exploration to Serranía de las Quinchas we have identified a few undescribed species, one of the stands out because it has peltate leaves, a characteristic rarely observed in the genus.

“Serranía de las Quinchas” is a small mountain spur west of the Cordillera Oriental in the middle Magdalena Valley, in Colombia. The region’s flora is particularly interesting as it combines its own floristic elements, mixed with taxa from Mesoamerica, the Chocó Region, and Amazonia (Balcázar-Vargas et al. 2000). Additionally, various endemic species and genera occur in the area (Rodríguez-Duque et al. 2021). The middle Magdalena Valley deserves close examination and more fieldwork to uncover the history of its biological richness. For centuries, the forest has been preserved because of its scarp ridges and high rainfall. However, the construction of a pipeline and road in the 1990s opened the area to colonization (Stiles and Bohórquez 2000). Today the region is threatened by the extension of agriculture and mining activities. Fortunately, Serranía de las Quinchas Regional Park was created in 2008 by Corpoboyacá (the regional environmental protection agency) to protect the remaining forest (Bohorquez-Osorio et al. 2020). Despite the deforestation in the region, new species are still being encountered. Here we describe a new species of Neotropical *Piper* with peltate leaves that we detected during our expeditions in this interesting location.

Materials and methods

Specimens were collected in Serranía de las Quinchas, located in the middle Magdalena Valley in Colombia, in the department of Boyacá. The sites visited range from 800–1200 m. a. s. l., the locality is dominated by humid montane forest (Fig. 1). Annual rainfall is 3333 mm on average; two dry seasons occur in January-February and June-August, and the average temperature is 26 °C (Instituto de Hidrología, Meteorología y Estudios Ambientales, IDEAM). Detailed observations in the field, combined with examination of available herbarium collections, allowed the description of growth habits and phenological stages. Voucher specimens were deposited in the following herbaria HUA, H-UPTC, and




Figure 1. Geographic distribution map of *Piper quinchasense* M.A.Jaram.

Table 1. Genbank accessions for the new species of *Piper*.

Taxon	Voucher	Genbank Accession #
<i>Piper quinchasense</i> M. A. Jaram.	MAJ1807	OQ354973
	MAJ1939	OQ354974

UMNG-H (herbarium acronyms according to Index Herbariorum [Thiers continuously updated], and Instituto Humboldt – Red Nacional de Colecciones [<http://rnc.humboldt.org.co/wp/>]). In addition, taxonomic literature on *Piper* was examined (Trelease and Yuncker 1950, Steyermark 1984, Callejas-Posada 2020). The measurements included in the description below are based on herbarium specimens collected by the authors. Conservation status assessments employ the categories and criteria of the IUCN (2012, 2022). We calculated the extent of occurrence and area of occupancy using the R package *ConR* (Dauby et al. 2017).

For preparing the key of Neotropical *Piper* species with peltate leaves we used the literature (Trelease and Yuncker 1950, Steyermark 1984, Callejas-Posada 2020, Carvalho-Silva et al. 2022) and type specimens were examined using digitized plant specimens available on-line from JSTOR Global Plants (<https://plants.jstor.org/>).

We extracted DNA from silica gel dried tissue, using the DNAeasy plant mini kit (Qiagen, Valencia, California, USA). The ITS region was amplified using one of two pairs of primers ITS5-ITS4, or LEU1-ITS4 (Baldwin 1992). Sequencing was contracted with GenCore (Universidad de los Andes, Bogotá, Colombia). The resulting sequences (Table 1) were manually aligned against previously obtained alignments (Jaramillo et al. 2008). We selected 43 ITS sequences from the large alignment to portray here. Three sequences from *Piper* species from Asia and the South Pacific were used as outgroups. Forty sequences of Neotropical *Piper*, comprising representatives of all clades, and 14 species with peltate leaves were selected. Maximum Likelihood (ML) phylogenetic and bootstrap (100 replicates) analyses were conducted using RAxML (Stamatakis 2014).

Taxonomic treatment

Piper quinchasense M.A.Jaram., sp. nov.

urn:lsid:ipni.org:names:77320207-1

Type. COLOMBIA. Boyacá: Otanche, vereda Las Quinchas, Sector La Y, Finca Lote Terreno, 5°48'17"N, 75°15'24"W, 1210 m, 17 Mayo 2022 [fl], M. A. Jaramillo et al. 1807 (holotype: HUA; isotypes: UPTC, UMNG-H). Figs 1–3.

Description. *Piper quinchasense* is similar to *P. parianum*, it differs from the latter in having all leaf blades peltate (vs. leaves deeply lobed to peltate), and inflorescence peduncle 4–5 cm long, (vs. peduncle 1–2.7 cm long).

Shrub, 3 m tall, branched in the upper portion only, exhibiting stilt roots (Fig. 2C). **Internodes** 5–10 × 2.5–4.6 cm, green when young and becoming brown when maturing, tomentose, idioblasts not evident. **Prophylls** not seen. **Petioles** 4.5–8.5 cm long, vaginate the entire length (Figs 2D, 3J), tomentose, idioblasts evident. **Leaf** blades (28) 35–49 × 7.5–19 cm, oblong-lanceolate, base obliquely-peltate, asymmetric to truncate, peltate, petiole inserting 2.5–10.5 cm from the margin, on the adaxial surface leaf blade is depressed and umbonate above




Figure 2. *Piper quinchaense* M. A. Jaram. **A** habit **B** magnified view of inflorescence **C** stilt roots **D** sheathing petiole **E** adaxial surface of leaves **F** abaxial surface of leaves and inflorescence. Photographs by D. Rodríguez-Duque.

petiole insertion (Fig. 2E), blade medially asymmetric, apex long attenuate, green on the adaxial surface and green-silver on the abaxial surface when alive (Fig. 2E, F), coriaceous, chartaceous when dry, drying dark maroon on the adaxial surface and ochre on the abaxial surface, glabrous with visible idioblasts on the adaxial surface (Fig. 3K), sparsely tomentose on the blade and veins densely tomentose on the abaxial surface (Fig. 3I), eciliate, margin folds towards the abaxial surface appearing to form irregular spaced teeth; pinnately nerved to the distal third, 4–6 pairs of secondary veins, curved and ascending, diverging in angles that decrease towards the apex (from 80–30 degrees) and spacing slightly decreasing towards the base, tertiary veins forming areoles 1.2–1.8 × 0.5–1 cm, rectangular and not uniform in size, perpendicular to secondary nerves, nerves not impressed on the adaxial surface, elevated on the abaxial surface. **Inflorescences** a simple spike, terminal, pendulous; peduncle 4.5–5 cm long, tomentose, green, idioblasts not evident; rachis length in flower 12–19 cm × 3–4 mm, rachis length in fruit 25–27 cm. **Floral bracts** cucullate, sagitate from above (Fig. 3F), 1.8–2.3 × 0.9–1.0 mm, pellucid dotted, pedicel fimbriate on the distal portion (Fig. 3E), forming bands around the spike (Fig. 3B). **Flowers** sessile with 4 stamens (Fig. 3C), filaments 0.25–0.40 mm long, anthers 0.3–0.6 × 0.16–0.26 long, with connective glabrate, longitudinally dehiscent, ditheous (Fig. 3D), ovary four carpellate, four stigmas sessile, not persistent in fruit, 0.4–0.5 mm long




Figure 3. *Piper quinchasense* M. A. Jaram. **A** sympodial branch, showing both the abaxial and adaxial surface of the leaves **B** magnified view of inflorescence **C** floral diagram **D** anther **E** floral bract, abaxial view **F** floral bract view from above **G** fruit **H** seed **I** magnified view of leaf abaxial surface **J** sheathing petiole **K** magnified view of leaf adaxial surface. Illustration by Ariadna Valenzuela, based on M. Escobar-Alba 764, and photographs by D. Rodríguez-Duque.

(Figs 2B, 3G). **Fruits** obpyriform, 2.1–2.4 × 1.0–1.6 mm, glabrous, green when alive, brown when dry (Fig. 3G). **Seeds** smooth, pellucid dotted, obpyriform, 1.9–2.1 × 1.0–1.3 mm, glabrous, brown when dry (Fig. 3H).

Phylogenetic relationships. *Piper quinchasense* belongs to the *Macrostachys* clade (Fig. 4). A group of shrubs and treelets reaching 7–8 (15) m tall, petioles sheathing above the middle or throughout their length, pinnately nerved leaves, mostly larger than 30cm long, inflorescences mostly pendulous (erect in some species), and flowers forming bands around the spike (Jaramillo et al. 2008). Species with peltate leaves have evolved independently in clades *Macrostachys*, *Pothomorphe*, *Oxodium* (= *Schilleria*, Callejas 2020) and *Ottonia*. Peltate leaves are known to occur in the two large genera of Piperaceae: *Piper* and *Peperomia*. They are more common in plants that grow in the shade of humid tropical forests (Wunnenberg et al. 2021). There is still much to learn about the functional morphology and anatomy of *Piper* species with peltate leaves.

Distribution and habitat. The species is only known from the type locality Serranía de las Quinchas (Fig. 1). *Piper quinchasense* is a tall shrub in the understory of very humid forests; it often grows near streams. The occurrence of stilt roots (Fig. 2C), a character not commonly seen in *Piper*, suggests it is adapted for flooded areas near watercourses. It also is a very resilient plant that will produce adventitious roots if tumbled and it will continue growing, or even resprout, from a fallen leaf. Resprouting is common among shade-tolerant *Piper* (Lasso et al. 2009).

Phenology. Flowering specimens were collected in March, and May. Fruiting specimens were collected in October.

Etymology. The epithet *quinchasense*, refers to Serranía de las Quinchas, the type locality for this species. According to locals, Quinchas derives from the indigenous groups “Quinchos” that inhabited the region.

Conservation status. This species is known only from one population in the type locality (Fig. 1). The extent of occurrence (EOO) of 8 km² and area of occupancy (AOO) of 8 km² are very small. The locality is under threat of disappearing for the extension of agricultural and mining activities (Rodríguez-Duque et al. 2021), which suggests it is Endangered [EN B1a]. Serranía de las Quinchas was declared a Regional Park in 2008; since its creation, the park extension has been reduced by 20% as the buffer zones were eliminated. Furthermore, the constant threat of coal mining makes the locality highly exposed to deforestation.

Additional specimens examined. COLOMBIA. —Boyacá: Otanche, vereda Las Quinchas, Sector La Y, Finca Lote Terreno, 5°48'17"N, 75°15'24"W, 26 October 2022, [fr] M. A. Jaramillo et al 1939 (HUA, UMNG-H); Boyacá: Otanche, Parque Regional Natural, Serranía de las Quinchas, 5°48'45.5"N, 75°15'22.2"W, [st] 14 June 2021, Magda Escobar-Alba et al. 489 (UPTC); Parque Regional Natural Serranía de las Quinchas, vereda las Quinchas, Finca Chorro Negro, 5°49'7.2"N, 75°14'57.3"W, [fl] 3 March 2022, Magda Escobar-Alba et al. 762 (UPTC).

Notes. *Piper quinchasense* is a handsome species that differs from related *Macrostachys* taxa in having long lanceolate-oblong leaves. It is morphologically similar to *P. parianum* from which it differs in having monomorphic leaves (all of them peltate) vs. leaves dimorphic, some peltate and others are deeply lobed. *P. parianum* is only known in the isolated cloud forests of “Península de Paria”, a region located in the eastern portion




Figure 4. Phylogenetic relationships of *Piper quinchasense* M. A. Jaram. and 14 species of Neotropical *Piper* with peltate leaves (in bold). Topology based on Maximum likelihood analysis of nrITS sequences. Numbers on branches are maximum likelihood bootstrap support values (>90%).

of Coastal Cordillera in the extreme northeast of Venezuela. The flora of Paria Peninsula is characterized by the high occurrence of endemic species (Steyermark and Agostini 1966; Meier 2011) and several types of vegetation communities (Beard 1946). The flora of this region has many affinities to plants of Trinidad and Tobago (*Selaginella hartii* Hieron.; – Selaginellaceae; 5 Turrill – Acanthaceae: sensu Baksh-comeau et al. 2016), and the forests

of the Northeastern sector of Guayana Shield in Venezuela and the Guianas (e.g., *Besleria insolita* C.V. Morton – Gesneriaceae; Meier 2011). Serranía de las Quinchas, on the other hand, has the most floristic affinities with the Chocó region (e. g. *Dimerocostus cryptocalyx* N. R. Salinas & Betancur, Salinas and Betancur 2004). The regions have similar habitats driven by high precipitations despite their divergent biogeographical affinities.

A key to species of Neotropical *Piper* with peltate leaves is presented below.

New status

While preparing the key for peltate Neotropical *Piper*, we realized some species deserve new status. New status and new names are proposed for two species.

***Piper neovenezuelense* M.A.Jaram., stat. nov et nom. nov.**

[urn:lsid:ipni.org:names:77320208-1](https://nomenclature.ipni.org/names/77320208-1)

Basionym. *Piper veraguense* C. DC. var. *venezuelense* Steyerem. Fl. Venez. 2(2): 590 (1984). Type: Venezuela, Edo. Trujillo, 16 km de Boconó a lo largo de la carretera a Biscucuy, 1850 m, 11 aug 1964, F. Breteler 4082 (Holotype; VEN; Isotype: MER). Non *Piper venezuelense* C.DC., J. Bot. 4: 216 (1866).

Note. The epithet *P. neovenezuelense* is proposed here to replace *Piper veraguense* C. DC. var. *venezuelense* Steyerem. Because *Piper venezuelense* C. DC. (De Candolle 1866: 216) is already in use.

Etymology. The new epithet *neovenezuelense* honors the intention of J. Steyermark to highlight the occurrence of this species in Venezuela.

Steyermark (1984) synonymized under *P. veraguense* C. DC. various distinct species: *P. albert-smithii* Trel. & Yunck., and *P. mutisii* Trel. & Yunck., and proposed that *P. veraguense* C. DC. has three varieties: *P. veraguense* var. *veraguense*, *P. veraguense* var. *mutisii* (Trel. & Yunck.) Steyermark, and he added *P. veraguense* var. *venezuelense* Steyermark (Steyermark 1984). These species are similar in their overall morphology and potentially form a species complex. Unfortunately, we do not have sequence data to test the latter hypothesis. Here we consider them separate species. *P. veraguense* has glabrous leaves with nerves puberulent on the abaxial surface, not pellucid dotted (Callejas-Posada 2020); *P. albert-smithii* is distinct because it has fleshy warts below the nodes, and leaves are glabrous (Trelease and Yuncker 1950); and *P. mutisii* has fleshy warts below the nodes, nerves are dense velvety puberulent on abaxial surface; *P. neovenezuelense* has glabrous internodes without warty outgrowths, and nerves pilose on abaxial surface.

***Piper andersii* M.A.Jaram., stat. nov. et nom. nov.**

[urn:lsid:ipni.org:names:77320209-1](https://nomenclature.ipni.org/names/77320209-1)

Piper mikanianum Steud. var. *peltatum* Yunck., Bol. Inst. Bot. (São Paulo) no. 3: 54 (1966). Type: Brazil, Minas Gerais, Caldas, A. F. Regnell II 256*, 9 Jul 1864. Non *Piper peltatum* L.

Note. The epithet *P. andersii* is proposed here to replace *Piper mikanianum* Steud. var. *peltatum* Yunck. because *Piper peltatum* L., (Sp. Pl. 1: 30 1753) is already in use.

Etymology. The new epithet *andersii* honors Anders Fredrik Regnell (1807–1884), Swedish physician and botanist who established himself in Minas Gerais (Brazil) and collected the type specimen for this species.

T. Yuncker provided a key to four varieties of *Piper mikanianum* (Kunth) Steud.: *P. mikanianum* (Kunth) Steud var. *mikanianum*; *P. mikanianum* f. *clausum* characterized by the closed sinus and overlapping lobes; *P. mikanianum* var. *pilosius* C. DC. with leaves and stems strongly pilose with hairs up to 1 mm long; and *P. mikanianum* var. *peltatum* Yunck with peltate leaves and nerves minutely hirtellous on the abaxial surface (Yuncker 1972). Yuncker had mentioned that this latter variety might deserve species rank, we propose to make the change suggested and propose the new name *P. andersii*. Furthermore, *P. mikanianum* occurs in the states of Minas Gerais, Paraná and Rio Grande do Sul in Brazil and in Argentina, *P. andersii*, with peltate leaves has only been registered from the state of Minas Gerais.

Key to Neotropical *Piper* species with peltate leaves

- 1 Leaves subpeltate, petiole inserted slightly inside the leaf margin2
- Leaves peltate, petiole inserted 0.5–11 cm from the leaf margin12
- 2 Leaves ovate-lanceolate or elliptic, up to 18 cm long, petioles terete3
- Leaves ovate, ovate –oblong or elliptic, more than 20 cm long, petioles sheathing5
- 3 Inflorescence a raceme with glabrous to sparsely pubescent rachis.....4
- Inflorescence a spike, with pubescent rachis.....
.....*P. klotzschianum* (Kunth) C. DC.
- 4 Leaves elliptic and glabrous without hirtellous intramarginal nerve
.....*P. brumadinense* M. Carv.-Silva & E. Guim.
- Leaves ovate, intramarginal nerve hirtellous on abaxial surface
.....*P. ovatum* Vahl
- 5 Petioles sheathing only at the base, nerves puberulent on the abaxial surface.....6
- Petioles sheathing over half their length, leaves and nerves glabrous or pubescent7
- 6 Leaves 35–20 cm long, base lobate *P. omega* Trel.
- Leaves 10–20 cm long, base barely cordate.....
.....*P. marginecontinuum* Callejas
- 7 Leaf base rounded or cordate, inflorescences erect.....8
- Leaf base lobate, inflorescences pendulous9
- 8 Leaf ovate, leaf base rounded, peduncle up to 0.5 cm long
.....*P. palenquense* Callejas
- Leaf broadly ovate, leaf base deeply cordate, sinus open, peduncle 0.7–1 cm long*P. vallicola* C. DC.
- 9 Leaves more than 35 cm long, inflorescences 50–60 long.....10
- Leaves up to 26 cm long, inflorescences up to 20 cm long11

- 10 Leaves up to 50cm long, sinus closed, longer lobe overlapping the petiole, peduncle 1.5–2.7cm long ***P. caracasenum* Bredem. ex Link**
- Leaves up to 35cm long, sinus open, peduncle 7 cm long ***P. gualeanum* C. DC.**
- 11 Petioles shortly pubescent, inflorescences 10–20 cm long ***P. calcariforme* Tebbs**
- Petioles with trichomes forming lines, inflorescences 6–9 cm long..... ***P. hebetifolium* W. C. Burger**
- 12 Plants herbaceous, inflorescences spikes, arranged in umbels ***P. peltatum* L.**
- Shrubs, suffrutex, or climbers, inflorescences solitary racemes or spikes... **13**
- 13 Leaves 10–30 cm long, inflorescences erect..... **14**
- Leaves 35 cm long or longer, inflorescences erect or pendulous..... **31**
- 14 Lianescent vines..... **15**
- Shrubs or suffrutex..... **17**
- 15 Internodes, petioles and leaves glabrous, leaf blade ovate or oblong-elliptic, smooth **16**
- Internodes, petioles and abaxial leaf surface pilose, leaf blade obovate, slightly bullate..... ***P. parmatum* Dressler**
- 16 Leaf blades with evident idioblasts on abaxial surface, spikes 1–2 cm long, with obtuse apices ***P. foreroi* Gentry**
- Leaf blades without visible idioblasts, spikes 5–6 cm long with mucronate apices..... ***P. peltifolium* Callejas**
- 17 Leaf base lobed or cordate..... **18**
- Leaf base rounded, obtuse or scutellate **24**
- 18 Leaves rounded –ovate, leaf base deeply cordate..... ***P. andersii* M.A.Jaram., stat nov. nom nov.**
- Leaves ovate, leaf base cordate or lobed **19**
- 19 Shrubs densely crisp –villous..... ***P. copeyanum* (C.DC) Trel.**
- Shrubs glabrous **20**
- 20 Idioblasts visible on both surfaces ***P. subscutatum* (Miq.) C. DC.**
- Idioblasts not visible **21**
- 21 Petioles vaginate at the base, flowers not forming bands on the spikes ... **22**
- Petioles vaginate half the length or more, flowers forming bands on the spikes ***P. maxonii* C. DC.**
- 22 Leaves 5–10 cm wide, petiole inserted ca. 1 cm from the margin..... ***P. jacaleapaense* Callejas**
- Leaves 15 –20 cm wide, petiole inserted 5–6 cm from the margin **23**
- 23 Leaves broadly ovate, glabrous on both surfaces..... ***P. veraguense* C. DC.**
- Leaves elliptic-oblong to ovate, nerves on the abaxial surface pilose..... ***P. neovenezuelense* M.A.Jaram., stat. nov. nom nov.**
- 24 Leaves glabrous **25**
- Leaves pubescent at least abaxially on veins **27**
- 25 Leaves oblong-lanceolate, leaf base rounded ***P. imberbe* Trel.**
- Leaves ovate, leaf base obtuse or spatulate **26**
- 26 Leaf base narrowly spatulate ***P. indiwassii* W. Trujillo & M.A.Jaram.**
- Leaf base broadly obtuse ***P. scutilimum* C. DC.**

- 27 Apex short acuminate, inflorescence obtuse28
 – Apex long attenuate, inflorescences mucronate.....***P. tuerckheimii* C. DC.**
 28 Flowers forming bands around the spike ***P. hammelii* Callejas**
 – Flowers laxly arranged in a spike or raceme29
 29 Plant glabrous, inflorescences a spike***P. scutifolium* Yunck.**
 – Villous shrub, inflorescence a raceme30
 30 Petiole inserted ca. 1 cm from margin, with visible idioblasts on leaf lamina ***P. cariacicaense* M. Carv.-Silva & E.F.Guim.**
 – Petiole inserted 0.5 cm from margin, without visible idioblasts on leaf lamina..... ***P. carautensei* E.F.Guim. & M. Carv.-Silva**
 31 Leaves broadly ovate, petiole vaginate to the middle.....32
 – Leaves ovate, lanceolate or oblong, petiole vaginate above the middle...33
 32 Leaf base rounded, inflorescence short-apiculate...***P. mutisii* Trel. & Yunck.**
 – Leaf base cordate, inflorescence obtuse.....***P. grandilimbus* C. DC.**
 33 Internodes warty above nodes, leaves glabrous
 ***P. albert-smithii* Trel. & Yunck.**
 – Internodes smooth, leaves pubescent or tomentose, at least on the abaxial surface34
 34 Leaf strongly bullate..... ***P. hartwegianum* (Benth.) C. DC.**
 – Leaves not bullate, or occasionally softly bullate35
 35 Leaves elliptic-ovate, apex acute or short acuminate.....36
 – Leaves oblong-lanceolate, apex gradually acuminate or long attenuate...37
 36 Heteromorphic trichomes of intermixed on the abaxial surface, inflorescence up to 43 cm long.....***P. peltilimbus* Yunck.**
 – Trichomes of uniform length on the abaxial surface, inflorescences 50–60 cm long ***P. candollei* Sodiro**
 37 Leaves 14–26 cm wide, leaf base deeply cordate sub-peltate to obliquely peltate ***P. parianum* Yunck.**
 – Leaves 7.5–19 cm wide, leaf base obliquely peltate
 ***P. quinchasense* M.A.Jaram., sp. nov.**

Acknowledgements

We thank Juan E. Carvajal Cogollo for his assistance, Ariadna Valenzuela Zúñiga for preparing the illustration, A. F. Majin for field assistance, and Jenny Peña Varon for preparing the distribution map. We thank G. Aymard-Corredor, M. Carvalho-Silva, E. Tepe, E. J. Lirio and one anonymous reviewer for comments on previous versions of the manuscript. We thank K. Gandhi for his expert advice on nomenclatural issues regarding the new names proposed here. Special thanks to the Galvis Family: Don Lucindo, Doña Edilsa and Leidy for hosting our field expeditions in Serranía de las Quinchas. Don Lucindo's expert guidance in the forest was key to finding the plants. This work was supported by project "BPIN No. 2020000100003 – Investigación de la biodiversidad de Boyacá: complementación y síntesis a través de gradientes altitudinales e implicaciones de su incorporación en proyectos de apropiación social de conocimiento y de efectos de cambio climático." Funds for field expeditions were also provided by Universidad Militar Nueva Granada as part of the Plant Taxonomy and Systematics course. Gencore (Sequencing Center) at Los Andes University partially subsidized the sequencing analysis.

Additional information

Conflict of interest

No conflict of interest was declared.

Ethical statement

No ethical statement was reported.

Funding

No funding was reported.

Author contributions

Conceptualization: MAJ. Investigation: DRD, MEA. Writing – original draft: MAJ.

Author ORCIDs

M. Alejandra Jaramillo  <https://orcid.org/0000-0002-6539-4149>

Dayro Rodríguez-Duque  <https://orcid.org/0000-0002-3829-3377>

Magda Escobar-Alba  <https://orcid.org/0000-0002-3756-9132>

Data availability

All of the data that support the findings of this study are available in the main text or Supplementary Information.

References

- Baksh-Comeau YS, Maharaj SS, Adams CD, Hawthorne WD (2016) An annotated checklist of the vascular plants of Trinidad and Tobago with analysis of vegetation types and botanical “hotspots.” *Phytotaxa* 250: 1–431. <https://doi.org/10.11646/phytotaxa.250.1.1>
- Balcázar-Vargas M, Rangel-Ch JO, Linares E (2000) Diversidad florística de la Serranía de Las Quinchas, Magdalena Medio (Colombia). *Caldasia* 22: 191–224. <http://www.revistas.unal.edu.co/index.php/cal/article/viewFile/17571/18415>
- Baldwin BG (1992) Phylogenetic utility of the internal transcribed spacers of nuclear ribosomal DNA in plants: An example from the Compositae. *Molecular Phylogenetics and Evolution* 1(1): 3–16. [https://doi.org/10.1016/1055-7903\(92\)90030-K](https://doi.org/10.1016/1055-7903(92)90030-K)
- Beard JS (1946) Notes on the vegetation of the Paria Peninsula, Venezuela. *Caribbean Forester* 7: 37–46.
- Bohlender EE, Pérez-Torres J, Borray-Escalante NA, Stevens RD (2018) Dietary variation during reproduction in Seba’s short-tailed fruit bat. *Journal of Mammalogy* 99(2): 440–449. <https://doi.org/10.1093/jmammal/gyx189>
- Bohorquez-Osorio AF, Ulián T, Díazgranados M (2020) Guía de plantas útiles de la Serranía de las Quinchas. *Ecología, Economía y Ética, Rutas Turísticas por los Bosques y la Paz*. Royal Botanic Gardens, Kew, E3, UK.
- Bornstein AJ, Smith JF, Tepe EJ (2014) Two new species of *Piper* from the greater antilles. *Systematic Botany* 39(1): 10–16. <https://doi.org/10.1600/036364414X678206>
- Callejas-Posada R (2020) Piperaceae. In: Davidse G, Ulloa Ulloa C, Hernández Macías HM, Knapp S (Eds) *Flora Mesoamericana* 2. Missouri Botanical Garden Press, Saint Louis.
- Carvalho-Silva M, Eduardo P, Saraiva A, Guimarães EF (2022) Synopsis of the *Piper* subgenus *Ottonia* Spreng. (Piperaceae) from Brazil, 2: 1–25. <https://doi.org/10.1590/1677-941x-abb-2021-0360>

- Dauby G, Stévant T, Droissart V, Cosiaux A, Deblauwe V, Simo-Droissart M, Sosef M, Lowry P, Schatz GE, Gereau RE, Couvreur TL (2017) *ConR*: An R package to assist large-scale multispecies preliminary conservation assessments using distribution data. *Ecology and Evolution* 7(24): 11292–11303. <https://doi.org/10.1002/ece3.3704>
- De Candolle C (1866) Piperaceae Novae. *Journal of Botany, British and Foreign* 4: 210–219.
- Draper FC, Costa FRC, Arellano G, Phillips OL, Duque A, Macía MJ, ter Steege H, Asner GP, Berenguer E, Schiatti J, Socolar JB, de Souza FC, Dexter KG, Jørgensen PM, Tello JS, Magnusson WE, Baker TR, Castilho CV, Monteagudo-Mendoza A, Fine PVA, Ruokolainen K, Coronado ENH, Aymard G, Dávila N, Sáenz MS, Paredes MAR, Engel J, Fortunel C, Paine CET, Goret JY, Dourdain A, Petronelli P, Allie E, Andino JEG, Brienen RJW, Pérez LC, Manzatto ÂG, Zambrana NYP, Molino JF, Sabatier D, Chave J, Fauset S, Villacorta RG, Réjou-Méchain M, Berry PE, Melgaço K, Feldpausch TR, Sandoval EV, Martinez RV, Mesones I, Junqueira AB, Roucoux KH, de Toledo JJ, Andrade AC, Camargo JL, del Aguila Pasquel J, Santana FD, Laurance WF, Laurance SG, Lovejoy TE, Comiskey JA, Galbraith DR, Kalamandeen M, Aguilar GEN, Arenas JV, Guerra CAA, Flores M, Llampazo GF, Montenegro LAT, Gomez RZ, Pansonato MP, Moscoso VC, Vleminckx J, Barrantes OJV, Duivenvoorden JF, de Sousa SA, Arroyo L, Perdiz RO, Cravo JS, Marimon BS, Junior BHM, Carvalho FA, Damasco G, Disney M, Vital MS, Diaz PRS, Vicentini A, Nascimento H, Higuchi N, Van Andel T, Malhi Y, Ribeiro SC, Terborgh JW, Thomas RS, Dallmeier F, Prieto A, Hilário RR, Salomão RP, Silva RC, Casas LF, Vieira ICG, Araujo-Murakami A, Arevalo FR, Ramírez-Angulo H, Torre EV, Peñuela MC, Killeen TJ, Pardo G, Jimenez-Rojas E, Castro W, Cabrera DG, Pipoly J, de Sousa TR, Silvera M, Vos V, Neill D, Vargas PN, Vela DM, Aragão LEOC, Umetso RK, Sierra R, Wang O, Young KR, Prestes NCCS, Massi KG, Huaymacari JR, Gutierrez GAP, Aldana AM, Alexiades MN, Baccaro F, Céron C, Muelbert AE, Rios JMG, Lima AS, Lloyd JL, Pitman NCA, Gamarra LV, Oroche CJC, Fuentes AF, Palacios W, Patiño S, Torres-Lezama A, Baraloto C (2021) Amazon tree dominance across forest strata. *Nature Ecology & Evolution* 5(6): 757–767. <https://doi.org/10.1038/s41559-021-01418-y>
- Fleming TH (2004) Dispersal ecology of Neotropical *Piper* shrubs and treelets. In: Dyer L, Palmer AN (Eds) *Piper: A Model Genus for Studies of Phytochemistry, Ecology, and Evolution*. Kluwer Academic/Plenum Publishers, Boston, 58–77. https://doi.org/10.1007/978-0-387-30599-8_4
- Frodin DG (2004) History and concepts of big plant genera. *Taxon* 53(3): 753–776. <https://doi.org/10.2307/4135449>
- Gentry AH (1990) Floristic similarities and differences between southern Central America and upper Central Amazonia. In: Gentry AH (Ed.) *Four Neotropical rainforests*. Yale University Press, New Haven, Connecticut, USA., 141–157.
- IUCN (2012) Guidelines for application of IUCN Red List criteria at regional and national levels: Version 4.0. IUCN, Gland, Switzerland, 41 pp.
- IUCN (2022) 1 Guidelines for Using the IUCN Red List Categories and Criteria.
- Jaramillo MA, Callejas-Posada R (2004) A reappraisal of *Trianaeopiper* Trelease: Convergence of dwarf habit in some *Piper* species of the Chocó. *Taxon* 53(2): 269–278. <https://doi.org/10.2307/4135607>
- Jaramillo MA, Manos PS (2001) Phylogeny and the patterns of floral diversity in the genus *Piper* (Piperaceae). *American Journal of Botany* 88(4): 706–716. <https://doi.org/10.2307/2657072>
- Jaramillo MA, Callejas-Posada R, Davidson C, Smith JF, Stevens AC, Tepe EJ (2008) A phylogeny of the tropical genus *Piper* using ITS and the chloroplast intron psbJ – petA. *Systematic Botany* 33(4): 647–660. <https://doi.org/10.1600/036364408786500244>

- Kunth K (1839) Bemerkungen über die Familie der Piperaceen. *Linnaea* 13: 562–726. <https://doi.org/10.5962/bhl.title.110118>
- Lasso E, Engelbrecht BMJ, Dalling JW (2009) When sex is not enough: Ecological correlates of asexual reproductive ability in eight co-occurring *Piper* species in Panama. *Oecologia* 161: 43–56. <https://doi.org/10.1007/s00442-009-1353-6>
- Massad TJ, Richards LA, Philbin C, Yamaguchi LF, Kato MJ, Jeffrey CS, Oliveira C, Ochsenrider K, de Moraes MM, Tepe EJ, Cebrian Torregón G, Sandivo M, Dyer LA (2022) The chemical ecology of tropical forest diversity: Environmental variation, chemical similarity, herbivory, and richness. *Ecology* 103(9): 1–12. <https://doi.org/10.1002/ecy.3762>
- Meier W (2011) Los bosques nublados de la Cordillera de La Costa de Venezuela. *Biollania* 10: 106–121.
- Miquel FAG (1843) *Systema Piperacearum*. Rotterdam, 304 pp. <https://doi.org/10.5962/bhl.title.151>
- Perez Gutierrez RM, Neira Gonzalez AM, Hoyo-Vadillo C (2013) Alkaloids from *Piper*: A Review of its Phytochemistry and Pharmacology. *Mini-Reviews in Medicinal Chemistry* 13(2): 163–193. <https://doi.org/10.2174/138955713804805148>
- Rodríguez-Duque DL, Allende JRG, García-González JD, Escobar-Alba M, Hernández-Avendaño P, Aymard-Corredor GA (2021) Rediscovering *Ternstroemia killipiana* (Pentapetalaceae, S.L.), a Colombian Andes species not collected since 1926: Its geographic distribution and current conservation status. *Harvard Papers in Botany* 26(2): 449–454. <https://doi.org/10.3100/hpib.v26iss2.2021.n11>
- Salazar D, Jaramillo MA, Marquis RJ (2016a) Chemical similarity and local community assembly in the species rich tropical genus *Piper*. *Ecology* 97(11): 3176–3183. <https://doi.org/10.1002/ecy.1536>
- Salazar D, Jaramillo MA, Marquis RJ (2016b) The impact of plant chemical diversity on plant-herbivore interactions at the community level. *Oecologia* 181(4): 1199–1208. <https://doi.org/10.1007/s00442-016-3629-y>
- Salehi B, Zakaria ZA, Gyawali R, Ibrahim SA, Rajkovic J, Shinwari ZK, Khan T, Sharifi-Rad J, Ozleyen A, Turkdonmez E, Valussi M, Tumer TB, Fidalgo LM, Martorell M, Setzer WN (2019) *Piper* species: A comprehensive review on their phytochemistry, biological activities and applications. *Molecules* 24(7): 1364 <https://doi.org/10.3390/molecules24071364>
- Salinas NR, Betancur J (2004) Una nueva especie de *Dimerocostus* (costaceae) de Colombia. *Revista de la Academia Colombiana de Ciencias Exactas. Físicas y Naturales* 28: 465–470.
- Slinn HL, Richards LA, Dyer LA, Hurtado PJ, Smilanich AM (2018) Across multiple species, phytochemical diversity and herbivore diet breadth have cascading effects on herbivore immunity and parasitism in a tropical model system. *Frontiers in Plant Science* 9: 1–12. <https://doi.org/10.3389/fpls.2018.00656>
- Stamatakis A (2014) RAxML version 8: A tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* 30: 1312–1313. <https://doi.org/10.1093/bioinformatics/btu033>
- Steyermark JA (1984) *Flora de Venezuela. Piperaceae. Vol. 2 Segunda Parte*. Ediciones Fundación Educación Ambiental, Caracas, Venezuela, 619 pp.
- Steyermark JA, Agostini G (1966) Exploración Botánica del Cerro Patao y zonas adyacentes a Puerto Hierro en la Península de Paria, Edo. Sucre. *Acta Botanica Venezuelica* 1: 7–80.
- Stiles F, Bohórquez CI (2000) Evaluando el estado de la biodiversidad: el caso de la avifauna de la Serranía de las Quinchas, Boyacá, Colombia. *Caldasia* 22: 61–92.

- Tebbs MC (1990) Revision of *Piper* (Piperaceae) in the New World. 1. Review of characters and taxonomy of *Piper* section *Macrostachys*. Bulletin of the Natural History Museum of London 19: 117–158.
- Tebbs MC (1993) Revision of *Piper* (Piperaceae) in the New World, 3. The taxonomy of *Piper* sections *Lepianthes* and *Radula*. Bulletin of the Natural History Museum of London 23: 1–50.
- Tepe EJ, Rodríguez-Castañeda G, Glassmire AE, Dyer LA (2014) *Piper kelleyi*, a hotspot of ecological interactions and a new species from Ecuador and Peru. PhytoKeys 2014: 19–32. <https://doi.org/10.3897/phytokeys.34.6376>
- Thiers B (2023) Index Herbariorum A global directory of public herbaria and associated staff. <http://sweetgum.nybg.org/ih/>
- Trelease W (1928) *Trianaeopiper*, a new genus of Piperaceae. Proceedings of the American Philosophical Society 67: 47–50.
- Trelease W, Yuncker TG (1950) The Piperaceae of Northern South America. University of Illinois Press, Indiana.
- Trujillo W, Trujillo ET, Ortiz-Morea FA, Toro DA, Jaramillo MA (2022) New *Piper* species from the eastern slopes of the Andes in northern South America. PhytoKeys 206: 25–48. <https://doi.org/10.3897/phytokeys.206.75971>
- Wilson JS, Forister ML, Dyer LA, Connor JMO, Burls K, Feldman CR, Jaramillo MA, Miller JS, Rodri G, Tepe EJ, Whitfield JB, Young B (2012) Host conservatism, host shifts and diversification across three trophic levels in two Neotropical forests. Journal of Evolutionary Biology 25(3): 1–15. <https://doi.org/10.1111/j.1420-9101.2011.02446.x>
- Wunnenberg J, Rjosk A, Neinhuis C, Lautenschläger T (2021) Strengthening structures in the petiole-lamina junction of peltate leaves. Biomimetics 6: 25. <https://doi.org/10.3390/biomimetics6020025>
- Yohe LR, Leiser-Miller LB, Kaliszewska ZA, Donat P, Santana SE, Dávalos LM (2021) Diversity in olfactory receptor repertoires is associated with dietary specialization in a genus of frugivorous bat. G3: Genes, Genomes. G3 (Bethesda, Md.) 11(10): jkab260. <https://doi.org/10.1093/g3journal/jkab260>
- Yuncker TG (1972) The Piperaceae of Brazil, I. *Piper*-group I, II, III and IV. Hoehnea 19: 336.