

On the identity and typification of *Viola tenuis* Benth. (Violaceae)

Yan-Shuang Huang¹, Stephan W. Gale², Jinlong Zhang², Qiang Fan¹

¹ State Key Laboratory of Biocontrol and Guangdong Provincial Key Laboratory of Plant Stress Biology, School of Life Sciences, Sun Yat-sen University, Guangzhou, China

² Flora Conservation Department, Kadoorie Farm and Botanic Garden, Tai Po, New Territories, Hong Kong SAR, China

Corresponding author: Qiang Fan (fanqiang@mail.sysu.edu.cn)

Abstract

Viola tenuis Benth. was first described by George Benth. in 1842, without any type specimen designated, but with a collection record by Richard Brinsley Hinds in 1841. Currently, *Viola tenuis* is considered a synonym of another species according to Plants of the World Online (POWO). However, based on a combination of morphological and molecular evidence, we propose reinstating *Viola tenuis* as a distinct species. Our analysis places it within *Viola* sect. *Plagiostigma* subsect. *Diffusae*, where it shows morphological similarities to five other species endemic to Guangdong Province, China. We designated the type of *V. tenuis* here and assigned additional KFBG specimens to the species.

Key words: Conservation, nomenclature, typification, *Viola* sect. *Plagiostigma* subsect. *Diffusae*

Introduction

Viola sect. *Plagiostigma* subsect. *Diffusae*, which is mainly distributed in Southeast Asia, has undergone a long and controversial history in classification. Wilhelm Becker first published this subsection as an unranked group *Diffusae* (Becker 1923) and it was then reclassified as *V. sect. Nominium* ser. *Diffusae* (Van Steenis 1934). Thereafter, Wang (1991) elevated the rank of this taxon as *V. sect. Diffusae* (W.Becker) C.J.Wang in *Flora Reipublicae Popularis Sinicae*. Recently, Marcussen et al. (2022) placed it in *V. sect. Plagiostigma* as subsect. *Diffusae* (W.Becker) C.C.Chang. The diversity of *Diffusae* species in China has long been underestimated, as 19 species have been discovered recently, 10 of which are new to science (Huang 2023). Amongst them, there is an unresolved name, *V. tenuis* Benth., published by George Benth. in 1842, whose identity we resolved here.

Viola tenuis, a species having typical characters of *V. subsect. Diffusae*, with long peduncles, ovate-oblong leaf blades, thin rhizomes and flowers with short spurs and marginated style, was first collected in Hong Kong by Mr. Hinds and described by George Benth. in 1842 (Benth. 1842). Wilhelm Becker (1907) described a later homonym, *V. tenuis* W.Becker, from Brazil and reclassified the Hong Kong species as *V. diffusa* subsp. *tenuis* (Benth.) W.Becker (Becker 1921). The name *V. tenuis* Benth. was also accepted as a valid species in *Taiwania* (Wang and Huang 1990), but treated as a synonym of *V. diffusa* Ging.



Academic editor: Alexander Sennikov

Received: 11 October 2024

Accepted: 22 January 2025

Published: 20 February 2025

Citation: Huang Y-S, Gale SW, Zhang J, Fan Q (2025) On the identity and typification of *Viola tenuis* Benth. (Violaceae). *PhytoKeys* 252: 197–206. <https://doi.org/10.3897/phytokeys.252.138994>

Copyright: © Yan-Shuang Huang et al.
This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0).

in “Flora Reipublicae Popularis Sinicae” (Wang 1991), “Flora of China” (Chen et al. 2007) and Plants of the World Online (Govaerts 2023).

Based on field surveys, molecular and morphological analyses, we provide evidence supporting the recognition of *V. tenuis* Benth. as a distinct species and its name should not be treated as a synonym. In this paper, we present updated descriptions and illustrations for the species.

Materials and methods

We conducted field investigations and observations in Guangdong and Hong Kong during the flowering and fruiting periods of *Viola tenuis*. Leaf material of the species and its relatives were collected and stored in zip-lock plastic bags with silica gel for comparisons and taxonomical treatment. We used a micrometer and a stereomicroscope to observe and measure morphological features to compile detailed descriptions, based on both fresh and dry specimens. Voucher specimens were deposited in the Herbarium of Sun Yat-sen University (SYS) and the Herbarium of Kadoorie Farm and Botanic Garden (KFBG).

Total DNA was extracted by using the modified CTAB method (Doyle and Doyle 1987). Previously reported primers ITS1 and ITS4 (White et al. 1990) were used to amplify the regions of partial internal transcribed spacer 1, 5.8S ribosomal RNA gene and partial internal transcribed spacer 2. PCR amplifications were performed following Fan et al. (2015). We downloaded the sequences of the species and related ones from NCBI and aligned them using MEGA 6.0 (Tamura et al. 2013) with ClusterW, then manually adjusted the alignment. Phylogenetic inferences were carried out with Maximum Likelihood (ML) using IQ-TREE 2.0.3 (Minh 2020). SYM+I+I+R2 was selected as the best-fit model and 2,000 bootstraps were conducted to evaluate the confidence.

Results

The aligned length of our ITS sequences was 724 bps in total. *Viola tenuis* belongs to the monophyletic group *V. subsect. Diffusae* (BS = 100%, Fig. 1) and specimens of *V. tenuis* from Shenzhen and Hong Kong form a sister group with *V. heyuanensis* Yan S.Huang, Q.L.Ye & Q.Fan (BS = 90%).

Taxonomic treatment

Viola tenuis Benth. in London J. Bot. 1: 482 (1842)

Fig. 2

≡ *Viola diffusa* subsp. *tenuis* W.Becker in Philipp. J. Sci. 19(6): 714 (1921).

Type. CHINA • Hong Kong, 1841, *Hinds s.n.* (holotype: K000370141 [photo!]).

Additional specimens examined. CHINA • Hong Kong, Sai Kung, 06 May 2016, *Danial Hang HKY0012* (KFBG, Fig. 3c); • Hong Kong, Sai Kung, 13 Mar 2018, *Jingang Liu JG0615* (KFBG, Fig. 3b); • Hong Kong, Sai Kung, 21 Jul 2023, *Jinlong Zhang JL1488* (KFBG); • Zhuhai, Dawanshan Island, 07 Nov 1973, *Yue73 73-3160* (IBSC); • Zhuhai, Dawanshan Island, 04 Nov 1973, *Yue73 73-3055* (IBSC); • Zhuhai, Dangan Island, 12 Jul 2017, *Ming Tan Tu TY3390* (CDBI);




Figure 1. Maximum Likelihood (ML) tree of the new species and related species. Bootstrap values of the Maximum Likelihood are shown along the branches. Outgroups: *Melicytus obovatus*. *Viola tenuis* are marked in grey.

- Shenzhen, Qiniangshan Mountain, 18 Mar 2005, *Shouzhou Zhang & Liangqian Li 0259* (SZG);
- Shenzhen, Qiniangshan Mountain, 17 May 2005, *Shouzhou Zhang & Liangqian Li 1528* (SZG);
- Shenzhen, Qiniangshan Mountain, 16 Apr 2005, *Shouzhou Zhang & Liangqian Li 0196* (SZG);
- Shenzhen, Qiniangshan Mountain, 16 Apr 2019, *Q. Fan 18910* (SYS);

Etymology. It is hypothesised that the specific epithet refers to the plant's thin rhizomes.

Vernacular name. We propose a Chinese name, *Xì Jīng Jǐn Cài* (细茎堇菜), to reflect the thin rhizomes of this species.

Description. Perennial, stoloniferous herb with basal leaves rosulate, 5–8.5 cm tall. Rhizome erect or obliquely erect, rather slender, dark purple, with dense remains of stipules and petioles. Lateral stolons purple or light green, spreading and elongating, usually producing adventitious roots, with an apical rosette of leaves. Stipules ca. 8 mm, adnate to petioles for about 1/3 of their length at base, linear-lanceolate, acute at apex, with margins pinnatifid, glabrous or rarely puberulous. Petioles 2–4 cm, with wings narrow, glabrous




Figure 2. *Viola tenuis* Benth. **a** habitat and habit (photograph in Hong Kong, J. Zhang) **b** flower, front view **c** flower, side view **d** leaf and stipules **e** bracteoles **f** ovary and stigma **g** stamens **h** seeds.

or sparsely puberulous along the margin. Leaf blades ovate or ovate-oblong, 1.0–2.4 × 1.5–3.3 cm, glabrous or densely puberulous; margin finely crenate; base wedge-shaped to truncated; apex obtuse, rarely acuminate. Chasmogamous flowers ca. 1.5 cm in diam.; peduncles slender, 6–7 cm long, glabrous, much higher above the basal leaves, with two opposite bracteoles above middle; bracteoles linear-lanceolate, ciliate, 5.5–6 mm long, apex acuminate or obtuse. Sepals green, ciliate, lanceolate, 1.5–1.7 × 4–5.5 mm, with entire margin, acuminate apex and truncate, with extremely short semicircular appendages. Petals purple, anterior one with apparent violet lines, posterior and lateral ones with a yellow to green patch at base; posterior petals, narrowly ovate to oblong, 4.5 × 12 mm, glabrous, with entire margin and obtuse apex; lateral petals with straight to slightly clavate hairs at the base, oblong, ca. 1.3 cm long, with entire margin, apex obtuse; anterior petal spatulate, 8 mm long, with a short, round spur, interior side of the spur puberulent. Stamens 5, unequal, puberulent;



Figure 3. **a** holotype of *Viola tenuis* Benth. (K000370141) **b, c** specimen of *V. tenuis* in KFBG (JG0615, HKY0012).

anther thecae ca. 0.9 mm long, with terminal appendages ca. 0.6–0.85 mm long; posterior appendages (nectar spurs) of the two anterior stamens ca. 0.95 mm long, triangular. Ovary ovoid, ca. 1.25 × 1.49 mm, glabrous; style clavate, ca. 1.35 mm long, conspicuously geniculate at base; stigma glabrous, with thickened lateral margins, shortly beaked at apex. Cleistogamous flowers ca. 3.2 mm long; peduncles ca. 6.5 mm long; bracteoles linear-lanceolate, puberulous, ca. 2.8 mm long, apex acute. Sepals green, glabrous or puberulous, lanceolate, ca. 0.8 × 3.2 mm, with entire margin, acuminate obtuse. Capsules glabrous, light green to yellow, ovoid to oblong, ca. 9 mm long. Seeds brown, oblongoid, ca. 1.4–1.55 mm long, with tubercles and inconspicuous elaiosomes.

Phenology. Chasmogamous flowers from April to May, rarely October, cleistogamous flowers from May to September and fruits from April to October.

Distribution and habitat. *Viola tenuis* is currently known from Hong Kong, Shenzhen and Zhuhai in China (Fig. 4). The species grows on rocks or in rocky crevices at altitudes of 100–700 m a.s.l.




Figure 4. Distribution of *Viola tenuis* Benth.

Conservation status. As its area of occupancy is estimated less than 20 km², the conservation status of *Viola tenuis* should be considered as Vulnerable (VU) according to IUCN Red List criteria (D2; IUCN 2024).

Discussion

Becker (1921), when merging *Viola tenuis* into *V. diffusa* as a subspecies, described its leaves as roundish or ovate with a distinctly cordate base and long petioles. While accepting the subspecies, he suggested it might simply be a small-leaved variety of *V. diffusa*. However, Bentham (1842), when publishing this species, described it as having ovate-oblong leaves with a truncate base, along with rhizomes and stolons; he specifically mentioned that the leaves are similar to those of *V. patrinii* Ging. Although the original description did not include a type designation, there is only one specimen collected by Richard Brinsley Hinds in Hong Kong in 1841 (Fig. 3a). As the individuals on it also have narrow and truncate leaves, the distinct morphological characteristics clearly distinguish it from *V. diffusa* and other species of the *Diffusae* group. Based on our field surveys, *V. tenuis* Benth. is endemic to Guangdong and Hong Kong (Figs. 3 and 4) and forms a paraphyletic group with *V. diffusa* (Fig. 1), so it is more appropriate to be treated as a separate species. Both the name *Viola tenuis* proposed by Becker in 1907 and *Viola × tenuis* proposed by Karl Richter in 1891 are illegitimate later homonyms, based on Shenzhen Code Art. 11.4 and 53.1 (Turland et al. 2017).

In our phylogenetic tree, *V. tenuis*, *V. huizhouensis* Yan S.Huang & Q.Fan, *V. chaozhouensis* Yan S.Huang, J.H.Ding & Q.Fan, *V. guangzhouensis* A.Q.Dong, J.S.Zhou & F.W.Xing, *V. longissima* Yan S.Huang & Q.Fan and *V. heyuanensis* form a monophyletic group (Fig. 1). The latter three species have aerial stems, while the former three do not. The colour and hairiness of the leaf surface, the thickness of the rhizome and the size of the flowers can distinguish the first three species, but the relationship between morphological characters and the phylogeny still needs more investigation and evidence. These species are dis-

Table 1. Morphological comparisons between *Viola tenuis*, *V. diffusa* and *V. heyuanensis*.

Species name	<i>Viola tenuis</i>	<i>Viola diffusa</i>	<i>Viola heyuanensis</i>
Leaf blade	Ovate to ovate-oblong, 1.0–2.4 × 1.5–3.3 cm, glabrous or puberulous; finely crenate margin; wedge-shaped to truncated base; obtuse apex, rarely acuminate	Ovate to oblong, 1.7–2.5 × 3.0–4.9 cm, nearly glabrous; crenate margin; wedge-shaped base, gradually tapering into petiole	Narrowly ovate to triangular, 1.8–3.2 × 1.3–1.8 cm, densely pubescent; coarsely serrate margin; cuneate to truncate base, rarely cordate; obtuse or acuminate apex
Petioles	2–4 × ca. 0.1–0.2 cm, glabrous or sparsely puberulent, narrow wings	1.6–2.0 × 0.35–0.47 cm, sparsely pubescent, broad wings	2–4.5 cm, densely pubescent, narrow wings
Aerial stem	Absent	Absent	Aerial stems solitary or several and fasciculate, erect, slender
Rhizome	Erect or obliquely erect, rather slender, dark purple, with dense remains of stipules and petioles	Absent, with fibrous root	Erect or obliquely erect, rather stout, reddish-brown rarely branched
Flower	Chasmogamous flowers ca. 1.5 cm in diam.	Chasmogamous flowers ca. 0.8 cm in diam.	Chasmogamous flowers ca 1.2 cm diam.
Seeds	Brown, oblongoid, with tubercles, 1.4–1.55 mm long	Yellow, obovoid, with smooth surface, 1–1.2 mm long	Brown, obovoid, with sparse tubercles, 1–1.5 mm long

tributed in the mountainous area of Guangdong Province and appear to have undergone adaptive radiation (Huang et al. 2023). We hypothesise that sea level change might have influenced their diversification (Holbourn et al. 2018).

Based on our field surveys, molecular and morphological analyses (Table 1), we provide strong evidence supporting the recognition of *V. tenuis* Benth. as a distinct species and, thus, that this name should not be treated as a synonym.

There is no type specimen of *V. tenuis* Benth. designated in the protologue (Bentham 1842), which was entitled as ‘Enumeration of plants collected from Hong Kong, by RB Hinds, determined and described by George Bentham, ESQ’. According to the publication, we determined K000370142 (Fig. 2a) in the Royal Botanic Garden Kew as the holotype of *V. tenuis* (Shenzhen Code, Art. 7.9, Art. 8.1), as it is the only specimen collected by Hinds and identified as *Viola tenuis* by G. Bentham.

Key to *Viola* subsect. *Diffusae*

- 1 Leaf base conspicuously decurrent to the petiole 2
- Leaf base inconspicuously or not decurrent 3
- 2 Seeds with a conspicuous elaisome..... ***Viola pendulipes***
- Elaisome inconspicuous or absent 4
- 3 Leaf blades oblong to oblong–lanceolate, distinctly longer than broad
..... ***V. yunnanensis***
- Leaf blades ovate or orbicular, about as long as broad 5
- 4 Flowers 20–35 mm across 6
- Flowers 4–20 mm across 7
- 5 Flowers with a yellow to green patch at the base ***V. qingruui***
- Flowers without a yellow to green patch at the base 8
- 6 Plants with distinct aerial stems 9
- Plants with short or no aerial stem 10
- 7 Leaf blades abaxially purple ***V. jinggangshanensis***
- Leaf blades light green 11
- 8 Stipules triangular; flowers white ***V. lucens***
- Stipules linear to linear-lanceolate; flowers light pink ***V. changii***

9	Leaf blades abaxially purple	<i>V. guangzhouensis</i>
–	Leaf blades light green.....	12
10	Leaves papery when dried	<i>V. nanlingensis</i>
–	Leaves thinly leathery when dried	13
11	Seeds yellow to light brown, with smooth surface.....	14
–	Seeds dark brown, with tubercles	15
12	Posterior petals shorter than lateral ones; aerial stems 5–7 cm tall.....	<i>V. heyuanensis</i>
–	Posterior petals longer than lateral ones; aerial stems up to 40 cm tall	<i>V. longissima</i>
13	Leaves abaxially green; rhizomes slender	<i>V. tenuis</i>
–	Leaves abaxially dark purple	16
14	Lateral petals bearded	<i>V. diffusa</i>
–	Lateral petals glabrous	<i>V. amamiana</i>
15	Lateral petals bearded	<i>V. pricei</i>
–	Lateral petals glabrous	<i>V. nagasawae</i>
16	Leaf apex acute; leaf blade glabrous or pubescent only along veins	<i>V. chaozhouensis</i>
–	Leaf apex obtuse; leaf blade densely covered with white pubescence	<i>V. huizhouensis</i>

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

This work was supported by the project of the Research on Coevolution of Vegetation and Geological Environment of Shenzhen Dapeng Peninsula Geopark (2020F36).

Author contributions

All authors have contributed equally.

Author ORCIDs

Jinlong Zhang  <https://orcid.org/0000-0002-1161-5460>

Qiang Fan  <https://orcid.org/0000-0003-4254-6936>

Data availability

All of the data that support the findings of this study are available in the main text or Supplementary Information.

References

- Becker W (1907) Systematische Bearbeitung der Violen-Section Leptidium. Beihefte zum Botanischen Centralblatt 22(2): 78–96.
- Becker W (1921) Die Violen der Philippinen. Philippine Journal of Science 19(6): 707–722.

- Becker W (1923) *Violae Asiaticae et Australenses VI*. Beihefte zum Botanischen Centralblatt Abteilung 2 40: 119–171.
- Bentham G (1842) Vegetation of Hong Kong. *London Journal of Botany* 1: 482–494.
- Chen YS, Yang QE, Ohba H, Nikitin VV (2007) *Violaceae*. In: Wu ZY, Raven PH, Hong DY (Eds) *Flora of China*. Vol. 13. Science Press, Beijing and Missouri Botanical Garden Press, St. Louis, 74–111.
- Doyle JJ, Doyle JL (1987) A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochemical Bulletin* 19: 11–15.
- Fan Q, Chen SF, Wang LY, Chen ZX, Liao WB (2015) A new species and new section of *Viola* (*Violaceae*) from Guangdong, China. *Phytotaxa* 197(1): 15–26. <https://doi.org/10.11646/phytotaxa.197.1.2>
- Govaerts R (2023) The World Checklist of Vascular Plants (WCVP). Royal Botanic Gardens, Kew. [Checklist dataset accessed via GBIF.org on 2024-01-03]
- Holbourn AE, Kuhnt W, Clemens SC, Kochhann KG, Jöhnck J, Lübbers J, Andersen N (2018) Late Miocene climate cooling and intensification of southeast Asian winter monsoon. *Nature Communications* 9(1): 1584. <https://doi.org/10.1038/s41467-018-03950-1>
- Huang YS, Ding JH, Ye QL, Dai JM, Zhong ZM, Fan Q (2023) Four new species of *Viola* (*Violaceae*) from southern China. *Nordic Journal of Botany* 2023(6): e03941. <https://doi.org/10.1111/njb.03941>
- IUCN (2024) The IUCN Red List of Threatened Species (Version 2024-2). <https://www.iucnredlist.org> [January, 27, 2025]
- Marcussen T, Ballard HE, Danihelka J, Flores AR, Nicola MV, Watson JM (2022) A revised phylogenetic classification for *Viola* (*Violaceae*). *Plants* 11(17): 2224. <https://doi.org/10.3390/plants11172224>
- Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, Von Haeseler A, Lanfear R (2020) IQ-TREE 2: New models and efficient methods for phylogenetic inference in the genomic era. *Molecular Biology and Evolution* 37(5): 1530–1534. <https://doi.org/10.1093/molbev/msaa015>
- Tamura K, Stecher G, Peterson D, Filipowski A, Kumar S (2013) MEGA6: Molecular evolutionary genetics analysis version 6.0. *Molecular Biology and Evolution* 30(12): 2725–2729. <https://doi.org/10.1093/molbev/mst197>
- Turland NJ, Wiersema JH, Barrie FR, Greuter W, Hawksworth DL, Herendeen PS, Knapp S, Kusber W-H, Li D-Z, Marhold K, May TW, McNeill J, Monro AM, Prado J, Price MJ, Smith GF (Eds) (2017) International code of nomenclature for algae, fungi, and plants (Shenzhen code) adopted by the nineteenth international botanical congress, Shenzhen, China, July 2017. *Regnum Vegetabile* 159. Koeltz Botanical Books, Glashütten. <https://doi.org/10.12705/Code.2018>
- Van Steenis CGGJ (1934) *Bulletin du Jardin Botanique de Buitenzorg, série 3*, 13: 260–263.
- Wang QR (1991) *Violaceae*. In: Wang QR (Ed.) *Flora Reipublicae Popularis Sinicae*. Vol. 51. Science Press, Beijing, 1–148.
- Wang CJ, Huang TC (1990) Notes on the Flora of Taiwan (9) The *Viola* L. (*Violaceae*). *Taiwania* 35(1): 12–56.
- White TJ, Bruns T, Lee SJWT, Taylor J (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: *PCR Protocols: A Guide to Methods and Applications*. 18: 315–322. <https://doi.org/10.1016/B978-0-12-372180-8.50042-1>

Supplementary material 1

GenBank accessions for phylogenetic analysis

Authors: Yan-Shuang Huang, Stephan W. Gale, Jinlong Zhang, Qiang Fan

Data type: docx

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/phytokeys.252.138994.suppl1>