



First report of the genus *Conostigmus* Dahlbom (Hymenoptera: Ceraphronoidea: Megaspilidae) from India with description of a new species

C Bijoy[†], K Rajmohana[†], Ramesh Kumar[‡]

[†] Western Ghats Regional Centre, Zoological Survey of India, Calicut-673006, Kerala, India

[‡] National Bureau of Agriculturally Important Insects, Hebbal, Bangalore-560024, Karnataka, India

Corresponding author: C Bijoy (bejoyaaryan@gmail.com)

Academic editor: Lyubomir Penev

Received: 30 Aug 2013 | Accepted: 26 Dec 2013 | Published: 02 Jan 2014

Citation: Bijoy C, Rajmohana K, Kumar R (2014) First report of the genus *Conostigmus* Dahlbom (Hymenoptera: Ceraphronoidea: Megaspilidae) from India with description of a new species. Biodiversity Data Journal 2: e991. doi: [10.3897/BDJ.2.e991](https://doi.org/10.3897/BDJ.2.e991)

ZooBank: urn:lsid:zoobank.org:pub:7B5C7A33-F8F5-489A-9A3C-A736B5FB1AB6

Abstract

The genus *Conostigmus* Dahlbom (Hymenoptera: Ceraphronoidea: Megaspilidae) is reported for the first time from India, along with description and illustration of *C. neotubifer* sp. n. A comparative discussion on the morphological affinities of the new species with its putative sister *C. tubifer* Dessart is provided. An intermixing of character states of genera *Conostigmus* as well as *Dendrocerus* Ratzeburg is observed.

Keywords

Conostigmus, Megaspilidae, new species, India

Introduction

Superfamily Ceraphronoidea (Hymenoptera) comprises two extant families of parasitoid wasps, Ceraphronidae and Megaspilidae (Johnson and Musetti 2004). Megaspilidae

includes 305 valid species in 13 genera (<http://hol.osu.edu/index.html?id=91>). The species of Megaspilidae are generally ectoparasitoids attacking cocoons and puparia, primarily on Diptera, Neuroptera, Coleoptera, Mecoptera and also as hyperparasitoids of Hymenoptera (Dessart 1995).

Among the 13 genera of Megaspilidae, the most speciose and widely distributed are *Dendrocerus* Ratzeburg 1852 and *Conostigmus* Dahlbom 1858. The genus *Conostigmus* was erected by Dahlbom (1858), as a subgenus of *Megaspilus* Westwood, with *Megaspilus alutaceus* Thomson as type species. Later in 1951, Muesebeck and Walkley elevated the taxon to generic level. The genus is distributed widely in all geographical realms and has 168 species. Only 9 species have been reported from Oriental region so far (Johnson and Musetti 2004, Dessart 1997) (catalogued as 8 species by Johnson and Musetti (2004), missing out *C. occipitalis* Dessart 1997).

In continuation with our pioneer taxonomic studies on the superfamily Ceraphronoidea of India, a new species of *Conostigmus* viz. *C. neotubifer* sp. n. is hereby described and illustrated. A comparative discussion on the morphological affinities of the new species with *C. tubifer* Dessart 1997, the most resembling species is also provided. Both the species are remarkable for their intermixing of characters seen in two genera, *Conostigmus* and *Dendrocerus*.

Materials and methods

The specimens under study were collected using malaise trap set among the grassy patches of Port Blair, Andaman and Nicobar Islands, India. The specimens were card mounted on point-card tips. Description and imaging works were carried out employing Leica M205A stereomicroscope, with 1 × objective and Leica DFC-500 digital camera. Morphological terminology follows Fergusson (1980), Dessart (1997) and Miko and Deans (2009). Based on Dessart (1997), we have attempted to define the various states of ocellar triangle in Megaspilidae. If OOL > POL, the ocellar triangle is 'narrow-based' and if OOL POL.

Both the Holotype and the Paratypes are deposited in the National Zoological Collection at Zoological Survey of India, Calicut.

Abbreviations: DFIm – Minimal Interocular Facial Distance; OOL – Ocellocular Length; POL – Postocellar Length; LOL – Lateral Ocellar Length; T3 – Metasomal tergite 3.

NHM – Natural History Museum, London.

Taxon treatment

Conostigmus neotubifer Bijoy & Rajmohana, 2014, sp. n.

- ZooBank urn:lsid:zoobank.org:act:4BF76167-8B66-4FB5-B4C1-DEB4C9F8F302

Materials

Holotype:

- a. scientificNameID: *Conostigmus neotubifer*; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Hymenoptera; family: Megaspilidae; genus: *Conostigmus*; specificEpithet: *neotubifer*; scientificNameAuthorship: Bijoy and Rajmohana; continent: Asia; island: Andaman and Nicobar island; country: India; stateProvince: Andaman and Nicobar island; locality: Port Blair; verbatimLocality: Bathu Barthi; decimalLatitude: 11.768 N; decimalLongitude: 92.737 E; habitat: grassy field; eventRemarks: collected in malasia trap; individualID: ZSI/WGRC/IR/INV.2652; individualCount: 1; sex: male; lifeStage: adult; preparations: Card mount; recordedBy: Ramesh Kumar; identifiedBy: Bijoy and Rajmohana; dateIdentified: 06/27/2013; institutionID: ZSIC

Paratypes:

- a. scientificNameID: *Conostigmus neotubifer*; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Hymenoptera; genus: *Conostigmus*; specificEpithet: *neotubifer*; scientificNameAuthorship: Bijoy and Rajmohana; continent: Asia; island: Andaman and Nicobar island; country: India; stateProvince: Andaman and Nicobar island; locality: Port Blair; verbatimLocality: Bathu Barthi; decimalLatitude: 11.768 N; decimalLongitude: 92.737 E; habitat: grassy field; eventRemarks: collected in malasia trap; individualID: ZSI/WGRC/IR/INV.2653; individualCount: 1; sex: female; lifeStage: adult; preparations: Card mount; recordedBy: Ramesh Kumar; identifiedBy: Bijoy and Rajmohana; dateIdentified: 06/27/2013; institutionID: ZSIC
- b. scientificNameID: *Conostigmus neotubifer*; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Hymenoptera; genus: *Conostigmus*; specificEpithet: *neotubifer*; scientificNameAuthorship: Bijoy and Rajmohana; continent: Asia; island: Andaman and Nicobar island; country: India; stateProvince: Andaman and Nicobar island; locality: Port Blair; verbatimLocality: Bathu Barthi; decimalLatitude: 11.768 N; decimalLongitude: 92.737 E; habitat: grassy field; eventRemarks: collected in malasia trap; individualID: ZSI/WGRC/IR/INV.2654; individualCount: 1; sex: female; lifeStage: adult; preparations: Card mount; recordedBy: Ramesh Kumar; identifiedBy: Bijoy and Rajmohana; dateIdentified: 06/27/2013; institutionID: ZSIC

Description

Holotype: Male (Fig. 1).



Figure 1.
Male habitus, lateral view.



Figure 2.
Male antenna, lateral view.



Figure 3.
Male head, dorsal view.



Figure 4.
Male head, anterior view.




Figure 5.
Male head and mesosoma, dorsal view.



Figure 6.
Male mesosoma, lateral view.



Figure 7.
Male clypeus, anterior view.

Coloration: Body blackish brown with head more darker; eyes and ocelli silvery with a bronze tinge; fore wing clear at base, with large infuscate area below pterostigma and radial vein extending to its posterior margin; scape, pedicel and A3 brownish yellow turning darker towards segments from A4 onwards; legs brownish yellow with coxae and posterior half of hind tibia blackish brown (Fig. 1); mandible light brown with darker distal extreme; pterostigma brown getting darker to wing margin; radial vein and costal vein pale brown (Fig. 8); body pubescence white.




Figure 8.
Male forewing.

Body: length: 1.52 mm.

Head: (length/width/height: 230/580/460); eye (length/width: 260/210); preoccipital crescent well-separated from posterior ocelli merging laterally into eye margin (Fig. 3); temple strongly carinated; preoccipital furrow present; ocellar triangle isosceles, narrow, short and raised; LOL

Antenna: (Fig. 2); very long (subequal to the body length) and slender; scape nearly 4 × longer than wide; pedicel small and almost globular; A3 highly slender, nearly 4.8 ×

longer than wide, subequal to scape; flagellar segments quasi cylindrical with very slight serrations basally; pubescence shorter than breadth of segments; length/width measurements of antennal segments: scape (252/66), pedicel (60/48), A3 (255/53), A4 (147/54), A5 (140/53), A6 (122/53), A7 (128/61), A8 (117/64), A9 (108/57), A10 (82/70), A11 (127/54).

Mesosoma: (length/width/height: 610/450/460); mesosoma robust with very angular shoulders; alutaceous in sculpture; mesoscutum: (length/width: 230/440); mesoscutum nearly 2 × wider than long; lateral margin of mesoscutum posterior to anterior margin of notaulus convex; coarsely foveolate notaulices curve smoothly and converging posteriorly, meeting median furrow at transscutal articulation (Fig. 5); scutellum broadly rounded at apex with long hairs; axilliluar carinae present; metanotal-propodeal sulcus placed in regular intervals medially and with some longitudinal carina remote laterally (Fig. 9); propodeum unarmed, smooth and bare with sculpture effaced and with irregular carinae (Fig. 9) but stretched posteriorly into a inverted 'U' shaped narrow apex. Lateral propodeal carinae present. Propodeal spiracle large and conspicuous (Fig. 1); sternaulus absent (Fig. 6); anterior mesopleural sulcus and meso-metapleural sulcus distinct and foveolate; metapleuron bordered by incomplete carina; lower margin of mesopleuron and metapleuron with dense hairs.



Figure 9.
Male propodeum.

Forewing: (Fig. 8); total wing length 1.25 mm. Pterostigma (length/width: 200/80) semielliptical, 2.4 × longer than wide. Radius (0.27 mm), slightly curved and 1.36 × the length of pterostigma; maximal width of fused costal and subcostal vein wider than radius; basal part with less hairs.

Metasoma: (Fig. 10); (length/width/height: 484/390/340); metasoma smooth; gastral collar well developed and widened more than that of propodeal strip, with syntergal translucent patch. Four strong transverse gastral carinae present on basal portion of

metasoma. A pair of paler gastrocoeli seen quite below gastral collar; largest tergite, T3 occupying 72% of metasomal length. Metasoma held in an elevated manner, above propodeal plane (Fig. 1) (best visible in lateral view). Genitalia with short basal ring, volsella widely separated basally with long terminal setae on each parossiculi (Fig. 15) (harpe damaged).



Figure 10.
Male, metasoma dorsal view.



Figure 11.
Female habitus.



Figure 12.
Female antenna.



Figure 13.
Female forewing.




Figure 14.
Female, metasoma dorsal view.



Figure 15.
Male genitalia ventral view.

Female (Paratype): Figs 11, 12, 13, 14.

Body length 1.68 mm. Body colouration and the morphological features of head, mesosoma and metasoma are the same as holotype, except the measurements of antennal segments, proportion of radius and pterostigma and length of metasoma.

Scape more than 4 × as long as wide and A3 more than 3 × as long as wide. Pedicel slender to succeeding antennal segments. Length/width measurements of antennal segments: scape (321/73), pedicel (103/51), A3 (170/51), A4 (97/65), A5 (86/79), A6 (83/90), A7 (90/87), A8 (79/84), A9 (80/85), A10 (87/78), A11 (124/68).

DFIm 52% of head width and fore wing infuscation much darker compared to paler infuscation in male fore wing. T3 occupying 57% of metasomal length.

Diagnosis

Conostigmus neotubifer sp. n. can be diagnosed by the following features.

Head transverse, wider than mesosoma. Male antenna subequal to body length with A3 highly slender, 4.8 × as long as wide and subequal to scape. Flagellar segments in male quasi cylindrical with very slight serrations basally. Female antenna with scape more than 4 × as long as wide and A3 more than 3 × as long as wide. Ocellar triangle isosceles, narrow based, short and raised in both sexes. Facial sulcus extending to intertorular carina in both sexes. Preoccipital furrow distinct. Supraclypeal furrow absent. DFIm 0.57% of head width in male and 0.52% of head width in female. Eyes densely pubescent. Dorsal margin of propodeal foramen 'U' shaped in dorsal view; median propodeal projection absent. Sternaulus absent. Metasoma dorsally elevated from the propodeal axis. Forewing infuscus, darker in female and paler in male.

Etymology

The species is named '*neotubifer*', since this species resembles *Conostigmus tubifer* ('Neo' in Latin = 'new').

Distribution

INDIA, Andaman Nicobar island, Port Blair.

Biology

Unknown.

Discussion

The proposed new species is placed under *Conostigmus*, since its putative sister *C. tubifer* belongs also to *Conostigmus* and can be justified mainly by the presence of independent parossiculi in male genitalia, one of the diagnostic characters for the subgenus *Conostigmus* s.str. in Dessart and Cancemi (1989). *C. tubifer* is unique among the Oriental species because of its affinity to *Dendrocerus* due to the quasi cylindrical appearance of male antennal segments, broad based nature of ocellar triangle (Dessart 1997), along with characters as mentioned (Table 1). Except in the ocellar ratio *C. neotubifer* resembles *C. tubifer*, in aspects regarding head, mesosoma and metasoma (Table 1) and also in having an upwardly elevated posture of metasoma. This intermixing of character states of the two genera *Conostigmus* and *Dendrocerus* in *C. tubifer* and *C. neotubifer* sp. n. emphasizes the uncertain state of the current classification of Ceraphronoidea.

Table 1.

Characters for distinguishing *Conostigmus* and *Dendrocerus*.

Characters	<i>Conostigmus</i>	<i>Dendrocerus</i>	<i>C. neotubifer</i> sp. n.	<i>C. tubifer</i>
Ocellar triangle	Usually with narrow base	Usually with broad base	<i>Conostigmus</i>	<i>Dendrocerus</i>
Shape of head	Usually globular	Usually transverse	<i>Dendrocerus</i>	<i>Dendrocerus</i>
Intertorular carina	Complete and with a median projection	Complete or incomplete but without a median projection	<i>Conostigmus</i>	<i>Conostigmus</i>
Notauli	Smoothly curved towards anterior angles of mesoscutum	Sharply angulate towards anterior angles of mesoscutum	<i>Conostigmus</i>	<i>Conostigmus</i>
Mesoscutum	Anteriorly narrowed	Quadrate	<i>Dendrocerus</i>	<i>Dendrocerus</i>
Basal antennal segments of male	Cylindrical	Serrate, triangular, ramose or quasi cylindrical	Quasi cylindrical as in <i>Dendrocerus</i>	Quasi cylindrical as in <i>Dendrocerus</i>
Shape of scutellum	Flat	Convex or raised	<i>Conostigmus</i>	<i>Conostigmus</i>
Parossiculi	Independent	Fused	<i>Conostigmus</i>	<i>Conostigmus</i>

In *C. neotubifer* sp. n., OOL > POL (ocellar triangle is narrow based), while in *C. tubifer* OOL < POL (ocellar triangle is broad based) in Dessart (1997), regarding ocelli, is a mismatch. In the description of the holotype male, Dessart clearly states, supported by numerical values, that the ocellar triangle is broad based, which is in contradiction to the medial placement of ocellar triangle in the illustration. As the holotype of *C. tubifer* could not be traced, we had no means than to go by the descriptive part of the original literature, since it is supported by numerical values. Though NHM is mentioned as the type depository of *C. tubifer*, in Johnson and Musetti (2004) and Dessart (1997), on enquiry at NHM, we were informed that the types were never deposited there in reality.

In addition, the differences in the following characters also ensure that *tubifer* and *neotubifer* are not conspecific.

1. Forewing infuscation extending further to the distal margins beyond pterostigma and radius in *C. neotubifer* sp. n. (restricted under stigma and radius in *C. tubifer*).
2. Facial groove distinct in both sexes in *C. neotubifer* sp. n. (in *C. tubifer* facial groove present in male and absent in female).
3. Pterostigma is 2.4 × longer than wide in *C. neotubifer* sp. n. (only 2 × longer than wide in *C. tubifer*).
4. DFlm value is 52% in female in *C. neotubifer* sp. n. (46% in female in *C. tubifer*).
5. Scape and A3 of male subequal in length in *C. neotubifer* sp. n. (in *C. tubifer*, A3 is shorter (only 0.85 ×) than scape).

6. In *C. neotubifer* sp. n., A3 of female antenna is 3 × longer than wide and scape 2 × longer than A3 (in *C. tubifer*, A3 is only two 2 × longer than wide and scape is 4 × longer than A3).

Acknowledgements

The authors are grateful to the Director of the Zoological Survey of India, Kolkata and the Officer-in-Charge, ZSI, Western Ghat Regional Centre, Calicut, Kerala, for providing facilities and encouragement. Thanks are also due to Istvan Miko for his valuable help in interpreting the generic limits of *Conostigmus* and *Dendrocerus* and also for literature support. We are grateful to David Notton, for checking the availability of the types at the Natural History Museum, London. The authors profusely thank Dr. J Poorani, for sharing specimens for this study.

Author contributions

Bijoy C: Identification, species concept and imaging

Rajmohana K: Identification and species concept

Ramesh Kumar: Specimen collection and manuscript preparation

References

- Dahlbom AG (1858) Svenska sma-ichneumonernas familjer och slagten. Ofvers. K. Vet. Akad. Forh. 14: 289-298.
- Dessart P (1995) Megaspilidae. In: Hanson P, Gauld ID (Eds) Hymenoptera of Costa Rica. Oxford University Press, Oxford, 893 pp.
- Dessart P (1997) Les Megaspilidae ni europeens, ni americains 1. Le genre *Conostigmus* Dahlbom, 1858 (Hym. Ceraphronoidea Megaspilidae). Mem. Soc. R. Belge Entomol. 37: 144.
- Dessart P, Cancemi P (1989) Trois especes nouvelles de *Ceraphron* a feutrage propodeal (Hymenoptera Ceraphronoidea Ceraphronidae). Bull. Inst. R. Sci. Nat. Belg. 58: 95-100.
- Fergusson NDM (1980) A revision of the British species of *Dendrocerus* Ratzburg (Hymenoptera: Ceraphronoidea) with a review of their biology as aphid hyperparasites. Bull. Brit. Mus. (Nat. Hist.) Entomol. 41: 255-314.
- Johnson NF, Musetti L (2004) Catalog of systematic literature of the super family Ceraphronoidea (Hymenoptera). 33(2). Contributions of the American Entomological institute, 149 pp.
- Miko I, Deans AR (2009) *Masner*, a new genus of Ceraphronidae (Hymenoptera, Ceraphronoidea) described using controlled vocabularies. Zookeys 20: 127-153. DOI: [10.3897/zookeys.20.119](https://doi.org/10.3897/zookeys.20.119)
- Ratzburg JTC (1852) Die Ichneumonien der Forstinsecten in forstlicher und entomologischer Beziehung. 3. Nicolaischen Buchhandlung, Berlin, 272 pp.