



Single Taxon Treatment

Heilipus squamosus (LeConte, 1824): clarification of the correct name for the “avocado tree girdler” with updates on its biology and distribution (Coleoptera, Curculionidae, Molytinae, Hylobiini)

Salvatore S Anzaldo[‡], Valentina Díaz-Grisales[§]

[‡] Arizona State University, Tempe, United States of America

[§] Programa de Entomología y Acarología, Colegio de Postgraduados, Campus Montecillo, Montecillo, Mexico

Corresponding author: Salvatore S Anzaldo (sanzaldo@asu.edu)

Academic editor: Jennifer C. Girón Duque

Received: 18 Apr 2022 | Accepted: 30 Aug 2022 | Published: 30 Sep 2022

Citation: Anzaldo SS, Díaz-Grisales V (2022) *Heilipus squamosus* (LeConte, 1824): clarification of the correct name for the “avocado tree girdler” with updates on its biology and distribution (Coleoptera, Curculionidae, Molytinae, Hylobiini). Biodiversity Data Journal 10: e85499. <https://doi.org/10.3897/BDJ.10.e85499>

Abstract

Background

A single species of the genus *Heilipus* Germar, 1824 is native to the south-eastern United States and was a pest of avocado in Florida in the mid-20th century. Two names—*Heilipus apiatus* (Olivier, 1807) and *H. squamosus* (LeConte, 1824)—have both recently been used as the valid name for this species, with *H. apiatus* also being recently used as the valid name for a species from French Guiana. Confusion surrounding the correct application of these names and the distribution of these species traces back to an erroneous distribution given in Olivier's 1807 description of *H. apiatus* and, although some authors clarified this previously, there continues to be confusion concerning the correct name. Outside of avocado-producing areas, this species was rarely collected and little was known about its

biology. Recent observations on iNaturalist and BugGuide suggest the species is more widely distributed in the United States and less rare than it was previously thought to be.

New information

Heilipus squamosus (LeConte, 1824) is recognised as the valid name for the sole species of *Heilipus* occurring in the United States, while *Heilipus apiatus* (Olivier, 1807) is a very different species from French Guiana. *Heilipus squamosus* was previously recorded from eight States in the south-eastern United States and, after examining records from natural history collections, iNaturalist, BugGuide and literature sources, the species is newly recorded from an additional seven States: Arkansas, Kansas, Kentucky, Louisiana, Missouri, Oklahoma and Texas. Though native host plants have been unconfirmed by rearing records, the evidence indicating the possible host plants in the plant family Lauraceae is reviewed.

Keywords

weevil, Lauraceae, host associations, new state record, *Persea*

Introduction

The genus *Heilipus* Germar, 1824 (Germar 1824) contains 88 species in the New World (O'Brien and Wibmer 1982, Wibmer and O'Brien 1986, Díaz et al. 2021), with a single species occurring in the United States (O'Brien and Wibmer 1982, Anderson 2002). The species described as *Rhynchaenus apiatus* Olivier, 1807 and *Pissodes squamosus* LeConte, 1824 were both originally indicated to be from the south-eastern United States (Olivier 1807, LeConte 1824). The introduction of South American countries to the published distribution of *H. apiatus* (Schoenherr 1836, Berg 1881) added confusion to the identity of this species which persists today. The two species were synonymised by Blatchley and Leng (1916) in the genus *Heilipus*, with Olivier's name (*H. apiatus*) having precedence. Since this act, the valid name for the species of *Heilipus* occurring in the United States has been inconsistently used, with *H. squamosus* (LeConte) and *H. apiatus* (Olivier) both being used in the 20th and 21st centuries.

On three occasions, the confusion surrounding the distribution and valid names for these species appeared to have been resolved, but was overlooked by subsequent authors. First, Schoenherr (1836) includes both Cayenna and Florida in the distribution for *H. apiatus*, but only Cayenna is listed in a later volume (Schoenherr 1843). This possibly indicates awareness of an erroneous North American distribution originally given by Olivier, but no clarification was given. Later authors (e.g. Gemminger and Harold 1871, Berg 1881) include both North and South America in the distribution of *H. apiatus* again. Second, Wolfenbarger (1950) references the synonymy of Blatchley and Leng (1916) and states, "This synonymy is accepted as an error since Dr. [William H.] Anderson reported that,

‘Olivier’s species is quite different and occurs in South America’”. *Heilipus apiatus* continued to be used by Woodruff (1963) and O’Brien and Wibmer (1982) for the U.S. species without reference to Wolfenbarger’s statement (Wolfenbarger 1950). Third, O’Brien and Wibmer (1984) correct their previous use of *H. apiatus* (O’Brien and Wibmer 1982) in the often overlooked supplement to their checklist, stating that the type locality of *H. apiatus* as Florida is an error and that it is a South American species, making *H. squamosus* (LeConte) the valid name for the species from the United States. In their later treatment on South American weevils, Wibmer and O’Brien (1986) accordingly list *H. apiatus* (Olivier) as a species from French Guiana.

Since the latest clarification by O’Brien and Wibmer (1984), however, the name *H. apiatus* has persisted as the valid name for the species in the United States in published literature and online resources. The name *H. apiatus* has also been used correctly recently, following Wibmer and O’Brien (1986), for the species from French Guiana (Gombauld 2004, Rheinheimer 2010, Rheinheimer 2017). Examination of images of Olivier’s *H. apiatus* syntypes (Fig. 1) reconfirms the positions taken by Schoenherr (1836), Wolfenbarger (1950) and O’Brien and Wibmer (1984)—*H. apiatus* and *H. squamosus* are distinct species and *H. apiatus* is not the species found in the United States.




Figure 1. [doi](#)

Heilipus apiatus syntype (MNHN EC13569). Scale bars = 2 mm. Photographs copyright Maxime Le Cesne/MNHN. A: lateral habitus. B: dorsal habitus. C: labels.

Materials and methods

One hundred and thirty-five occurrence records of *H. squamosus* were analysed (Suppl. material 1). Fifty-seven of those records are preserved specimens, only 10 of which are digitised records available online (SCAN 2022, Ecdysis 2022, GBIF.org 2022b) and are housed in the following natural history collections:

- **ABS** – Archbold Biological Station Arthropod Collection, Lake Placid, FL
- **ASUCOB** – Arizona State University Charles W. O’Brien Collection, Tempe, AZ
- **AUMNH** – Auburn University Museum of Natural History, Auburn, AL

The remaining 47 records of preserved specimens were not previously available online; information for these records is available in Suppl. material 1 and GBIF.org (2022a). The

following collections provided specimen data and/or photographs of their *H. squamosus* material:

- **CMNC** – Canadian Museum of Nature Collection, Ottawa, Canada
- **FSCA** – Florida State Collection of Arthropods, Gainesville, FL
- **LSAM** – Louisiana State Arthropod Museum, Baton Rouge, LA
- **MEM** – Mississippi Entomological Museum, Mississippi State, MS
- **NCSU** – North Carolina State University Insect Collection, Raleigh, NC
- **UGCA** – University of Georgia Collection of Arthropods, Athens, GA
- **VMNH** – Virginia Museum of Natural History, Martinsville, VA

In addition to specimens from natural history collections, 43 observations posted on the websites BugGuide (VanDyk 2021) and iNaturalist (iNaturalist 2022) that could be verified as being *H. squamosus* (all identified as *H. apiatus*) were also analysed (Suppl. materials 1, 2). Of these records, 21 are published to GBIF, including 14 research-grade iNaturalist observations (GBIF.org 2022b).

Literature sources citing *H. squamosus* occurrences from additional localities (Blatchley and Leng 1916, Woodruff 1963, Hoffman 2003, Ciegler 2010, Fraedrich et al. 2011) were also included. Records of any type without at least county-level locality data (fewer than 10 overall) were excluded from further analysis.

Botanical names used are the accepted name from Tropicos.org (2021) at the time of access. Photographs of the syntype series of *H. apiatus* were provided by the Muséum National d'Histoire Naturelle (MNHN; Fig. 1). Habitus images of authoritatively identified *H. squamosus* (Fig. 2A-B) were taken with a BK Lab imaging system with a Canon EOS 7D camera equipped with a 65 mm macro lens. Images were stacked with Zerene Stacker. Distribution maps were created with SimpleMapp (Shorthouse 2010; Fig. 2C).

Taxon treatment

Heilipus squamosus (LeConte, 1824)

Nomenclature

Original combination: *Pissodes squamosus* LeConte, 1824: 161

Description: <https://www.biodiversitylibrary.org/page/15913340>

Holotype: <https://mczbase.mcz.harvard.edu/guid/MCZ:Ent:5176>

Synonyms:

Heilipus squamosus Boheman, 1836: 171 (not LeConte, 1824).

Description: <https://www.biodiversitylibrary.org/page/4109434>

Distribution

The previously-published state distribution of *Heilipus squamosus* is in the southeastern United States—Georgia (LeConte 1824), Florida (LeConte and Horn 1876), Tennessee (Blatchley and Leng 1916), South Carolina, Alabama and Mississippi (Wolfenbarger 1950), North Carolina (O'Brien and Wibmer 1982) and Virginia (Hoffman 2003). The species was considered rare (LeConte 1824, LeConte and Horn 1876, Barber 1912) and, apart from specimens from Florida where the species was a pest on avocado, relatively few specimens of this large and distinctive weevil have been collected and deposited in natural history collections.




Figure 2. [doi](#)

Heilipus squamosus. Scale bars = 2 mm. A: lateral habitus. B: dorsal habitus. C: distribution map (circles = iNaturalist/BugGuide records, stars = specimen data, squares = literature records; red symbols = 20th century records, black symbols = 21st century records).

In the past 20 years, citizen science initiatives (e.g. BugGuide, iNaturalist) have yielded observations of *H. squamosus* from all States in its previously reported range plus seven additional States, expanding the distribution of this species to the northern and western regions of the United States:

Alabama, Arkansas (**new State record**), Florida, Georgia, Kansas (**new State record**), Kentucky (**new State record**), Louisiana (**new State record**), Missouri (**new State record**), Mississippi, North Carolina, Oklahoma (**new State record**), South Carolina, Tennessee, Texas (**new State record**) and Virginia (Fig. 2C).

Biology

The biology of *H. squamosus* is still incompletely known. Before it became a pest of non-native avocado in Florida, the published natural history information was limited to a record from “under pine bark” (LeConte and Horn 1876: 141) and from “budding shoots of sassafras” (Blatchley and Leng 1916: 187). The avocado industry in Florida traces back to the early 20th century, though *Persea americana* Mill. was first brought to and became naturalised in Florida before 1835 (Wolfe et al. 1942). Wolfenbarger (1948) reports the earliest evidence of *H. squamosus* causing damage to avocado to be from the late 1930s and, by the late 1940s, the weevil became “one of the most dangerous pests of the avocado” (Wolfe et al. 1949). Larvae feed under the bark near ground level and can girdle trees and cause an 8–10% loss of avocado trees in a grove (Wolfenbarger 1948). The status of *H. squamosus* as a pest appears to have continued through the 1950s, but the species was rarely observed after (Wolfenbarger 1971).

The native hosts for the weevil have been suspected to be other members of the Lauraceae, namely *Lindera melissifolia* (Walter) Blume (pondberry), *Litsea aestivalis* (L.) Fernald (pondspice), *Persea borbonia* (L.) Spreng. (redbay) and *Sassafras albidum* (Nutt.) Nees (sassafras), though no definitive rearing records are known. Of these species, sassafras is widely distributed throughout the eastern United States; the other three are restricted to coastal plains and swampy areas, with pondberry and pondspice being listed as a federally endangered and threatened species, respectively.

Adult weevils have been found on pondberry in North Carolina (U.S. Fish and Wildlife Service 1993) and on pondspice in Georgia (Cameron et al. 2008, Fraedrich et al. 2011). An unidentified weevil larva described as being a similar size to the adult *H. squamosus* was found causing feeding damage to the root collar at the base of a pondberry plant in South Carolina (Fraedrich et al. 2011). Redbay was suggested as a possible host due to the weevil's distribution along the East Coast (Barber 1912, Hoffman 2003); Cameron et al. (2008) observed debarking damage to redbay branches which was noted to be very similar to the damage observed on pondspice branches at a locality in Georgia where *H. squamosus* was also found, but no more direct associations are known with redbay. Adult weevils have been collected on sassafras in Tennessee (Blatchley and Leng 1916) and Alabama [AUMNH].

Additional records: Wolfenbarger (1948) reported specimens collected “on cotton, sassafras and Satsuma orange” and, in a subsequent paper (Wolfenbarger 1950), he adds “camphor” to the list. Camphor, *Cinnamomum camphora* (L.) J. Presl, is a non-native species of Lauraceae naturalised in parts of south-eastern United States. No additional records were found indicating an association of *H. squamosus*. Three specimens have been collected from Isle of Wight Co., Virginia on *Diospyros virginiana* L. (wild persimmon; Ebenaceae) (CMNC; Art Evans, *pers. comm.*). One specimen from Spotsylvania Co., Virginia was photographed “on dogbane hemp”, *Apocynum cannabinum* L. (Apocynaceae) (iNaturalist, BugGuide).

Discussion

Heilipus squamosus was previously known from eight States in the south-eastern United States and is herein reported from an additional seven States, expanding the distribution in the country to the west and north. The reason for the poorly-documented distribution is likely related to the weevil's rarity, as well as a possible recent range expansion. The species has been historically reported to be rare and this is supported by only 57 adult specimens of this large and distinctive species being located in natural history collections. Removing records from Florida, where *H. squamosus* is known to be a pest on avocado, 33 specimens were found from only nine of the other 14 States where the species is now recorded. The rarity of the species can possibly be attributed to the rarity of the potential host plants: pondberry and pondspice. Pondberry is thought to have been uncommon even before recent habitat destruction and extant populations are known from only a few counties in each State in its range (Devall 2013). Of the newly-recorded States, two of three records from Arkansas and the only record from Missouri occur either in the same county (Craighead Co., AR) or a neighbouring county (White Co., AR; Carter Co., MO) to where the few extant populations of pondberry are known in these States (Devall 2013). The distribution of pondberry and pondspice alone do not account for the full distribution of *H. squamosus* as both plant species are absent from Kansas, Kentucky, Oklahoma, Tennessee and Texas (Gramling 2010, Devall 2013). Redbay ranges into east Texas and, while sassafras is present in each State, it apparently does not occur as far east as Tulsa Co., OK, where *H. squamosus* was observed (Randolph 2017). Other native Lauraceae, such as *Lindera benzoin* (L.) Blume, do occur in the full range of *H. squamosus*, but no associations of the weevil with these plants are known.

Another explanation for the poorly-documented distribution is that the range of *H. squamosus* has recently expanded. In the 21st century, citizen science initiatives like BugGuide and iNaturalist have provided an easy way for users to contribute observation data. Forty-three additional records of *H. squamosus* came from these sources from all 15 States from where the weevil is currently recognised, including the only known observations from five of the seven newly-recorded States (Fig. 2C, black circles). Such a large and distinctive species would be expected to be present in natural history collections if it had occurred in a region for a long time. So far, no specimens collected in Arkansas, Kansas, Missouri, Oklahoma or Texas were found in collections, suggesting that the weevil has not been present in these States for very long. All known records for those five States plus Kentucky have occurred in the 21st century (Fig. 2C, black circles). The iNaturalist observation data also indicate that *H. squamosus* may not actually be rare in the newly-recorded States. Seven total specimens were observed from Tulsa County, Oklahoma and Madison County, Kentucky since July 2020—the same number of records that have been found from all sources from the State of Georgia, from which *H. squamosus* has been recorded for nearly two centuries!

Native Lauraceae, including the possible host plants discussed here, are threatened by the spread of laurel wilt disease (Cameron et al. 2008, Gramling 2010, Fraedrich et al. 2011, Devall 2013, Randolph 2017, Best and Fraedrich 2018). It is currently unclear whether this,

or other factors such as climate change, are driving changes to the distribution and/or abundance of the weevil historically known only in the southeast. While the only avocado-producing region in which *H. squamosus* is currently known to occur is Florida, the ability of this species to live in 14 other States indicates that the species might also be able to continue spreading to and thrive in other avocado-producing regions (e.g. California). Locating more natural history collection specimens and making new observations will reveal more about the biology and distribution of this incompletely known species.

Conclusion

- *Heilipus squamosus* (LeConte, 1824) is the valid name for the single species of *Heilipus* occurring in the United States. Much of the prior literature and current internet resources refer to this species as *H. apiatus* (Olivier, 1807), which is an incorrect name for this species.
- *Heilipus squamosus* was previously recorded from Alabama, Florida, Mississippi, North Carolina, South Carolina, Tennessee and Virginia and is newly recorded from Arkansas, Kansas, Kentucky, Louisiana, Missouri, Oklahoma and Texas.
- Possible host plants of *H. squamosus* include *Lindera melissifolia* (Walter) Blume (pondberry), *Litsea aestivalis* (L.) Fernald (pondspice), *Persea borbonia* (L.) Spreng. (redbay) and *Sassafras albidum* (Nutt.) Nees (sassafras), which are all species of Lauraceae, although larval associations with these plants are currently limited.
- *Heilipus apiatus* (Olivier, 1807) is a valid name that applies to a species known only from French Guiana.

Acknowledgements

This research was made possible, in part, by a PPA 7721-funded Cooperative Agreement (AP21PPQS&T00C035) from the United States Department of Agriculture's Animal and Plant Health Inspection Service (APHIS). It may not necessarily express APHIS' views. We would like to thank the following collections personnel for looking for specimens in their collection and providing records if specimens were present: Robert Anderson (CMNC), Thomas Atkinson (University of Texas), Victoria Bayless (LSAM), Bob Blinn (NCSU), Janet Braun (University of Oklahoma), Melissa Callahan (AUMNH), Eric Chapman (University of Kentucky), Art Evans (VMNH), Rick Hoebeke (UGCA), Terence Schiefer (MEM), Kyle Schnepf (FSCA) and Kristin Simpson (University of Missouri). Additionally, thank you to iNaturalist users Thomas Shahan and John Abrams for providing more information on their observations in Oklahoma and Kentucky, respectively and Antoine Mantilleri, H el ene Perrin and Maxime Le Cesne for locating and providing images of *H. apiatus* syntypes at the MNHN.

References

- Anderson RS (2002) Molytinae Schoenherr 1823. In: Arnett Jr. RH, Thomas MC, Skelley PE, Frank JH (Eds) American beetles - Polyphaga: Scarabaeoidea through Curculionoidea. Volume 2. CRC Press, Boca Raton, 786-792 pp.
- Barber HS (1912) Note on the avocado weevil (*Heilipus lauri* Boheman). Proceedings of the Entomological Society of Washington 14: 181-183. URL: <https://www.biodiversitylibrary.org/item/20238#page/217/mode/1up>
- Berg C (1881) Entomologisches aus dem Indianergebiet der Pampa. Stettiner Entomologische Zeitung 42 (1-3): 36-72. URL: <https://www.biodiversitylibrary.org/item/35945#page/42/mode/1up>
- Best GS, Fraedrich SW (2018) An assessment of the potential impact of laurel wilt on clonal populations of *Lindera melissifolia* (Pondberry). Southeastern Naturalist 17 (4): 616-628. <https://doi.org/10.1656/058.017.0409>
- Blatchley WS, Leng CW (1916) Rhynchophora or weevils of North Eastern America. The Nature Publishing Company, Indianapolis <https://doi.org/10.5962/bhl.title.1557>
- Cameron RS, Bates C, Johnson J (2008) Distribution and spread of laurel wilt disease in Georgia: 2006-08 Survey and Field Observations. Georgia Forestry Commission.
- Ciegler JC (2010) Weevils of South Carolina (Coleoptera: Nemonychidae, Attelabidae, Brentidae, Ithyceridae, and Curculionidae). Biota of South Carolina. Vol. 6. Clemson University Public Service Publishing, Clemson, South Carolina, 276 pp. [ISBN 9780979877742]
- Devall MS (2013) The endangered pondberry (*Lindera melissifolia* [Walter] Blume, Lauraceae). Natural Areas Journal 33 (4): 455-465. <https://doi.org/10.3375/043.033.0409>
- Díaz V, Equihua-Martínez A, Romero-Nápoles J, Valdes-Carrasco J, Vallejo-Espinosa LF (2021) Nuevos registros de especies y un hospedero vegetal del género *Heilipus* Germar (Curculionidae: Molytinae: Hylobiini) para Colombia. Acta Zoológica Mexicana (nueva serie) 37: 1-12. <https://doi.org/10.21829/azm.2021.3712334>
- Ecdysis (2022) Ecdysis. A portal for live-data arthropod collections. <https://serv.biokic.asu.edu/ecdysis/index.php>. Accessed on: 2022-1-05.
- Fraedrich SW, Harrington TC, Bates CA, Johnson J, Reid LS, Best GS, Leininger TD, Hawkins TS (2011) Susceptibility to laurel wilt and disease incidence in two rare plant species, pondberry and pondspice. Plant Disease 95 (9): 1056-1062. <https://doi.org/10.1094/PDIS-11-10-0841>
- GBIF.org (2022a) GBIF Occurrence Dataset. Release date: 2022-8-24. URL: <https://doi.org/10.15468/e3gyvh>
- GBIF.org (2022b) GBIF Occurrence Download. Release date: 2022-8-22. URL: <https://doi.org/10.15468/dl.dsx827>
- Gemminger M, Harold Ev (1871) Catalogus Coleopterorum hucusque descriptorum synonymicus et systematicus. Vol. 8. Curculionidae. E. H. Gummi (G. Beck), Monachii, 2181-2668 pp. URL: <https://www.biodiversitylibrary.org/item/37367#page/9/mode/1up>
- Germar EF (1824) Insectorum species novae aut minus cognitae, descriptionibus illustratae. Vol. 1. Coleoptera. J.C. Hendelii et filii, Halae, xxiv + 624 pp. <https://doi.org/10.5962/bhl.title.130964>

- Gombauld P (2004) Introduction aux Charançons de Guyane. In: Gombauld P (Ed.) Insectes de Guyane: Beauté et Diversité. SEPANGUY, Collection Nature Guayanaise, Cayenne, Pp. 95–104 pp.
- Gramling JM (2010) Potential effects of laurel wilt on the flora of North America. *Southeastern Naturalist* 9 (4): 827-836. <https://doi.org/10.1656/058.009.0417>
- Hoffman RL (2003) *Heilipus apiatus*, a striking large weevil new to the Virginia fauna (Coleoptera: Curculionidae). *Banisteria* 22: 58-59. URL: <https://www.biodiversitylibrary.org/page/57144986#page/60/mode/1up>
- iNaturalist (2022) <https://www.inaturalist.org>. Accessed on: 2021-8-20.
- LeConte J (1824) New Coleopterous insects of North America. *Annals of the Lyceum of Natural History of New York* 1: 169-173 + pl. XI. URL: <https://www.biodiversitylibrary.org/item/54048#page/179/mode/1up>
- LeConte JL, Horn GH (1876) The *Rhynchophora* of America North of Mexico. *Proceedings of the American Philosophical Society* 15 (96): i-xvi + 1-455. URL: <https://www.biodiversitylibrary.org/item/85593#page/8/mode/1up>
- O'Brien CW, Wibmer GJ (1982) Annotated checklist of the weevils (Curculionidae sensu lato) of North America, Central America, and the West Indies (Coleoptera: Curculionidae). *Memoirs of the American Entomological Institute* 34: 1-382.
- O'Brien CW, Wibmer GJ (1984) Annotated checklist of the weevils (Curculionidae sensu lato) of North America, Central America, and the West Indies - Supplement I. *Southwestern Entomologist* 9 (3): 286-307.
- Olivier AG (1807) Entomologie, ou Histoire Naturelle des Insectes, avec leurs caractères génériques et spécifiques, leur description, leur synonymie, et leur figure enluminée. Coléoptères. Vol. 5. Desray, Paris, 612 pp. URL: <https://www.biodiversitylibrary.org/item/125992#page/11/mode/1up>
- Randolph KC (2017) Status of *Sassafras albidum* (Nutt.) Nees in the presence of laurel wilt disease and throughout the eastern United States. *Southeastern Naturalist* 16 (1): 37-58. <https://doi.org/10.1656/058.016.0104>
- Rheinheimer J (2010) Les Hylobiini de Guyane (Coleoptera, Curculionidae). In: Tourout J (Ed.) Contribution à l'étude des coléoptères de Guyane. Tome I. – Supplément au Bulletin de liaison d'ACOREP-France "Le Coléoptériste". 59-72 pp.
- Rheinheimer J (2017) Neue Arten der Tribus Hylobiini und Cryptorhynchini aus Französisch Guayana. *Koleopterologische Rundschau* 87: 297-324.
- SCAN (2022) The Symbiota Collections of Arthropods Network (SCAN) serves specimen occurrence records and images from North American arthropod collections. <https://scan-bugs.org/>. Accessed on: 2022-1-05.
- Schoenherr CJ (1836) Genera et species curculionidum, cum synonymia hujus familiae. Species novae aut hactenus minus cognitae, descriptionibus a Dom. Leonardo Gyllenhal, C.H. Boheman, et entomologis aliis illustratae. Vol. 3 (1). Roret, Paris, 1-505 pp. URL: <https://www.biodiversitylibrary.org/item/24767#page/5/mode/1up>
- Schoenherr CJ (1843) Genera et species curculionidum, cum synonymia hujus familiae. Species novae aut hactenus minus cognitae, descriptionibus a Dom. Leonardo Gyllenhal, C.H. Boheman, et entomologis aliis illustratae. Vol. 7 (2). Roret, Paris, 1–461 pp. URL: <https://www.biodiversitylibrary.org/item/24771#page/499/mode/1up>
- Shorthouse DP (2010) SimpleMappr, an online tool to produce publication-quality point maps. <https://www.simplemappr.net>. Accessed on: 2021-11-30.

- Tropicos.org (2021) Missouri Botanical Garden. <http://www.tropicos.org>. Accessed on: 2021-11-30.
- U.S. Fish and Wildlife Service (1993) Recovery plan for pondberry (*Lindera melissifolia*). U.S. Fish and Wildlife Service, Atlanta, Georgia, 56 pp.
- VanDyk J (Ed.) (2021) BugGuide.Net: Identification, images, & information for insects, spiders & their kin for the United States & Canada. Iowa State University. <https://bugguide.net/>. Accessed on: 2021-11-30.
- Wibmer GJ, O'Brien CW (1986) Annotated checklist of the weevils (Curculionidae sensu lato) of South America (Coleoptera: Curculionoidea). *Memoirs of the American Entomological Institute* 39: i–xvi + 1-563.
- Wolfe HS, Toy LR, Stahl AL (1942) Avocado production in Florida. Cooperative Extension Work in Agriculture and Home Economics, Gainesville, Florida. Bulletin 112, 111 pp.
- Wolfe HS, Toy LR, Stahl AL (1949) Avocado production in Florida. Cooperative Extension Work in Agriculture and Home Economics, Gainesville, Florida. Bulletin 141, 110 pp.
- Wolfenbarger DO (1948) *Heilipus squamosus* Lec., a new enemy of the avocado. *Proceedings of the Florida State Horticultural Society* 61: 260-264.
- Wolfenbarger DO (1950) On the distribution of *Heilipus squamosus* (Lec.) a pest of the avocado. *The Florida Entomologist* 33 (4): 139-141. <https://doi.org/10.2307/3492736>
- Wolfenbarger DO (1971) Updating of changes in pests, pesticides and other factors affecting subtropical insect pest control. *Proceedings of the Florida State Horticultural Society* 84: 318-320.
- Woodruff RE (1963) An avocado weevil (*Heilipus apiatus* Oliv.) (Coleoptera: Curculionidae). Florida Department of Agriculture. Contribution No. 19, Entomology Circular No. 11.

Supplementary materials

Suppl. material 1: *Heilipus squamosus* records observed in this study, separated by record type [doi](#)

Authors: Salvatore Anzaldo. Data provenance: cited natural history collections, repositories and authors.

Data type: Excel spreadsheet

Brief description: An Excel spreadsheet with three tables: 1) iNaturalist and BugGuide observations, 2) natural history collection specimens and 3) literature records.

[Download file](#) (23.99 kb)

Suppl. material 2: iNaturalist records of *H. squamosus* [doi](#)

Authors: iNaturalist

Data type: Observation data in a .csv file.

Brief description: The file includes all records (not only research grade) on iNaturalist at the time of access pertaining to the species *Heilipus squamosus*.

[Download file](#) (11.49 kb)