



# Species conservation profile and revision of *Salvia korolkowii* (Lamiaceae, Lamiales), a narrow endemic of the Western Tian-Shan

Alexander Sennikov<sup>‡</sup>, Obidjon Turdiboev<sup>§</sup>

<sup>‡</sup> University of Helsinki, Helsinki, Finland

<sup>§</sup> Institute of Botany, Tashkent, Uzbekistan

Corresponding author: Alexander Sennikov ([alexander.sennikov@helsinki.fi](mailto:alexander.sennikov@helsinki.fi))

Academic editor: Dimitrios Koureas

Received: 23 Jun 2022 | Accepted: 28 Jul 2022 | Published: 12 Aug 2022

Citation: Sennikov A, Turdiboev O (2022) Species conservation profile and revision of *Salvia korolkowii* (Lamiaceae, Lamiales), a narrow endemic of the Western Tian-Shan. Biodiversity Data Journal 10: e89437. <https://doi.org/10.3897/BDJ.10.e89437>

## Abstract

### Background

*Salvia korolkowii* (Lamiaceae) has been considered a species of highest conservation priority due to its narrow distribution and isolated taxonomic position. The species has been known from Uzbekistan and, questionably, Kyrgyzstan and treated as endemic to the Western Tian-Shan. Its modern conservation status according to the IUCN Criteria has not been established.

### New information

The taxonomic position of *Salvia korolkowii* is evaluated; the species is treated as the sole member of *S.* sect. *Odontochilus* (Pobed.) Sennikov, comb. nov. because of its morphological differences and phylogenetic isolation. The herbarium collections are completely revised and the species is proven to occur mainly in Uzbekistan with a single locality (new country record) in Kazakhstan; its former report from Kyrgyzstan (one locality) is confirmed and documented by herbarium specimens. The species occurrences are

mapped and its conservation status is assessed as Vulnerable due to the restricted size and continuous decline of its populations because of the ongoing degradation and destruction of its primary habitat (variegated outcrops) at lower elevations. This study highlights the importance of a thorough revision of herbarium collections in Central Asia for conservation purposes.

## Keywords

Central Asia, conservation, IUCN assessment, Kazakhstan, Kyrgyzstan, Labiatae, Mountains of Central Asia biodiversity hotspot, Tian-Shan, Uzbekistan

## Introduction

The genus *Salvia* L. (Lamiaceae Martinov, Mentheae Dumort., Salviinae Endl.), encompassing nearly 1000 species (Kriebel et al. 2019), has undergone marked species radiations in three regions of the world: Central and South America, Central Asia/Mediterranean and Eastern Asia, where it colonised different biomes and adapted to different pollinators (Drew et al. 2017, Hu et al. 2018, Kriebel et al. 2019). A broad-scale phylogeny of the genus was developed and demonstrated that the genus comprises three major clades, which largely correspond to the Mediterranean and south-western Asian, eastern Asian and New World centres of its species diversity (Kriebel et al. 2019). Over a half of its total species diversity falls into the New World, which is a secondary radiation area, whereas the ancestral area for the genus radiation was placed to the Irano-Turanian floristic region, which is the second great centre of its species diversity (Walker et al. 2004, Kriebel et al. 2019).

Recent phylogenetic studies (Drew et al. 2017, Hu et al. 2018) argued that a broad circumscription of *Salvia* is preferable, although an opposite view has been advocated by some researchers (Will and Claßen-Bockhoff 2017). In a broad sense, *Salvia* is characterised by two anterior fertile stamens, with two thecae on each stamen (one theca may be sterile) separated by a variably elongated connective tissue, whereas the other genera of Mentheae possess four fertile stamens (Drew et al. 2017 and references therein). In this circumscription, the broad *Salvia* includes some small genera traditionally recognised in Central Asia and the former USSR (e.g. *Perovskia* Kar., *Rosmarinus* L.), which are deeply embedded within *Salvia* on the phylogenetic tree (Drew et al. 2017) and, therefore, abandoned in our latest synoptic revisions (Tojibaev et al. 2021). Morphological differences between major phylogenetic clades of *Salvia* are largely limited to the characters of the elongated connective tissue that separates thecae in stamens, are at least partly driven by changes between different pollinators (Drew et al. 2017) and evolved parallel variants of the staminal architecture independently in each part of the world (e.g. Fragoso-Martínez et al. 2018).

*Salvia* is well represented in the mountainous regions of Central Asia. Its species grow on mountain slopes and outcrops of variegated beds, in foothills and the lower mountain belt.

According to Makhmedov (1984a), 21 and 11 species of *Salvia* in its traditional circumscription occur in the Pamir-Alai Mountains and in the Western Tian-Shan, respectively. The expanded *Salvia* counted 41 species in Central Asia (Turdiboev et al. 2022), although this figure did not take into account the synonymy proposed by Lazkov (2016) and Tojibaev et al. (2021) and should, therefore, be corrected. *Salvia korolkowii* Regel & Schmalh. (Regel 1879), *S. tianschanica* Makhm. (Makhmedov 1980) and *S. vvedenskii* Nikitina (Nikitina 1962) of the Western Tian-Shan are currently listed in the Red Data Books of Kyrgyzstan (Mosolova 2006) and Uzbekistan (Tojibaev 2009) as rare and subendemic plants.

Being part of the Mountains of Central Asia biodiversity hotspot (Körner 2004, Spehn et al. 2005), the Western Tian-Shan is rich in plant diversity (3440 species and subspecies) and endemic plants (634 taxa) (Tojibaev et al. 2021). *Salvia korolkowii* is endemic to the Western Tian-Shan (Tojibaev et al. 2020) and occurs in rare and scattered populations. The species has potential ornamental and medicinal values (Mamadaliyeva et al. 2017).

While working on the Flora of Uzbekistan project (Sennikov et al. 2016), we revised herbarium specimens of *Salvia* from Central Asia stored at TASH and LE. During this revision, we have found that the published data on *S. korolkowii* are highly incomplete and its conservation assessment is badly outdated. Although this species has been legally protected for nearly 50 years, its distribution remains poorly researched and its taxonomic position has not been updated since Pobedimova (1954). Its latest regional revision (Lazkov 2016) highlighted a number of open questions, which should be resolved for a modern conservation assessment under the IUCN.

The aim of the present contribution is to resolve historical and nomenclatural problems connected with *S. korolkowii* and to re-assess the distribution and conservation status of this species, which has been under the highest conservation priority in Kyrgyzstan (Sharashova 1985, Mosolova 2006) and Uzbekistan (Makhmedov 1984b, Tojibaev 2009) since the times of the USSR (Takhtajan 1975, Takhtajan 1981, Makhmedov and Denisova 1984), due to its restricted distribution area and isolated taxonomic position.

## Material and methods

The data about *Salvia korolkowii* (history of discovery, distribution, ecology, morphology, taxonomic position) were collected from literature, herbarium collections and online depositories of published observations.

The nomenclature was re-assessed on the basis of the protologue (Regel 1879) and historical literature (Lipsky 1905), according to the International Code of Nomenclature of algae, fungi and plants (Turland et al. 2018). Original collections were examined at the Komarov Botanical Institute of Russian Academy of Sciences (LE).

Morphological characters were examined from taxonomic revisions (Pobedimova 1954, Lazkov 2016). The phylogenetic position and taxonomy was assessed on the basis of

phylogenetic (Drew et al. 2017, Will and Claßen-Bockhoff 2017, Hu et al. 2018) and taxonomic (Kudryashev 1937, Pobedimova 1954, Makhmedov 1984a) literature.

A distributional dataset (Turdiboev and Sennikov 2022) is based on our comprehensive *de visu* revision of herbarium specimens kept at LE and TASH; several smaller collections of Central Asian plants (at E, H, MW, NL, NSK, P, PE, TALL and ZT) were examined online. The specimen data were complemented with documented observations (Plantarium 2022). The occurrences were georeferenced with the help of printed maps adjusted with Google Maps, using the utmost accuracy available; historical records from the same locality were linked to a single occurrence when no further precision was available (Usmonov et al. 2021). Place names were recorded according to the national languages (Kazakh, Kyrgyz, Uzbek); their spelling adopted here may deviate from the international tradition which is based on the Russian spelling.

To assess the threat status of the species, IUCN Red List Categories and Criteria were employed (IUCN 2012). The geospatial analysis was performed using the Geospatial Conservation Assessment Web Tool GeoCAT (Bachman et al. 2011). The AOO was based on a recommended cell width of 2 km.

## Results

### Historical background

The first difficulty connected with the name *Salvia korolkowii* is bibliographic. This species was described by Eduard Regel in a series of his numerous contributions of new species described on the basis of dried and living plant collections of the Imperial Botanical Garden in Saint-Petersburg. His contribution (fasciculus) 7 appeared in part 2 of volume 6 of the main periodical of the garden, *Acta Horti Petropolitani* (Regel 1880). At that time, due to the massive size of contributions, the journal appeared in bulky volumes consisting of a few large-sized parts, usually with three or more papers in each. In such cases, production and printing was rather slow and the volume parts were inconvenient in private circulation; for this reason, many significant papers were printed separately and distributed privately by the authors, whereas the journal volumes were intended for library use. For practical reasons, individual papers were typically preprinted immediately after typesetting by the same commercial printer (brothers Schumacher for volume 6), rather than reprinted from journal volumes, thus pre-dating the journal publication and distribution. The preprints were separately paginated, omitting the journal pagination, whereas the typeset remained identical. For this reason, bibliographically, the instalments of Regel's *Descriptiones plantarum novarum* may be treated as books rather than articles in periodicals and the nomenclatural novelties published in that series may be cited from those books rather from the journals. The time difference between preprints and journal volumes (or their parts) was sufficiently short to fall within the same calendar year, making the distinction rather impractical. However, the situation was different for Regel's fasciculus 7 (Regel 1879),

which was printed and distributed in 1879, whereas its volume part was dated 1880. This difference is significant and should be taken into account.

*Salvia korolkowii* Regel & Schmalhausen (Regel 1879: 70) was described with an indication of its provenance in Latin: "In Turkestan occidentali prope Chodschent 3000' alt.; inter Ak-tag-tau et Ak Cagoti, 5–8000' alt. (Korolkow)". The protologue states that the plants were collected by a Russian military officer in Turkestan, Colonel (thereafter General) Nikolai I. Korolkov near Xuçand (also spelled Khodzhent in Russian) Town, now in northern Tajikistan, probably implying the southern part of the Kurama Range. Kudryashev (1937) suggested that this indication is a spelling mistake and the locality should rather read Xo'jakent (also spelled Khodzhakent), now in eastern Uzbekistan, situated along Chirchiq River between the Qorjontov Range and the Chatqol Range. This conclusion is confirmed by the analysis of Korolkov's herbarium collections published by Lipsky (1905), who found that Korolkov collected near Burchmullo Village at the stated elevation, and by the other collections cited in Regel (1879), where "Chodschent" is clearly linked with Burchmullo. Climatic conditions in the two locations differ (Kudryashev 1937) and, according to the published data, *S. korolkowii* does not occur at Xuçand (Zhogoleva and Kochkareva 1986).

The other locality mentioned in the protologue is situated near Chimgon (Chimyon), very close to Xo'jakent and Burchmullo. As noted by Lipsky (1905), Korolkov collected near Chimgon and Xo'jakent in June of 1872, travelling between Chimgon (Chimyon) Mt. and Oqsoqotasoy River. The toponyms mentioned in the protologue can be interpreted as follows: Ak-tag-tau = Oqtaxta Pass, Ak Cagoti = Oqsoqotasoy River; this apparently denotes the upper course of the river, which starts under that pass.

This means that all the original localities of *S. korolkowii* are situated in the mountains surrounded by the Chirchiq, Chatqol, Oqsoqotasoy and Teraklisoy Rivers (Chatqol Range, Western Tian-Shan), within the area of ca. 30 km in diameter. Among the collections examined at LE, only one gathering (collected between Oqsoqotasoy and Oqtaxta) was found. The species nomenclature is as follows.

***Salvia korolkowii*** Regel & Schmalh., Trudy Imp. S.-Peterburgsk. Bot. Sada 6: 356 [repr. 70] (1879) ≡ *Schraderia korolkowii* (Regel & Schmalh.) Pobed. in Schischkin, Fl. URSS 21: 373 (1954) ≡ *Arischrada korolkowii* (Regel & Schmalh.) Pobed., Novit. Syst. Pl. Vasc. 9: 247 (1972) ≡ *Stiefia korolkowii* (Regel & Schmalh.) Soják, Čas. Nár. Mus., Odd. Přír. 152(1): 22 (1983).

Type: UZBEKISTAN. "Inter Ak-tag-tau et Ak Cagoti, 5–8000' alt." [between Oqsoqotasoy River and Oqtaxta Pass, 1500–2400 m a.s.l.], June 1872, *Korolkoff s.n.* (lectotype LE0051698, designated by Lazkov (2016); isolectotypes LE0051697, LE01072812, K000929822 and P02864044).

The species name was spelled "Korolkowi" in the protologue. This spelling is correctable to "korolkowii" according to Art. 60.8(b). In modern databases, the correct spelling was adopted in the Leipzig Catalogue of Vascular Plants (Freiberg et al. 2020, Freiberg 2020),

whereas the World Checklist of Vascular Plants (Govaerts et al. 2021, The Royal Botanic Gardens, Kew 2021) used the wrong spelling "korolkovii".

When designating the lectotype, Lazkov (2016) did not specify the particular sheet because of the lack of appropriate identifiers at that time. The lectotype specimen was clearly labelled as "Type" and, therefore, Lazkov's lectotype designation was unambiguous and effective.

### Taxonomic classification

The taxonomic position of *Salvia korolkowii* varied with time, when researchers realised that the early attempts to circumscribe and subdivide the genus *Salvia* were unnatural and, therefore, unsatisfactory. This species was originally assigned to *S. sect. Hymenospace* Benth. (Regel 1879), which was treated very broadly in those times. The most extreme splitting of *Salvia* was upheld by Nevski (1937), who accepted many minor generic segregates including the genus *Schraderia* Medik. Pobedimova (1954) followed this approach and classified *Salvia korolkowii* in *Schraderia*. When *Schradera* Vahl was conserved against its near-homonym *Schraderia* Medik., Pobedimova (1972) renamed this segregate to *Arischrada* Pobed. At the next twist, Soják (1983) returned to the old nomenclature of Medikus (1791) and found another suitable generic name, thus renaming the species as *Stiefia korolkowii* (Regel & Schmalh.) Soják.

The phylogeny of *Salvia* (Dizkirici et al. 2015, Will and Claßen-Bockhoff 2017) revealed a broad paraphyly of the traditional *S. sect. Hymenospace*, in which *S. korolkowii* was retained by Makhmedov (1984a). Bentham (1833b) originally included 12 species into this section, of which *S. pomifera* L., *S. canariensis* L., *S. aurea* L. and *S. paniculata* L. were listed in its protologue (Bentham 1833a).

Pobedimova (Pobedimova 1954, Pobedimova 1972) understood that the broad *S. sect. Hymenospace* was too diverse and artificial. She subdivided this taxon (as *Schraderia*) into smaller, more natural groups. The species of this affinity occurring in Central Asia, *S. bucharica* Popov and *S. korolkowii*, she placed in separate sections, which were subsequently downgraded to subsections (Makhmedov 1982, Makhmedov 1984a, Lazkov 2016, Turdiboev and Turginov 2021). The relationship and separate positions of these sections were recently confirmed (Will and Claßen-Bockhoff 2017).

The sections differ in the following morphological characters (Pobedimova 1954): *Holochilus* (*S. bucharica*) – upper lip nearly entire, flowers pink or purple, floral leaves persistent, leaves pinnatisect, upper thecae connected and fertile; *Odontochilus* (*S. korolkowii*): upper lip tridentate, flowers yellow, floral leaves caducous, leaves simple, upper thecae free and sterile.

The correct type species of *S. sect. Hymenospace* is *S. pomifera*, which was designated by Makhmedov (1982). It is clear that the Central Asian species previously placed into *S. sect. Hymenospace* are not closely related to any species originally assigned to this section (Will and Claßen-Bockhoff 2017). For this reason, the system of narrowly defined

sections proposed by Pobedimova (Pobedimova 1954, Pobedimova 1972) can be retained and transferred to the nomenclature of *Salvia* as elaborated here.

***Salvia* sect. *Holochilus*** (Pobed.) Sennikov, **comb. nov.**  $\equiv$  *Schraderia* sect. *Holochilus* Pobed. in Schischkin, Fl. URSS 21: 664 (1954)  $\equiv$  *Arischrada* sect. *Holochilus* (Pobed.) Pobed., Novit. Syst. Pl. Vasc. 9: 247 (1972). Type: *Salvia bucharica* Popov. Species: *Salvia bucharica* Popov, *S. hydrangea* Benth., *S. maymanica* Hedge.

***Salvia* sect. *Odontochilus*** (Pobed.) Sennikov, **comb. nov.**  $\equiv$  *Schraderia* sect. *Odontochilus* Pobed. in Schischkin, Fl. URSS 21: 664 (1954)  $\equiv$  *Arischrada* sect. *Odontochilus* (Pobed.) Pobed., Novit. Syst. Pl. Vasc. 9: 247 (1972)  $\equiv$  *Salvia* subsect. ***Odontochilus*** (Pobed.) Makhm., Bot. Mater. Gerb. Inst. Bot. Akad. Nauk Uzbebsk. S.S.R. 20: 30 (1982). Type: *Salvia korolkowii* Regel & Schmalh. Monotypic section.

### Distribution area

The species was considered endangered due to its narrow distribution area (Makhmedov and Denisova 1984, Makhmedov 1984b). For this reason, it was mapped already in the 1980s (Makhmedov 1984a), but that mapping was extremely coarse and incomplete.

The species distribution is apparently centred in Uzbekistan, where the greatest majority of its localities are situated (Turdiboev and Sennikov 2022). Its occurrence in Kyrgyzstan (in "Chatkal River basin") was postulated (Sharashova 1960) seemingly because of the close proximity of localities in Uzbekistan to the border with Kyrgyzstan, but no specimens have been known to date (Mosolova 2006, Lazkov 2016). We found two specimens of *Salvia korolkowii* at LE which were collected in Kyrgyzstan by R. Kamelin in 1972, who unknowingly entered the territory from the neighbouring Uzbekistan along the Chatqol River and collected the plants within 0.5-2 km from the border. These are the first documented records of the species from Kyrgyzstan.

During our inventory of herbarium collections, quite unexpectedly we found one overlooked specimen at TASH, which was collected in 1924 near Trbat Village in Trkstan Region of Kazakhstan. This specimen documents the first record of *S. korolkowii* from the country.

According to our data, the localities of *S. korolkowii* are concentrated along tributaries of the Chirchiq River: Oqsoqotasoy, G'alvasoy, Qorango'Isoy, Chatqol, Oqbuloq, Ko'ksuv, Piskom and Ugom, with isolated localities along the Boshqizilsoy River and at Trbat Village. This area is situated within the Chatqol, Ugom, Ko'ksuv, Piskom and Qorjontov mountain ranges.

*Salvia korolkowii* occurs at lower elevations, mostly between 900 and 1200 m, but sometimes up to 1500-1700 m above sea level. In the Western Tian-Shan, lower elevations are very hot and highly arid; in this territory, the species prefers open stony substrates of red clay and sandstone mixed with gypsaceous sediments, which are called variegated denudations due to their alternation of red sandstone and white gypsum patches. This landscape is very rich in narrowly distributed plants (Popov 1923, Kamelin

2017) and seems to be a major contributor to the highest plant endemism recorded in the Western Tian-Shan (Tojibaev et al. 2021).

Makhmedov (1984b) noted that many populations of *S. korolkowii* are small but some may be significant in size. Due to its availability, the species was distributed to international scientific herbarium collections as exsiccata, collected along the Ko'ksuv (Makhmedov 1988) and Piskom (Pobedimova 1967) Rivers in 50 duplicates.

## Species Conservation Profile

### *Salvia korolkowii* Regel & Schmalh.

#### Species information

##### Synonyms

*Schraderia korolkowii* (Regel & Schmalh.) Pobed., *Arischrada korolkowii* (Regel & Schmalh.) Pobed., *Stiefia korolkowii* (Regel & Schmalh.) Soják.

##### Common names

Корольков мармараги (Uzbek Cyrillic), Шалфей Королькова (Russian), Корольков кек башы (Kyrgyz), Корольков сэлбен (Kazakh Cyrillic).

##### Taxonomy

Kingdom	Phylum	Class	Order	Family
Plantae	Spermatophyta	Magnoliopsida	Lamiales	Lamiaceae

##### Taxonomic notes

The species belongs to a presumably monotypic section, *Salvia* sect. *Odontochilus* (Pobed.) Sennikov. Its closest relative in Central Asia is *S. bucharica* Popov. (*S.* sect. *Holochilus* (Pobed.) Sennikov).

##### Region for assessment:

- Global

##### Figure(s) or Photo(s):

Figs 1, 2





Figure 1. [doi](#)

A plant of *Salvia korolkowii* at Xumson, Uzbekistan. Photographed by A. Gaziev, 1 June 2008. Source: <https://www.plantarium.ru/page/image/id/48037.html>.



Figure 2. [doi](#)

A large cushion of *Salvia korolkowii* along the Beldorsoy River, Uzbekistan. Photographed by A. Gaziev, 8 June 2014. Source: <https://www.plantarium.ru/page/image/id/246310.html>.

## Reviewers

Lazkov, G.A.

**Editor**

Sennikov, A.N. & Turdiboev, O.A.

**Geographic range****Biogeographic realm:**

- Palearctic

**Countries:**

- Uzbekistan
- Kazakhstan
- Kyrgyzstan

**Map of records (image):**

Fig. 3




Figure 3. [doi](#)

Distribution area of *Salvia korolkowii* according to the specimens examined and documented observations. Symbols: triangle - lectotype locality, dots - other records.

**Map of records (Google Earth):**

Suppl. material 1

**Basis of EOO and AOO:** Observed

**Basis (narrative)**

The species is known from several populations. It was estimated that ca. 200 local populations may exist in Uzbekistan (Makhmedov 1984b), where the species forms a compact core distribution area ca. 65 km in diam. with many documented localities, from which it has been repeatedly sampled for 150 years (Turdiboiev and Sennikov 2022). The species is highly conspicuous, easy to detect and identify and tends to occur in easy-to-access places; for this reason, we assume that the major part of its distribution has been revealed and most of its populations are known. Our data inventory has expanded the total known distribution area (Tojibaev 2009) by ca. 30 km only.

**Min Elevation/Depth (m):** 850

**Max Elevation/Depth (m):** 1700

**Range description**

The species habitat is mountainous, occurring along the Chirchik River and its tributaries (Oqsoqotasoy, G'alvasoy, Qorangko'lsoy, Chatqol, Oqbuloq, Ko'ksuv, Piskom and Ugom), in the Chatqol, Ugom, Ko'ksuv, Piskom and Qorjontov mountain ranges. The distribution area lies mostly in Uzbekistan, with a minor penetration into Kyrgyzstan and one isolated locality in Kazakhstan.

The species occurs in highly arid territories, mostly between 900 and 1200 m, but sometimes up to 1500-1700 m above sea level.

**Extent of occurrence**

**EOO (km<sup>2</sup>):** 2800

**Trend:** Stable

**Justification for trend**

Although the species is under high anthropogenic pressure because of many of its populations being situated in the immediate vicinity of populated places or roads, which may lead to destruction or impoverishing of some populations, there is no evidence that any part of its distribution area is lost.

**Causes ceased?:** No

**Causes understood?:** Yes

**Causes reversible?:** No

**Area of occupancy**

**AOO (km<sup>2</sup>):** 192

**Trend:** Decline (observed)

**Justification for trend**

Most of the localities of *Salvia korolkowii* in Uzbekistan are situated in the nearest proximity to populated places, roads and other areas of human activity. The species occurs mostly in the lower mountain belt, in the lower part of river valleys. Such locations are most easily accessible and susceptible to residential and recreational development, road construction and land use, especially close to the Chirchiq River; as an example, several populations were destroyed when the territory was inundated by the Chorvoq water reservoir (Makhmedov 1984b).

The most significant populations of the species are known along the Chatqol, Piskom and Ugom Rivers (Makhmedov 1984b, Turdiboev and Sennikov 2022). The species is still regularly encountered in accessible places along the Chirchiq and Chatqol Rivers (Plantarium 2022, Turdiboev and Sennikov 2022) despite the anthropogenic pressure.

**Causes ceased?:** No

**Causes understood?:** Yes

**Causes reversible?:** No

**Locations**

**Number of locations:** 25

**Justification for number of locations**

The species is documented from a number of localities, which can be treated as about 25 locations, based on the most significant threatening event, i.e. habitat destruction by large-scale actions (inundation, road construction, mining, development).

**Trend:** Decline (observed)

**Justification for trend**

At least one location was nearly destroyed under the Chorvoq water reservoir.

**Population**

**Number of individuals:** Estimated as less than 10000 mature individuals.

**Trend:** Decline (observed)

**Justification for trend**

The species is under constant pressure due to large-scale activities that lead to irreversible habitat destruction and degradation, and some of these activities (inundation, road

construction, development) are known to reduce the number of individuals, although the extent of this reduction cannot be estimated with certainty.

**Basis for decline:**

- (a) direct observation

**Causes ceased?:** No

**Causes understood?:** Yes

**Causes reversible?:** No

**Population Information (Narrative)**

The exact population size and the number of individuals are unknown. The count provided by Makhmedov (1984b) is based on extrapolations.

**Subpopulations**

**Abundance largest subpopulation:** 1000

**Number of subpopulations:** 25

**Trend:** Decline (inferred)

**Justification for trend**

Some subpopulations situated at lower elevations close to populated places are under ongoing threat because of urban development and road construction and may disappear or noticeably degrade in the future.

**Habitat**

**System:** Terrestrial

**Habitat specialist:** Yes

**Habitat (narrative)**

The species often grows on variegated beds, which are situated at elevations of ca. 900-1200 m a.s.l. This habitat is highly arid, much insolated and very warm; its vegetation cover is naturally very sparse.

More seldom, the species occurs also at higher elevations of 1500-1700 m a.s.l., in sparse steppoid plant communities on rocky slopes.

**Trend in extent, area or quality?:** Decline (observed)

**Justification for trend**

Variogated beds are highly prone to destruction because of their situation at lower elevations and close proximity to populated places. Habitat loss occurs due to expansion of populated places, road construction, human changes in watercourses etc.

**Habitat importance:** Major Importance

**Habitats:**

- 6. Rocky areas (e.g. inland cliffs, mountain peaks)

**Ecology**

**Size:** 30-50 cm

**Generation length (yr):** 30

**Dependency of single sp?:** No

**Ecology and traits (narrative)**

The plants are subshrubs forming a long vertical root and a strong branching caudex which is capable to last for several decades. Shoots bicyclic, developing a rosette in the first year and a leafy stem in the second year. Leaves long-petiolate, blades narrowly oblong in rosettes and ovate-oblong on stems, white-tomentose on the lower side. Flowering stems are very showy because of conspicuous yellow flowers 2.5-3 cm long, which are collected in large thyrsoïd inflorescences. Reproduction exclusively by seed.

**Threats**

**Threat type:** Ongoing

**Threats:**

- 1.1. Residential & commercial development - Housing & urban areas
- 1.3. Residential & commercial development - Tourism & recreation areas
- 2.3.2. Agriculture & aquaculture - Livestock farming & ranching - Small-holder grazing, ranching or farming
- 4.1. Transportation & service corridors - Roads & railroads

**Threat type:** Past

**Threats:**

- 7.2. Natural system modifications - Dams & water management/use

### Justification for threats

The species populations are situated at lower elevations and in the lower part of river ravines. For this reason, many populations are situated next to populated places and along roads and have been in danger because of continuous residential development and road construction. Some complementary threat comes from grazing, sports and recreation, which may damage individual plants.

In the past (1960s), some populations were destroyed by the construction of the Chorvoq water reservoir.

### Conservation

**Conservation action type:** In Place

**Conservation actions:**

- 1.1. Land/water protection - Site/area protection
- 5.1.2. Law & policy - Legislation - National level

### Justification for conservation actions

The species is not protected in any strict nature reserve, but its distribution area is largely covered by the Ugom-Chatqol National Park (Uzbekistan), in which some lands are protected, whereas economic activities and recreation are regulated. The only locality in Kazakhstan is situated within the Sairam-Ögem National Park.

Cultivated in the Tashkent Botanical Garden. Seeds have not been submitted to the Millennium Seed Bank or any similar institution.

The species is included in the Red Data Books of Uzbekistan (Tojibaev 2009) and Kyrgyzstan (Mosolova 2006) as a rare and vulnerable plant and is, therefore, under legal protection even outside strictly protected areas.

Using the IUCN Criteria (rather small population size and continuous decline), we suggest the global conservation status of *Salvia korolkovii* to be assessed as Vulnerable (criteria C1+2a(i)).

### Other

**Use type:** National

**Use and trade:**

- 5. Manufacturing chemicals
- 13. Pets/display animals, horticulture
- 16. Establishing ex-situ production \*

### Justification for use and trade

Despite its limited distribution, the species was considered as a source of essential and drying oils tested as a component of varnish (Zuckerwanik and Grachev 1948). The actual use is prevented by legal protection.

Although the plants are not picked up by local people and tourists, the species has a potential value as ornamental for plant enthusiasts. Such actual use is unknown.

There has been no noticeable harvest of the species, except for scientific purposes. The species is no longer collected for scientific distribution either.

The species is cultivated ex-situ in the Tashkent Botanical Garden.

**Ecosystem service type:** Less important

### Research needed:

- 1.2. Research - Population size, distribution & trends
- 3.1. Monitoring - Population trends

### Justification for research needed

Although we believe that the major part of the distribution area of *Salvia korolkowii* has been already detected, its exact localities and population size have not been properly studied and documented.

## IUCN Red List assessment

Due to its restricted extent of occurrence (2800 km<sup>2</sup>) and area of occupancy (190 km<sup>2</sup>), the species may qualify for Endangered, but we estimate the number of its locations as ca. 25. Its total population size is estimated as less than 10000 individuals and the projected loss is estimated at 10% within the future 100 years; for this reason, we assess its global threatened category as Vulnerable (criteria C1+2a(i)).

## Conclusions

The endemic flora of the Western Tian-Shan is very rich (Tojibaev et al. 2021) but has never been assessed for conservation purposes in its entirety. Modern inventories have been recently started with selected species of *Cousinia* (Usmonov et al. 2021) and *Tulipa* (Wilson et al. 2021). This effort should be continued to cover other species of the area with restricted distributions.

*Salvia korolkowii* has been legally protected since the 1970s in the USSR, based on the draft assessment of R.V.Kamelin who considered its isolated taxonomic position, showy habit and very narrow distribution area (Takhtajan 1975). This protection continues in



present-day Kyrgyzstan (Mosolova 2006) and Uzbekistan (Tojibaev 2009). Although the species does not occur in any strictly protected area, its distribution is largely covered by national parks in Kazakhstan and Uzbekistan, thus ensuring some protection for its native landscapes from large-scale destruction. However, many of its populations are under continuous threat of reduction or even extinction due to their proximity to populated places and communications, which makes the species vulnerable as assessed here.

Our study highlights the importance of herbarium collections in the botanical research of Central Asia. Even in cases of such well-known and unambiguous species as *Salvia korolkowii*, which have been in conservation focus for decades, historical data may contain some important but cryptic information which have never been evaluated and included in scientific works. Given the small size of distribution areas of many plants of the Western Tian-Shan, precise and comprehensive data inventory and mobilisation is required to overcome the obscurity and complexity of historical collections, which are often dispersed among various countries and require a special effort in their deciphering and evaluation (e.g. Sennikov 2021).

## Acknowledgements

Denis Melnikov (Saint-Petersburg) is thanked for communication about the type material at L.E. Open Access publication was funded by the Library of the University of Helsinki.

## References

- Bachman S, Moat J, Hill AW, de Torre J, Scott B (2011) Supporting Red List threat assessments with GeoCAT: geospatial conservation assessment tool. *ZooKeys* 150: 117-26. <https://doi.org/10.3897/zookeys.150.2109>
- Bentham G (1833a) Account of Indian Labiatae, in the collection of J. F. Royle. In: Hooker WJ (Ed.) *Botanical Miscellany*. 3. J. Murray, London, 370-384 pp.
- Bentham G (1833b) *Labiatarum genera et species*. 3. J. Ridgway & Sons, London, 189-323 pp.
- Dizkirici A, Celep F, Kansu C, Kahraman A, Dogan M, Kaya Z (2015) A molecular phylogeny of *Salvia euphratica* sensu lato (*Salvia* L., Lamiaceae) and its closely related species with a focus on the section Hymenosphace. *Plant Systematics and Evolution* 301: 2313-2323. <https://doi.org/10.1007/s00606-015-1230-1>
- Drew BT, González-Gallegos JG, Xiang C, Kriebel R, Drummond CP, Walked JB, Sytsma KJ (2017) *Salvia* united: The greatest good for the greatest number. *Taxon* 66 (1): 133-145. <https://doi.org/10.12705/661.7>
- Fragoso-Martínez I, Martínez-Gordillo M, Salazar GA, Sazatornil F, Jenks AA, del Rosario García Peña M, Barrera-Aveleida G, Benitez-Vieyra S, Magallón S, Cornejo-Tenorio G, Granados Mendoza C (2018) Phylogeny of the Neotropical sages (*Salvia* subg. *Calosphace*; Lamiaceae) and insights into pollinator and area shifts. *Plant Systematics and Evolution* 304: 43-55. <https://doi.org/10.1007/s00606-017-1445-4>

- Freiberg M (2020) The Leipzig catalogue of vascular plants. German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig <https://doi.org/10.15468/9qxm3>
- Freiberg M, Winter M, Gentile A, Zizka A, Muellner-Riehl AN, Weigelt A, Wirth C (2020) LCVP, The Leipzig catalogue of vascular plants, a new taxonomic reference list for all known vascular plants. *Scientific Data* 7 (1): 416. <https://doi.org/10.1038/s41597-020-00702-z>
- Govaerts R, Nic Lughadha E, Black N, Turner R, Paton A (2021) The World checklist of vascular plants, a continuously updated resource for exploring global plant diversity. *Scientific Data* 8 (1): 215. <https://doi.org/10.1038/s41597-021-00997-6>
- Hu GX, Takano A, Drew BT, Liu ED, Soltis DE, Soltis PS, Peng H, Xiang CL (2018) Phylogeny and staminal evolution of *Salvia* (Lamiaceae, Nepetoideae) in East Asia. *Annals of Botany* 122: 649-668. <https://doi.org/10.1093/aob/mcy104>
- IUCN (2012) IUCN Red List Categories and Criteria. Version 3.1. Gland, 32 pp. URL: [http://jr.iucnredlist.org/documents/redlist\\_cats\\_crit\\_en.pdf](http://jr.iucnredlist.org/documents/redlist_cats_crit_en.pdf)
- Kamelin RV (2017) Flora of variegated denudations of Central Asia (brief analysis and genesis). *Turczaninowia* 20 (4): 125-151. [In Russian]. <https://doi.org/10.14258/turczaninowia.20.4.14>
- Körner C (2004) Mountain biodiversity, its causes and function. *AMBIO* 33 (sp13): 11-17. <https://doi.org/10.1007/0044-7447-33.sp13.11>
- Kriebel R, Drew BT, Drummond CP, González Gallegos JG, Celep F, Mahdjoub MM, Rose JP, Xiang CL, Hu GX, Walker JB, Lemmon EM, Lemmon AR, Sytsma KJ (2019) Tracking temporal shifts in area, biomes, and pollinators in the radiation of *Salvia* (sages) across continents: leveraging anchored hybrid enrichment and targeted sequence data. *American Journal of Botany* 106 (4): 573-597. <https://doi.org/10.1002/ajb2.1268>
- Kudryashev SN (1937) On the sages of Central Asia. Committee for science of the Uzbek SSR, Tashkent, 35 pp. [In Russian].
- Lazkov GA (2016) The family Labiatae Juss. in the flora of Kyrgyzstan. Korea National Arboretum, Pocheon, 384 pp. [ISBN 9791187031444]
- Lipsky VI (1905) Flora of Central Asia, i.e. Russian Turkestan and the khanates of Bukhara and Khiva 3. Botanical collections from Central Asia. *Trudy Tiflisskogo Botanicheskogo Sada* 7: 339-841. [In Russian].
- Makhmedov AM (1980) A new species of *Salvia* from the Western Tian-Shan. *Novosti Sistematiki Vysshikh Rastenii* 17: 215-218. [In Russian].
- Makhmedov AM (1982) New taxa of *Salvia* L. *Botanicheskie Materialy Gerbariya Instituta Botaniki Akademii Nauk Uzbekskoi SSR* 20: 30-31. [In Russian].
- Makhmedov AM (1984a) Sages of Central Asia and Kazakhstan (taxonomy, geography and rational use). Science Publishers, Tashkent, 112 pp. [In Russian].
- Makhmedov AM (1984b) *Salvia korolkowii* Regel et Schmalh., *Salvia tianschanica* Makhmedov. In: Nabiev MM (Ed.) *Red Data Book of Uzbek SSR*. Science Publishers, Tashkent, 48-49 pp. [In Russian].
- Makhmedov AM, Denisova LV (1984) *Arischrada korolkowii* (Regel et Schmalh.) Pobed. In: Borodin AM, Bannikov AG, Sokolov VE (Eds) *Red Data Book of the USSR*. 2 ed., 2. Forestry Publishers, Moscow, 223-224 pp. [In Russian].
- Makhmedov AM (1988) *Salvia korolkowii* Regel et Schmalh. In: Ivanina LI (Ed.) *Schedae ad Herbarium florum URSS ab Instituto Botanico Academiae Scientiarum*

- URSS editum. 26. Science Publishers, Leningrad, 107-107 pp. [In Russian]. [ISBN 5-02-026616-7].
- Mamadalieva NZ, Akramov DK, Ovidi E, Tiezzi A, Nahar L, Azimova SS, Sarker SD (2017) Aromatic medicinal plants of the Lamiaceae family from Uzbekistan: ethnopharmacology, essential oils composition, and biological activities. *Medicines* 4 (1). <https://doi.org/10.3390/medicines4010008>
  - Medikus F (1791) *Philosophische Botanik*. 2. Hof- und Akademischen Buchhandlung, Mannheim, 112 pp.
  - Mosolova SL (2006) *Salvia korolkowii* Regel et Schmalh., *Salvia vvedenskii* E. Nikit. In: Shukurov EJ (Ed.) *Red Data Book of Kyrgyz Republic*. 2 ed. State agency on environment protection and forestry under the Government of Kyrgyz Republic, Bishkek, 194-197 pp. [ISBN 9967-23-367-2].
  - Nevski SA (1937) Contributions to the flora of the Kuhitang-Tau and its foothills. *Acta Instituti Botanici Academiae Scientiarum USSR* (ser. 1) 4: 199-346. [In Russian].
  - Nikitina EV (1962) *Salvia Vvedenskii* E. Nik. sp. n. In: Vvedensky AI (Ed.) *Flora of the Kirghizian SSR. Manual of vascular plants of the Kirghizian SSR*. 10. Academy of Sciences of the Kirghizian SSR, Frunze, 377-378 pp. [In Russian].
  - Plantarium (2022) *Plantarium: open on-line atlas and key to plants and lichens of Russia and neighbouring countries*. <http://www.plantarium.ru/>. Accessed on: 2022-6-01.
  - Pobedimova EG (1954) *Schraderia* Medik. In: Schischkin BK (Ed.) *Flora of the USSR*. 21. Academy of Sciences of the USSR, Moscow & Leningrad, 363-374 pp. [In Russian].
  - Pobedimova EG (1967) *Schraderia korolkovii* (Regel et Schmalh.) Pobed. In: Czerepanov SK (Ed.) *Schedae ad Herbarium florum URSS ab Instituto Botanico Academiae Scientiarum URSS editum*. 17. Science Publishers, Leningrad, 82-82 pp. [In Russian].
  - Pobedimova EG (1972) *Taxa nova Florae URSS*, 2. *Novosti Sistematiki Vysshikh Rastenii* 9: 247-250. [In Russian].
  - Popov MG (1923) *Flora of variegated sediments (red sandstone foothills) of Buxoro*. *Trudy Turkestanского nauchnogo obshchestva* 1: 27-64. [In Russian].
  - Regel E (1879) *Descriptiones plantarum novarum et minus cognitatarum*. 7. Brüder Schumacher, St. Petersburg, 263 pp.
  - Regel E (1880) *Descriptiones plantarum novarum et minus cognitatarum, fasciculus VII*. *Acta Horti Petropolitani* 6 (2): 287-538.
  - Sennikov AN, Tojibaev KS, Khassanov FO, Beshko NY (2016) The flora of Uzbekistan project. *Phytotaxa* 282 (2): 107-118. <https://doi.org/10.11646/phytotaxa.282.2.2>
  - Sennikov AN (2021) Botanical expeditions of Boris K. Schischkin and Vasily V. Sapozhnikov in Turkey. *Annales Botanici Fennici* 58 (1-3): 171-180. <https://doi.org/10.5735/085.058.0119>
  - Sharashova VS (1960) *Salvia* L. In: Vvedensky AI (Ed.) *Flora of the Kirghiz SSR*. 9. Academy of Sciences of the Kirghiz SSR, Frunze, 131-143 pp. [In Russian].
  - Sharashova VS (1985) *Salvia korolkowii* Regel et Schmalh., *Salvia vvedenskii* E. Nikit. In: Vorobiev GG (Ed.) *Red Data Book of the Kyrgyz SSR*. Forestry Publishers, Frunze, 101-103 pp. [In Russian].
  - Soják J (1983) *Fragmenta phytotaxonomica et nomenclatorica*, 3. *Časopis Národního muzea v Praze* 152 (1): 22.
  - Spehn EM, Liberman M, Körner C (Eds) (2005) *Land use change and mountain biodiversity*. CRC Press, Boca Raton, 376 pp. [ISBN 0-8439-3523-X]

- Takhtajan AL (Ed.) (1975) Red Data Book. Native plant species to be protected in the USSR. Science Publishers, Leningrad, 204 pp. [In Russian].
- Takhtajan AL (Ed.) (1981) Rare and vanishing plants of the USSR to be protected. 2 ed. Science Publishers, Leningrad, 264 pp. [In Russian].
- The Royal Botanic Gardens, Kew (2021) The World checklist of vascular plants (WCVP). The Catalogue of Life Partnership <https://doi.org/10.15468/k8u286>
- Tojibaev KS (2009) *Salvia korolkowii* Regel et Schmalh., *Salvia tianschanica* Makhm. In: Prатов O, Khassanov FO (Eds) Red Data Book of the Republic of Uzbekistan. 3 ed., 1. Chinor ENK, Tashkent, 270-273 pp.
- Tojibaev KS, Jang CG, Lazkov GA, Chang KS, Sitpayeva GT, Safarov N, Beshko N, Muktabaeyeva S, Vesselova PV, Turakulov I, Abdurakhmanova YY, Na N, Park M, Choi K, Choi H, Oh BU, Oh SH (2020) An annotated checklist of endemic vascular plants of the Tian-Shan Mountains in Central Asian countries. Phytotaxa 464 (2): 117-158. <https://doi.org/10.11646/phytotaxa.464.2.1>
- Tojibaev KS, Sennikov AN, Lazkov GA, Jang CG, Choi HJ, Chang KS, Jung SY, Kim A, Choi K (2021) Checklist of vascular plants of the Tian-Shan Mountain System. Korea National Arboretum, Pocheon, 607 pp. [ISBN 979-11-90509-72-5 (93480)]
- Turdiboev OA, Turginov OT (2021) The taxonomy of *Salvia* L. in Uzbekistan. Uzbekistan Biological Journal 2021 (1): 34-38. [In Russian].
- Turdiboev OA, Sennikov AN (2022) Distribution of *Salvia korolkowii*. Version 1.4. Komarov Botanical Institute, Russian Academy of Sciences, St. Petersburg. <https://doi.org/10.15468/7j3uer>
- Turdiboev OA, Shormanova AA, Sheludyakova MB, Akbarov F, Drew BT, Celep F (2022) Synopsis of the Central Asian *Salvia* species with identification key. Phytotaxa 543 (1): 1-20. <https://doi.org/10.11646/phytotaxa.543.1.1>
- Turland N, Wiersema JH, Barrie FR, Greuter W, Hawksworth DL, Herendeen PS, Knapp S, Kusber WH, Li DZ, Marhold K, May TW, McNeill J, Monro AM, Prado J, Price MJ, Smith GF (Eds) (2018) International code of nomenclature for algae, fungi, and plants (Shenzhen Code) adopted by the Nineteenth International Botanical Congress Shenzhen, China, July 2017. Koeltz Botanical Books, Glashütten. [ISBN 978-3-946583-16-5] <https://doi.org/10.12705/Code.2018>
- Usmonov M, Tojibaev K, Jang CG, Sennikov AN (2021) Species conservation profile and amended distribution of *Cousinia knorringiae* (Asteraceae), a narrow endemic of the Western Tian-Shan. Biodiversity Data Journal 9: e64115. <https://doi.org/10.3897/BDJ.9.e64115>
- Walker JB, Sytsma KJ, Treutlein J, Wink M (2004) *Salvia* (Lamiaceae) is not monophyletic: implications for the systematics, radiation, and ecological specializations of *Salvia* and tribe Mentheae. American Journal of Botany 91 (7): 1115-1125. <https://doi.org/10.3732/ajb.91.7.1115>
- Will M, Claßen-Bockhoff R (2017) Time to split *Salvia* s.l. (Lamiaceae) - New insights from Old World *Salvia* phylogeny. Molecular Phylogenetics and Evolution 109: 33-58. <https://doi.org/10.1016/j.ympev.2016.12.041>
- Wilson B, Dolotbakov A, Burgess BJ, Clubbe C, Lazkov GA, Shalpykov K, Ganybaeva M, Sultangaziev O, Brockington SF (2021) Central Asian wild tulip conservation requires a regional approach, especially in the face of climate change. Biodiversity and Conservation 30: 1705-1730. <https://doi.org/10.1007/s10531-021-02165-z>

- Zhogoleva EP, Kochkareva TF (1986) *Salvia* L. In: Kochkareva TF (Ed.) Flora of Tajikistan. 8. Science Publishers, Leningrad, 241–257 pp. [In Russian].
- Zuckerwanik IP, Grachev TF (1948) An analysis of *Salvia korolkowii* Rgl. et Schmalh. Trudy Instituta Khimii (AN UzbSSR) 1: 41-43. [In Russian].

## Supplementary material

### Suppl. material 1: Distributional data for *Salvia korolkowii* [doi](#)

**Authors:** Turdiboev, O.A. & Sennikov, A.N.

**Data type:** occurrences

**Brief description:** Distribution data based on the comprehensive revision of herbarium specimens and published human observations, as used in GeoCAT. Detailed information: <https://doi.org/10.15468/7j3uer>.

[Download file](#) (189.34 kb)