



Unveiling hidden diversity: new records of *Chaitophorus* (Hemiptera, Aphididae) in Korea from historical specimens

Hwalran Choi[‡], Yeyeun Kim[‡], Seunghwan Lee[‡]

[‡] Insect Biosystematics Laboratory, Research Institute of Agricultural and Life Sciences, Department of Agricultural Biotechnology, Seoul National University, Seoul, Republic of Korea

Corresponding author: Seunghwan Lee (seung@snu.ac.kr)

Academic editor: Colin Favret

Received: 19 Feb 2025 | Accepted: 24 Apr 2025 | Published: 07 May 2025

Citation: Choi H, Kim Y, Lee S (2025) Unveiling hidden diversity: new records of *Chaitophorus* (Hemiptera, Aphididae) in Korea from historical specimens. Biodiversity Data Journal 13: e150852.

<https://doi.org/10.3897/BDJ.13.e150852>

Abstract

Background

The aphid genus *Chaitophorus* Koch, 1854 (Hemiptera, Aphididae, Chaitophorinae) has been studied in Korea from historical specimens.

New information

Korean *Chaitophorus* aphids have been confirmed, including four new records: *C. horii* Takahashi, 1939; *C. leucomelas* Koch, 1854; *C. salijaponicus* Essig and Kuwana, 1918; and *C. tremulae* Koch, 1854. These four newly-recorded species are described with biometric measurements and illustrations. A modified taxonomic key for Korean *Chaitophorus* spp. is presented.

Keywords

aphids, Salicaceae, East Asia

Introduction

The aphid genus *Chaitophorus* Koch, 1854 is a species-rich group within the subfamily Chaitophorinae. These aphids are distributed across the Holarctic Region, with approximately 90 species recorded (Blackman and Eastop 2024, Favret and Aphid Taxon Community, eds. 2025). *Chaitophorus* species are well-known for their tree-dwelling habit, with no host alternation and they exhibit host specificity at the genus level, feeding exclusively on either *Salix* or *Populus* (Salicaceae), but never both (Blackman and Eastop 2024). Distinct morphological traits of the genus include a body covered with long setae, short siphunculi and knobbed cauda, all of which are particularly useful for distinguishing these aphids from related groups.

Taxonomic studies on *Chaitophorus* have been conducted in various countries. Pintera (1987) revised the Palaearctic fauna, covering 58 species. Additionally, research has been conducted in China and Japan by Tao (1964) and Higuchi (1972), respectively. Nieto Nafria and Mier Durante (1998) and Wieczorek and Osiadacz (2007) reviewed the European species, while Chakrabarti (1977) and Ghosh (1980) studied the species found in India. In North America, Hille Ris Lambers (1960) and Richards (1972) conducted reviews.

Systematic and co-evolutionary studies on the genus *Chaitophorus* have also been undertaken. Shingleton and Stern (2003) were the first to conduct a molecular phylogenetic study on this genus, using mitochondrial DNA (mtDNA) from 15 species. Wieczorek and Kajtoch (2011) and Wieczorek et al. (2017) examined the tribal relationships within Chaitophorinae and their closely-related groups using both mtDNA and nuclear genes. Yang et al. (2016) investigated the co-evolution of *Chaitophorus* with the symbiont *Buchnera* and Liu et al. (2022) explored macroevolution in relation to host plants.

Taxonomic research on Korean *Chaitophorus* species has been conducted by several researchers, including Shinji (1941), Paik (1972), Szelegiewicz (1981), Pintera (1987) and Lee and Seo (1992). In this study, we newly record four species of *Chaitophorus* collected from Korea.

Materials and methods

Historical specimens of this study are deposited in the Consortium of Korea Biodiversity and Sustainable Use, College for Agriculture and Life Sciences, Seoul National University (CALS SNU, Korea).

Illustrations for each species were taken by digital camera (14.2 Color Mosaic, Diagnostic Instruments, Sterling Heights, MI, USA) attached to the microscope (DM 400B, Leica Microsystems, Wetzlar, Germany) at a resolution of 600 dpi. Measurements for each specimen are taken from the digital images by Image Laboratory v.2.2.4.0 software (MCM Design Ltd, Hillerod, Denmark).

Abbreviations used for descriptions are as follows: ap., apterous viviparous female; Ant., antennae; Ant. I, Ant. II, Ant. III, Ant. IV, Ant. V, Ant. VI and Ant. VIb, antennal segments I, II, III, IV, V, VI and base of VI, respectively; BD Ant. III, basal diameter of antennal segment III; AbdT. III, Abdominal Tergite III; GP, genital plate; 2HT, second segment of hind tarsus; PT, processus terminalis of antennal segment VI; SIPH, siphunculus; URS, ultimate rostral segments (segment IV+V). For the localities of distribution, provincial abbreviations are also used: South Korea: CB, Chungcheongbuk-do; GG, Gyeonggi-do; JB, Jeollabuk-do; North Korea: HB, Hwanghaebuk-do; PB, Pyeonganbuk-do; PN: Pyeongannam-do; RG: Ryanggang-do.

Taxon treatments

Chaitophorus horii Takahashi, 1939

Materials

- a. scientificName: *Chaitophorus horii* Takahashi, 1939; namePublishedIn: Takahashi, R. (1939) Some Aphididae from Hokkaido (Hemiptera). *Insecta Matsumurana*, 13(4), 114–128.; namePublishedInID: <http://hdl.handle.net/2115/9419>; taxonomicStatus: accepted; taxonID: 2074388; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Hemiptera; family: Aphididae; nomenclaturalCode: ICZN; genus: *Chaitophorus*; specificEpithet: *horii*; continent: Asia; country: North Korea; countryCode: KP; stateProvince: Ryanggang Province; municipality: Samjiyeon; locality: Mt. Baekdu; decimalLatitude: 41.899; decimalLongitude: 128.394; geodeticDatum: WGS84; coordinateUncertaintyInMeters: 100; samplingProtocol: hand collected; eventDate: 1988-06-24; year: 1988; month: 6; day: 24; individualCount: 3; sex: female; lifeStage: adult; recordedBy: J. Havelka; occurrenceStatus: present; disposition: in collection; associatedOccurrences: host: *Salix herbacea*; identifiedBy: Yeyun Kim; dateIdentified: 2025-02-18; type: specimen; modified: 2025-03-28; language: en; rights: Seoul National University; rightsHolder: Content licensed under Seoul National University; accessRights: not for profit use only; institutionID: NPRI; collectionID: SNU; basisOfRecord: PreservedSpecimen; occurrenceID: 19819F5C-4912-5340-A02F-DFD62934B644
- b. scientificName: *Chaitophorus horii* Takahashi, 1939; namePublishedIn: Takahashi, R. (1939) Some Aphididae from Hokkaido (Hemiptera). *Insecta Matsumurana*, 13(4), 114–128.; namePublishedInID: <http://hdl.handle.net/2115/9419>; taxonomicStatus: accepted; taxonID: 2074388; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Hemiptera; family: Aphididae; nomenclaturalCode: ICZN; genus: *Chaitophorus*; specificEpithet: *horii*; continent: Asia; country: North Korea; countryCode: KP; stateProvince: Ryanggang Province; municipality: Samjiyeon; locality: Mt. Baekdu; decimalLatitude: 41.899; decimalLongitude: 128.394; geodeticDatum: WGS84; coordinateUncertaintyInMeters: 100; samplingProtocol: hand collected; eventDate: 1988-06-17; year: 1988; month: 6; day: 17; individualCount: 2; sex: female; lifeStage:

adult; recordedBy: J. Havelka; occurrenceStatus: present; disposition: in collection; associatedOccurrences: host: *Salix herbacea*; identifiedBy: Yeyeun Kim; dateIdentified: 2025-02-18; type: specimen; modified: 2025-03-28; language: en; rights: Seoul National University; rightsHolder: Content licensed under Seoul National University; accessRights: not for profit use only; institutionID: NPRI; collectionID: SNU; basisOfRecord: PreservedSpecimen; occurrenceID: F12B8F44-2F65-551B-8C05-8C7EE953F41E

Description

Apterous viviparous female (Fig. 1, Table 1). **Colour (macerated specimens).** Body and appendages wholly pale. **Morphology.** Body elongated and oval-shaped; its length twice as long as width, about 1.36 mm. **Head:** length of antenna slightly over 0.5 times of the body length; PT almost 2 times longer than Ant. Vlb.; longest seta on Ant. III same as long as basal width of segment. URS beak-shaped, with 6-7 setae; nearly 1.1 times as long as 2HT. **Thorax:** Legs comparatively shorter than other *Chaitophorus* spp., hind femur and tibia about 0.29 and 0.45 mm long, respectively. **Abdomen:** abdominal segments fused from I to VII; most setae on tergite relatively long, stunted or furcate; nearly 10 setae each on tergite VI and VIII. SIPH slightly long and stump-shaped. Cauda knobbed with 5-6 setae.

Host plants: *Salix caprea*, *S. herbacea*, *S. rorida*, *S. viminalis*, *Salix* spp. (Blackman and Eastop 2024).

Part		Apterous vivipara (n = 5)	
		Average	(Min-Max)
Length (mm)	Body (from the most forward point of the frons to end of Cauda)	1.36	(1.35-1.38)
	Whole Ant.	0.71	(0.69-0.72)
	Ant. I	0.05	(0.05-0.05)
	Ant. II	0.04	(0.04-0.04)
	Ant. III	0.14	(0.13-0.14)
	Ant. IV	0.10	(0.09-0.10)
	Ant. V	0.10	(0.10-0.11)
	Ant. Vlb	0.08	(0.08-0.08)
	PT	0.17	(0.16-0.17)
	URS	0.09	(0.09-0.10)

Part		Apterous vivipara (n = 5)	
		Average	(Min-Max)
	Hind femur	0.29	(0.29-0.30)
	Hind tibia	0.45	(0.45-0.47)
	2HT	0.09	(0.08-0.09)
	SIPH	0.02	(0.02-0.03)
	Knob of Cauda	0.05	(0.05-0.05)
	Width of the knob of Cauda	0.05	(0.05-0.05)
	Longest seta on Ant. III	0.02	(0.01-0.02)
	Longest seta on AbdT. III	0.14	(0.14-0.15)
No. of setae on	Mandibular lamina	6	(6-7)
	Ant. I	3	(3-4)
	Ant. II	3	(3-3)
	Ant. III	2	(2-3)
	URS (subsidiary)	6	(6-7)
	Tergite VI between SIPH	10	(9-11)
	Tergite VIII	11	(10-11)
	Median of GP	4	(3-6)
	Posterior margin of GP	13	(12-15)
	Knob of Cauda	7	(6-8)
No. of secondary rhinaria on	Ant. III	0	
	Ant. IV	0	
	Ant. V	0	
Ratio (times)	Whole antennae / Body	0.52	(0.51-0.53)
	PT / Ant. VIb	1.97	(1.94-1.97)
	PT / Ant. III	1.21	(1.21-1.21)
	URS / 2HT	1.10	(1.10-1.11)
	URS / Ant. VIb	1.14	(1.14-1.15)
	SIPH / Body	0.02	(0.02-0.02)
	SIPH / Ant. III	0.21	(0.20-0.21)
	SIPH / Hind femur	0.10	(0.10-0.10)

Part		Apterous vivipara (n = 5)	
		Average	(Min-Max)
	SIPH / Cauda	0.49	(0.47-0.51)
	Knob of Cauda / Width of the knob of Cauda	1.00	(1.00-1.02)
	Seta on Ant. III / Ant. III BD	1.00	(0.90-1.00)
	Seta on AbdT. III / Ant. III BD	7.35	(7.14-7.37)




Figure 1. [doi](#)

Apterous viviparous female of *Chaitophorus horii* Takahashi, 1939. **A** whole body; **B** hind tibia and tarsus; **C** antenna; **D** head; **E** cauda; **F** siphunculus; **G** ultimate rostral segments. Scale bar: A, 0.5 mm; B,C, 0.1 mm; D, 0.1 mm; E,F,G, 0.05 mm.

Diagnosis

Chaitophorus horii morphologically resembles *C. purpureae*. However, it is distinguished by its beak-shaped URS and long dorsal setae.

Distribution

Korea (new record), South Europe, Japan, Thailand (Blackman and Eastop 2024).

Chaitophorus leucomelas Koch, 1854

Materials

- a. scientificName: *Chaitophorus leucomelas* Koch, 1854; namePublishedIn: Koch, C.L. (1854) In *Die Pflanzenläuse Aphiden getreu nach dem Leben abgebildet und beschrieben*. J.L. Lotzbeck, Nürnberg. Vol. 1, 1–36 pp.; namePublishedInID: <https://doi.org/10.5281/zenodo.14387414>; taxonomicStatus: accepted; taxonID: 2074434;

higherClassification: Animalia; kingdom: Animalia; phylum: Arthropod; class: Insects; order: Hemiptera; family: Aphididae; taxonRank: species; nomenclaturalCode: ICZN; genus: *Chaitophorus*; specificEpithet: *leucomelas*; scientificNameAuthorship: Koch, 1854; continent: Asia; country: North Korea; countryCode: KP; stateProvince: South Hwanghae; municipality: Kaesong; locality: Gongmin Wang Tomb; decimalLatitude: 37.982; decimalLongitude: 126.473; geodeticDatum: WGS84; coordinateUncertaintyInMeters: 100; samplingProtocol: hand collected; eventDate: 1987-06-01; year: 1987; month: 06; day: 01; individualCount: 24; sex: females; lifeStage: adult; catalogNumber: 87HA1644; recordedBy: J. Havelka; occurrenceStatus: present; disposition: in collection; associatedOccurrences: host: *Populus alba*; identifiedBy: Yeyeun Kim; dateIdentified: 2025-03-28; type: specimen; modified: 2025-03-28; language: en; rights: Seoul National University; rightsHolder: Seoul National University; accessRights: not for profit use only; institutionID: NPRI; collectionID: SNU; occurrenceID: 71388D05-EB1D-5BF5-A4D7-514768BE84CB

- b. scientificName: *Chaitophorus leucomelas* Koch, 1854; namePublishedIn: Koch, C.L. (1854) In *Die Pflanzenläuse Aphiden getreu nach dem Leben abgebildet und beschrieben*. J.L. Lotzbeck, Nürnberg. Vol. 1, 1–36 pp.; namePublishedInID: <https://doi.org/10.5281/zenodo.14387414>; taxonomicStatus: accepted; taxonID: 2074434; higherClassification: Animalia; kingdom: Animalia; phylum: Arthropod; class: Insects; order: Hemiptera; family: Aphididae; taxonRank: species; nomenclaturalCode: ICZN; genus: *Chaitophorus*; specificEpithet: *leucomelas*; scientificNameAuthorship: Koch, 1854; continent: Asia; country: South Korea; countryCode: KR; stateProvince: North Jeolla Province; municipality: Muju; decimalLatitude: 36.019; decimalLongitude: 127.658; geodeticDatum: WGS84; coordinateUncertaintyInMeters: 100; samplingProtocol: direct collection; eventDate: 1969-10-29; year: 1969; month: 10; day: 29; individualCount: 4; sex: females; lifeStage: adult; catalogNumber: #5783; recordedBy: J. Havelka; occurrenceStatus: present; disposition: in collection; associatedOccurrences: host: *Populus* sp.; identifiedBy: Yeyeun Kim; dateIdentified: 1989-07-10; type: specimen; modified: 1969-10-29; language: en; rights: Seoul National University; rightsHolder: Seoul National University; accessRights: not for profit use only; institutionID: NPRI; collectionID: #5783; basisOfRecord: PreservedSpecimen; occurrenceID: D29D7CDD-DF15-5196-87CE-EEE7A14FDD6F

Description

Apterous viviparous female (Fig. 2, Table 2). **Colour** (macerated specimens). Body pale, leg, cauda and antenna faintly grey. Head, mandibular lamina, whole Ant. except Ant. III through apical half of Ant. V and SIPH more deeply pigmented. URS dark in peak. **Morphology**. Body ovoid, about 1.98 mm long. **Head**: flat with stout seta on frons; mandibular lamina with eight or more setae. Antennal length almost 0.6 times as long as body length; PT almost 3 times longer than Ant. VIb. URS short with eight or more setae. **Thorax**: pronotum smooth. Hind femur and tibia length about 0.51 mm and 0.69 mm, respectively; hind tibia entirely with long and fine setae. **Abdomen**: abdominal segments fused from II to VII; dorsal cuticles with nodules and faintly pigmented, with fine, long and pointed setae. SIPH 0.02 times the body length. Cauda knobbed with eight setae.

Host plants: *Populus alba*, *P. deltoides*, *P. nigra*, *P. simonii*, *Populus* spp. (Blackman and Eastop 2024).

Table 2. Biometric data of <i>Chaitophorus leucomelas</i> .			
Part		Apterous vivipara (n = 17)	
		Average (Min-Max)	
Length (mm)	Body (from the most forward point of the frons to end of Cauda)	1.98	(1.96-2.39)
	Whole Ant.	1.17	(1.14-1.41)
	Ant. I	0.07	(0.07-0.08)
	Ant. II	0.05	(0.05-0.06)
	Ant. III	0.31	(0.30-0.37)
	Ant. IV	0.17	(0.17-0.20)
	Ant. V	0.14	(0.13-0.16)
	Ant. VIb	0.10	(0.09-0.13)
	PT	0.30	(0.30-0.39)
	URS	0.11	(0.11-0.12)
	Hind femur	0.51	(0.51-0.52)
	Hind tibia	0.69	(0.68-0.70)
	2HT	0.12	(0.11-0.13)
	SIPH	0.04	(0.04-0.05)
	Knob of Cauda	0.06	(0.05-0.06)
	Width of the knob of Cauda	0.06	(0.05-0.06)
	Longest seta on Ant. III	0.06	(0.06-0.07)
Longest seta on AbdT. III	0.13	(0.12-0.14)	
No. of setae on	Mandibular lamina	8	(7-10)
	Ant. I	5	(4-5)
	Ant. II	4	(4-5)
	Ant. III	15	(12-17)
	URS (subsidiary)	8	(7-8)
	Tergite VI between SIPH	10	(9-12)
	Tergite VIII	7	(7-8)
	Median of GP	10	(9-14)
	Posterior margin of GP	19	(17-21)
	Knob of Cauda	8	(7-8)

Part		Apterous vivipara (n = 17)	
		Average (Min-Max)	
No. of secondary rhinaria on	Ant. III	0	
	Ant. IV	0	
	Ant. V	0	
Ratio (times)	Whole Ant. / Body	0.59	(0.59-0.61)
	PT / Ant. VIb	3.01	(3.00-3.10)
	PT / Ant. III	0.98	(0.98-0.98)
	URS / 2HT	0.97	(0.93-1.01)
	URS / Ant. VIb	1.17	(1.15-1.18)
	SIPH / Body	0.02	(0.02-0.03)
	SIPH / Ant. III	0.16	(0.15-0.17)
	SIPH / Hind femur	0.10	(0.09-0.10)
	SIPH / Knob of Cauda	0.80	(0.78-0.87)
	Knob of Cauda / Width of the knob of Cauda	1.02	(1.00-1.04)
	Longest seta on Ant. III / Ant. IIIBD	2.62	(2.60-2.73)
Longest seta on AbdT. III / Ant. IIIBD	5.15	(5.14-5.27)	




Figure 2. [doi](#)

Apterous viviparous female of *Chaitophorus leucomelas* Koch, 1854. **A** whole body; **B** hind tibia and tarsus; **C** antenna; **D** head; **E** cauda; **F** siphunculus; **G** ultimate rostral segments. Scale bar: A, 0.5 mm; B,C, 0.1 mm; D, 0.1 mm; E,F,G, 0.05 mm.

Diagnosis

This species typically has pigmentation on the dorsum, frons and SIPH, but the pigmentation on the dorsum varies depending on the specimen, whereas the pigmentation on the frons and SIPH is always present. Korean *Chaitophorus leucomelas* has faintly pigmented dorsum, but the frons and SIPH distinctly have pigmentation. Additionally, the pigmentation of antennae and their ratio and dorsal cuticles with nodules follow the referred description (Pintera 1987).

Distribution

Korea (new record), China, Kazakhstan, Mongolia, Siberia, Europe, Canada, USA, Chile, Africa (Blackman and Eastop 2024).

Chaitophorus salijaponicus Essig and Kuwana, 1918

Materials

- a. scientificName: *Chaitophorus salijaponicus* Essig and Kuwana, 1918; namePublishedIn: Essig, E.O. & Kuwana, S.I. (1918) Some Japanese Aphididae. *Proceedings of the California Academy of Sciences*, 8(3), 35–112.; namePublishedInID: 10.1163/187631204788920158; taxonID: 2074456; higherClassification: Animalia; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Hemiptera; family: Aphididae; taxonRank: species; nomenclaturalCode: ICZN; genus: *Chaitophorus*; specificEpithet: *salijaponicus*; continent: Asia; country: North Korea; countryCode: KP; stateProvince: Ryanggang Province; locality: Hyesan; decimalLatitude: 41.387; decimalLongitude: 128.173; geodeticDatum: WGS84; coordinateUncertaintyInMeters: 100; samplingProtocol: hand collected; eventDate: 1988-06-28; year: 1988; month: 06; day: 28; individualCount: 8; sex: females; lifeStage: adult; catalogNumber: #88HA3688; recordedBy: J. Havelka; occurrenceStatus: present; disposition: in collection; associatedOccurrences: host: *Salix purpurea*; identifiedBy: Yeyeun Kim; dateIdentified: 2025-03-28; type: specimen; modified: 2025-03-28; language: en; rights: Seoul National University; rightsHolder: Seoul National University; accessRights: not for profit use only; institutionID: NPRI; collectionID: SNU; occurrenceID: C7336A4E-0FF3-5014-B231-3CA2EEC96C4B
- b. scientificName: *Chaitophorus salijaponicus* Essig and Kuwana, 1918; namePublishedIn: Essig, E.O. & Kuwana, S.I. (1918) Some Japanese Aphididae. *Proceedings of the California Academy of Sciences*, 8(3), 35–112.; namePublishedInID: 10.1163/187631204788920158; taxonID: 2074456; higherClassification: Animalia; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Hemiptera; family: Aphididae; taxonRank: species; nomenclaturalCode: ICZN; genus: *Chaitophorus*; specificEpithet: *salijaponicus*; continent: Asia; country: South Korea; countryCode: KR; stateProvince: North Jeolla Province; municipality: Muju; decimalLatitude: 36.014; decimalLongitude: 127.656; geodeticDatum: WGS84; coordinateUncertaintyInMeters: 100; samplingProtocol: hand collected; eventDate: 1969-10-29; year: 1969; month: 10; day: 29; individualCount: 2; sex: females; lifeStage: adult; catalogNumber: #5718; recordedBy: J. Havelka; occurrenceStatus: present; disposition: in collection; associatedOccurrences: host: *Salix* sp.; identificationID: Yeyeun Kim; identifiedBy: Yeyeun Kim; dateIdentified: 2025-02-19; type: specimen; modified: 2025-03-28; language: en;

rights: Seoul National University; rightsHolder: Seoul National University; accessRights: not for profit use only; institutionID: SNU; collectionID: #5718; occurrenceID: 4A417E0F-2855-5528-B85B-9E47C1522690

Description

Apterous viviparous female (Fig. 3, Table 3). **Colour** (macerated specimens). Body blackish and oval-shaped, intersegmentally pale on unfused tergites. Antenna and leg slightly grey; hind tibia, Ant. VIb and PT darker than body. Abdominal tergite with a pale ring around base of SIPH. **Morphology**. Body elongated oval-shaped; about 1.52 mm long; marginal setae around body distinctly long and fine. **Head**: flat with long pointed setae on frons; mandibular lamina with eight setae. Antenna slender; whole antennae almost 0.55 times as long as body length; PT 2.55 times as long as Ant. VIb.; Ant. III with seven long setae; Ant. V with relatively stunt setae. URS with eight setae. **Thorax**: hind femur and tibia about 0.31 mm and 0.49 mm long, respectively; length of leg a little shorter than other species in comparison with length of body. **Abdomen**: abdominal tergites from II to VII fused; more sclerotised on marginal tergites of each segment. SIPH short and reticulate. Cauda knobbed with 6 setae.

Table 3.

Biometric data of *Chaitophorus salijaponicus*.

Part		Apterous vivipara (n = 20)	
		Average (Min-Max)	
Length (mm)	Body (from the most forward point of the frons to end of Cauda)	1.52	(1.37-1.69)
	Whole Ant.	0.83	(0.75-0.91)
	Ant. I	0.05	(0.04-0.06)
	Ant. II	0.05	(0.04-0.05)
	Ant. III	0.17	(0.14-0.19)
	Ant. IV	0.12	(0.10-0.13)
	Ant. V	0.10	(0.09-0.11)
	Ant. VIb	0.09	(0.08-0.10)
	PT	0.23	(0.21-0.24)
	URS	0.08	(0.08-0.09)
	Hind femur	0.31	(0.28-0.33)
	Hind tibia	0.49	(0.45-0.51)
	2HT	0.12	(0.11-0.12)
	SIPH	0.05	(0.05-0.06)

Part		Apterous vivipara (n = 20)	
		Average (Min-Max)	
	Knob of Cauda	0.03	(0.03-0.03)
	Width of the knob of Cauda	0.04	(0.04-0.04)
	Longest seta on Ant. III	0.07	(0.05-0.10)
	Longest seta on AbdT. III	0.15	(0.12-0.18)
No. of setae on	Mandibular lamina	8	(7-8)
	Ant. I	6	(5-6)
	Ant. II	4	(4-5)
	Ant. III	7	(5-8)
	URS (subsidiary)	8	(7-9)
	Tergite VI between SIPH	12	(10-15)
	Tergite VIII	9	(8-10)
	Median of GP	5	(4-7)
	Posterior margin of GP	15	(14-18)
No. of secondary rhinaria on	Knob of Cauda	6	(4-6)
	Ant. III	0	
	Ant. IV	0	
Ratio (times)	Ant. V	0	
	Whole Ant. / Body	0.55	(0.51-0.58)
	PT / Ant. VIb	2.55	(2.49-2.61)
	PT / Ant. III	1.32	(1.30-1.35)
	URS / 2HT	0.68	(0.62-0.71)
	URS / Ant. VIb	0.89	(0.84-0.92)
	SIPH / Body	0.03	(0.03-0.03)
	SIPH / Ant. III	0.29	(0.22-0.35)
	SIPH / Hind femur	0.16	(0.14-0.17)
	SIPH / Knob of Cauda	1.50	(1.47-1.52)
	Knob of Cauda / Width of the knob of Cauda	0.77	(0.75-0.78)
	Longest seta on Ant. III / Ant. III BD	3.04	(2.94-3.10)
Longest seta on AbdT. III / Ant. III BD	6.04	(5.87-6.24)	

Host plants: *S. babylonica*, *S. caprea*, *S. integra*, *S. koriyanagi*, *Salix* spp. (Blackman and Eastop 2024).




Figure 3. [doi](#)

Apterous viviparous female of *Chaitophorus salijaponicus* Essig and Kuwana, 1918. **A** whole body; **B** hind tibia and tarsus; **C** antenna; **D** head; **E** cauda; **F** siphunculus; **G** ultimate rostral segments. Scale bar: A, 0.5 mm; B,C, 0.1 mm; D, 0.1 mm; E,F,G, 0.05 mm.

Diagnosis

C. salijaponicus closely resembles *C. hokkaidensis* morphologically. However, it can be distinguished by the state of abdominal tergite I: in *C. salijaponicus*, it is free, whereas in *C. hokkaidensis*, it is completely fused with the preceding segment.

Distribution

Korea (new record), China, Siberia, Mongolia, Japan (Blackman and Eastop 2024).

Chaitophorus tremulae Koch, 1854

Materials

- a. scientificName: *Chaitophorus tremulae* Koch, 1854; namePublishedIn: Koch, C.L. (1854) In *Die Pflanzenläuse Aphiden getreu nach dem Leben abgebildet und beschrieben*. J.L. Lotzbeck, Nürnberg. Vol. 1, 1–36 pp.; namePublishedInID: 10.5281/zenodo.14387432; taxonID: 2074419; higherClassification: Animalia; kingdom: Animalia; phylum: Arthropod; class: Insects; order: Hemiptera; family: Aphididae; taxonRank: species; nomenclaturalCode: ICZN; genus: *Chaitophorus*; specificEpithet: *tremulae*; continent: Asia; country: North Korea; countryCode: KP; stateProvince: Ryanggang Province; municipality: Hyesan; decimalLatitude: 41.387; decimalLongitude: 128.174; geodeticDatum: WGS84; coordinateUncertaintyInMeters: 100; samplingProtocol: hand collected; eventDate: 1988-06-28; year: 1988; month: 06; day: 28; individualCount: 6; sex: female; lifeStage: adult; catalogNumber: #88HA3711; recordedBy: J. Havelka; occurrenceStatus: present; disposition: in collection; associatedOccurrences: host: *Populus davidiana*; identifiedBy: yeyeun Kim; dateIdentified: 2025-02-19; type: specimen; modified: 2025-03-28; language: en; rights: Seoul National University; rightsHolder: Seoul

National University; accessRights: not for profit use only; institutionID: NPRI; collectionID: SNU; basisOfRecord: PreservedSpecimen; occurrenceID: 045CC865-6020-5684-B585-D80F05169086

- b. scientificName: *Chaitophorus tremulae* Koch, 1854; namePublishedIn: Koch, C.L. (1854) *In Die Pflanzenläuse Aphiden getreu nach dem Leben abgebildet und beschrieben*. J.L. Lotzbeck, Nürnberg. Vol. 1, 1–36 pp.; namePublishedInID: 10.5281/zenodo.14387432; taxonomicStatus: accepted; taxonID: 2074419; higherClassification: Animalia; kingdom: Animalia; phylum: Arthropod; class: Insecta; order: Hemiptera; family: Aphididae; taxonRank: species; nomenclaturalCode: ICZN; genus: *Chaitophorus*; specificEpithet: *tremulae*; continent: Asia; country: South Korea; countryCode: KR; municipality: Seoul; decimalLatitude: 37.475; decimalLongitude: 126.948; geodeticDatum: WGS84; coordinateUncertaintyInMeters: 100; samplingProtocol: hand collected; eventDate: 1970-05-06; year: 1970; month: 05; day: 06; individualCount: 2; sex: female; lifeStage: adult; catalogNumber: #5895; recordedBy: J. Havelka; occurrenceStatus: present; disposition: in collection; associatedOccurrences: host: *Populus davidiana*; identifiedBy: yeyeun Kim; dateIdentified: 2025-02-19; type: specimen; modified: 2025-03-28; language: en; rights: Seoul National University; rightsHolder: Seoul National University; accessRights: not for profit use only; institutionID: NPRI; collectionID: SNU; basisOfRecord: PreservedSpecimen; occurrenceID: B1911DEB-ECE7-5D4C-B3C1-B6F695873087

Description

Apterous viviparous female (Fig. 4, Table 4). **Colour** (macerated specimens). Body extremely blackish, slightly bright in the spinal part of abdomen. Antennae pale, except Ant. I, distal part of Ant. IV, Ant. V., Ant. VIb and PT. Legs pale, except mid- and hind femora, basal part of hind tibia and 2HT. Cauda and anal plate distinctly pale. SIPH dark as body colour. **Morphology**. Body elongated oval-shaped; about 2.20 mm long. **Head**: smooth, seta very long and spine. Antenna slender; whole antennae almost 0.57 times as long as body length; PT 2.73 times as long as Ant. VIb. URS short and slightly blunt, with eight or more setae. **Thorax**: pronotum smooth. Hind femur and tibia about 0.55 mm and 0.69 mm long, respectively; **Abdomen**: abdominal tergites fused from I to VI, with blunt or furcate seta; dorsum with pore-like microstructure in a row of each segment. SIPH short and finely reticulate. Cauda distinctly knobbed with six setae.

Table 4.

Biometric data of *Chaitophorus tremulae*.

Part		Apterous vivipara (n = 7)	
		Average (Min-Max)	
Length (mm)	Body (from the most forward point of the frons to end of Cauda)	2.20	(1.94-2.34)
	Whole Ant.	1.25	(1.09-1.32)
	Ant. I	0.06	(0.05-0.07)

Part		Apterous vivipara (n = 7)	
	Ant. II	0.05	(0.04-0.05)
	Ant. III	0.35	(0.31-0.38)
	Ant. IV	0.20	(0.17-0.21)
	Ant. V	0.16	(0.15-0.17)
	Ant. VIb	0.11	(0.10-0.12)
	PT	0.30	(0.25-0.32)
	URS	0.09	(0.08-0.10)
	Hind femur	0.55	(0.49-0.58)
	Hind tibia	0.69	(0.64-0.74)
	2HT	0.12	(0.11-0.14)
	SIPH	0.06	(0.06-0.07)
	Knob of Cauda	0.06	(0.05-0.06)
	Width of the knob of Cauda	0.06	(0.06-0.06)
	Longest seta on Ant. III	0.08	(0.06-0.09)
	Longest seta on AbdT. III	0.17	(0.15-0.20)
No. of setae on	Mandibular lamina	8	(7-10)
	Ant. I	8	(7-10)
	Ant. II	5	(4-6)
	Ant. III	11	(9-13)
	URS (subsidiary)	8	(7-8)
	Tergite VI between SIPH	16	(15-18)
	Tergite VIII	12	(11-13)
	Median of GP	8	(4-10)
	Posterior margin of GP	16	(14-22)
	Cauda	6	(5-7)
No. of secondary rhinaria on	Ant. III	0	
	Ant. IV	0	
	Ant. V	0	
Ratio (times)	Whole Ant. / Body	0.57	(0.51-0.60)
	PT / Ant. VIb	2.73	(2.59-2.84)
	PT / Ant. III	0.85	(0.76-0.91)
	URS / 2HT	0.72	(0.68-0.75)

Part	Apterous vivipara (n = 7)	
URS / Ant. VIb	0.83	(0.79-0.85)
SIPH / Body	0.03	(0.03-0.03)
SIPH / Ant. III	0.18	(0.17-0.19)
SIPH / Hind femur	0.11	(0.10-0.12)
SIPH / Knob of Cauda	1.19	(1.18-1.21)
Knob of Cauda/ Width of the knob of Cauda	0.82	(0.75-0.85)
Longest seta on Ant. III / Ant. III BD	3.00	(2.86-3.14)
Longest seta on AbdT. III / Ant. III BD	6.07	(5.87-6.24)




Figure 4. [doi](#)

Apterous viviparous female of *Chaitophorus tremulae* Koch, 1854. **A** whole body; **B** hind tibia and tarsus; **C** antenna; **D** head; **E** cauda; **F** siphunculus; **G** ultimate rostral segments. Scale bar: A, 0.5 mm; B,C, 0.1 mm; D, 0.1 mm; E,F,G, 0.05 mm.

Host plants: *P. maximowiczii*, *P. sieboldii*, *P. simonii*, *P. tremula* (Blackman and Eastop 2024).

Diagnosis

C. tremulae closely resembles *C. salijaponicus* morphologically. However, it can be distinguished by body length and marginal tergites: in *C. tremulae*, body length is longer than 1.8 mm and has no sclerotised marginal tergites, whereas in *C. salijaponicus*, body length is shorter than 1.8 mm and sclerotised marginal tergites are developed.

Distribution

Korea (new record), Great Britain, Netherlands, Germany, Denmark, Spain, Norway, Finland, Poland, Hungary, Turkey, Russia, Caucasus, Kazakhstan, Siberia, China, Mongolia, Japan (Blackman and Eastop 2024).

Identification keys

Key to the species of the genus <i>Chaitophorus</i> in the Korean Peninsula (apterous viviparous female) (modified from Albert Pintera (1987) and Blackman and Eastop (2024)).		
1	Pseudosensoria present on hind tibia	2
–	Pseudosensoria absent on hind tibia	3
2	Pseudosensoria scattered over nearly whole hind tibia. Live on <i>Populus</i> spp.	<i>C. populeti</i> (Panzer, 1801)
–	Pseudosensoria (approximately 12) on the enlarged part near the knees of the hind tibia. Live on <i>Salix</i> spp.	<i>C. saliniger</i> Shinji, 1924
3	Body wholly pale	4
–	Body wholly or partly pigmented	11
4	Abdominal tergite I fused with the following tergite. URS with beak-shaped or U-shaped	5
–	Abdominal tergite I not fused with the following tergite. URS with U-shaped	6
5	URS with beak-shaped. URS/2HT 1.1. PT/Ant. Vlb 2.0. Dorsal setae usually furcated. Live on <i>Salix</i> spp.	<i>C. horii</i> Takahashi, 1939
–	URS with U-shaped. URS/2HT 0.8. PT/Ant. Vlb 2.0-2.5. Dorsal setae long, not furcated. Live on <i>Salix purpurea</i> var. <i>smithiana</i>	<i>C. purpureae</i> Lee and Seo, 1992
6	Abdominal tergite VII completely fused with the preceding tergite. SIPH dark. Live on <i>Populus</i> spp.	<i>C. leucomelas</i> Koch, 1854
–	Abdominal tergite VII mostly free with the preceding tergite. SIPH pale	7
7	Longest seta on Ant. III and AbdT. III almost 3-6 times as long as the basal diameter of Ant. III, respectively. Live on <i>Populus</i> spp.	<i>C. populialbae</i> (Boyer de Fonscolombe, 1841)

–	Longest seta on Ant. III and AbdT. III less than 1.2 times and 3.2 times as long as the basal diameter of Ant. III, respectively	8
8	Body elongate, oval without pigmentation. Microstructure consisting of blunt nodules, very sparsely distributed in the middle of dorsum and head, often poorly visible. Live on <i>Salix</i> spp.	<i>C. saliapterus</i> Shinji, 1924
–	Body rather elongate, oval and fuscous. Microstructure not developed or, at most, with small granules on the frons. Live on <i>Salix</i> spp. or <i>Populus</i> spp.	9
9	Seta on Ant. III short, not more than twice as long as basal diameter of Ant. III. Ant. V with 2-3 setae. SIPH pigmented and reticulated on all their length. Live on <i>Salix bakko</i>	<i>C. matsumurai</i> Hille Ris Lambers, 1960
–	Seta on Ant. III more than 4 times as long as basal diameter of Ant. III. PT more than 4 times as long as basal part of Ant. VI. Microstructure developed only with small granules on the frons. Live on <i>Populus koreanus</i>	<i>C. variegatus</i> Szelegiewicz, 1981
10	Body length longer than 1.8 mm, usually over 2 mm. Live on <i>Populus</i> spp.	<i>C. tremulae</i> Koch, 1854
–	Body length 1.3-1.9 mm. Live on <i>Salix</i> spp.	11
11	Abdominal tergite I free. Body length shorter than 1.8 mm, usually about 1.5 mm. Live on <i>Salix</i> spp.	<i>C. salijaponicus</i> Essig and Kuwana, 1918
–	Abdominal tergite I completely fused with the preceding one. Body length shorter than 1.8 mm, usually about 1.4 mm. Live on <i>Salix</i> spp.	<i>C. hokkaidensis</i> Higuchi, 1972

Discussion

Chaitophorus species have been studied from 311 historically old slide-mounted specimens collected across North and South Korea and stored at SNU. Amongst these, we confirmed 10 valid species: *C. horii* Takahashi, 1939 (new record); *C. leucomelas* Koch, 1854 (new record); *C. salijaponicus* Essig and Kuwana, 1918 (new record); *C. tremulae* Koch, 1854 (new record); *C. hokkaidensis* Higuchi, 1972; *C. populeti* (Panzer, 1801); *C. populiaebae* (Boyer de Fonscolombe, 1841); *C. purpureae* Lee and Seo, 1992; *C. saliapterus* Shinji, 1924; and *C. saliniger* Shinji, 1924 (Fig. 5).

In 1941, Shinji reported *C. narae* Shinji, 1941 for the first time from a Korean specimen. However, this species was associated with a host plant from the order Fagales — an atypical host for *Chaitophorus*, which primarily feeds on *Salix* and *Populus* species. Additionally, Shinji's morphological description of the species lacked sufficient detail, complicating subsequent taxonomic efforts. The combination of an unusual host plant,

the limited number of specimens and inadequate morphological characterisation has hindered further research on this species. As a result, *C. narae* Shinji, 1941 has been excluded from the taxonomic key of Korean *Chaitophorus* species.




Figure 5. [doi](#)

Apterous viviparous females of Korean *Chaitophorus* spp. from this study. 1. *C. populeti* 2. *C. saliniger* 3. *C. horii* 4. *C. purpureae* 5. *C. leucomelas* 6. *C. populiabae* 7. *C. salipterus* 8. *C. salijaponicus* 9. *C. tremulae* 10. *C. hokkaidensis*

C. variegatus and *C. matsumurai* were recorded by Szelegiewicz (1981) and Paik (1972), respectively (with the original record of *C. salicicolus*). However, due to a lack of specimens and the absence of any subsequent records, further confirmation of these species is required.

For future studies, an integrated classification system, based on morphological characteristics and DNA barcode data is necessary, along with a comparative analysis with global datasets.

Acknowledgements

We are grateful to Dr. Hong-yul Seo of the National Institute of Biological Resources (NIBR), Incheon 22689, Republic of Korea, for the loan of the *C. purpureae* specimen. We also thank an anonymous reviewer, Dr. Colin Favret and Daisuke Sasaki for their dedicated reviews, which greatly improved this manuscript. This work was supported by the National Research Foundation of Korea (NRF) grant funded by the Korean government (MSIT) (No. RS-2024-00405751) and the National Institute of Biological

Resources (NIBR), funded by the Ministry of Environment (MOE) of the Republic of Korea (NIBR202402202).

References

- Blackman RL, Eastop VF (2024) Aphids on the world's plants: an online identification and information guide. <http://www.aphidinworldsplants.info/>. Accessed on: 2024-8-30.
- Chakrabarti S (1977) Chaitophorinae (Aphididae: Homoptera) of India with descriptions of three new species from north west India. *Oriental Insects* 11: 205-224. <https://doi.org/10.1080/00305316.1977.10434535>
- Favret C, Aphid Taxon Community, eds. (2025) Blackman & Eastop's Aphids on the World's Plants, version 1.0. <https://aphidsonworldsplants.info/>. Accessed on: 2025-3-26.
- Ghosh AK (1980) Part 1. General Introduction and sub-family Chaitophirinae. In: Ghosh AK (Ed.) *The fauna of India and the adjacent countries. Homoptera, Aphidoidea*. Calcutta
- Higuchi H (1972) A taxonomic study of the subfamily Callipterinae in Japan (Homoptera: Aphididae). *Insecta Matsumurana* 35: 19-126.
- Hille Ris Lambers D (1960) The genus *Chaitophorus* Koch in North America (Homoptera, Aphididae). *Tijdschrift voor Entomologie* 103: 1-30.
- Lee W, Seo H (1992) Description of two new species and notes on two unrecorded species (Drepanosiphidae: Chaitophorinae) from Korea. *Korean Journal of Applied Entomology* 22: 251-262.
- Liu T, Chen J, Zhu X, Li C, Jiang L, Wiczorek K, Qiao G (2022) Molecular phylogeny and macroevolution of Chaitophorinae aphids (Insecta: Hemiptera: Aphididae). *Systematic Entomology* 47: 303-314. <https://doi.org/10.1111/syen.12531>
- Nieto Nafria JM, Mier Durante MP (1998) Hemiptera, Aphididae I. *Fauna Iberica*. 11. Museo Nacional de Ciencias Naturales, CSIC, Madrid, Spain, 424 pp.
- Paik WH (1972) *Illustrated Encyclopedia of Fauna and Flora of Korea*. 13. Insecta (V), 165–179 pp.
- Pintera A (1987) Taxonomic revision of the species of genus *Chaitophorus* Koch in Palaearctis (Homoptera: Aphidoidea). *Deutsche Entomologische Zeitschrift* 34: 219-340. <https://doi.org/10.1002/mmnd.19870340402>
- Richards WR (1972) The Chaitophorinae of Canada (Homoptera: Aphididae). *Memoirs of the Entomological Society of Canada* 87: 1-109. <https://doi.org/10.4039/entm10487fv>
- Shingleton AW, Stern DL (2003) Molecular phylogenetic evidence for multiple gains or losses of ant mutualism within the aphid genus *Chaitophorus*. *Molecular Phylogenetics and Evolution* 26: 26-35. [https://doi.org/10.1016/S1055-7903\(02\)00328-7](https://doi.org/10.1016/S1055-7903(02)00328-7)
- Shinji O (1941) *Monograph of Japanese Aphididae*. Tokyo, 137–139 pp. [In Japanese].
- Szelegiewicz H (1981) Two new chaitophorids from the Korean Peninsula. *Annales Zoologici* 36: 37-43.
- Tao CC (1964) Revision of Chinese Callipterinae (Aphidae, Homoptera). *Quarterly Journal of the Taiwan Museum* 17: 209-226.
- Wiczorek K, Osiadacz B (2007) The faunistic review of the Polish species of the subfamily Chaitophorinae (Hemiptera, Aphidoidea). Part 1: genus *Chaitophorus* Koch, 1854. 14-15. *Acta Entomologica Sinica*, 5–19 pp.

- Wieczorek K, Kajtoch Ł (2011) Relationships within Siphini (Hemiptera, Aphidoidea: Chaitophorinae) in light of molecular and morphological research. *Systematic Entomology* 36: 164-174. <https://doi.org/10.1111/j.1365-3113.2010.00550.x>
- Wieczorek K, Lachowska-Cierlik D, Kajtoch Ł, Kanturski M (2017) The relationships within the Chaitophorinae and Drepanosiphinae (Hemiptera, Aphididae) inferred from molecular-based phylogeny and comprehensive morphological data. *PLOS One* 12 (e0173608). URL: <https://doi.org/10.1371/journal.pone.0173608>
- Yang X, Jiang L, Chen J, Qiao G (2016) Study of the evolutionary relationship between *Chaitophorus* aphids and their primary endosymbiont *Buchnera* (Hemiptera: Aphididae: Chaitophorinae). *Chinese Journal of Applied Entomology* 53: 1274-1287.