



# Woody flora of the Prof. Dr. Karl Arens Reserve, Corumbataí, São Paulo, Brazil

Lilian Silva Santos<sup>‡</sup>, Pedro Luís Rodrigues de Moraes<sup>§</sup>

<sup>‡</sup> Universidade Estadual Paulista (UNESP), Instituto de Biociências de Rio Claro, Programa de Pós-Graduação em Biologia Vegetal (Interunidades), Rio Claro, Brazil

<sup>§</sup> Universidade Estadual Paulista (UNESP), Instituto de Biociências de Rio Claro, Departamento de Biodiversidade, Rio Claro, Brazil, Rio Claro, Brazil

Corresponding author: Lilian Silva Santos ([lilian.s.santos@unesp.br](mailto:lilian.s.santos@unesp.br))

Academic editor: Gianmarco Tavilla

Received: 19 Nov 2024 | Accepted: 25 Jan 2025 | Published: 03 Feb 2025

Citation: Santos L, de Moraes PL (2025) Woody flora of the Prof. Dr. Karl Arens Reserve, Corumbataí, São Paulo, Brazil. Biodiversity Data Journal 13: e142217. <https://doi.org/10.3897/BDJ.13.e142217>

## Abstract

## Background

This study was conducted in the Prof. Dr. Karl Arens Reserve, Corumbataí, São Paulo. The area is a *Cerrado* fragment, with no records of fire since 1962 and is characterised by the predominance of *Cerradão* phytophysiology. Although several studies have been conducted in the Reserve, there was no testimony material for some species in the Herbarium Rioclarense (HRCB), nor a list of vouchers for any taxon in the main publications. Thus, the objective of this work was to undertake a floristic survey of the woody species in the Reserve, based on herbarium specimens and fieldwork.

## New information

The survey in HRCB recorded 510 exsiccatae belonging to 160 species and 44 families. We also conducted 24 fieldwork expeditions between 2019 and 2021 and recorded 118 species in 39 families. In total, 193 woody species were recorded for the Reserve in Corumbataí. Species such as *Copaifera langsdorffii* and *Myrcia neoclusiifolia* had already been cited for the area, but there was no testimony material in the herbarium.

*Machaerium nyctitans* and *Alchornea glandulosa* are examples of new records for the Reserve. Fabaceae, Asteraceae, Myrtaceae, Melastomataceae and Rubiaceae were the families with the greatest species diversity.

## Keywords

*Cerradão*, *Cerrado*, floristics, herbarium, species

## Introduction

The Prof. Dr. Karl Arens Reserve was obtained in 1962 by the São Paulo Research Foundation (Fundação de Amparo à Pesquisa do Estado de São Paulo - FAPESP) and is currently managed by the Institute of Bioscience, Rio Claro, of the São Paulo State University (UNESP). It is a fragment of the *Cerrado*, isolated by sugarcane plantations and cattle pastures and has no records of fire since its acquirement (Camargo and Arens 1967, Piccolo et al. 1971).

The *Cerrado* is the second largest phytogeographic domain in Brazil (Ab'Sáber 1983, Batalha 2011) and occupies 23% of the country (Ribeiro and Walter 2008). This important hotspot (Myers et al. 2000, Strassburg et al. 2017) represents 30% of the biodiversity in the country (Françoso et al. 2015). In the State of São Paulo, the *Cerrado* vegetation is severely fragmented (Eiten 1994, Cassavan 2002, Nalon et al. 2022), being part of the most threatened *Cerrado* ecoregion (Sano et al. 2019). According to the last inventory of the native vegetation cover (Nalon et al. 2022), of the 8,106,085 ha of original cover, only 239,312 ha remain.

The vegetation in the *Cerrado* is not uniform (Coutinho 2002) and includes field, savannah and forest physiognomies (Ribeiro and Walter 2008). Fire can change *Cerrado* physiognomies, impacting the vegetation (Eiten 1972). The absence of fire allows a savannah formation to become denser and form *Cerradão*, which is already the dominant formation in São Paulo State (Durigan et al. 2004a).

*Cerradão* is the tallest physiognomy in the *Cerrado* (Eiten 1994), characterised by a forest formation with trees 8–15 m tall, mostly continuous canopy and distinct layers of shrubs and herbs in the understorey due to variation in light reaching the ground. Individuals that occur in *cerrado sensu stricto* and other forest formations are common in this physiognomy (Ribeiro and Walter 2008). The Reserve in Corumbataí comprises *Cerradão* and *Cerrado sensu stricto* (Pinheiro et al. 2010, Pinheiro et al. 2021). The change in the proportion of the area occupied by each physiognomy is notable over more than 60 years. Due to the expansion and dominance of *Cerradão* in the fragment, only 3% of the *Cerrado sensu stricto* remains out of more than 34 initial hectares (Pinheiro et al. 2021). This phenomenon is mainly associated with the absence of fire (Pinheiro et al. 2010, Pinheiro et al. 2021), which changes the landscape, provokes a decrease in biodiversity and mostly harms non-arboreal species (Durigan 2020).

Since the Institute of Biosciences at UNESP in Rio Claro obtained custody of the Reserve, studies have been conducted in different areas of plant biology, mainly phytosociological and floristic works (Camargo and Arens 1967, Piccolo et al. 1971, Cesar et al. 1988, Campos 1989, Pagano et al. 1989, Paula and Monteiro 2000, Andena et al. 2005, Pinheiro 2013, Pinheiro et al. 2021).

Despite the studies and large species lists made for the Reserve, gaps were observed, such as the absence of testimony material of several taxa in the Herbarium Rioclarense (HRCB) and lack of citations of vouchers in those works. Thus, the objective of the present study was to identify and record the woody species in the Prof. Dr. Karl Arens Reserve, Corumbataí, São Paulo, based on a floristic survey and herbarium specimens mainly deposited at HRCB and from a set housed at SPFR, with images available in speciesLink (2024).

## Materials and methods

### Study area

The Prof. Dr. Karl Arens Reserve (22°14'30"S, 47°41'10"W) is located in Corumbataí, São Paulo, approximately 30 km from the city of Rio Claro (Fig. 1) and 200 km from the state capital. The study area is in the Paraná Sedimentary Basin, in a transition region between the Basalt Cuestas and Peripheral Depression geomorphological Provinces (Troppmair 2000, Marques e Amorim et al. 2005), more specifically on the Residual Plateau of Brotas-Itirapina (IBGE 2023).

The fragment is in the *Cerrado* domain and has an area of almost 39 ha (Pinheiro et al. 2010). The area is surrounded by a monoculture of sugarcane and pasture and ranges from 793–866 m. *Cerradão* is predominant in relation to *Cerrado sensu strictu* (Pinheiro et al. 2021) and there are fragments of semi-deciduous seasonal forest on nearby hills (Pinheiro et al. 2010, Pinheiro et al. 2021).

The soil is a dystrophic, Red-Yellow/Red Latosol with a medium texture (Rossi 2017) and, despite the presence of clay, is predominantly sandy (Cesar et al. 1988, Pinheiro et al. 2010). The climate in the region is the Cwa type: subtropical humid with a cold and dry period between April and September and a hot and rainy period between October and March (Peel et al. 2007). The average temperature from 2015–2023 was 21.8°C, with a minimum of 12.8°C and a maximum of 30.9°C (CIIAGRO 2023). The average annual precipitation was 1492 mm between 1987 and 2017 (ANA 2021) and 1197 mm between 2018 and 2023 (CIIAGRO 2023).

### Obtaining and analysing the material

Exsiccatae from the Reserve in Corumbataí were selected from the Herbarium Rioclarense (HRCB), which belongs to the Institute of Biosciences at UNESP in Rio Claro. HRCB has the largest and most important collection of specimens from the

Reserve of Corumbataí, with duplicates that were distributed to many herbaria i.e.: ALCB, BHCB, CEN, CESJ, CPAP, ESA, FLOR, FUEL, HUEM, HUESB, HUFJSJ, HUFU, HUPG, HXBH, IAC, INPA, MBM, MBML, MO, PAMG, PEUFR, SJRP, SP, SPF, SPFR, SPSF, UEC and UPCB. We manually checked all the fertile material with subshrub, shrub and tree habits in HRCB. We did not include herbs and vines because we lack adequate reference collections and we had limited time available for more extensive sampling efforts. Information, such as voucher number, identification, collector and collection date, were taken from the labels.




Figure 1. [doi](#)

Study Area: location and natural coverage of *Cerrado* in São Paulo State, including the Municipality of Corumbataí, its border with Itirapina and Rio Claro and limits from Prof. Dr. Karl Arens Reserve. Prepared by: Lilian S. Santos.

Additionally, we conducted rapid floristic walkover surveys on foot (Figueiras et al. 1994) in the fragment to collect woody and subwoody material, except for vines. However, facultative climbers were considered in this study, as well as "woody" palms. During these field expeditions, we walked along the entire perimeter of the Reserve, along the trail that crosses the fragment, as well as into the middle of the fragment from this trail. There were no predetermined transects; the walking was random and focused on scanning as much area as possible in search of fertile material. A drone was used to take images of the area and we took photographs *in situ* from the specimens to be obtained. The collected material was processed in the herbarium (Fidalgo and Bononi 1989) and exsiccatae were prepared and incorporated into HRCB.

All the material obtained in the field and herbarium was checked and determined using studies conducted in the area (Camargo and Arens 1967, Piccolo et al. 1971, Cesar et al. 1988, Pagano et al. 1989, Andena et al. 2005, Pinheiro et al. 2010, Pinheiro 2013, Pinheiro et al. 2021), floras for the State, for example, Flora Fanerogâmica do Estado de São Paulo (Romão and Souza 2016) and field guides (Durigan et al. 2004b, Souza et al. 2018). The specimens were analysed using a Leica M80 stereomicroscope and the software Leica LAS EZ.

Morphological comparisons were made with material in HRCB and on virtual herbaria (e.g. MNHN (2023), speciesLink (2024)). Dictionaries and glossaries were used to help interpret the terminology and morphology (Font Quer 1953, Harris and Harris 1994, Hickey and King 2000, Gonçalves and Lorenzi 2011). Species life-form was verified in the Flora e Funga do Brasil (2024) platform and specialised literature of the different taxonomic groups. The families follow APG IV (2016), the epithets and authors were updated (IPNI 2024, speciesLink 2024) and the herbaria acronyms follow Thiers [continuously updated] (2024).

The software QGIS version 3.36.1 was used to create the maps and georeference the satellite image. The remote sensing image and coordinates were obtained with the software Google Earth Pro version 7.3.6. All figures were edited with the software Adobe Photoshop CS5.

## Data resources

We found 510 exsiccatae in HRCB, distributed in 160 species, 111 genera and 44 families of angiosperms (Table 1). Nearly half of the collections are from 1981–1985, especially 1984 with 104 exsiccatae (Fig. 2). During that period, most collections were gathered by O. Cesar & S.N. Pagano and M.J.O. Campos, resulting, respectively, in a publication on the phytosociological structure of the Reserve's tree stratum (Cesar et al. 1988) and a doctoral thesis (Campos 1989). Smaller collection peaks occurred in 1962, led by H. Amaral and, in 2000, with collections conducted by C.E. Carneiro, L. Cordeiro, V.B. Ziparro and by V.F.O. Miranda. However, the only work potentially associated with those collections is Piccolo et al. (1971). The complete list of the specimens analysed and their respective vouchers are in the Supplementary Material (Suppl. material 1).

Table 1.

Checklist of the taxa from the Prof. Dr. Karl Arens Reserve; specimens obtained from the HRCB collection and field floristic survey in 2019, 2020, 2021 and 2023. Number of vouchers in brackets. Taxa organised, based on APG IV.

### Checklist of the taxa from the Prof. Dr. Karl Arens Reserve

PHYLUM > CLASS > ORDER > Family > Species	HRCB	SURVEY
TRACHEOPHYTA		
MAGNOLIOPSIDA		

<b>Checklist of the taxa from the Prof. Dr. Karl Arens Reserve</b>		
<b>LAURALES</b>		
<b>Lauraceae</b>		
<i>Ocotea corymbosa</i> (Meisn.) Mez	(1)	(6)
<i>Ocotea pulchella</i> (Nees & Mart.) Mez	(8)	(8)
<i>Persea willdenovii</i> Kosterm.	(3)	
<b>Siparunaceae</b>		
<i>Siparuna guianensis</i> Aubl.	(4)	(3)
<b>MAGNOLIALES</b>		
<b>Annonaceae</b>		
<i>Annona coriacea</i> Mart.		(7)
<i>Duguetia furfuracea</i> (A.St.-Hil.) Saff.		(3)
<i>Guatteria australis</i> A.St.-Hil.		(4)
<i>Xylopia aromatica</i> (Lam.) Mart.	(3)	(4)
<b>Myristicaceae</b>		
<i>Virola sebifera</i> Aubl.	(8)	(5)
<b>LILIOPSIDA</b>		
<b>ARECALES</b>		
<b>Areceaceae</b>		
<i>Syagrus flexuosa</i> (Mart.) Becc.		(1)
<b>MAGNOLIOPSIDA</b>		
<b>PROTEALES</b>		
<b>Proteaceae</b>		
<i>Roupala montana</i> Aubl.	(4)	(5)
<b>DILLENIALES</b>		
<b>Dilleniaceae</b>		
<i>Davilla elliptica</i> A.St.-Hil.	(1)	(1)
<i>Davilla rugosa</i> Poir.	(4)	(2)
<b>CELASTRALES</b>		
<b>Celastraceae</b>		
<i>Monteverdia evonymoides</i> (Reissek) Biral	(1)	
<i>Peritassa campestris</i> (Cambess.) A.C.Sm.	(5)	
<i>Plenckia populnea</i> Reissek	(1)	(3)
<b>OXALIDALES</b>		
<b>Connaraceae</b>		
<i>Connarus suberosus</i> Planch.	(4)	(1)
<b>MALPIGHIALES</b>		

<b>Checklist of the taxa from the Prof. Dr. Karl Arens Reserve</b>		
<b>Calophyllaceae</b>		
<i>Kielmeyera coriacea</i> Mart. & Zucc.	(2)	
<b>Caryocaraceae</b>		
<i>Caryocar brasiliense</i> Cambess.	(8)	(5)
<b>Chrysobalanaceae</b>		
<i>Couepia grandiflora</i> (Mart. & Zucc.) Benth.	(3)	
<i>Leptobalanus humilis</i> (Cham. & Schtdl.) Sothers & Prance	(1)	
<b>Erythroxylaceae</b>		
<i>Erythroxylum campestre</i> A.St.-Hil.	(2)	
<i>Erythroxylum cuneifolium</i> (Mart.) O.E.Schulz	(4)	(4)
<i>Erythroxylum daphnites</i> Mart.	(2)	
<i>Erythroxylum deciduum</i> A.St.-Hil.	(1)	
<i>Erythroxylum pelleterianum</i> A.St.-Hil.	(3)	(1)
<i>Erythroxylum suberosum</i> A.St.-Hil.	(5)	
<b>Euphorbiaceae</b>		
<i>Alchornea glandulosa</i> Poepp. & Endl.		(1)
<i>Croton gnaphaloides</i> Schrad.	(4)	(2)
<i>Microstachys serrulata</i> (Mart.) Müll.Arg.	(3)	
<i>Sapium glandulosum</i> (L.) Morong	(5)	(1)
<b>Lacistemataceae</b>		
<i>Lacistema hasslerianum</i> Chodat	(7)	(4)
<b>Malpighiaceae</b>		
<i>Banisteriopsis campestris</i> (A.Juss.) Little	(1)	
<i>Banisteriopsis stellaris</i> (Griseb.) B.Gates		(4)
<i>Byrsonima coccolobifolia</i> Kunth	(4)	(1)
<i>Byrsonima intermedia</i> A.Juss.	(8)	(8)
<i>Byrsonima verbascifolia</i> (L.) DC.	(4)	
<i>Heteropterys umbellata</i> A.Juss.	(3)	
<i>Peixotoa tomentosa</i> A.Juss.	(1)	(1)
<b>Ochnaceae</b>		
<i>Ouratea spectabilis</i> (Mart.) Engl.	(3)	(4)
<b>Peraceae</b>		
<i>Pera glabrata</i> (Schott) Baill.	(14)	(10)
<b>Salicaceae</b>		
<i>Casearia sylvestris</i> Sw.	(7)	(5)
<b>FABALES</b>		

<b>Checklist of the taxa from the Prof. Dr. Karl Arens Reserve</b>		
<b>Fabaceae</b>		
<i>Anadenanthera peregrina</i> (L.) Speg.	(4)	(5)
<i>Andira humilis</i> Benth.	(2)	
<i>Andira vermifuga</i> (Mart.) Benth.	(4)	(1)
<i>Bauhinia brevipes</i> Vogel	(3)	(4)
<i>Bauhinia holophylla</i> (Bong.) Steud.	(3)	(2)
<i>Cerradicola</i> cf. <i>decumbens</i> (Benth.) L.P.Queiroz	(3)	
<i>Chamaecrista cathartica</i> (Mart.) H.S.Irwin & Barneby	(4)	
<i>Chamaecrista desvauxii</i> (Collad.) Killip	(3)	
<i>Chamaecrista flexuosa</i> (L.) Greene	(6)	(3)
<i>Chamaecrista rotundifolia</i> (Pers.) Greene	(3)	
<i>Clitoria densiflora</i> (Benth.) Benth.	(2)	
<i>Copaifera langsdorffii</i> Desf.		(6)
<i>Crotalaria maypurensis</i> Kunth	(2)	
<i>Crotalaria micans</i> Link		(2)
<i>Crotalaria pallida</i> Aiton var. <i>obovata</i> (G.Don) Polhill	(1)	
<i>Dahlstedtia muehlbergiana</i> (Hassl.) M.J.Silva & A.M.G.Azevedo	(1)	
<i>Dalbergia frutescens</i> (Vell.) Britton var. <i>frutescens</i>	(1)	
<i>Dalbergia miscolobium</i> Benth.	(1)	(3)
<i>Desmodium tortuosum</i> (Sw.) DC.	(1)	
<i>Dimorphandra mollis</i> Benth.	(2)	(4)
<i>Indigofera hirsuta</i> L.	(1)	
<i>Leptolobium dasycarpum</i> Vogel	(4)	(4)
<i>Leptolobium elegans</i> Vogel		(8)
<i>Machaerium acutifolium</i> Vogel	(3)	(4)
<i>Machaerium nyctitans</i> (Vell.) Benth.		(3)
<i>Machaerium villosum</i> Vogel	(1)	
<i>Mimosa debilis</i> Willd.	(2)	(1)
<i>Mimosa xanthocentra</i> Mart. var. <i>subsericea</i> (Benth.) Barneby	(3)	(1)
<i>Senegalia lowei</i> (L.Rico) Seigler & Ebinger	(2)	
<i>Senna pendula</i> (Willd.) H.S.Irwin & Barneby	(1)	(2)
<i>Senna rugosa</i> (G.Don) H.S.Irwin & Barneby	(4)	(5)
<i>Stryphnodendron adstringens</i> (Mart.) Coville	(5)	
<i>Stryphnodendron rotundifolium</i> Mart.	(4)	(8)
<i>Stylosanthes scabra</i> Vogel	(1)	
<i>Stylosanthes viscosa</i> (L.) Sw.	(1)	



<b>Checklist of the taxa from the Prof. Dr. Karl Arens Reserve</b>		
<b>Polygalaceae</b>		
<i>Bredemeyera floribunda</i> Willd.	(2)	(3)
<i>Polygala poaya</i> Mart.	(6)	
<b>ROSALES</b>		
<b>Moraceae</b>		
<i>Ficus guaranitica</i> Chodat		(3)
<b>MYRTALES</b>		
<b>Lythraceae</b>		
<i>Cuphea thymoides</i> Cham. & Schtdl.	(1)	
<i>Diplusodon virgatus</i> Pohl	(6)	(4)
<i>Lafoensia pacari</i> A.St.-Hil.	(1)	(3)
<b>Melastomataceae</b>		
<i>Leandra aurea</i> (Cham.) Cogn.	(2)	
<i>Miconia albicans</i> (Sw.) Triana	(5)	(1)
<i>Miconia fallax</i> DC.	(2)	
<i>Miconia flammea</i> Casar.	(3)	(3)
<i>Miconia ligustroides</i> (DC.) Naudin	(9)	(9)
<i>Miconia paucidens</i> DC.	(1)	
<i>Miconia rubiginosa</i> (Bonpl.) DC.	(9)	(14)
<i>Miconia sellowiana</i> Naudin	(2)	(4)
<i>Miconia stenostachya</i> DC.	(4)	(2)
<i>Microlicia polystemma</i> Naudin	(1)	
<i>Pleroma stenocarpum</i> (DC.) Triana	(4)	
<b>Myrtaceae</b>		
<i>Blepharocalyx salicifolius</i> (Kunth) O.Berg	(2)	(1)
<i>Campomanesia pubescens</i> (DC.) O.Berg	(6)	(2)
<i>Eugenia aurata</i> O.Berg	(2)	(1)
<i>Eugenia bimarginata</i> DC.	(4)	(3)
<i>Eugenia</i> cf. <i>hiemalis</i> Cambess.		(3)
<i>Eugenia puniceifolia</i> (Kunth) DC.	(4)	(5)
<i>Myrcia bela</i> Cambess.	(5)	(1)
<i>Myrcia guianensis</i> (Aubl.) DC.	(4)	(7)
<i>Myrcia neoclusiifolia</i> A.R.Lourenço & E.Lucas		(1)
<i>Myrcia splendens</i> (Sw.) DC.	(2)	(1)
<i>Myrcia tomentosa</i> (Aubl.) DC.	(2)	
<i>Myrciaria floribunda</i> (Willd.) O.Berg	(3)	(1)

<b>Checklist of the taxa from the Prof. Dr. Karl Arens Reserve</b>		
<i>Psidium grandifolium</i> DC.	(2)	
<b>Vochysiaceae</b>		
<i>Qualea cordata</i> Spreng.	(1)	(2)
<i>Qualea grandiflora</i> Mart.	(3)	(5)
<i>Qualea multiflora</i> Mart.	(3)	(5)
<i>Vochysia tucanorum</i> Mart.	(10)	(6)
<b>SAPINDALES</b>		
<b>Meliaceae</b>		
<i>Cedrela fissilis</i> Vell.		(3)
<i>Guarea guidonia</i> (L.) Sleumer		(1)
<b>Sapindaceae</b>		
<i>Serjania erecta</i> Radlk.	(2)	
<i>Talisia angustifolia</i> Radlk.	(1)	
<b>MALVALES</b>		
<b>Bixaceae</b>		
<i>Cochlospermum regium</i> (Schrank) Pilg.	(1)	
<b>Malvaceae</b>		
<i>Byttneria sagittifolia</i> A.St.-Hil.	(1)	
<i>Eriotheca gracilipes</i> (K.Schum.) A.Robyns	(2)	(3)
<i>Luehea grandiflora</i> Mart.		(1)
<i>Pavonia malacophylla</i> (Link & Otto) Garcke	(1)	
<b>Thymelaeaceae</b>		
<i>Daphnopsis fasciculata</i> (Meisn.) Nevling	(5)	
<b>CARYOPHYLLALES</b>		
<b>Nyctaginaceae</b>		
<i>Guapira noxia</i> (Netto) Lundell	(8)	(3)
<i>Neea theifera</i> Oerst.	(5)	(3)
<b>ERICALES</b>		
<b>Ebenaceae</b>		
<i>Diospyros lasiocalyx</i> (Mart.) B.Walln.	(1)	
<b>Primulaceae</b>		
<i>Myrsine cf. coriacea</i> (Sw.) Roem. & Schult.	(1)	
<i>Myrsine guianensis</i> (Aubl.) Kuntze	(4)	
<i>Myrsine lancifolia</i> Mart.	(1)	(1)
<i>Myrsine umbellata</i> Mart.	(4)	(2)
<b>Sapotaceae</b>		

<b>Checklist of the taxa from the Prof. Dr. Karl Arens Reserve</b>		
<i>Pouteria ramiflora</i> (Mart.) Radlk.	(1)	
<i>Pouteria torta</i> (Mart.) Radlk.	(3)	
<b>Styracaceae</b>		
<i>Styrax camporum</i> Pohl	(7)	(6)
<i>Styrax ferrugineus</i> Nees & Mart.	(4)	
<b>Symplocaceae</b>		
<i>Symplocos oblongifolia</i> Casar.	(3)	
<i>Symplocos pubescens</i> Klotzsch ex Benth.	(4)	(1)
<b>SOLANALES</b>		
<b>Solanaceae</b>		
<i>Cestrum mariquitense</i> Kunth	(3)	(2)
<i>Solanum</i> cf. <i>didymum</i> Dunal		(1)
<i>Solanum granuloseprosum</i> Dunal	(1)	(3)
<i>Solanum lycocarpum</i> A.St.-Hil.	(5)	(4)
<b>GENTIANALES</b>		
<b>Apocynaceae</b>		
<i>Aspidosperma tomentosum</i> Mart.	(1)	
<i>Tabernaemontana catharinensis</i> A.DC.	(1)	
<b>Loganiaceae</b>		
<i>Strychnos bicolor</i> Progel	(3)	(1)
<b>Rubiaceae</b>		
<i>Amaioua intermedia</i> Mart.	(8)	(7)
<i>Cordia obtusa</i> (K.Schum.) Kuntze	(2)	
<i>Cordia sessilis</i> (Vell.) Kuntze		(2)
<i>Declieuxia fruticosa</i> (Roem. & Schult.) Kuntze	(1)	
<i>Palicourea hoffmannseggiana</i> (Schult.) Borhidi	(2)	(1)
<i>Palicourea marcgravi</i> A.St.-Hil.	(3)	(4)
<i>Palicourea rigida</i> Kunth	(6)	
<i>Palicourea sessilis</i> (Vell.) C.M. Taylor	(5)	(3)
<i>Palicourea violacea</i> (Aubl.) A.Rich	(5)	(4)
<i>Tocoyena formosa</i> (Cham. & Schldl.) K.Schum.	(4)	(2)
<b>LAMIALES</b>		
<b>Bignoniaceae</b>		
<i>Adenocalymma peregrinum</i> (Miers) L.G.Lohmann		(2)
<i>Anemopaegma acutifolium</i> DC.	(1)	
<i>Anemopaegma arvense</i> (Vell.) J.F.Souza	(1)	

<b>Checklist of the taxa from the Prof. Dr. Karl Arens Reserve</b>		
<i>Handroanthus</i> cf. <i>ochraceus</i> (Cham.) Mattos	(1)	
<i>Jacaranda caroba</i> (Vell.) DC.	(3)	(2)
<i>Jacaranda decurrens</i> Cham.	(1)	
<i>Zeyheria montana</i> Mart.	(1)	
<b>Lamiaceae</b>		
<i>Aegiphila integrifolia</i> (Jacq.) Moldenke		(2)
<i>Aegiphila verticillata</i> Vell.		(1)
<i>Cyanocephalus lippoides</i> (Benth.) Harley & J.F.B.Pastore	(5)	
<i>Medusantha eriophylla</i> (Benth.) Harley & J.F.B.Pastore	(3)	(1)
<i>Medusantha</i> sp.		(2)
<b>Verbenaceae</b>		
<i>Lantana camara</i> L.		(2)
<i>Lantana fucata</i> Lindl.	(1)	(1)
<i>Lippia origanoides</i> Kunth	(2)	(3)
<i>Stachytarpheta cayennensis</i> (Rich.) Vahl	(1)	
<b>AQUIFOLIALES</b>		
<b>Aquifoliaceae</b>		
<i>Ilex cerasifolia</i> Reissek	(3)	(4)
<b>ASTERALES</b>		
<b>Asteraceae</b>		
<i>Achyrocline</i> cf. <i>satureioides</i> (Lam.) DC.	(2)	(1)
<i>Aldama arenaria</i> (Baker) E.E.Schill. & Panero	(4)	
<i>Baccharis dracunculifolia</i> DC.	(4)	(1)
<i>Chresta sphaerocephala</i> DC.	(3)	
<i>Chromolaena odorata</i> (L.) R.M.King & H.Rob.		(1)
<i>Chromolaena squalida</i> (DC.) R.M.King & H.Rob.	(2)	(1)
<i>Chrysolaena cognata</i> (Less.) Dematt.	(2)	
<i>Grazielia dimorpholepis</i> (Baker) R.M.King & H.Rob.	(1)	
<i>Heterocondylus alatus</i> (Vell.) R.M.King & H.Rob.	(5)	
<i>Hoehnephytum trixoides</i> (Gardner) Cabrera	(1)	
<i>Lepidaploa</i> sp.1		(2)
<i>Lepidaploa</i> sp.2		(1)
<i>Lessingianthus bardanoides</i> (Less.) H.Rob.	(1)	
<i>Moquiniastrum barrosoae</i> (Cabrera) G.Sancho	(3)	(3)
<i>Moquiniastrum polymorphum</i> (Less.) G.Sancho		(4)
<i>Moquiniastrum pulchrum</i> (Cabrera) G.Sancho	(4)	(1)

Checklist of the taxa from the Prof. Dr. Karl Arens Reserve		
<i>Piptocarpha macropoda</i> (DC.) Baker		(1)
<i>Piptocarpha rotundifolia</i> (Less.) Baker	(8)	
<i>Pterocaulon lanatum</i> Kuntze		(2)
<i>Trixis</i> cf. <i>antimenorrhoea</i> (Schrank) Kuntze		(1)
<i>Vernonanthura ferruginea</i> (Less.) H. Rob.	(1)	
<i>Vernonanthura polyanthes</i> (Spreng.) Vega & Dematteis		(1)
<i>Vernonanthura</i> cf. <i>rubiramea</i> (DC.) Loeuille & P.N. Soares		(2)
<b>APIALES</b>		
<b>Araliaceae</b>		
<i>Schefflera vinosa</i> (Cham. & Schldl.) Frodin & Fiaschi	(6)	(5)




Figure 2. [doi](#)

Specimens obtained from the HRCB collection: most collected periods in the Prof. Dr. Karl Arens Reserve.

Fabaceae, Asteraceae, Melastomataceae, Myrtaceae and Rubiaceae were the families with the greatest species diversity in the HRCB collection (Fig. 3). Nevertheless, Annonaceae, Aquifoliaceae, Araliaceae, Bixaceae, Calophyllaceae, Caryocaraceae, Connaraceae, Ebenaceae, Lacistemataceae, Loganiaceae, Myristicaceae, Ochnaceae, Peraceae, Proteaceae, Salicaceae, Siparunaceae and Thymelaeaceae were represented by only one species each (Table 1).

Between May 2019 and July 2021, 24 fieldwork expeditions were undertaken. The interval between the expeditions varied due to rainy and dry periods which affected finding the same fertile species and lockdown because of the Covid-19 pandemic. During this time, 364 individuals distributed in 116 species, 82 genera and 39 families of angiosperms were collected. In October and November of 2023, two new collections were made, totalling 366 individuals, 118 species, 84 genera and 39 families (Table 1). A complete list of the vouchers is presented in Supplementary Material (Suppl. material 1).

Fabaceae (66), Asteraceae (22), Myrtaceae (26), Rubiaceae (23) and Melastomataceae (33) also had the greatest species diversity (Fig. 4). Aquifoliaceae, Araliaceae, Arecaceae, Caryocaraceae, Celastraceae, Connaraceae, Lacistemataceae, Loganiaceae, Moraceae, Myristicaceae, Ochnaceae, Peraceae, Polygalaceae,

Proteaceae, Salicaceae, Siparunaceae, Styracaceae and Symplocaceae had only one taxon each (Table 1).




Figure 3. [doi](#)

Specimens obtained from the HRCB collection: families with the largest diversity in species.

For the HRCB and floristic surveys, there was a total of 192 species, 126 genera and 47 families (Table 1). However, 193 species were considered for the Reserve in Corumbataí, since *Syagrus romanzoffiana* (Cham.) Glassman occurs in this area, but was not collected. Popularly known as *jerivá*, this is a common species in the fragment and has been cited for the area (Pinheiro et al. 2021). The families with the greatest species diversity can be seen in Fig. 5. Exotic specimens of *Syzygium jambos* (L.) Alston (*jambo*) and *Mangifera indica* L. (*manga*) were collected, but are not on the list of taxa in the present study. Sterile individuals of *Pouteria* sp. and *Aspidosperma* sp. were also observed.

## Checklist of taxa from the Prof. Dr. Karl Arens Reserve

### Class Liliopsida and Magnoliopsida

**Notes:** The list of taxa recorded in this study is presented in Table 1. Details of the examined material from the Dr. Karl Arens Reserve, Corumbataí (SP), are provided in Supplementary Material (Suppl. material 1).




Figure 4. [doi](#)

Field floristic surveys conducted in 2019, 2020, 2021 and 2023: families with the largest diversity in species.

## Discussion

Species such as *Xylopia aromatica* (Lam.) Mart. (HRCB 17668), *Schefflera vinosa* (Cham. & Schlttdl.) Frodin & Fiaschi (HRCB 32188), *Caryocar brasiliense* Cambess. (HRCB 32249), *Sapium glandulosum* (L.) Morong (HRCB 55742), *Bauhinia brevipes* Vogel (HRCB 5516), *Leptolobium dasycarpum* Vogel (HRCB 27646) and *Machaerium acutifolium* Vogel (HRCB 17667) have been cited for the Reserve (Camargo and Arens 1967, Piccolo et al. 1971, Cesar et al. 1988, Pagano et al. 1989, Andena et al. 2005, Pinheiro 2013, Pinheiro et al. 2010, Pinheiro et al. 2021). However, those authors did not present their respective vouchers, even though there is herbarium material of them existing at HRCB and SPFR. Furthermore, no vouchers are cited for any other taxa in these works.

*Annona crassiflora* Mart., *Syagrus romanzoffiana*, *Protium heptaphyllum* (Aubl.) Marchand, *Monteverdia gonoclada* (Mart.) Biral, *Terminalia glabrescens* Mart., *Nectandra megapotamica* (Spreng.) Mez, *Ocotea acutifolia* (Nees) Mez, *Trichilia hirta* L., *Campomanesia guaviroba* (DC.) Kiaersk., *Savia dictyocarpa* Müll.Arg., *Esenbeckia grandiflora* Mart., *Zanthoxylum rhoifolium* Lam. and *Copaifera langsdorffii* Desf. are cited as present in the Reserve (Pinheiro et al. 2021); however, there is no testimony material in HRCB; however, with the exception of *Syagrus* and *Copaifera*, none of the cited species was found in our floristic survey. This may indicate either a change in the flora of

the Reserve or incorrect identification. For example, the cited *Ocotea acutifolia* is likely HRCB 5448, later recognised as *Ocotea corymbosa* (Meisn.) Mez.




Figure 5. [doi](#)

HRCB collection and field floristic survey: the families with the largest diversity in species.

*Annona cacans* Warm., *Croton floribundus* Spreng. and *Piptocarpha axillaris* (Less.) Baker are other species cited for the Reserve that have no testimony material and, according to Flora e Funga do Brasil (2024), occur in ombrophilous forests. On the other hand, *Persea willdenovii* Kosterm. (HRCB 59275) was collected in 2012, but is not on any list in the works published. According to Moraes (personal observation), the only known individual in the Reserve died in 2013. *Ocotea corymbosa* (Meisn.) Mez (HRCB 5448) was collected in 1984 and is also not mentioned in the recent studies.

Some exsiccatae in the collection have inconsistent localities, with two municipalities (Corumbataí and Itirapina), for example, HRCB 1490, HRCB 1325, HRCB 1328, HRCB 36700, HRCB 1310, HRCB 1591, HRCB 1592, HRCB 1388, HRCB 1465, HRCB 1477, HRCB 1497, HRCB 1598, HRCB 1102, HRCB 1103, HRCB 1104, HRCB 1105, HRCB 1106, HRCB 1121, HRCB 1122 and HRCB 1123). *Anacardium humile* A.St.-Hil., *Annona*



*cornifolia* A.St.-Hil, and *Duguetia furfuracea* (A.St.-Hil.) Saff. are amongst these vouchers and are species cited in a study only made with herbarium material (Pinheiro 2013). The specimens with a conflicting locality on their labels are not on the species list of the present study.

Pinheiro et al. (2021) determined 510 individuals from the Reserve, which were distributed in 103 species and 43 families. However, we only found two collections in HRCB made by the first author (HRCB 45190, *Pera glabrata* (Schott) Poepp. ex Baill. and HRCB 45191, *Daphnopsis fasciculata* (Meisn.) Nevling), both from 2003 and without a collection number (Table 1). Providing a voucher from analysed specimens is important for increasing the credibility of studies and enabling the location of the examined material. According to Baker et al. (2017), a reliable identification is linked to a well-prepared and cited voucher. Adequate documentation is fundamental to allow comparisons, improvements (Schilthuizen et al. 2015) or corrections. This practice forms the basis of collections and scientific reproducibility (Funk et al. 2005); however, the studies with the specimens from the Reserve have been neglected over the years.

*Copaifera langsdorffii*, *Syagrus flexuosa* (Mart.) Becc., *Leptolobium elegans* Vogel, *Eugenia* cf. *hiemalis* Cambess. and *Myrcia neoclusiifolia* A.R.Lourenço & E.Lucas (Fig. 6) were added to the HRCB collection for the first time, although they were already cited for the Reserve. According to Pinheiro et al. (2021), *Copaifera langsdorffii* is one of the most abundant species in the Reserve, as observed by the authors in the present study, particularly along the northern border of the fragment.



Figure 6. [doi](#)

Some of the species mentioned in previous studies that did not have testimonial material deposited in the HRCB: (a–b) *Syagrus flexuosa*; (c–e) *Copaifera langsdorffii*; (f–h) *Leptolobium elegans*. Photos: Lilian S. Santos (a–f).

On the other hand, *Annona coriacea* Mart., *Duguetia furfuracea*, *Guatteria australis* A.St.-Hil., *Moquiniastrum polymorphum* (Less.) G.Sancho, *Piptocarpha macropoda* (DC.) Baker, *Alchornea glandulosa* Poepp. & Endl., *Machaerium nyctitans* (Vell.) Benth., *Aegiphila integrifolia* (Jacq.) Moldenke, *Aegiphila verticillata* Vell., *Cedrela fissilis* Vell., *Ficus guaranitica* Chodat, *Luehea grandiflora* Mart. and *Guarea guidonia* (L.) Sleumer did not have testimony material, were not on species lists for the Reserve (Pinheiro et al. 2021) and were added to HRCB for the first time (Fig. 7). Despite *A. coriacea* being added to the HRCB collection for the first time, there are records of this species from Corumbataí in other herbaria (ESA 023328 and PEL 20336).




Figure 7. [doi](#)

Some of the species not mentioned in previous studies and without testimonial material deposited in the HRCB: (a–b) *Annona coriacea*; (c–d) *Duguetia furfuracea*; (e–f) *Piptocarpha macropoda*; (g–h) *Alchornea glandulosa*; (i–j) *Machaerium nyctitans*; (k–l) *Aegiphila verticillata*. Photos: Lilian S. Santos (a,b,e–l); Alessandra I. Coan (c–d).

The floristic survey recorded fewer species and families than what was found in the herbarium. This could be related to the phenology of the plants and their fertile periods (Baker et al. 2017), as well as to the chance of finding plants with flowers or fruits (e.g. *Pouteria* sp. that was only observed in a vegetative state). The field of vision of the collector is impaired in denser vegetation and this may have influenced the lower number of taxa collected. The lower number of species may also be related to the difficulty some plants have in competing for light, since the *Cerradão* grows over the *Cerrado sensu stricto*. Although most of the collected specimens were found on the border and not in the middle of the fragment, the denser vegetation still harbours species that are not found elsewhere within the Reserve, such as *Andira vermifuga* (Mart.) Benth. and *Piptocarpha macropoda* (DC.) Baker. Despite being modest, differences were observed between the species diversity of the floristic survey and the HRCB collection. The family Annonaceae, for example, was represented by only one species in the herbarium and is now represented by four taxa (Table 1).

According to Durigan and Ratter (2006), the change in the physiognomy is accompanied by changes in the floristic composition. The densification of the vegetation makes it difficult for sunlight to enter, which affects the species from *cerrado sensu stricto* in the understorey (Pinheiro and Durigan 2012, Pinheiro et al. 2021). In the most closed areas in the middle of the fragment, dead individuals were observed with very suberised trunks. The absence of fires can be associated with a decrease in species in more open areas and an increase in large generalist species (Pinheiro et al. 2010, Pinheiro et al. 2021). Additionally, the absence of disturbances, such as fire and deforestation, contribute to an increase in tree cover to the detriment of grasses (Scholes and Archer 1997) and other plants more dependent on the sun. The single or few exsiccatae of *Aspidosperma tomentosum* Mart., *Cochlospermum regium* (Mart. ex Schrank) Pilg., *Kielmeyera coriacea* Mart. & Zucc., *Couepia grandiflora* (Mart. & Zucc.) Benth., *Leptobalanus humilis* (Cham. & Schltdl.) Sothers & Prance and *Diospyros lasiocalyx* (Mart.) B.Walln., encountered in the HRCB collection, are old specimens that were not collected in the floristic inventory.

Images taken by drones (Fig. 8) allowed the collector on the ground to have another angle of observation. However, during the period when the photographs were taken, only a predominance of vines in the families Malpighiaceae and Bignoniaceae, as well as crowns without apparent flowers or fruits, were observed. Most of the collections were made on the edge of the Reserve, where there is more sunlight. Many shrubs and treelets are notable on the edge, as well as trees of *Anadenanthera peregrina* (L.) Speg. and *Copaifera langsdorffii* (Fig. 9), which are generalists and common in the Reserve (Pinheiro et al. 2021). In the interior of the fragment, many individuals also found along the border were observed, for example, *Pera glabrata*, *Ocotea pulchella* (Nees & Mart.) Mez, *Amaioua intermedia* Mart. and *Virola sebifera* Aubl.

This study highlights the relevance of floristic surveys in the field and of herbarium material as sources of biological information, as well as the importance of documenting and publishing analysed vouchers. Although previous studies have been conducted in the Reserve, it was possible to collect species that have not been listed or lack testimony material, which reinforces the importance of floristic studies in already known areas. The



influence of the lack of disturbances, such as fire, is noticeable, based on the change in the physiognomy and flora over the years.

In addition to the floristic diversity, the Reserve has a fauna comprised of various birds, for example, *Cariama cristata* (seriema), *Mimus saturninus* (sabiá-do-campo), *Taraba major* (choró-boi) and *Caracara plancus* (carcará), as well as reptiles, arthropods, cervids and other small mammals (Santos, personal observation). The Prof. Dr. Karl Arens Reserve maintains its importance as an area used for research and to preserve species and it is one of the few *Cerrado* remnants in the State of São Paulo. An updated floristic list, now with the respective vouchers, is also relevant regarding academic training because there are plant biology field classes taught to undergraduates in the Reserve.



Figure 8. [doi](#)

Aerial images from the Prof. Dr. Karl Arens Reserve: (a) May 2021; (b) December 2019; (c) July 2019. Photos: Bruno D. Borges (a–c).




Figure 9. [doi](#)

Some of the more common species from the Prof. Dr. Karl Arens Reserve: (a–b) *Anadenanthera peregrina*; (c–d) *Miconia rubiginosa*; (e–f) *Pera glabrata*; (g–h) *Xylopia aromatica*; (i–j) *Virola sebifera*; (k–l) *Ocotea pulchella*; (m–n) *Amaioua intermedia*; (o–p) *Qualea grandiflora*. Photos: Lilian S. Santos (a–p).

## Acknowledgements

We are grateful to everyone who helped with the fieldwork, especially to Aline Sayuri Tajima, Luis Antonio Letizio and Lucas Pellegrini Elias during the less severe period of the pandemic. Special thanks to Monica Midori Suemitsu for her assistance in the herbarium and to Prof. Alessandra Ike Coan for kindly providing the photographs. We also thank the Coordination for the Improvement of Higher Education Personnel (Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - CAPES) for the

scholarship awarded to the first author during the development of her Masters dissertation.

## Author contributions

Lilian Silva Santos: Conceptualisation, Fieldwork, Data Collection, Data Analysis, Taxa Identification, Writing - Original Draft, Review and Editing.

Pedro Luís Rodrigues de Moraes: Conceptualisation, Supervision, Fieldwork, Data Collection, Taxa Identification, Translation and Manuscript Review.

## References

- Ab'Sáber AN (1983) O domínio dos cerrados: introdução ao conhecimento. *Revista do Serviço Público (RSP)* 40 (4): 41-55. <https://doi.org/10.21874/rsp.v40i4.2144>
- ANA (2021) Agência Nacional de Águas. Portal Hidroweb: Séries Históricas de Estações. <https://snirh.gov.br/hidroweb/serieshistoricas>. Accessed on: 2021-2-01.
- Andena SR, Bego LR, Mechi MR (2005) A comunidade de abelhas (Hymenoptera, Apoidea) de uma área de cerrado (Corumbataí, SP) e suas visitas às flores. *Revista Brasileira de Zoociências* 7 (1): 47-54.
- APG IV (2016) An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG IV. *Botanical journal of the Linnean Society* 181 (1): 1-20. <https://doi.org/10.1111/boj.12385>
- Baker TR, Pennington RT, Dexter KG, Fine PVA, Fortune-Hopkins H, Honorio EN, Huamantupa-Chuquimaco I, Klitgård BB, Lewis G, Lima HC, Ashton P, Baraloto C, Davies S, Donoghue MJ, Kaye M, Kress WJ, Lehmann CER, Monteagudo A, Phillips OL, Vasquez R (2017) Maximising synergy among tropical plant systematists, ecologists, and evolutionary biologists. *Trends in Ecology & Evolution* 32 (4): 258-267. <https://doi.org/10.1016/j.tree.2017.01.007>
- Batalha MA (2011) O cerrado não é um bioma. *Biota Neotropica* 11 (1): 21-24. <https://doi.org/10.1590/S1676-06032011000100001>
- Camargo PN, Arens K (1967) Observações sobre uma reserva de cerrado. *Rev Agricultura* 42: 3-9.
- Campos MJO (1989) Estudo das interações entre comunidade de Apoidea, na procura de recursos alimentares, e a vegetação de cerrado da Reserva de Corumbataí-SP. Thesis (PhD in Ecology). Federal University of São Carlos, SP, São Carlos, 114 pp.
- Cassavan O (2002) O cerrado do Estado de São Paulo. In: Klein AL (Ed.) *Eugen Warming e o cerrado brasileiro: um século depois*. Ed. da UNESP: Imprensa Oficial do Estado, São Paulo, 156 pp. [ISBN 8571393540].
- Cesar O, Pagano SN, Leitão Filho HF, Monteiro R, Silva OA, Marinis G, Shepherd GJ (1988) Estrutura fitossociológica do estrato arbóreo de uma área de vegetação de cerrado no município de Corumbataí (Estado de São Paulo). *Naturalia* 13: 91-101.
- CIIAGRO (2023) Centro integrado de informações agrometeorológicas. Portal Agrometeorológico e Hidrológico do Estado de São Paulo. <http://ciiagro.org.br/diario/cperiodo>. Accessed on: 2023-12-12.

- Coutinho LM (2002) O bioma cerrado. In: Klein AL (Ed.) Eugen Warming e o cerrado brasileiro: um século depois. Ed. da UNESP: Imprensa Oficial do Estado, São Paulo, 77-91 pp. [ISBN 8571393540].
- Durigan G, Franco GADC, Siqueira MF (2004a) A vegetação dos remanescentes de cerrado no Estado de São Paulo. In: Bitencourt MD, Mendonça RR (Eds) Viabilidade de conservação dos remanescentes de cerrado no estado de São Paulo. Annablume: FAPESP, São Paulo, 29-56 pp. [ISBN 8574194751].
- Durigan G, Baitello JB, Franco GADC, Siqueira MF (2004b) Plantas do cerrado paulista: imagens de uma paisagem ameaçada. Páginas & Letras, São Paulo, 475 pp. [ISBN 8586508322]
- Durigan G, Ratter JA (2006) Successional changes in cerrado and cerrado/forest ecotonal vegetation in western São Paulo State, Brazil, 1962–2000. *Edinburgh Journal of Botany* 63 (1): 119-130. <https://doi.org/10.1017/S0960428606000357>
- Durigan G (2020) Zero-fire: Not possible nor desirable in the Cerrado of Brazil. *Flora* 268: 151612. <https://doi.org/10.1016/j.flora.2020.151612>
- Eiten G (1972) The cerrado vegetation of Brazil. *The Botanical Review* 38 (2): 201-341. <https://doi.org/10.1007/BF02859158>
- Eiten G (1994) Vegetação do cerrado. In: Pinto MN (Ed.) Cerrado: caracterização, ocupação e perspectivas. 2. ed. Editora Universidade de Brasília, Brasília, 681 pp. [ISBN 8523003444].
- Fidalgo O, Bononi VL (1989) Técnicas de coleta, preservação e herborização de material botânico. Instituto de Botânica, São Paulo, 62 pp. [ISBN 8585131071]
- Filgueiras TS, Nogueira PE, Brochado AL, Guala GF (1994) Caminhamento: um método expedito para levantamentos florísticos qualitativos. *Cadernos de geociências* 12 (1): 39-43.
- Flora e Funga do Brasil (2024) Jardim Botânico do Rio de Janeiro. <http://floradobrasil.jbrj.gov.br>. Accessed on: 2024-4-03.
- Font Quer P (1953) Diccionario de botánica. Labor, Barcelona, 1244 pp.
- Françoso RD, Brandão R, Nogueira CC, Salmona YB, Machado RB, Colli GR (2015) Habitat loss and the effectiveness of protected areas in the Cerrado Biodiversity Hotspot. *Natureza & conservação* 13 (1): 35-40. <https://doi.org/10.1016/j.ncon.2015.04.001>
- Funk VA, Hoch PC, Prather LA, Wagner WL (2005) The importance of vouchers. *Taxon* 54 (1): 127-129. <https://doi.org/10.2307/25065309>
- Gonçalves EG, Lorenzi H (2011) Morfologia vegetal: organografia e dicionário ilustrado de morfologia das plantas vasculares. 2. ed. Instituto Plantarum de Estudos da Flora, São Paulo. [ISBN 9788586714382]
- Harris JG, Harris MW (1994) Plant identification terminology: an illustrated glossary. Spring Lake Publishing, Spring Lake, 197 pp. [ISBN 096402215X]
- Hickey M, King C (2000) The Cambridge illustrated glossary of botanical terms. Cambridge University Press, Cambridge, 222 pp. [ISBN 0521794013]
- IBGE (2023) Instituto Brasileiro de Geografia e Estatística. Banco de Dados e Informações Ambientais (BDiA). <https://bdiaweb.ibge.gov.br/#/home>. Accessed on: 2023-12-12.
- IPNI (2024) International Plant Names Index. The Royal Botanic Gardens, Kew, Harvard University Herbaria & Libraries and Australian National Herbarium. <https://ipni.org>. Accessed on: 2024-4-03.



- Marques e Amorim G, Ebert HD, Horst R (2005) Integração de informações geológicas para o geoturismo na Bacia do Rio Corumbataí e sua divulgação na Web através do Mapserv. *Geociências* 24 (3): 221-238.
- MNHN (2023) Muséum national d'Histoire naturelle. The vascular plants collection (P) at the Herbarium of the Muséum national d'Histoire Naturelle. <https://science.mnhn.fr/institution/mnhn/search>. Accessed on: 2023-12-12.
- Myers N, Mittermeier RA, Mittermeier CG, Fonseca GAB, Kent J (2000) Biodiversity hotspots for conservation priorities. *Nature* 403 (6772): 853-858. <https://doi.org/10.1038/35002501>
- Nalon MA, Matsukuma CK, Pavão M, Ivanauskas NM, Kanashiro MM (2022) Inventário da cobertura vegetal nativa do Estado de São Paulo. SIMA/IPA, São Paulo, 238 pp. [ISBN 9786599641725]
- Pagano SN, Cesar O, Leitão Filho HF (1989) Estrutura fitossociológica do estrato arbustivo-arbóreo da vegetação de cerrado da Área de Proteção Ambiental (APA) de Corumbataí-Estado de São Paulo. *Revista brasileira de biologia* 49 (1): 49-59.
- Paula CC, Monteiro R (2000) The genus *Tillandsia* Lindley (Bromeliaceae) in a "cerrado" reserve (savanna) in Corumbataí, state of São Paulo, Brazil. *Naturalia* 25: 127-138.
- Peel MC, Finlayson BL, McMahon TA (2007) Updated world map of the Köppen-Geiger climate classification. *Hydrology and earth system sciences* 11 (5): 1633-1644. <https://doi.org/10.5194/hess-11-1633-2007>
- Piccolo ALG, Thamazini L, Massa CS, Cesar O, Pagano SN, Moraes JAPV, Amaral H (1971) Aspecto fitossociológico de uma reserva de cerrado. *Revista de Agricultura* 46: 81-92.
- Pinheiro ES, Durigan G (2012) Diferenças florísticas e estruturais entre fitofisionomias do cerrado em Assis, SP, Brasil. *Revista Árvore* 36 (1): 181-193. <https://doi.org/10.1590/S0100-67622012000100019>
- Pinheiro MHO, Azevedo TS, Monteiro R (2010) Spatial-temporal distribution of fire-protected savanna physiognomies in Southeastern Brazil. *Anais da Academia Brasileira de Ciências* 82 (2): 379-395. <https://doi.org/10.1590/S0001-37652010000200014>
- Pinheiro MHO (2013) Fenologia reprodutiva de comunidade savânica através de acervo de herbário. *Brazilian Geographical Journal: Geosciences and Humanities research medium* 4 (1): 233-254.
- Pinheiro MHO, Azevedo TS, Ferreira FL, Monteiro R (2021) Vegetation patterns and the influence of rainfall after long-term fire suppression on a woody community of a Brazilian savanna. *Anais da Academia Brasileira de Ciências* 93 (1): e20191405. <https://doi.org/10.1590/0001-3765202120191405>
- Ribeiro JF, Walter BMT (2008) Fitofisionomias do bioma cerrado. In: Sano SM, Almeida SP, Ribeiro JF (Eds) *Cerrado: ecologia e flora*. 2. v. Embrapa Informação Tecnológica, Brasília, 151-199 pp. [ISBN 9788573833973].
- Romão GO, Souza VC (2016) *Senna* P. Miller. In: Wanderley MGL, Shepherd GJ, Melhem TS, Giulietti AM, Martins SE (Eds) *Flora Fanerogâmica do Estado de São Paulo*. v. 8. Instituto de Botânica, São Paulo, 51-62 pp. [ISBN 8575230514].
- Rossi M (2017) Mapa pedológico do Estado de São Paulo: revisado e ampliado. Instituto Florestal, São Paulo, 118 pp. [ISBN 9788564808171]
- Sano EE, Rodrigues AA, Martins ES, Bettioli GM, Bustamante MMC, Bezerra AS, Couto Jr AF, Vasconcelos V, Schüller J, Bolfe EL (2019) Cerrado ecoregions: A spatial framework to assess and prioritize Brazilian savanna environmental diversity for



conservation. *Journal of environmental management* 232: 818-828. <https://doi.org/10.1016/j.jenvman.2018.11.108>

- Schilthuizen M, Vairappan CS, Slade EM, Mann DJ, Miller JA (2015) Specimens as primary data: museums and 'open science'. *Trends in ecology & evolution* 30 (5): 237-238. <https://doi.org/10.1016/j.tree.2015.03.002>
- Scholes RJ, Archer SR (1997) Tree-grass interactions in savannas. *Annual Review of Ecology and Systematics* 28 (1): 517-544. <https://doi.org/10.1146/annurev.ecolsys.28.1.517>
- Souza VC, Flores TB, Colletta GD, Coelho RLG, Mattos MHDC, Mattos FDC (2018) Guia das plantas do cerrado. Taxon Brasil, Piracicaba, 583 pp. [ISBN 9788554312008]
- speciesLink (2024) Simple search. Centro de Referência em Informação Ambiental, CRIA. <https://specieslink.net/search/>. Accessed on: 2024-4-03.
- Strassburg BBN, Brooks T, Feltran-Barbieri R, Iribarrem A, Crouzeilles R, Loyola R, Latawiec AE, Oliveira Filho FJB, Scaramuzza CAM, Scarano FR, Soares-Filho B, Balmford A (2017) Moment of truth for the Cerrado hotspot. *Nature Ecology & Evolution* 1: 0099. <https://doi.org/10.1038/s41559-017-0099>
- Thiers [continuously updated] (2024) Index Herbariorum: a global directory of public herbaria and associated staff. New York Botanical Garden's Virtual Herbarium. <http://sweetgum.nybg.org/science/ih/>. Accessed on: 2024-4-02.
- Troppmair H (2000) Geossistemas e geossistemas paulistas. UNESP, Instituto de Geociências e Ciências Exatas, Departamento de Geografia, Rio Claro, 105 pp.

## Supplementary material

### Suppl. material 1: List of the examined vouchers from Dr. Karl Arens Reserve, Corumbataí (SP) [doi](#)

**Authors:** Lilian Silva Santos & Pedro Luís Rodrigues de Moraes

**Data type:** Taxa

**Brief description:** List of the examined vouchers from Dr. Karl Arens Reserve, Corumbataí (SP), deposited in the Herbário Rioclarense (HRCB) of the Universidade Estadual Paulista (UNESP), Câmpus de Rio Claro.

[Download file](#) (335.38 kb)