



# First discovery of the genus *Enicopus* Stephens (Coleoptera, Dasytidae) from China

Jialin Miao<sup>‡</sup>, Haoyu Liu<sup>‡§</sup>, Junbo Tong<sup>‡</sup>, Xingke Yang<sup>‡</sup>, Yuxia Yang<sup>‡§</sup>

<sup>‡</sup> Key Laboratory of Zoological Systematics and Application, School of Life Sciences, Hebei University, Baoding, China

<sup>§</sup> Hebei Basic Science Center for Biotic Interaction, Hebei University, Baoding, China

| Key Laboratory of Zoological Systematics and Evolution, Institute of Zoology, Chinese Academy of Sciences, Beijing 100101, China, Beijing, China

Corresponding author: Haoyu Liu ([liuhy@hbu.edu.cn](mailto:liuhy@hbu.edu.cn)), Yuxia Yang ([yuxia0305@126.com](mailto:yuxia0305@126.com))

Academic editor: Cheng-Bin Wang

Received: 12 Dec 2024 | Accepted: 22 Jan 2025 | Published: 28 Jan 2025

Citation: Miao J, Liu H, Tong J, Yang X, Yang Y (2025) First discovery of the genus *Enicopus* Stephens (Coleoptera, Dasytidae) from China. Biodiversity Data Journal 13: e144281.

<https://doi.org/10.3897/BDJ.13.e144281>

## Abstract

## Background

*Enicopus* Stephens, 1830, is classified within the subfamily Dasytinae of the family Dasytidae (Coleoptera, Cleroidea). Currently, it comprises two subgenera and 27 known species. This genus primarily inhabits south-central Europe, with certain species extending into west Asia; however, it has not been reported in China until now.

## New information

The genus *Enicopus* Stephens, 1830 is reported for the first time from China, following the discovery of *E. (Enicopus) ater* (Fabricius, 1787) in Xinjiang Autonomous Region. This species is thoroughly re-described and illustrated, including detailed depictions of the ultimate abdominal tergites and sternites, genitalia of both sexes, as well as tarsomeres 1 from the right front and hind legs of male.

## Keywords

New faunistic record, alpha taxonomy, China, Cleroidea, Dasytidae

## Introduction

The genus *Enicopus* was first proposed by Stephens 1830, with *Lagria atra* Fabricius, 1787 designated as the type species. This genus is classified within the subfamily Dasytinae of the beetle family Dasytidae (Bahillo de la Puebla and López-Colón 2004). Adult specimens can be readily distinguished from other dasytid beetles by their unique tarsal appendages found in males (Liberti and Constantin 2009). *Enicopus* comprises two subgenera (Bahillo de la Puebla and López-Colón 2020), namely *E. (Enicopus)* Stephens, 1830 and *E. (Parahenicopus)* Portevin, 1931. These subgenera are differentiated, based on the characteristics of male tarsi. Specifically, tarsomeres 1 exhibit appendages on both front and hind tarsi in *E. (Enicopus)*, whereas such appendages are present only on the hind tarsi in *E. (Parahenicopus)*.

So far, it has been reported that there are 27 species (Bahillo de la Puebla and López-Colón 2020). Most of the species are distributed in south-central Europe, with one species spreading to Central Asia (Liberti and Constantin 2009). Amongst them, *E. (Enicopus) ater* (Fabricius, 1787) is the most widely distributed, extending from Western Europe to Central Asia (Liberti and Constantin 2009). It has been predicted to spread eastwards throughout Mongolia (Liberti and Constantin 2009), but has never been reported in China.

In this study, some materials of *E. (Enicopus) ater* were discovered in Xinjiang Autonomous Region, northwest China. They have been deposited in the Institute of Zoology, Chinese Academy of Sciences, Beijing since 1960. Due to the insufficient study of this beetle group in the Chinese fauna, they have been overlooked for a long time. Therefore, it is necessary to re-describe and illustrate it here to enhance its recognition and also enrich the beetle diversity of China.

## Materials and methods

In this study, we adhere to the conventional taxonomic classification of dasytid beetles as a separate family, Dasytidae (Majer 1994, Majer 2002, Mayor 2007, Bocakova et al. 2012, Constantin 2021), rather than regarding them as a subfamily within Melyridae (Constantin and Bahillo de la Puebla 2019, Gimmel and Mayor 2019, Gimmel and Mayor 2024). This study primarily focuses on the species descriptions of *Enicopus*, despite the ongoing debates regarding its higher classification, which falls outside the scope of this research. The specimens examined in this study are deposited in the Institute of Zoology, Chinese Academy of Sciences, Beijing, China (IZAS).

The specimens were initially soaked in water for softening, followed by the separation of their abdomens. The separated abdomens were then immersed in a 10% sodium hydroxide (NaOH) solution and heated at a constant temperature for several minutes using a metal bath. Once the fat had dissolved, they were transferred to a Nikon SMZ1500 stereomicroscope for the dissection of the pygidium, ultimate abdominal ventrite and genitalia. To facilitate observation, the spiculum gastrale, tegmen and median lobe were isolated respectively. The ovipositor was stained with haematoxylin. Subsequently, the dissected genitalia were placed on a glass slide with glycerol and photographed using a Leica M205A stereomicroscope before being stored in glycerol for preservation. A Canon EOS 80D digital camera was used to capture images of habitus which were later processed using Helicon Focus 7 software. Adobe Photoshop CC 2019 version 20.0.4 was utilised for editing in plate preparation. The body length was measured from the anterior margin of the head to the elytral apices and the width at the humeri. The terminology of genital segments follows Lawrence et al. (2010) and that of genitalia follows Gimmel and Mayor (2019).

## Taxon treatments

### *Enicopus* Stephens, 1830

#### Nomenclature

*Enicopus* (*Enicopus*) Stephens, 1830 - Stephens 1830: 318.

*Enicopus* (*Parahenicopus*) Portevin, 1931 - Portevin 1931: 453.

#### Type species

*Lagria ater* Fabricius, 1787

#### Diagnosis

Body black and covered with long hairs. Pronotum with two sinuous lateral grooves on disc. Front tibiae terminating in a short hooked appendage, tarsi with at least one pair of spines or appendages on tarsomeres 1 in males, while simple in female, all tarsal claws fitted with a membrane almost as long as the claws themselves. Tegmen with two well-developed and symmetrical lobes at apex, densely covered with long hairs; internal sac of median lobe with more than two pairs of large spines at apical part.

#### Distribution

China (new record: Xinjiang), Spain, Portugal, Albania, Algeria, Armenia, Austria, Azerbaijan, Bosnia-Herzegovina, Bulgaria, Caucasus, Croatia, Czechia, France, Germany, Georgia, Greece, Hungary, Iran, Italy, Kazakhstan, Kyrgyzstan, Macedonia,

Mongolia, Montenegro, Romania, Russia, Serbia, Slovakia, Sweden, Switzerland, Turkey, Ukraine, Uzbekistan.

## ***Enicopus (Enicopus) ater* (Fabricius, 1787)**

### **Nomenclature**

*Lagria atra* Fabricius, 1787 - Fabricius 1787: 94.

*Melyris ater* (Fabricius, 1787) - Olivier 1790: 9; Olivier 1808: plate 2, figs. 8a-8e.

*Dasytes ater* (Fabricius, 1787) - Germar 1817: 209; Stephens 1829: 136; Küster 1849: 15.

*Enicopus ater* (Fabricius, 1787) - Stephens 1830: 318; Stephens 1833: 46; Stephens 1839: 195; Spry and Shuckard 1840: 75; Mayor 2007: 407 (syn. of *E. pilosus* Scopoli, 1763); Liberti and Constantin 2009: 298.

### **Material**

- a. country: China; stateProvince: Xinjiang; county: Qinghe; verbatimElevation: 1450 m; year: 1960; month: 7; day: 3; individualCount: 8; sex: 4 males, 4 females; lifeStage: adult; recordedBy: Shuyong Wang; institutionID: Institute of Zoology, Chinese Academy of Sciences; institutionCode: IZAS; occurrenceID: 88CA14BD-017B-5FDF-8245-6F0B22E842FE

### **Description**

**Male** (Fig. 1A). Body length 7.4–9.4 mm, width 2.3–2.9 mm.

Body black with lustre, except for antennomeres 3 and 4 yellow at apices. Body densely and coarsely punctate on surface, as well as covered with short and recumbent black pubescence, grey on antennomeres 3–11, tibiae and tarsi. Vertex behind eyes densely covered with long erect black hairs. Antennomeres 1 and 2 covered with a few erect black hairs. Pronotum densely covered with long erect black hairs around all margins, of which that on lateral margins longer than those on anterior and posterior margins. Elytra covered with long erect black hairs along margins, which are progressively shortened towards apices.

Head width across eyes narrower than anterior margin of pronotum, present with a pair of shallow depressions on vertex. Antennae quite short, extending to posterior margin of pronotum when inclined, antennomeres 1 nearly ellipsoid, 1.5 times as long as wide, 2 short, apical half expanded, 3–10 triangular with truncate apex, 11 elongate and fusiform, twice as long as wide. Ultimate maxillary palpomere fusiform, about 1.7 times longer than the penultimate one.



Figure 1. [doi](#)

Habitus of *Enicopus (Enicopus) ater* (Fabricius, 1787), dorsal view: **A** male; **B** female. Scale bars: 1.0 mm.

Pronotum transverse and 1.3 times as wide as long, widest near middle, anterior margin slightly arcuate, lateral margins slightly diverging posteriorly and regularly rounded with distinct edges, posterior margin slightly bisinuate.

Elytra elongate and parallel-sided, 2.1–2.3 times longer than wide at humeri, 2.3–2.5 times longer than pronotum, rounded at humeri, acute at apices.

Front femora slightly swollen, each tibia with a pair of small and sharply-hooked spurs at apex, tarsomeres 1 with a large and sharply hooked appendage on outer side, which bear a large gable-shaped tooth on inner base (Fig. 2A–B); middle tarsomeres 1 short, with a pair of small sharp teeth; hind femora slightly swollen, tibiae slightly swollen, distinctly curved, tarsomeres 1 with a large spoon-shaped appendage on outer base, which is bent inwards and slightly wider at apical part than basal part, external angle acute and pointing upwards, tarsomeres 2 very long and twice longer than combined length of tarsomeres 3–5 and terminating into a small tooth at apex; tarsal claws symmetrical, with a pair of broad membranous appendages, which slightly shorter than the claws (Fig. 2C–E).

Ultimate abdominal ventrite (Fig. 3A) saddle-form, narrowed posteriorly, 1.2–1.3 times as wide as long, shallowly and trapezoidally emarginate in middle of posterior margin, present with a slender central process at anterior margin, which does not extend over antero-lateral angles, surface densely covered with long black hairs along lateral margins. Pygidium (Fig. 3B) shield-form, 1.0–1.1 times longer than wide, feebly narrowed posteriorly, hardly emarginate in middle of posterior margin, largely

and triangularly protuberant in middle of anterior margin, with antero-lateral angles obviously protruding, surface covered with long black hairs in centre and along lateral margins. Aedeagus: tegmen (Fig. 3C and D) nearly bullet-shaped, present with a pair of symmetrical and nearly semicircular lobes at apex, wider than long, where it is densely covered with long hairs; median lobe strongly bent ventrally, dorso-ventrally flattened apically in lateral view (Fig. 3E), with apex slightly hooked ventrally, slightly elongated at apex in ventral view (Fig. 3F); internal sac (Fig. 3H) long and membranous, fitted with four types of spines: the first type (I) located at base, a dozen or so, yellow and slender, arranged in parallel, the second type (II) dozens, brown, short and nearly triangular, the third type (III) dozens, black and slender and the fourth type (IV) located at the apical part, a dozen or so, black, stout and long. Spiculum gastrale Y-shaped (Fig. 3G).



Figure 2. [doi](#)

*Enicopus (Enicopus) ater* (Fabricius, 1787), male: **A** front tarsi, ventral view; **B** front tarsi, dorsal view; **C** hind tarsi, dorsal view; **D** hind tarsi, dorso-lateral view; **E** hind tarsi, lateral view. Scale bars: 1.0 mm.

**Female** (Fig. 1B). Similar to male, but body larger, length 8.7–9.6 mm, width 3.1–3.4 mm. Eyes less prominent. Front tarsomeres 1 without any appendage; middle tarsi simple; hind tibiae normally slender, straight, tarsomeres 1 without any appendage, tarsomeres 2 much shorter than that of male. Ultimate abdominal ventrite (Fig. 4A) trapezoidal, strongly narrowed posteriorly, posterior margin nearly truncate, present with a very long central process at anterior margin, which distinctly extends over postero-lateral angles. Pygidium (Fig. 4B) strongly narrowed posteriorly, semicircularly emarginate in middle of posterior margin, arcuate at anterior margin, antero-lateral angles obviously protruding. Ovipositor (Fig. 4C) slender and membranous, gonostylus feebly long and nearly cylindrical.

### Diagnosis

This species can be distinguished from all other species of *E. (Enicopus)* by the characteristics of tarsal appendages and aedeagus. It looks more similar to *E.*

(*Enicopus pilosus* (Scopoli, 1763), but can be distinguished by the combination of the following characters: appendage on hind tarsomeres 1 of male flattened and spoon-shaped, with the external angle pointing upwards; median lobe strongly curved ventrally in lateral view; tegmen with a pair of nearly semicircular lobes at apex, which are separated and feebly wider than long. Unlike in *E. pilosus*, appendage on hind tarsomeres 1 of male is sickle-shaped, with apical part bent upwards (Liberti and Constantin 2009: fig. 23); median lobe is slightly curved dorsally in lateral view (Liberti and Constantin 2009: fig. 8); tegmen bears a pair of slender and parallel lobes at apex, which are much longer than wide (Liberti and Constantin 2009: fig. 12).



Figure 3. [doi](#)

*Enicopus* (*Enicopus*) *ater* (Fabricius, 1787), male: **A** ultimate abdominal ventrite (apical sternite), ventral view; **B** pygidium (apical tergite), dorsal view; **C** tegmen, ventral view; **D** tegmen, lateral view; **E** median lobe, lateral view; **F** median lobe, ventral view; **G** spiculum gastrale, ventral view; **H** internal sac; **I-IV** the four types of spines covered on the internal sac. Scale bars: 0.5 mm.

### Distribution

China (new record: Xinjiang), Albania, Armenia, Austria, Azerbaijan, Bosnia-Herzegovina, Bulgaria, Caucasus, Croatia, southern France, Georgia, Greece, Hungary, Iran, Italy, Kazakhstan, Kyrgyzstan, Macedonia, Montenegro, Romania, Russia (incl. South Siberia), Serbia, Turkey, Ukraine, Uzbekistan.

## Notes

Liberti and Constantin (2009) presented a detailed list of all synonyms for this species, which are omitted here. Meanwhile, they depicted the internal sac of the median lobe with 3-5 spines at the apex (Liberti and Constantin 2009: fig. 11), but there are many more shown in our material (Fig. 3H). Later, they provided an illustration of the adult male, whose hind tarsomer 2 is slightly longer than the combined length of the last three tarsomer (Constantin and Liberti 2011: fig. 27), while it is much longer in our material (Fig. 1A). Nevertheless, we identify them as *E. (Enicopus) ater* because more or less variations can occur in such a widespread species.



Figure 4. [doi](#)

*Enicopus (Enicopus) ater* (Fabricius, 1787), female: **A** ultimate abdominal ventrite (apical sternite), ventral view; **B** pygidium (apical tergite), dorsal view; **C** ovipositor, ventral view. Scale bars: 0.5 mm.

## Acknowledgements

This study is financially supported by the National Natural Science Foundation of China (No. 32270491), the Natural Science Foundation of Hebei Province (No. C2022201005) and the Interdisciplinary Research Program of Natural Science of Hebei University (No. DXK202302).

## References

- Bahillo de la Puebla P, López-Colón J (2004) *Enicopus sarae*, nueva especie de *Enicopus* Stephens, 1830 de España (Coleoptera, Dasytidae). Boletín de la Sociedad Entomológica Aragonesa 34: 55-58. URL: [www.researchgate.net/journal/Boletin-de-la-](http://www.researchgate.net/journal/Boletin-de-la-)



[Sociedad-Entomologica-Aragonesa-1134-6094?\\_tp=eyJjb250ZXh0Ijp7ImZpcnN0UGFnZSI6InB1YmxpY2F0aW9uIiwicGFnZSI6InB1YmxpY2F0aW9uIn19](https://doi.org/10.1111/j.1096-0031.2011.00368.x)

- Bahillo de la Puebla P, López-Colón J (2020) Sobre el estatus taxonómico del subgénero *Dasysoma* Schilsky, 1896 (Coleoptera: Cleroidea: Melyridae: Dasytinae). *Heteropterus Revista de Entomología* 20 (2): 167-180.
- Bocakova M, Constantin R, Bocak L (2012) Molecular phylogenetics of the melyrid lineage (Coleoptera: Cleroidea). *Cladistics* 28 (2): 117-129. <https://doi.org/10.1111/j.1096-0031.2011.00368.x>
- Constantin R, Liberti G (2011) Coléoptères Dasytidae de France. Musée des Confluences, France, 144 pp.
- Constantin R, Bahillo de la Puebla P (2019) *Enicopus* (s. str.) tempranillo nov. sp. from southern Spain (Coleoptera: Melyridae: Dasytinae). *Heteropterus Revista de Entomología* 19 (2): 285-293.
- Constantin R (2021) Malachiidae Fleming, 1821. In: Tronquet M (Ed.) *Catalogue des Coléoptères de France*. Association Roussillonnaise d'Entomologie, Perpignan, 487–492 pp.
- Fabricius JC (1787) *Mantissa Insectorum sistens eorum species nuper detectas. Adiectis characteribus genericis, differentiis specificis, emendationibus, observationibus*. Tome I - Impensis Christ. Gotti. Proft., Hafniae, 348 pp. <https://doi.org/10.5962/bhl.title.36471>
- Germar EF (1817) *Reise nach Dalmatien und in das Gebiet von Ragusa*. Brockhaus FA, Leipzig, 325 pp.
- Gimmel ML, Mayor AJ (2019) On the north American species of *Dolichosoma* Stephens and *Psilothrix* Küster (Coleoptera: Melyridae: Dasytinae: Dasytini). *The Coleopterists Bulletin* 73 (2): 395-407. <https://doi.org/10.1649/0010-065X-73.2.395>
- Gimmel ML, Mayor AJ (2024) Taxonomic Overview of World Dasytinae (Coleoptera: Melyridae), with Emphasis on the North American Fauna of the Subfamily and Two Newly Established Tribes. *The Coleopterists Bulletin* 78 (4): 499-536. <https://doi.org/10.1649/0010-065X-78.4.499>
- Küster HC (1849) *Die Käfer Europas: nach der Natur beschrieben*, 19es Heft. Verlag von Bauer und Raspe, Nurnberg, 100 pp.
- Lawrence JF, Beutel RG, Leschen RA, Ślipiński SA (2010) Chapter 2. Glossary of Morphological Terms. In: Leschen RA, Beutel RG, Lawrence JF (Eds) *Handbook of Zoology, Coleoptera. Volume 2. Morphology and Systematics (Elateroidea, Bostrichiformia, Cucujiformia partim)*. Walter de Gruyter, Berlin, 9-20 pp. <https://doi.org/10.1515/9783110911213.9>
- Liberti G, Constantin R (2009) The *Enicopus* Stephens, 1830 species East of the Iberian Peninsula: *E. ater* (Fabricius, 1787) and *E. pilosus* (Scopli, 1763) (Coleoptera, Dasytidae). *Annali del Museo Civico di Storia Naturale Giacomo Doria* 100: 293-321.
- Majer K (1994) A review of the classification of the Melyridae and related families (Coleoptera, Cleroidea). *Entomologica Basiliensia* 17: 319-390.
- Majer K (2002) Subfamilial classification of the family Malachiidae (Coleoptera, Cleroidea). *Entomologica Basiliensia* 24: 179-244.
- Mayor AJ (2007) Dasytidae. In: Löbl I, Smetana A (Eds) *Catalogue of Palaearctic Coleoptera, Vol. 4. Elateroidea - Derodontoidea - Bostrichoidea - Lymexyloidea - Cleroidea - Cucujoidea*. Apollo Books, Stenstrup, 388–415 pp.

- Olivier AG (1790) *Entomologie, ou histoire naturelle des insectes, avec, leur caractères génériques et spécifiques, leur descriptions, leur synonymie, et leur figures enluminée. Coléoptères. Tome second*, Paris: de Baudouin, 485 pp.
- Olivier AG (1808) *Entomologie, ou histoire naturelle des insectes, avec leurs caractères génériques et spécifiques, leur description, leur synonymie, et leur enluminée. Coléoptères. Tome sixième*, Paris: Desray, 611-1104 pp.
- Portevin G (1931) *Histoire Naturelle des Coléoptères de France. Tome II. Polyphaga: Lamellicornia, Palpicornia, Diversicornia. Encyclopédies entomologiques, séries A, XIII éditions Paul Lechevalier, Paris, 542 pp.*
- Spry W, Shuckard WE (1840) *The British Coleoptera delineated: consisting of figures of all the genera of British beetles*. In: Shuckard WE, Lib RS (Eds) [Family [51] Melyridae, Pp. 42-43, 75, plate 51, and Supplement plate 5]. W. Crofts, 83 pp. + 86 plates + 8 Supplement plates.
- Stephens JF (1829) *A systematic Catalogue of British Insects; being an attempt to arrange all the hitherto discovered indigenous insects in accordance with their natural affinities*. Baldwin and Cradock, London, 388 pp.
- Stephens JF (1830) *Illustrations of British Entomology; or, a synopsis of indigenous insects: containing their generic and specific distinctions, with an account of their metamorphoses, times of appearance, localities, food, and economy, as far as practicable. Mandibulata, 3*. Baldwin and Cradock, London, 374 pp.
- Stephens JF (1833) *The Nomenclature of British Insects*. Baldwin and Cradock, London, 134 pp.
- Stephens JF (1839) *A manual of British Coleoptera, or beetles, containing a brief description of all the species of beetles hitherto ascertained to inhabit Great Britain and Ireland, together with a notice of their chief localities, times and places of appearances, etc*. Longman, Orme Brown, Green, and Longmans, London, 443 pp.