



Checklist of pioneer benthic taxa found on Autonomous Reef Monitoring Structures (ARMS) in Terra Nova Bay (Ross Sea, Antarctica)

Valentina Cometti^{‡,§}, Matteo Cecchetto^{l,§}, Alice Guzzi^{l,§,¶}, Marco Grillo^{‡,§}, Nicholas Francesco Noli^{‡,§}, Simonetta Corsolini^{‡,#}, Stefano Schiaparelli^{l,§,¶}

[‡] Department of Physical, Earth and Environmental Sciences, University of Siena, Siena, Italy

[§] Italian National Antarctic Museum (MNA, section of Genoa), Genoa, Italy

^l Department of Earth, Environmental and Life Sciences (DISTAV), University of Genoa, Genoa, Italy

[¶] National Biodiversity Future Center (NBFC), Palermo, Italy

[#] Institute of Polar Sciences, Italian National Research Council (ISP-CNR), Bologna, Italy

Corresponding author: Valentina Cometti (v.cometti@student.unisi.it)

Academic editor: Anton P. van de Putte

Received: 05 Feb 2025 | Accepted: 03 Apr 2025 | Published: 08 Apr 2025

Citation: Cometti V, Cecchetto M, Guzzi A, Grillo M, Noli N, Corsolini S, Schiaparelli S (2025) Checklist of pioneer benthic taxa found on Autonomous Reef Monitoring Structures (ARMS) in Terra Nova Bay (Ross Sea, Antarctica). Biodiversity Data Journal 13: e148863. <https://doi.org/10.3897/BDJ.13.e148863>

Abstract

Background

Benthic communities studies in the Southern Ocean highlight their potential for assessing climate and anthropogenic impacts. However, the lack of standardised methods limits result reliability and interpretation. This dataset presents the first checklist focus on the Antarctic pioneer benthic communities collected using a standardised approach such as Autonomous Reef Monitoring Structures (ARMS) located at 25 m depth in the surroundings of the Italian research station "Mario Zucchelli" (MZS) in the Terra Nova Bay (TNB) area of the Ross Sea, Antarctica. The data encompass ARMS time series corresponding to deployments of 1, 2, 3 and 5 years, from which 277 occurrence data corresponding to 12 phyla, 43 families, 49 genera and 39 species were obtained. All retrieved specimens are curated by the Italian National Antarctic Museum (MNA, section of Genoa). This dataset is a contribution to the Antarctic Biodiversity Portal, the thematic

Antarctic node for both the Ocean Biogeographic Information System (AntOBIS) and the Global Biodiversity Information Facility Antarctic Biodiversity Information Facility ([ANTABIF](#)). The dataset was uploaded and integrated with the SCAR-AntOBIS database under the licence CC-BY 4.0. Please follow the guidelines from the [SCAR Data Policy](#) (ISSN 1998-0337) when using the data. If you have any questions regarding this dataset, please contact us via the contact information provided in the metadata or via data-biodiversity-aq@naturalsciences.be. Issues with the dataset can be reported at the [biodiversity-aq GitHub project](#).

New information

We describe the biodiversity of the Antarctic pioneer benthic communities of TNB sampled using the ARMS installed at the Italian research station "Mario Zucchelli". ARMS is a standardised, reproducible and comparable method for quantifying biodiversity. This dataset provides essential baseline data on the occurrence and abundance of pioneer benthic communities in this study area, representing an important contribution for understanding the dynamics of benthic pioneer communities in an area where these structures have never been deployed and, in general, for an exposure time that largely exceed the standard one, which is usually of one year only.

The 277 occurrences reported here have been classified at the lowest possible taxonomic level and comprise 39 recognised species, 49 genera and 43 families. Approximately 98% of the samples are stored in 96% ethanol, while the others at -20°C, representing a potential resource for future genetic studies. To date, the entire ARMS collection has not been DNA barcoded, although preliminary metabarcoding analyses have already been published in Cecchetto et al. (2024). Outcomes of the barcoding activity will be the target of another future publication (Cometti et al., in prep). The publication of this data paper was funded by the Belgian Science Policy Office (BELSPO, contract n°FR/36/AN1/AntaBIS) in the framework of EU-Lifewatch as a contribution to the SCAR Antarctic Biodiversity Portal ([bio diversity.aq](#)).

Keywords

distributional occurrences, check-list, time series, ARMS, Southern Ocean, Ross Sea, Terra Nova Bay, Italian National Antarctic Museum (MNA), biodiversity

Introduction

The Southern Ocean has unique environmental conditions compared to other areas of the Planet and is characterised by a high degree of endemism (De Broyer et al. 2014), high levels of cryptic diversity at the species level (Wilson et al. 2009, Maroni and Wilson 2022, Maroni et al. 2022) and a Palaeozoic functional grade of organisation (Aronson and Blake 2001). To date, the available data on the development and dynamics of

benthic communities have shown interesting potential for understanding the impacts of anthropogenic and climate change on them, showing changes at both community and species levels (Barnes et al. 2021).

Antarctic hard substrates fouling communities of Terra Nova Bay (TNB) were initially studied during the expeditions 1987-88, 1989-90 and 1993-94, conducted by the Italian National Antarctic Research Program (PNRA) (Cattaneo-Vietti et al. 2000). During the same period, other areas of the continent, such as McMurdo Sound (Ross Sea), were investigated (e.g. Dayton (1989)).

Since these initial studies, there have been no new attempts to study pioneer benthic communities until recently, when new investigations into the colonisation of artificial substrates (e.g. Bowden et al. (2006), Dayton et al. (2016), Caruso (2019), Barnes et al. (2021)) and natural substrates (e.g. Barnes and Souster (2011), Fillinger et al. (2013), Barnes et al. (2014), Krzeminska and Kuklinski (2018)) were undertaken. To date, the data show a very critical and slow growth rate (Peck 2018) and reveal the presence of various taxa including Porifera, Cnidaria, Annelida, Mollusca, Bryozoa, Brachiopoda, Chordata, Algae (Arntz and Gallardo 1994, Bowden et al. 2006) and bacterial communities (Caruso et al. (2023), Papale et al. (2024)).

However, these studies have never employed reproducible and standardised techniques, limiting both the reliability of the results and the broader understanding of observed changes. A standardised approach is essential to ensure the replicability of analyses, a key feature of biological monitoring at regional and global scales.

This feature is provided by the Autonomous Reef Monitoring Structures (ARMS), designed by Leray and Knowlton (2015). The ARMS units, part of the [Global ARMS Programme](#), consists of ten 22.5 × 22.5-cm PVC plates stacked on top of each other, simulating a complex 3D environment. Furthermore, these structures present alternating layers of crevices open and closed to the flow of the current. ARMS provide a quantitative, reproducible, standardised and cost-effective method that enables reliable comparisons across studies (Leray and Knowlton 2015, David et al. 2019, Pearman et al. 2020).

The simplified design of these structures provides an easily quantifiable sampling methodology that, combined with the study of fouling organisms with High Throughput Sequencing (HTS), became a standard in monitoring activities at sea (Valentini et al. 2016) allowing communities to be inspected in different time intervals and environmental conditions (Cecchetto et al. 2024). ARMS are a powerful tool to obtain information on the likely resilience of the benthic fauna in relation to possible changes and to significantly improve the accuracy and feasibility of monitoring efforts (Fonseca et al. 2017), especially in the Ross Sea, where they have never been adopted.

This dataset presents the first checklist of pioneer benthic organisms in TNB, which have never been studied using these structures and over such a long monitoring time. Previous MNA contributions focused on Mollusca, Tanaidacea, Fungi, Ophiuroidea,

Porifera, Bryozoa, Rotifera, Asteroidea, Copepoda and Isopoda (Ghiglione et al. 2013, Piazza et al. 2014, Selbmann et al. 2015, Cecchetto et al. 2017, Ghiglione et al. 2018, Garlasché et al. 2019, Cecchetto et al. 2019, Bonello et al. 2020, Guzzi et al. 2022, Grillo et al. 2024, Noli et al. 2024).

The special issue that included this publication contains additional articles that centre on specific marine animals, such as Holothurians (Guzzi et al., in prep), Amphipods (Cecchetto et al., in prep) and fish (La Mesa et al., in prep). This dataset also represents another Italian contribution to the CCAMLR CONSERVATION MEASURE 91-05 (2016) for the Ross Sea region Marine Protected Area, specifically addressing Annex 91-05/C (“long-term monitoring of benthic ecosystem functions”).

Project description

Title: Checklist of pioneer benthic taxa found on Autonomous Reef Monitoring Structures (ARMS) in Terra Nova Bay (Ross Sea, Antarctica)


Personnel: Valentina Cometti, Matteo Cecchetto, Alice Guzzi, Marco Grillo, Nicholas Francesco Noli, Simonetta Corsolini, Stefano Schiaparelli

Study area description: The occurrence data of the pioneer benthic communities studied in this data paper derives from the XXXII, XXXIII, XXXIV, XXXVII and XXXVIII Expeditions of the Italian National Antarctic Program (PNRA). Samples were collected from ARMS located at a depth of 25 m at the ‘Zecca’ site in Tethys Bay (-74.690°, 164.103°), approximately 500 m from the Mario Zucchelli Station (Fig. 1), in the TNB area. The sampling period spanned from 1 December 2015 to 8 November 2022. The seabed surrounding the ARMS consists of heterogeneous, unsorted sediments including sand, gravel and small cobbles mainly colonised by Corallinales. The area is characterised by a high abundance of *Sterechinus neumayeri* (Meissner, 1900) and *Odontaster validus* Koehler, 1906 (Piazza et al. 2019, Piazza et al. 2020), which were frequently observed at the same site during the retrieval of the structures.

Funding: Data originated in the framework of six different PNRA (Italian National Antarctic Program) expeditions carried out from 2015 to 2022. The deployment, recovery and analyses of the ARMS deployed in TNB were funded by the Italian National Antarctic Program (PNRA) projects:

- “TNB-CODE - Terra Nova Bay barCODing and mEtabarcoding of Antarctic organisms from marine and limno-terrestrial environments” (Project code 2016/AZ1.17; PI Prof. Schiaparelli S.).
- “RosS-MODE – Ross Sea biodiversity Monitoring through barcoding, metabarcODing and e-DNA” (Project code PNRA18_00078, PI Prof. Ficetola F.).

The publication of this data paper was funded by the Belgian Science Policy Office (BELSPO, contract n°FR/36/AN1/AntaBIS) in the Framework of EU-Lifewatch as a contribution to the [SCAR Antarctic biodiversity portal](#).



Sampling methods

Sampling description: Samples were collected using [ARMS](#) (Fig. 2). Each ARMS consists of 10 PVC plates (22.5 x 22.5 x 0.5 cm) stacked one top of each other and separated by 1 cm nylon spacers at the corners of each plate, into which four stainless steel bolts are threaded, holding the entire structure together. This is then fixed on top of a large 45 x 35 cm PVC base plate, which allows the whole structure to be anchored to the seafloor. ARMS have been recovered thanks to the help of PNRA SCUBA divers, which covered each retrieved structure with a rigid plastic crate perforated on each side and internally lined with a 100 μ m nylon net, in order to avoid the escape of vagile benthic organisms. Other details available at [Global ARMS Program](#) site.

ARMS were deployed at sea for varying time ranges of 1, 2, 3 and 5 years. These six structures were recovered in different years: 2 in 2016 (1 year later), 2 in 2017 (2 years later), 2 in 2018 (3 years later). Additionally, one pair of ARMS was installed in 2016 and recovered in 2021 (5 years later) and another additional pair of ARMS was installed in 2017 and recovered in 2022 (5 years later).

Quality control: All records were visually checked, identified at the lowest possible taxonomic level, validated and assigned an MNA voucher. Throughout all phases, quality control and data cleaning ensured high-quality data and reliable identifications. Throughout sorting, classification and storage at the MNA, quality control and data

cleaning ensured high-quality data and reliable identifications. Coordinates were converted into decimal latitude and decimal longitude and plotted to verify the geographical location and locality. All scientific names were inspected for typos and were updated using the “WoRMS Taxon match” tool of the “[World Register of Marine Species](#)” (WoRMS) and AphiaID was assigned to each taxon as scientificNameID. The event dates and times were converted into ISO 8601 and verified with the field reports.




Figure 2. [doi](#)

Photo extracted from underwater video showing an ARMS deployed in the sampling area (A). The structure was recovered during the XXXVII PNRA expedition in 2021 (after 5 years of colonisation). After recovery, photographs were taken of the PVC panels. In detail, we see panel number 7 facing downwards (B) and panel number 5 facing upwards (C).

Step description: The structures deployed in 2015 and recovered in 2016 (1 year), 2017 (2 years) and 2018 (3 years) were stored entirely at -20°C to be transported to Italy, where they were disassembled and analysed. After retrieval, each individual plate was appropriately photographed, from both sides, scraped and homogenised to form subsamples that were subsequently stored in ethanol at -20°C . However, during the recovery of the first pair of structures (2016) and one of the second (2017), these crates malfunctioned and, thus, no quantitative analyses on the vagile component of the community inhabiting the ARMS could be performed. For the fifth-year structures, one pair deployed in 2016 and recovered in 2021 and the other pair deployed in 2017 and recovered in 2022, processing took place directly in Antarctica. The samples were sorted, acquired by the Italian National Antarctic Museum (MNA, Section Genoa) and directly stored in ethanol (96%) or at -20°C to be identified later. Most of the records were identified by one researcher, using original descriptions and taxonomic keys and the online WoRMS portal to confirm the acceptance of species names. The identification was often supported by scanning electron microscopy (SEM), combining stack images of the

analysed specimens, particularly for the bryozoan specimens (Fig. 3). High-resolution SEM images were taken, focusing on diagnostic traits such as details of the primary orifice, ovicell morphology and other morphological characteristics. The light and contrast parameters were carefully adjusted to enhance specimen features, ensuring clarity for accurate identification. For bryozoan, a small portion of the colony of less than 1 cm was taken and treated with sodium hypochlorite (NaClO) for 10 minutes and washed with ethanol (EtOH) at different concentrations 70%, 90% and 100%. When identification was inconclusive, only genus or family names were assigned. The samples were deposited in the biological collection of the MNA. All data were uploaded to the [GBIF portal](#).




Figure 3. [doi](#)

Sample of *Camptoplites tricornis* (Waters, 1904) (MNA-13822 voucher number) on PVC plate (Fig. 3A) identified using scanning electron microscopy (SEM). In detail, avicularia (Fig. 2B) and autozooid with spines (Fig. 2C). The sample is part of the material collected during the XXXIV PNRA Expedition (2018/2019).

Geographic coverage

Description: Samples were collected at one location, nominally "Zecca", from the Tethys Bay area. The sampling site is approximately 500 m distant from the MZS in TNB (Ross Sea, Antarctica) and at 25 m of depth (Fig. 1).

Coordinates of the deployment site: -74.690 Latitude, 164.104 Longitude

Taxonomic coverage

Description: This dataset focuses on pioneer benthic taxa collected using ARMS. A total of 277 occurrences were recorded, with the largest proportions in 2022 (30.68%) and 2018 (29.97%), followed by 2021 (27.07%) and 2017 (9.75%). The smallest percentage was in 2016, accounting for only 2.53% of the total occurrences (Fig. 4).




Figure 4. [doi](#)

Occurrence percentages over the years from 2016 to 2022.

After analysing the complete dataset of 277 occurrences, 43 families, 49 genera and 39 species were identified, along with 31 morphotypes. However, some taxa could not be identified to the species level and this uncertainty was indicated by 'sp.' or 'cf.' identification qualifiers in the dataset (Fig. 5). Furthermore, six specimens from the order Amphipoda were excluded from this dataset, as they are already listed in the checklist by Cecchetto et al. (in prep).

Of the 277 occurrences, 154 are sessile and 123 are vagile species.

Annelida is the most abundant phylum (73 occurrences), followed by Bryozoa (70 occurrences). While Echinodermata (45 occurrences) and Mollusca (29 occurrences) are also notable, they have fewer records compared to the top two phyla. Other phyla, such as Chordata and Porifera, have significantly fewer records (respectively 11 and 5), with the number of records gradually decreasing towards the least represented phyla.

The most representative families in the dataset are Polynoidae (30 occurrences), Echinidae (23 occurrences) and Serpulidae (22 occurrences). Amongst these, the most abundant genera were *Harmothoe* Kinberg, 1856 (27 occurrences) (POLYCHAETA, Polynoidae) and *Sterechinus* Koehler, 1901 (23 occurrences) (ECHINOIDEA, Echinidae). The species with the highest number of occurrences was *Sterechinus neumayeri* (Meissner, 1900), recorded 22 times.

After analysing the life stages of the specimens, a total of 277 occurrences were recorded, comprising 272 adults, three juveniles (corresponding to *Lanicides bilobata*

(Grube, 1877) and two specimens of *Adamussium colbecki* (E.A. Smith, 1902) and two egg masses (*Neobuccinum eatoni* (E.A. Smith, 1875)).

Species with the symbol (*) in the following table indicate that they are new records for the TNB area. All the records recorded in this dataset are based on physical museum vouchers (hereafter “MNA collection records”) curated by the Genoa section of the MNA.




Figure 5. [doi](#)

Number of records identified at the phylum, family, genus and species level, for each specific taxa.

Taxa included:

Rank	Scientific Name
kingdom	Animalia
kingdom	Chromista
kingdom	Plantae
phylum	Annelida
phylum	Arthropoda
phylum	Bryozoa
phylum	Cercozoa
phylum	Chordata
phylum	Cnidaria
phylum	Echinodermata
phylum	Heterokontophyta
phylum	Mollusca
phylum	Nemertea
phylum	Porifera
phylum	Rhodophyta
class	Ascidiacea
class	Asteroidea
class	Bacillariophyceae
class	Bivalvia
class	Demospongiae
class	Echinoidea
class	Gastropoda
class	Gromiidea
class	Gymnolaemata
class	Hexacorallia
class	Hydrozoa
class	Malacostraca
class	Octocorallia
class	Ophiuroidea

class	Ostracoda
class	Polychaeta
class	Pycnogonida
class	Stenolaemata
order	Actiniaria
order	Amphipoda
order	Anthoathecata
order	Arcida
order	Camarodonta
order	Cheilostomatida
order	Ctenostomatida
order	Cyclostomatida
order	Forcipulatida
order	Gromiida
order	Haplosclerida
order	Isopoda
order	Littorinimorpha
order	Malacalcyonacea
order	Mysida
order	Neogastropoda
order	Ophiurida
order	Pantopoda
order	Pectinida
order	Phlebobranchia
order	Phyllodocida
order	Podocopida
order	Poecilosclerida
order	Sabellida
order	Stolidobranchia
order	Suberitida
order	Tanaidacea

order	Terebellida
order	Trochida
order	Valvatida
family	Alcyoniidae
family	Alcyoniidae
family	Amphilochoidae
family	Arachnopusiidae
family	Asciidae
family	Asteriidae
family	Beaniidae
family	Bugulidae
family	Calliopiidae
family	Calliostomatidae
family	Capulidae
family	Celleporidae
family	Chalinidae
family	Clavulariidae
family	Crisiidae
family	Echinidae
family	Eudendriidae
family	Gromiidae
family	Janiridae
family	Laternulidae
family	Lichenoporidae
family	Microporidae
family	Munnidae
family	Mysidae
family	Myxillidae
family	Nototanaidae
family	Odontasteridae
family	Ophiopyrgidae

family	Orbiniidae
family	Pectinidae
family	Philobryidae
family	Philoporidae
family	Polynoidae
family	Prosiphonidae
family	Rissoidae
family	Romancheinidae
family	Serpulidae
family	Spirorbidae
family	Styelidae
family	Suberitidae
family	Terebellidae
family	Tubuliporidae
family	Xestoleberididae
genus	<i>Adamussium</i> Thiele, 1934
genus	<i>Alcyonidium</i> Lamouroux, 1813
genus	<i>Alcyonium</i> Linnaeus, 1758
genus	<i>Antarcticaetos</i> Hayward & Thorpe, 1988
genus	<i>Arachnopusia</i> Jullien, 1888
genus	<i>Ascidia</i> Linnaeus, 1767
genus	<i>Austrofilius</i> Hodgson, 1910
genus	<i>Barrukia</i> Bergström, 1916
genus	<i>Beania</i> Johnston, 1840
genus	<i>Camptoplites</i> Harmer, 1923
genus	<i>Clavularia</i> Blainville, 1830
genus	<i>Cnemidocarpa</i> Huntsman, 1913
genus	<i>Crisia</i> Lamouroux, 1812
genus	<i>Cryocapulus</i> Schiaparelli, Bouchet, Fassio & Oliverio, 2020
genus	<i>Diplasterias</i> Perrier, 1891
genus	<i>Disporella</i> Gray, 1848

genus	<i>Eucranta</i> Malmgren, 1865
genus	<i>Eudendrium</i> Ehrenberg, 1834
genus	<i>Exidmonea</i> David, Mongereau & Pouyet, 1972
genus	<i>Favosthimosia</i> Hayward & Winston, 2011
genus	<i>Filicrisia</i> d'Orbigny, 1853
genus	<i>Gitanopsilis</i> Rauschert, 1994
genus	<i>Gromia</i> Dujardin, 1835
genus	<i>Haliclona</i> Grant, 1841
genus	<i>Harmothoe</i> Kinberg, 1856
genus	<i>Helicosiphon</i> Gravier, 1907
genus	<i>Homaxinella</i> Topsent, 1916
genus	<i>Klugeflustra</i> Moyano, 1972
genus	<i>Lanicides</i> Hesse, 1917
genus	<i>Laternula</i> Röding, 1798
genus	<i>Leodamas</i> Kinberg, 1866
genus	<i>Margarella</i> Thiele, 1893
genus	<i>Micropora</i> Gray, 1848
genus	<i>Munna</i> Krøyer, 1839
genus	<i>Mysidetes</i> Holt & Tattersall, 1906
genus	<i>Neobuccinum</i> E. A. Smith, 1879
genus	<i>Nototanais</i> Richardson, 1906
genus	<i>Odontaster</i> Verrill, 1880
genus	<i>Ophioplinthus</i> Lyman, 1878
genus	<i>Oradarea</i> Walker, 1903
genus	<i>Philobrya</i> J. G. Cooper, 1867
genus	<i>Powellisetia</i> Ponder, 1965
genus	<i>Reteporella</i> Busk, 1884
genus	<i>Serpula</i> Linnaeus, 1758
genus	<i>Spirorbis</i> Daudin, 1800
genus	<i>Stelodoryx</i> Topsent, 1904
genus	<i>Sterechinus</i> Koehler, 1901

genus	<i>Subonoba</i> Iredale, 1915
genus	<i>Xestoleberis</i> Sars, 1866
species	<i>Adamussium colbecki</i> (E. A. Smith, 1902)
species	<i>Alcyonium antarcticum</i> Wright & Studer, 1889
species	<i>Alcyonium</i> cf. <i>antarcticum</i> Wright & Studer, 1889
species	<i>Antarcticaetos bubeccata</i> (Rogick, 1955)
species	<i>Arachnopusia decipiens</i> Hayward & Thorpe, 1988
species	<i>Arachnopusia</i> cf. <i>decipiens</i> Hayward & Thorpe, 1988
species	<i>Austrofillius furcatus</i> Hodgson, 1910
species	<i>Barrukia</i> cf. <i>cristata</i> (Willey, 1902)
species	<i>Beania erecta</i> Waters, 1904
species	<i>Camptoplites bicornis</i> (Busk, 1884)
species	<i>Camptoplites tricornis</i> (Waters, 1904)
species	<i>Clavularia frankliniana</i> Roule, 1902
species	<i>Clavularia</i> cf. <i>frankliniana</i> Roule, 1902
species	<i>Cnemidocarpa verrucosa</i> (Lesson, 1830)
species	<i>Cnemidocarpa</i> cf. <i>verrucosa</i> (Lesson, 1830)
species	<i>Cryocapulus subcompressus</i> (Pelseneer, 1903)
species	<i>Diplasterias brucei</i> (Koehler, 1907)
species	<i>Eudendrium scotti</i> Puce, Cerrano & Bavestrello, 2002
species	<i>Favosthimosia milleporoides</i> (Calvet, 1909)
species	<i>Gitanopsilis amissio</i> Rauschert, 1994*
species	<i>Gromia</i> cf. <i>melinus</i> Rothe, Gooday, Cedhagen, Fahrni, Hughes, Page, Pearce & Pawlowski, 2009
species	<i>Harmothoe</i> cf. <i>exanthema</i> (Grube, 1856)
species	<i>Harmothoe</i> cf. <i>spinosa</i> Kinberg, 1856
species	<i>Harmothoe antarctica</i> (McIntosh, 1885)
species	<i>Harmothoe crosetensis</i> (McIntosh, 1885)
species	<i>Harmothoe</i> cf. <i>crosetensis</i> (McIntosh, 1885)
species	<i>Harmothoe fuligineum</i> (Baird, 1865)
species	<i>Harmothoe</i> cf. <i>fuligineum</i> (Baird, 1865)
species	<i>Harmothoe fullo</i> (Grube, 1878)

species	<i>Homaxinella balfourensis</i> (Ridley & Dendy, 1886)
species	<i>Xestoleberis</i> cf. <i>meridionalis</i> Müller, 1908
species	<i>Klugeflustra</i> cf. <i>vanhoeffeni</i> (Kluge, 1914)
species	<i>Lanicides bilobata</i> (Grube, 1877)
species	<i>Laternula elliptica</i> (P. P. King, 1832)
species	<i>Leodamas marginatus</i> (Ehlers, 1897)
species	<i>Margarella crebrilirulata</i> (E. A. Smith, 1907)
species	<i>Margarella refulgens</i> (E. A. Smith, 1907)
species	<i>Micropora</i> cf. <i>notialis</i> Hayward & Ryland, 1993
species	<i>Munna antarctica</i> (Pfeffer, 1887)
species	<i>Mysidetes illigi</i> Zimmer, 1914*
species	<i>Neobuccinum eatoni</i> (E. A. Smith, 1875)
species	<i>Nototanais antarcticus</i> (Hodgson, 1902)
species	<i>Nototanais dimorphus</i> (Beddard, 1886)
species	<i>Odontaster roseus</i> Janosik & Halanych, 2010
species	<i>Odontaster validus</i> Koehler, 1906
species	<i>Ophioplinthus gelida</i> (Koehler, 1901)
species	<i>Oradarea acuminata</i> Thurston, 1974
species	<i>Philobrya sublaevis</i> Pelseneer, 1903
species	<i>Powellisetia deserta</i> (E. A. Smith, 1907)
species	<i>Stelodoryx cribrigera</i> (Ridley & Dendy, 1886)*
species	<i>Sterechinus neumayeri</i> (Meissner, 1900)
species	<i>Subonoba gelida</i> (E. A. Smith, 1907)

Temporal coverage

Data range: 2015-12-01 - 2022-11-08.

Collection data

Collection name: MNA – Biological Collections

Collection identifier: <https://www.gbif.org/grscicoll/collection/a57a1dc1-706c-42db-bb-ad-1e68d9685439>

Parent collection identifier: Italian National Antarctic Museum (section of Genoa)

Specimen preservation method: Specimens in 96% ethanol, slides with whole or dissected organisms (fixed in glycerol), part of organisms preserved in dry for SEM and frozen at -20°C.

Usage licence

Usage licence: Other

IP rights notes: The dataset was published under the licence CC-BY 4.0.

Data resources

Data package title: Checklist of pioneer benthic taxa found on Autonomous Reef Monitoring Structures (ARMS) in Terra Nova Bay (Ross Sea, Antarctica)

Resource link: <https://doi.org/10.15468/4c7zf8>

Alternative identifiers: <https://www.gbif.org/dataset/f024a3d3-7bb2-4985-be57-d890936769ff>

Number of data sets: 1

Data set name: Checklist of pioneer benthic taxa found on Autonomous Reef Monitoring Structures (ARMS) in Terra Nova Bay (Ross Sea, Antarctica)

Data format: Darwin Core

Description: The dataset comprises a total of 277 distributional records, each one corresponding to a voucher specimen stored at the MNA, Section Genoa (Cometti et al. 2024). These records originate from ARMS, at varying time ranges of 1, 2, 3 and 5 years in Tethys Bay (Ross Sea, Antarctica). The occurrences presented in this dataset represent an important contribution as a baseline of the taxonomic composition of pioneer benthic communities in the Ross Sea and it will be useful to study their future dynamics.

Column label	Column description
occurrenceID	A global unique identifier for the occurrence.
institutionCode	The name (or acronym) in use by the institution having custody of the object(s) or information referred to in the record.
institutionID	An identifier for the institution having custody of the objects or information referred to in the record.
collectionCode	The acronym identifying the collection or dataset from which the record was derived.

collectionID	An identifier for the dataset from which the record was derived.
catalogNumber	An identifier of any form assigned by the source within a physical collection or digital dataset for the record which may not be unique, but should be fairly unique in combination with the institution and collection code.
basisOfRecord	The specific nature of the data record (Preserved Specimen).
type	The genre of the resource (PhysicalObject).
scientificName	The full scientific name, with authorship and date information, if known.
taxonRank	The taxonomic rank of the most specific name in the scientificName.
kingdom	The full scientific name of the kingdom in which the taxon is classified.
phylum	The full scientific name of the phylum in which the taxon is classified.
class	The full scientific name of the class in which the taxon is classified.
order	The full scientific name of the order in which the taxon is classified.
family	The full scientific name of the family in which the taxon is classified.
genus	The full scientific name of the genus in which the taxon is classified.
specificEpithet	The name of the first or species epithet of the scientificName.
scientificNameAuthorship	The authorship information for the scientificName formatted according to the conventions of the applicable.
identificationQualifier	A controlled value to express the determiner's doubts about the Identification (sp. cf).
scientificNameID	An identifier for the nomenclatural (not taxonomic) details of a scientific name.
individualCount	The number of individuals present at the time of the Occurrence.
lifeStage	The life stage of organisms. In detail, juveniles and eggs.
occurrenceRemarks	Antarctic Expeditions in which the organisms were sampled.
eventDate	The date-time or interval during which an Event occurred.
sampleSizeValue	A numerical value indicating the time of colonisation.
sampleSizeUnit	The unit of measurement of the time duration. This term must have a corresponding to sampleSizeValue.
eventID	A global unique identifier for the set of information associated with an Event.
decimalLatitude	The geographic latitude (in decimal degrees, using the spatial reference system given in geodeticDatum).
decimalLongitude	The geographic longitude (in decimal degrees, using the spatial reference system given in geodeticDatum).
geodeticDatum	The spatial reference system (WGS84) upon which the geographic coordinates given in decimalLatitude and decimalLongitude are based.

minimumDepthInMetres	Minimum sampling depth during event in metres.
maximumDepthInMetres	Maximum sampling depth during event in metres.
coordinatePrecision	A decimal representation of the precision of the coordinates given in the decimalLatitude and decimalLongitude.
samplingProtocol	Gear used to collect specimens and relative DOI of manuscript in which the sampling method is described.
dynamicProperties	Concatenation of information, specifically: Movement (sessile, vagile), PVC plate number (lowest = 1 to highest = 10) and plate orientation (T = top, B = bottom).
preparations	Description of the tissue or the voucher specimen and preservation method.
occurrenceStatus	Statement about the presence or absence of a specimen.
locality	The specific description of the place.
continent	Continent where the organisms were sampled.
countryCode	The standard code for the country where the organisms were sampled.
recordedBy	Surname and name of the personnel who collected the samples.
recordedByID	ORCID of the personnel who collected the samples.
identifiedBy	Surname and name of the personnel who analysed and recognised the single species.
identifiedByID	ORCID of the personnel who analysed and recognised the single species.

Acknowledgements

This paper is an Italian National Antarctic Programme (PNRA) contribution to the [SCAR-ANTOS Expert Group](#) and to the CCAMLR CONSERVATION MEASURE 91-05 (2016) for the Ross Sea region Marine Protected Area, specifically, addressing Annex 91-05/C ('long-term monitoring of benthic ecosystem functions'). We are grateful to Blanca Figuerola and Paul D. Taylor for the valuable help to identify some Bryozoa's samples, Giorgio Bavestrello to identify some sample of Hydrozoa and Marco Bertolino to identify some samples of Porifera. The publication of this data paper was funded by the Belgian Science Policy Office (BELSPO, contract n°FR/36/AN1/AntaBIS) in the Framework of EU-Lifewatch as a contribution to the SCAR Antarctic biodiversity portal ([biodiversity.aq](#)). The authors are also immensely grateful to Dr. Anton Van De Putte and Dr. Yi Ming Gan, who provided much appreciated advice and help on the metadata standards and Darwin core archive format during the quality control of the dataset.

Author contributions

VC compiled and edited the dataset and wrote the original manuscript. VC, MC, AG, MG and NN identified and counted the specimens, compiled and edited the dataset, provided funding, lab equipment and reviewed the manuscript and SC wrote, reviewed and edited.

SS provided funding, resources, specimens and lab equipment, supervised the project and edited the manuscript. All authors have read and agreed to the published version of the manuscript.

References

- Arntz WE, Gallardo VA (1994) Antarctic benthos: present position and future prospects. In: Hempel G (Ed.) Antarctic science. 243-277 pp. https://doi.org/10.1007/978-3-642-78711-9_16
- Aronson R, Blake D (2001) Global climate change and the origin of modern benthic communities in Antarctica. *American Zoologist* 41 (1): 27-39. <https://doi.org/10.1093/icb/41.1.27>
- Barnes DA, Souster T (2011) Reduced survival of Antarctic benthos linked to climate-induced iceberg scouring. *Nature Climate Change* 1 (7): 365-368. <https://doi.org/10.1038/nclimate1232>
- Barnes DA, Fenton M, Cordingley A (2014) Climate-linked iceberg activity massively reduces spatial competition in Antarctic shallow waters. *Current Biology* 24 (12). <https://doi.org/10.1016/j.cub.2014.04.040>
- Barnes DA, Ashton G, Morley S, Peck L (2021) 1 °C warming increases spatial competition frequency and complexity in Antarctic marine macrofauna. *Communications Biology* 4 (1). <https://doi.org/10.1038/s42003-021-01742-w>
- Bonello G, Grillo M, Cecchetto M, Giallain M, Granata A, Guglielmo L, Pane L, Schiaparelli S (2020) Distributional records of Ross Sea (Antarctica) planktic Copepoda from bibliographic data and samples curated at the Italian National Antarctic Museum (MNA): checklist of species collected in the Ross Sea sector from 1987 to 1995. *ZooKeys* 969: 1-22. <https://doi.org/10.3897/zookeys.969.52334>
- Bowden D, Clarke A, Peck L, Barnes D (2006) Antarctic sessile marine benthos: colonisation and growth on artificial substrata over three years. *Marine Ecology Progress Series* 316: 1-16. <https://doi.org/10.3354/meps316001>
- Caruso G (2019) Microplastics as vectors of contaminants. *Marine Pollution Bulletin* 146: 921-924. <https://doi.org/10.1016/j.marpolbul.2019.07.052>
- Caruso G, Maimone G, Rappazzo AC, Dell'Acqua O, Laganà P, Azzaro M (2023) Microbial biofilm colonizing plastic substrates in the Ross Sea (Antarctica): first overview of community-level physiological profiles. *Journal of Marine Science and Engineering* 11 (7). <https://doi.org/10.3390/jmse11071317>
- Cattaneo-Vietti R, Chiantore M, Gambi MC, Albertelli G, Cormaci M, Di Geronimo I (2000) Spatial and vertical distribution of benthic littoral communities in Terra Nova Bay. In: Faranda FM, Guglielmo L, Ianora A (Eds) *Ross Sea ecology*. 503-514 pp. https://doi.org/10.1007/978-3-642-59607-0_36
- Cecchetto M, Alvaro MC, Ghiglione C, Guzzi A, Mazzoli C, Piazza P, Schiaparelli S (2017) Distributional records of Antarctic and sub-Antarctic Ophiuroidea from samples curated at the Italian National Antarctic Museum (MNA): check-list update of the group in the Terra Nova Bay area (Ross Sea) and launch of the MNA 3D model 'virtual gallery'. *ZooKeys* 705: 61-79. <https://doi.org/10.3897/zookeys.705.13712>
- Cecchetto M, Lombardi C, Canese S, Cocito S, Kuklinski P, Mazzoli C, Schiaparelli S (2019) The Bryozoa collection of the Italian National Antarctic Museum, with an updated

- checklist from Terra Nova Bay, Ross Sea. *ZooKeys* 812: 1-22. <https://doi.org/10.3897/zookeys.812.26964>
- Cecchetto M, Dettai A, Gallut C, Obst M, Kuklinski P, Balazy P, Chelchowski M, Małachowicz M, Poćwierz-Kotus A, Zbawicka M, Reiss H, Eléaume M, Ficetola GF, Pavlouli C, Exter K, Fontaneto D, Schiaparelli S (2024) Seasonality of primary production explains the richness of pioneering benthic communities. *Nature Communications* 15 (1). <https://doi.org/10.1038/s41467-024-52673-z>
 - Cometti V, Cecchetto M, Guzzi A, Grillo M, Noli N, Corsolini S, Schiaparelli S (2024) Checklist of pioneer benthic taxa found on Autonomous Reef Monitoring Structures (ARMS) in Terra Nova Bay (Ross Sea, Antarctica). Occurrence dataset. Version 1.2.. Italian National Antarctic Museum (MNA, Section of Genoa). URL: <https://doi.org/10.15468/4c7zf8>
 - David R, Uyerra M, Carvalho S, Anlauf H, Borja A, Cahill A, Carugati L, Danovaro R, De Jode A, Feral J, Guillemain D, Martire ML, D'Avray LTDV, Pearman J, Chenuil A (2019) Lessons from photo analyses of Autonomous Reef Monitoring Structures as tools to detect (bio-)geographical, spatial, and environmental effects. *Marine Pollution Bulletin* 141: 420-429. <https://doi.org/10.1016/j.marpolbul.2019.02.066>
 - Dayton P (1989) Interdecadal variation in an Antarctic sponge and its predators from oceanographic climate shifts. *Science* 245 (4925): 1484-1486. <https://doi.org/10.1126/science.245.4925.1484>
 - Dayton P, Jarrell S, Kim S, Thrush S, Hammerstrom K, Slattery M, Parnell E (2016) Surprising episodic recruitment and growth of Antarctic sponges: implications for ecological resilience. *Journal of Experimental Marine Biology and Ecology* 482: 38-55. <https://doi.org/10.1016/j.jembe.2016.05.001>
 - De Broyer C, Koubbi P, Griffiths HJ, Raymond B, Udekem d'Acoz Cd, Van de Putte AP, Danis B, David B, Grant S, Gutt J, Held C, Hosie G, Huettmann F, Post A, Ropert-Coudert Y (Eds) (2014) Biogeographic atlas of the Southern Ocean. Scientific Committee on Antarctic Research, Cambridge, 498 pp.
 - Fillinger L, Janussen D, Lundälv T, Richter C (2013) Rapid glass sponge expansion after climate-induced Antarctic ice shelf collapse. *Current Biology* 23 (14): 1330-1334. <https://doi.org/10.1016/j.cub.2013.05.051>
 - Fonseca VG, Sinniger F, Gaspar JM, Quince C, Creer S, Power DM, Peck LS, Clark MS (2017) Revealing higher than expected meiofaunal diversity in Antarctic sediments: a metabarcoding approach. *Scientific Reports* 7 (1): 6094. <https://doi.org/10.1038/s41598-017-06687-x>
 - Garlasché G, Karimullah K, Iakovenko N, Velasco-Castrillón A, Janko K, Guidetti R, Rebecchi L, Cecchetto M, Schiaparelli S, Jersabek C, De Smet W, Fontaneto D (2019) A data set on the distribution of Rotifera in Antarctica. *Biogeographia – The Journal of Integrative Biogeography* 35 <https://doi.org/10.21426/b635044786>
 - Ghigliione C, Schiaparelli S, Alvaro MC, Griffiths H, Linse K (2013) Ross Sea Mollusca from the Latitudinal Gradient Program: R/V *Italica* 2004 Rauschert dredge samples. *ZooKeys* 341: 37-48. <https://doi.org/10.3897/zookeys.341.6031>
 - Ghigliione C, Alvaro MC, Cecchetto M, Canese S, Downey R, Guzzi A, Mazzoli C, Piazza P, Tore Rapp H, Sarà A, Schiaparelli S (2018) Porifera collection of the Italian National Antarctic Museum (MNA), with an updated checklist from Terra Nova Bay (Ross Sea). *ZooKeys* 758: 137-156. <https://doi.org/10.3897/zookeys.758.23485>

- Grillo M, Bonello G, Cecchetto M, Guzzi A, Noli N, Cometti V, Schiaparelli S (2024) Planktonic, benthic and sympagic copepods collected from the desalination unit of Mario Zucchelli Research Station in Terra Nova Bay (Ross Sea, Antarctica). *Biodiversity Data Journal* 12 <https://doi.org/10.3897/bdj.12.e119633>
- Guzzi A, Alvaro MC, Danis B, Moreau C, Schiaparelli S (2022) Not all that glitters is gold: barcoding effort reveals taxonomic incongruences in iconic Ross Sea sea tars. *Diversity* 14 (6). <https://doi.org/10.3390/d14060457>
- Krzeminska M, Kuklinski P (2018) Biodiversity patterns of rock encrusting fauna from the shallow sublittoral of the Admiralty Bay. *Marine Environmental Research* 139: 169-181. <https://doi.org/10.1016/j.marenvres.2018.03.016>
- Leray M, Knowlton N (2015) DNA barcoding and metabarcoding of standardized samples reveal patterns of marine benthic diversity. *Proceedings of the National Academy of Sciences of the United States of America* 112 (7): 2076-81. <https://doi.org/10.1073/pnas.1424997112>
- Maroni P, Wilson N (2022) Multiple Doris “kerguelenensis” (Nudibranchia) species span the Antarctic Polar Front. *Ecology and Evolution* 12 (9). <https://doi.org/10.1002/ece3.9333>
- Maroni P, Baker B, Moran A, Woods HA, Avila C, Johnstone G, Stark J, Kocot K, Lockhart S, Saucède T, Rouse G, Wilson N (2022) One Antarctic slug to confuse them all: the underestimated diversity of Doris kerguelenensis. *Invertebrate Systematics* 36 (5): 419-435. <https://doi.org/10.1071/is21073>
- Noli N, Cecchetto M, Guzzi A, Grillo M, Cometti V, Schiaparelli S (2024) New distributional records of Southern Ocean Isopoda based on vouchers from the Italian National Antarctic Museum (MNA, Section of Genoa). *Biodiversity Data Journal* 12: e127689. <https://doi.org/10.3897/BDJ.12.e127689>
- Papale M, Fazi S, Severini M, Scarinci R, Dell'Acqua O, Azzaro M, Venuti V, Fazio B, Fazio E, Crupi V, Irrera A, Rizzo C, Giudice AL, Caruso G (2024) Structural properties and microbial diversity of the biofilm colonizing plastic substrates in Terra Nova Bay (Antarctica). *Science of The Total Environment* 943 <https://doi.org/10.1016/j.scitotenv.2024.173773>
- Pearman J, Chust G, Aylagas E, Villarino E, Watson J, Chenuil A, Borja A, Cahill A, Carugati L, Danovaro R, David R, Irigoien X, Mendibil I, Moncheva S, Rodríguez-Ezpeleta N, Uyarra M, Carvalho S (2020) Pan-regional marine benthic cryptobiome biodiversity patterns revealed by metabarcoding Autonomous Reef Monitoring Structures. *Molecular Ecology* 29 (24): 4882-4897. <https://doi.org/10.1111/mec.15692>
- Peck L (2018) Antarctic marine biodiversity: adaptations, environments and responses to change. In: Hawkins SJ, Evans AJ, Dale AC, Firth LB, Smith IP (Eds) *Oceanography and marine biology*. 1, 56. 518 pp. [ISBN 9780429454455]. <https://doi.org/10.1201/9780429454455>
- Piazza P, Błażewicz-Paszkowycz M, Ghiglione C, Alvaro M, Schnabel K, Schiaparelli S (2014) Distributional records of Ross Sea (Antarctica) Tanaidacea from museum samples stored in the collections of the Italian National Antarctic Museum (MNA) and the New Zealand National Institute of Water and Atmospheric Research (NIWA). *ZooKeys* 451: 49-60. <https://doi.org/10.3897/zookeys.451.8373>
- Piazza P, Cummings V, Guzzi A, Hawes I, Lohrer A, Marini S, Marriott P, Menna F, Nocerino E, Peirano A, Kim S, Schiaparelli S (2019) Underwater photogrammetry in Antarctica: long-term observations in benthic ecosystems and legacy data rescue. *Polar Biology* 42 (6): 1061-1079. <https://doi.org/10.1007/s00300-019-02480-w>

- Piazza P, Gattone SA, Guzzi A, Schiaparelli S (2020) Towards a robust baseline for long-term monitoring of Antarctic coastal benthos. *Hydrobiologia* 847 (7): 1753-1771. <https://doi.org/10.1007/s10750-020-04177-2>
- Selbmann L, Schiaparelli S, Onofri S, Zucconi L, Isola D, Rottigni M, Ghiglione C, Piazza P, Alvaro MC (2015) Distributional records of Antarctic fungi based on strains preserved in the Culture Collection of Fungi from Extreme Environments (CCFEE) Mycological section associated with the Italian National Antarctic Museum (MNA). *MycKeys* 10: 57-71. <https://doi.org/10.3897/mycokeys.10.5343>
- Valentini A, Taberlet P, Miaud C, Civade R, Herder J, Thomsen PF, Bellemain E, Besnard A, Coissac E, Boyer F, Gaboriaud C, Jean P, Poulet N, Roset N, Copp GH, Geniez P, Pont D, Argillier C, Baudoin J, Peroux T, Crivelli AJ, Olivier A, Acqueberge M, Le Brun M, Møller PR, Willerslev E, Dejean T (2016) Next-generation monitoring of aquatic biodiversity using environmental DNA metabarcoding. *Molecular Ecology* 25 (4): 929-42. <https://doi.org/10.1111/mec.13428>
- Wilson N, SchrodL M, Halanych K (2009) Ocean barriers and glaciation: evidence for explosive radiation of mitochondrial lineages in the Antarctic sea slug *Doris kerguelensis* (Mollusca, Nudibranchia). *Molecular Ecology* 18 (5): 965-984. <https://doi.org/10.1111/j.1365-294x.2008.04071.x>