



Two new species and one new record of *Ischnothyreus* Simon, 1893 (Araneae, Oonopidae) from China

Chenxue Song[‡], Yanfeng Tong[‡], Dongju Bian[§], Zhisheng Zhang^l

[‡] Life Science College, Shenyang Normal University, Shenyang 110034, China

[§] Key Laboratory of Forest Ecology and Management, Institute of Applied Ecology, Chinese Academy of Sciences, Shenyang 110016, China

^l Key Laboratory of Eco-environments in Three Gorges Reservoir Region (Ministry of Education), School of Life Sciences, Southwest University, Chongqing 400715, China

Corresponding author: Yanfeng Tong (tyf68@hotmail.com), Dongju Bian (biandongju@163.com), Zhisheng Zhang (zhangzs327@qq.com)

Academic editor: Alireza Zamani

Received: 02 Mar 2024 | Accepted: 03 Apr 2024 | Published: 11 Apr 2024

Citation: Song C, Tong Y, Bian D, Zhang Z (2024) Two new species and one new record of *Ischnothyreus* Simon, 1893 (Araneae, Oonopidae) from China. Biodiversity Data Journal 12: e122100.

<https://doi.org/10.3897/BDJ.12.e122100>

ZooBank: [urn:lsid:zoobank.org:pub:1E1758DC-BDF9-4A18-A234-73AB69A84ACE](https://zoobank.org/pub:1E1758DC-BDF9-4A18-A234-73AB69A84ACE)

Abstract

Background

Ischnothyreus Simon, 1893 is a large genus of oonopid spiders that currently contains 126 species, amongst which, 28 have been recorded in China.

New information

Two new *Ischnothyreus* species, *Ischnothyreus daheling* Tong & Zhang, **sp. nov.** and *Ischnothyreus longyang* Tong & Zhang, **sp. nov.**, are described, based on specimens collected from Yunnan Province and *Ischnothyreus velox* Jackson, 1908 is recorded in

China for the first time, based on material collected from Guangxi Province. All three species are illustrated.

Keywords

Asia, goblin spiders, new distribution, taxonomy

Introduction

Oonopidae is a diverse spider family with 1940 extant described species in 115 genera (WSC 2024). They have a nearly worldwide distribution, occurring mainly in the leaf litter, under bark and in the tree canopy (Jocqué and Dippenaar-Schoeman 2006, Ubick and Dupérré 2017).

The genus *Ischnothyreus* Simon, 1893 is one of the most speciose genera of Oonopidae, with 126 extant species mainly distributed in the Old World (Fu et al. 2023, Tong et al. 2023, WSC 2024). The genus *Ischnothyreus* Simon, 1893 can be recognised by the presence of leg spines, the usually small abdominal scutum, the strongly sclerotised male palps, the heavily sclerotised male endites and the winding genital tube in females (Kranz-Baltensperger 2011).

In this paper, two new species of *Ischnothyreus*, *Ischnothyreus daheling* **sp. nov.** and *Ischnothyreus longyang* **sp. nov.** are described from Yunnan, China. Furthermore, *Ischnothyreus velox* Jackson, 1908 is newly recorded from China (Guangxi).

Materials and methods

The specimens were examined using a Leica M205C stereomicroscope. Details were studied using an Olympus BX51 compound microscope. Photos were made with a Canon EOS 750D zoom digital camera (18 megapixels) mounted on an Olympus BX51 compound microscope. Endogynes were cleared in lactic acid. Scanning electron microscope (SEM) images were taken under high vacuum with a Hitachi S-4800 after critical point drying and gold-palladium coating. All measurements were taken using an Olympus BX51 compound microscope and are in millimetres. Material is deposited in Shenyang Normal University (SYNU) in Liaoning, China.

The following abbreviations are used in the text and figures: ALE = anterior lateral eyes; bss = bell-shaped structure; fp = flag-shaped process; PLE = posterior lateral eyes; PME = posterior median eyes; rl = retrolateral lobe; stp = strong, tooth-like projection.

Taxon treatments

Ischnothyreus daheling Tong & Zhang sp. nov.

- ZooBank [0457642A-703E-4A20-AC1B-FFF54BD7A6F1](https://doi.org/10.21203/rs.3.rs-10457642/v1)

Materials

Holotype:

- scientificName: *Ischnothyreus daheling*; order: Araneae; family: Oonopidae; genus: *Ischnothyreus*; country: China; stateProvince: Yunnan; county: Baoshan City, Tengchong County; locality: JietouTown, Datou Village, Dahelingganjiao; samplingProtocol: sifting leaf litter; eventDate: 23/02/2011; individualCount: 1; sex: male; lifeStage: adult; recordNumber: SYNU-707; recordedBy: Zongxu Li, Luyu Wang; identifiedBy: Yanfeng Tong; occurrenceID: A9E7AAC7-43DA-5861-B070-AA367E644B62

Paratypes:

- scientificName: *Ischnothyreus daheling*; order: Araneae; family: Oonopidae; genus: *Ischnothyreus*; country: China; stateProvince: Yunnan; county: Baoshan City, Tengchong County; locality: JietouTown, Datou Village, Dahelingganjiao; samplingProtocol: sifting leaf litter; eventDate: 23/02/2011; individualCount: 3; sex: female; lifeStage: adult; recordNumber: SYNU-708-710; recordedBy: Zongxu Li, Luyu Wang; identifiedBy: Yanfeng Tong; occurrenceID: 0FD72221-D529-5328-9438-36EDCECCEE10
- scientificName: *Ischnothyreus daheling*; order: Araneae; family: Oonopidae; genus: *Ischnothyreus*; country: China; stateProvince: Yunnan; county: Baoshan City, Tengchong County; locality: Jietou Town, Datang Village; samplingProtocol: sifting leaf litter; eventDate: 21/02/2011; individualCount: 4; sex: female; lifeStage: adult; recordNumber: SYNU-711-714; recordedBy: Zongxu Li, Luyu Wang; identifiedBy: Yanfeng Tong; occurrenceID: A39B8642-53DA-5493-82BD-1968E6544502
- scientificName: *Ischnothyreus daheling*; order: Araneae; family: Oonopidae; genus: *Ischnothyreus*; country: China; stateProvince: Yunnan; county: Baoshan City, Tengchong County; locality: Wuhe Town, Xiaodifang Village; samplingProtocol: sifting leaf litter; eventDate: 27/02/2011; individualCount: 3; sex: female; lifeStage: adult; recordNumber: SYNU-715-717; recordedBy: Zongxu Li, Luyu Wang; identifiedBy: Yanfeng Tong; occurrenceID: F489D23E-11EB-5FC1-ACD9-A3F0EBA8AB8B
- scientificName: *Ischnothyreus daheling*; order: Araneae; family: Oonopidae; genus: *Ischnothyreus*; country: China; stateProvince: Yunnan; county: Baoshan City, Tengchong County; locality: Jietou Town, Sabadi; samplingProtocol: sifting leaf litter; eventDate: 25/05/2011; individualCount: 1; sex: female; lifeStage: adult; recordNumber: SYNU-718; recordedBy: Zongxu Li, Luyu Wang; identifiedBy: Yanfeng Tong; occurrenceID: 6DFAC351-F3D7-5037-8495-2316D4E4E770
- scientificName: *Ischnothyreus daheling*; order: Araneae; family: Oonopidae; genus: *Ischnothyreus*; country: China; stateProvince: Yunnan; locality: Gaoligong Mountain, Nankang Protection Station; eventDate: 20/04/2011; individualCount: 1; sex: female; lifeStage: adult; recordNumber: SYNU-719; recordedBy: Ping Feng; identifiedBy: Yanfeng Tong; occurrenceID: AC048212-9E5A-59B5-8D18-EFB379F11EDE

Description

Male (Holotype). **Body:** habitus as in Fig. 1A–C; body length 1.64. **Carapace:** 0.88 long, 0.65 wide; yellow, oval in dorsal view, pars cephalica strongly elevated in lateral view, surface of elevated portion of pars cephalica reticulate, sides finely reticulate, lateral margin straight, smooth (Fig. 1C). **Clypeus:** straight in frontal view, ALE separated from edge of carapace by 1.5 times of their diameter (Fig. 1F). **Eyes:** ALE largest, ALE circular, PME circular, PLE oval; posterior eye row recurved from above; ALE separated by less than their radius, ALE and PLE touching. **Sternum:** as long as wide, pale orange (Fig. 1E). **Mouthparts:** chelicerae, endites and labium yellow; chelicerae straight, base of fangs with flag-like process, fang groove with a few small and one larger denticles (Fig. 2J–L); anteromedian tip of endites with one strong, tooth-like projection (Fig. 1G). **Abdomen:** 0.76 long, 0.44 wide; dorsal scutum well sclerotised, dark brown, covering 4/5 of the abdomen width and approximately 5/6 of the abdomen length, not fused to epigastric scutum; epigastric and postepigastric scutum pale orange, fused, covering 3/4 of the abdomen length. **Legs:** pale orange, femur I with 2 prolateral spines, tibia I with 4 pairs, metatarsus I with 2 pairs of long ventral spines. Leg II spination similar to leg I, except femur with only 1 prolateral spine. Legs III and IV spineless. **Palp:** trochanter without ventral projection; bulb with 1 ventral protuberance, distal end of bulb stout (Fig. 2A–I).

Female (paratype, SYNU-708). Same as male, except as noted. Habitus as in Fig. 3 A–C. **Body:** 1.82 long. **Carapace:** 0.81 long, 0.66 wide. **Mouthparts:** chelicerae and endites unmodified. **Abdomen:** 1.24 long, 0.79 wide; dorsal scutum approximately 1/3 of the abdomen length. **Epigastric area:** postepigastric scutum with inverted bell-shaped structure (Fig. 3G and H). **Endogyne:** winding tube simple, strongly convoluted; lateral apodemes present (Fig. 3I).

Diagnosis

This species is similar to *Ischnothyreus yueluensis* Yin & Wang, 1984 in the size of the abdominal scuta, but can be distinguished by the flag-shaped sclerotised process of cheliceral fang (Fig. 2J) vs. small stick-shaped sclerotised process (Huang et al. 2021: fig. 4G), the stout distal end of bulb (Fig. 2G and H) vs. elongated distal end, with several leaf-shaped membranes (Huang et al. 2021: figs. 5G and H) and the inverted bell-shaped structure of postepigastric scutum of female (Fig. 3G) vs. small, narrow bowl-shaped structure (Huang et al. 2021: fig. 6H).

Etymology

The specific epithet is a noun in apposition taken from the type locality.

Distribution

Known only from the type locality.




Figure 1. [doi](#)

Ischnothyreus daheling sp. nov., male holotype. **A** habitus, dorsal view; **B** habitus, ventral view; **C** habitus, lateral view; **D** prosoma, dorsal view; **E** prosoma, ventral view; **F** prosoma, anterior view; **G** left chelicera, anterior view; **H** left chelicera, lateral view; **I** left chelicera, posterior view. Abbreviation: stp = strong, tooth-like projection. Scale bars: A–F = 0.4 mm; G–I = 0.1 mm.




Figure 2. [doi](#)

Ischnothyreus daheling sp. nov., male holotype, light (A–C) and SEM (D–L) images. **A** left palp, prolateral view; **B** left palp, retrolateral view; **C** left palp, dorsal view; **D** left palp, prolateral view; **E** left palp, retrolateral view; **F** left palp, dorsal view; **G** distal part of palpal bulb, prolateral view; **H** distal part of palpal bulb, retrolateral view; **I** distal part of palpal bulb, dorsal view; **J** left chelicera, anterior view; **K** left chelicera, posterior view; **L** left chelicera, lateral view. Abbreviation: fp = flag-shaped process. Scale bars: A–F = 0.4 mm; G–L = 0.1 mm.




Figure 3. [doi](#)

Ischnothyreus daheling sp. nov., female paratype. **A** habitus, dorsal view; **B** habitus, ventral view; **C** habitus, lateral view; **D** prosoma, dorsal view; **E** prosoma, ventral view; **F** prosoma, anterior view; **G** epigastric region, ventral view **H** endogyne, ventral view; **I** endogyne, dorsal view. Abbreviation: bss = bell-shaped structure. Scale bars: A–F = 0.4 mm; G–I = 0.1 mm.

Ischnothyreus longyang Tong & Zhang sp. nov.

- ZooBank [E2BDA5A7-657E-4266-AF1B-F5A5EC00B662](https://doi.org/10.21203/rs.3.rs-10000000/v1)

Material

Holotype:

- scientificName: *Ischnothyreus longyang*; order: Araneae; family: Oonopidae; genus: *Ischnothyreus*; country: China; stateProvince: Yunnan; county: Baoshan City; locality: Longyang District, Lujiang Town, Nankang bealock,; samplingProtocol: sifting leaf litter; eventDate: 28/02/2011; individualCount: 1; sex: male; lifeStage: adult; recordNumber: SYNU-721; recordedBy: Zongxu Li, Luyu Wang; identifiedBy: Yanfeng Tong; occurrenceID: 57B4E8C2-B0A4-5835-8ED2-92C6C567A3F6

Description

Male (holotype). **Body:** habitus as in Fig. 4A–C; body length 1.91. **Carapace:** 0.91 long, 0.73 wide; yellow, oval in dorsal view, with brown egg-shaped patches behind eyes, pars cephalica strongly elevated in lateral view, surface of elevated portion of pars cephalica smooth, sides finely reticulate, lateral margin straight, smooth (Fig. 4C and D). **Clypeus:** straight in frontal view, ALE separated from edge of carapace less than their diameter (Fig. 4F). **Eyes:** ALE largest, ALE circular, PME circular, PLE oval; posterior eye row straight from above; ALE touching, ALE-PLE touching. **Sternum:** longer than wide, pale orange (Fig. 4E). **Mouthparts:** chelicerae, endites and labium yellow; chelicerae straight, base of fangs with flag-like process, fang groove with a few small and two larger denticles (Fig. 5K and L); anteromedian tip of endites with one strong, tooth-like projection (Fig. 4E). **Abdomen:** 1.00 long, 0.61 wide; dorsal scutum well sclerotised, dark brown, covering 1/2 of the abdomen width and approximately 1/2 of the abdomen length, not fused to epigastric scutum; epigastric and postepigastric scutum well sclerotised, yellow, fused; postepigastric scutum covering about 2/3 of the abdomen length. **Legs:** pale orange, femur I with 2 prolateral spines, tibia I with 4 pairs, metatarsus I with 2 pairs of long ventral spines. Leg II spination similar to leg I, except femur with only 1 prolateral spine. Legs III and IV spineless. **Palp:** trochanter without ventral projection; bulb with 1 ventral protuberance, distal end of bulb stout, with a retrolateral broad lobe (Fig. 5A–I).

Female. Unknown.

Diagnosis

The new species is similar to *Ischnothyreus qiuxing* Tong & Li, 2021 in the size of the abdominal scuta, but can be distinguished by the flag-shaped sclerotised process of cheliceral fang (Fig. 5K) vs. finger-shaped process (Tong et al. 2021: figs. 13H–J, 21F and G) and the broad retrolateral lobe of palpal bulb (Fig. 5H) vs. lack the lobe, but has a tuber-like projection (Tong et al. 2021: fig. 14H).




Figure 4. [doi](#)

Ischnothyreus longyang sp. nov., male holotype. **A** habitus, dorsal view; **B** habitus, ventral view; **C** habitus, lateral view; **D** prosoma, dorsal view; **E** prosoma, ventral view; **F** prosoma, anterior view; **G** left chelicera, anterior view; **H** left chelicera, lateral view; **I** left chelicera, posterior view. Abbreviation: stp = strong, tooth-like projection. Scale bars: A–F = 0.4 mm; G–I = 0.1 mm.

Etymology

The specific epithet is a noun in apposition taken from the type locality.




Figure 5. [doi](#)

Ischnothyreus longyang sp. nov., male holotype, light (A–C) and SEM (D–L) images. **A** left palp, prolateral view; **B** left palp, retrolateral view; **C** left palp, dorsal view; **D** left palp, prolateral view; **E** left palp, retrolateral view; **F** left palp, dorsal view; **G** palpal bulb, prolateral view; **H** distal part of palpal bulb, retrolateral view; **I** distal part of palpal bulb, dorsal view; **J** left chelicera, anterior view; **K** distal part of left chelicera, anterior view; **L** distal part of left chelicera, posterior view. Abbreviation: fp = flag-shaped process; rl = retrolateral lobe. Scale bars: A–F, J–L = 0.1 mm; G–I = 0.05 mm.

Distribution

Known only from the type locality.




Figure 6. [doi](#)

Ischnothyreus velox Jackson, 1908, female. **A** habitus, dorsal view; **B** prosoma, dorsal view; **C** habitus, ventral view; **D** prosoma, ventral view; **E** habitus, lateral view; **F** prosoma, anterior view; **G** endogyne, ventral view; **H** endogyne, dorsal view. Scale bars: A–F = 0.4 mm; G, H = 0.1 mm.

Ischnothyreus velox Jackson, 1908

Material

- a. scientificName: *Ischnothyreus velox* Jackson, 1908; order: Araneae; family: Oonopidae; genus: *Ischnothyreus*; country: China; stateProvince: Guangxi; county: Liuzhou City; locality: Hills behind Bus Master Station, Laohuchong, Bajiao cave; verbatimElevation: 85 m; verbatimCoordinates: 24°17'49.5"N, 109°24'10.3"E; eventDate: 10/12/2012; individualCount: 1; sex: female; lifeStage: adult; recordNumber: SYNU-700; recordedBy: Zhigang Chen & Zhe Zhao; identifiedBy: Yanfeng Tong; occurrenceID: AE872332-BE8E-5D62-900A-5AD15212C5FB

Description

See Platnick et al. (2012).

Diagnosis

Males differ from those of *Ischnothyreus peltifer* (Simon, 1892) in lacking a protuberance on the base of the fang; palp with rounded bulb and sperm pore larger (Platnick et al. 2012: figs. 104 and 107; Brescovit et al. 2019: figs. 26–30). Females can be recognised by the boat-shaped structure on the poststepigastric scutum (Fig. 6 G).

Distribution

Pantropical; newly recorded from China.

Acknowledgements

The manuscript benefitted greatly from comments by Alireza Zamani and an anonymous referee. This study was supported by the National Natural Science Foundation of China (NSFC-32370479, 31972867).

References

- Brescovit AD, Bonaldo AB, Ott R, Chavari JL (2019) To boldly go: on invasive goblin spiders in Brazil (Araneae, Oonopidae). *Iheringia, Série Zoologia* 109 (e2019033): 1-20. <https://doi.org/10.1590/1678-4766e2019033>
- Fu HJ, Wang ZX, Sun YF, Tong YF, Bian DJ (2023) A new species of *Ischnothyreus* Simon, 1893 (Araneae, Oonopidae) from Guangdong Province, China. *Biodiversity Data Journal* 11 (e105283): 1-8. <https://doi.org/10.3897/BDJ.11.e105283>
- Huang Y, Tong YF, Bian DJ, Li SQ (2021) One new species of the genus *Ischnothyreus* Simon, 1893 and re-description of *I. yueluensis* Yin & Wang, 1984 from China (Araneae, Oonopidae). *Biodiversity Data Journal* 9 (e66843): 1-11. <https://doi.org/10.3897/BDJ.9.e66843>

- Jocqué R, Dippenaar-Schoeman AS (2006) Spider families of the World. Musée Royal del'Afrique Central, Tervuren, 336 pp.
- Kranz-Baltensperger Y (2011) The oonopid spider genus *Ischnothyreus* in Borneo (Oonopidae, Araneae). Zootaxa 2939: 1-49. <https://doi.org/10.11646/zootaxa.2939.1.1>
- Platnick NI, Berniker L, Kranz-Baltensperger Y (2012) The goblin spider genus *Ischnothyreus* (Araneae, Oonopidae) in the New World. American Museum Novitates 3759: 1-32. <https://doi.org/10.1206/3759.2>
- Tong YF, Sun XC, Li SQ, Bian DJ (2021) Taxonomic study of the genus *Ischnothyreus* (Araneae, Oonopidae) from Xishuangbanna Rainforest, southwestern China. ZooKeys 1034: 165-197. <https://doi.org/10.3897/zookeys.1034.63388>
- Tong YF, Bian DJ, Li SQ (2023) Three new species of the genus *Ischnothyreus* Simon, 1893 and the discovery of the male of *I. linzhiensis* Hu, 2001 from Tibet, China (Araneae, Oonopidae). ZooKeys 1152: 119-131. <https://doi.org/10.3897/zookeys.1152.100341>
- Ubick D, Dupérré N (2017) Oonopidae. In: Ubick D, Paquin P, Cushing P, Roth V (Eds) Spiders of North America: an Identification Manual. 2nd Edition. American Arachnological Society, Keene, New Hampshire, 181-182.
- WSC (2024) World Spider Catalog. Version 25.0. Natural History Museum Bern. <http://wsc.nmbe.ch>. Accessed on: 2024-1-28.