



Short Communication

First record of leopard-spotted goby *Thorogobius ephippiatus* (Gobiiformes, Gobiidae) above the Arctic circle

Vsevolod Rudyi[‡], Cesc Gordó-Vilaseca[‡]

[‡] Nord University, Bodø, Norway

Corresponding author: Vsevolod Rudyi (vsevolodrudyi@gmail.com)

Academic editor: Paramasivam Kodeeswaran

Received: 20 May 2024 | Accepted: 23 Jun 2024 | Published: 26 Jul 2024

Citation: Rudyi V, Gordó-Vilaseca C (2024) First record of leopard-spotted goby *Thorogobius ephippiatus* (Gobiiformes, Gobiidae) above the Arctic circle. Biodiversity Data Journal 12: e127963.

<https://doi.org/10.3897/BDJ.12.e127963>

Brief communication

The leopard-spotted goby *Thorogobius ephippiatus* (Lowe, 1839) distribution extends from the Western edge of the Mediterranean to the Eastern Atlantic, from Madeira to the North Sea (Costello 1992, Gerovasileiou et al. 2017, Stern et al. 2018). This species inhabits crevices and caves, and they tend to hide and withdraw at the minimum disturbance by divers or light (Holm and Mattson 1981), making its detection particularly challenging and typically restricted to observations by scuba divers (Costello 1992).

In the last decades, new records of this species have been reported in Sweden and Norway, but never North of 64°22'24" (Kovačić and Svensen 2018, Pethon 2005) (Fig. 1). These new records at northern latitudes could either be the result of previous lack of knowledge of this species range limit, or the result of climate warming induced northward expansion of its range, as has been observed for several other species, including the Atlantic Mackerel (*Scomber scombrus*), a pelagic species, or the spotted dragonet (*Callionymus maculatus*), a demersal one (Gordó-Vilaseca et al. 2023). Here we document the first confirmed record of this species above the Arctic circle, with images from an individual observed during a recreational SCUBA dive in the eastern side of Saltfjorden, at 5 m depth. The coordinates of the place, where the fish was discovered, are 67°18'19.3464" N, 14°43'5.52" E (Fig. 2). The fish was observed in the upper subtidal

zone, on a mixed sand and shell rubble bottom, hiding under a large overhanging rock. This micro-habitat is typical for this cryptobenthic species (Costello 1992). This find proves a wider and more northern extent than previously considered, and places the leopard-spotted goby as the seventh species of goby living above the Arctic circle (Kovačić and Svensen 2019).




Figure 1. [doi](#)

Global records of *Thorogobius ephippiatus* (GBIF.org 2024). Green dots correspond to records in GBIF. Blue dots are the observations, not in GBIF, from (Gerovasileiou et al. 2017) and (Stern et al. 2018). The red dot represents our observation, the first one above the Arctic circle (black dashed line).

The morphological and coloration evidences should be provided if the record and identification are based on morphology and coloration following the recommended best practice for reporting new fish species records (Bello et al. 2014). We document this record with pictures, and without collecting specimens, because this species' coloration and general shape are sufficient for species identification from photographs of live specimens (Kovačić et al. 2022). Diagnosis by coloration of alive individuals: Head and body of alive specimens in water whitish, pale fawn to sandy or bluish grey (which is the case for this particular individual). Black, brown orange to brown purple blotches and spots on head and body, the largest along lateral midline and along dorsal fin bases, smaller ones on predorsal area and on head. We have accurately checked available regional fish diversity catalogues (Mecklenburg et al. 2018) and global and regional databases (GBIF.org 2024, Norwegian Biodiversity Information Centre and (2024) 2024 and OBIS 2024). The record has been documented in Artsdatabanken which in turn makes the data available in GBIF (<https://www.artsobservasjoner.no/Sighting/34568198>). We have regularly dived in the area

over the past three years and have consulted with other scuba divers who photograph and document species in the area, and nobody had observed it before. This observation, roughly 450 km away from the previously northernmost record (Kovačić and Svensen 2018), suggests a recent range extension for the species, as has been happening for other marine fishes (Gordó-Vilaseca et al. 2023), and following the positive temperature trend of the last decades in the area (Fig. 3). Further records in regularly dived localities may confirm the spread of the species.



Figure 2. [doi](#)

Pictures taken during the dive by CGV. The individual was photographed in a crevice at ~5 m depth on the 9th and 10th of April 2024.




Figure 3. [doi](#)

Annual mean sea surface temperature (SST) between 1994 and 2022, at the latitude of the previous northernmost record for the species, and at the latitude of this record.

Acknowledgements

We thank Professor Mark J. Costello for providing constructive comments on the manuscript and providing an exciting and stimulating discussion about the finding, and Nord&Ne for showing us the amazing underwater world around Bodø and Saltstraumen.

Conflicts of interest

The authors have declared that no competing interests exist.

References

- Bello G, Causse R, Lipej L, Dulčić J (2014) A proposed best practice approach to overcome unverified and unverifiable “first records” in ichthyology. *Cybum* 38 (1): 9-14.
- Costello MJ (1992) Abundance and spatial overlap of gobies (Gobiidae) in Lough Hyne, Ireland. *Environmental Biology of Fishes* 33 (3): 239-248. <https://doi.org/10.1007/BF00005868>
- GBIF.org (2024) GBIF Occurrence Download. <https://doi.org/10.15468/dl.6gg9q7>
- Gerovasileiou V, Akel EH, Akyol O, Alongi G, Azevedo F, Babali N, Bakiu R, Bariche M, Bennoui A, Castriota L, Chintiroglou CC, Crocetta F, Deidun A, Galinou-Mitsoudi S, Giovos I, Gökoğlu M, Golemaj A, Hadjioannou L, Hartingerova J, Insacco G, Katsanevakis S, Kleitou P, Korun J, Lipej L, Malegue M, Michailidis N, Mouzai Tifoura A, Ovalis P, Petović S, Piraino S, Rizkalla SI, Rousou M, Savva I, Şen H, Spinelli A, Vougioukalou KG, Xharahi E, Zava B, Zenetos A (2017) New mediterranean biodiversity records (July 2017). *Mediterranean Marine Science* 18 (2): 355-384. <https://doi.org/10.12681/mms.13771>
- Gordó-Vilaseca C, Stephenson F, Coll M, Lavin C, Costello MJ (2023) Three decades of increasing fish biodiversity across the northeast Atlantic and the Arctic Ocean. *Proceedings of the National Academy of Sciences of the United States of America* 120 (4). <https://doi.org/10.1073/pnas.2120869120>
- Holm T, Mattson S (1981) *Thorogobius ephippiatus* (Pisces) found on the west coast of Sweden. *Sarsia* 66 (1): 87-88. <https://doi.org/10.1080/00364827.1981.10414526>
- Kovačić M, Svensen R (2018) The confirmed and continuous northern distribution of *Thorogobius ephippiatus* (Teleostei: Gobiidae) with the scientific use of recreational fishing data. *Journal of Applied Ichthyology* 34 (3): 691-693. <https://doi.org/10.1111/jai.13584>
- Kovačić M, Svensen R (2019) Northern extension of *Lesueurigobius friesii* (Malm, 1874) (Pisces: Gobiidae) distribution and the gobiid diversity decline along the Norwegian coast. *Acta Adriatica* 60 (2): 147-156. <https://doi.org/10.32582/aa.60.2.4>
- Kovačić M, Renoult JP, Pillon R, Svensen R, Bogorodsky SV, Engin S, Louisy P (2022) Identification of Mediterranean marine gobies (Actinopterygii: Gobiidae) of the continental shelf from photographs of in situ individuals. *Zootaxa* 5144 (1): 1-103. <https://doi.org/10.11646/zootaxa.5144.1.1>
- Mecklenburg CW, Lynghammer A, Johannesen E, Byrkjedal I, Christiansen J, Dolgov A, Karamushko O, Mecklenburg TA, Møller PR, Steinke D, Wienerroither R (2018) Marine

fishes of the Arctic region. II. Conservation of Arctic Flora and Fauna [ISBN 978-9935-431-69-1]

- Norwegian Biodiversity Information Centre, (2024) HS (2024) Norwegian Biodiversity Information Centre - Other datasets. Version 13.281. The Norwegian Biodiversity Information Centre (NBIC). Occurrence dataset accessed via GBIF.org on 2024-05-16. <https://doi.org/10.15468/tm56sc>
- OBIS (2024) Ocean Biodiversity Information System.
- Pethon P (2005) Aschehougs store fiskebok: Norges fisker i farger. Aschehoug, 486 pp. [ISBN 9788203232473]
- Stern N, Weissman A, Makovsky Y (2018) East and deep: Range extension and depth record for the leopard-spotted goby *Thorogobius ephippiatus* (Lowe, 1839) (Osteichthyes: Gobiidae). Journal of Applied Ichthyology 34 (3): 681-683. <https://doi.org/10.1111/jai.13625>