



# Diversity of parasitoid wasps (Insecta, Hymenoptera) in oilseed rape fields in Serbia

Milan Plećaš<sup>‡</sup>, Vladimir Žikić<sup>§</sup>, Korana Kocić<sup>‡</sup>, Jelisaveta Čkrkić<sup>‡,||</sup>, Anđeljko Petrović<sup>‡</sup>,  
Željko Tomanović<sup>‡</sup>

<sup>‡</sup> University of Belgrade, Faculty of Biology, Studentski Trg 16, 11000 Belgrade, Serbia

<sup>§</sup> University of Niš, Faculty of Sciences and Mathematics, Department of Biology with Ecology, Višegradska 33, P.O. Box 224, 18000, Niš, Serbia

| Centre for Biodiversity Genomics, University of Guelph, 50 Stone Road, N1G 2W1, Guelph, Ontario, Canada

Corresponding author: Milan Plećaš ([mplecas@bio.bg.ac.rs](mailto:mplecas@bio.bg.ac.rs))

Academic editor: Daniel Silva

Received: 26 Jul 2023 | Accepted: 27 Nov 2023 | Published: 05 Dec 2023

Citation: Plećaš M, Žikić V, Kocić K, Čkrkić J, Petrović A, Tomanović Ž (2023) Diversity of parasitoid wasps (Insecta, Hymenoptera) in oilseed rape fields in Serbia. Biodiversity Data Journal 11: e110118.

<https://doi.org/10.3897/BDJ.11.e110118>

## Abstract

## Background

Oilseed rape is an important crop grown worldwide and used for various purposes, including oil extraction and animal feed. In Europe, there are six major pest species and several other minor pests that can significantly affect oilseed rape production, requiring growers to effectively control them in order to ensure crop yield. The host-parasitoid complexes of these pests have been studied in detail and recorded mainly in western, central and northern Europe. As an abundant source of pollen and nectar, oilseed rape may also be attractive to other parasitoids that do not have direct trophic interactions with oilseed rape pest species. The aim of this study is to fill the knowledge gap regarding the wider parasitoid community in oilseed rape fields, particularly in southern Europe.

## New information

During the two-year study, a total of 3135 specimens of primary and secondary parasitoids were sampled, of which 2855 were found in oilseed rape fields and 280 in semi-natural habitats. We found 153 taxa, of which 119 were found in oilseed rape fields and 87 in semi-natural habitats. We identified 31 genera (33 species) as parasitoids of oilseed rape pests, 54 genera (97 species) parasitising non-pest species and 10 genera (23 species) as possible parasitoids of oilseed rape pests. This study shows that the parasitoid community in oilseed rape fields is very diverse and that includes parasitoids of both oilseed rape pest and non-pest species.

## Keywords

oilseed rape pests, host species, Balkans, ecological interactions, biological control agents

## Introduction

Oilseed rape (*Brassica napus* L.) is a very important crop, grown on more than 36 million hectares worldwide and more than 8 million hectares in Europe in 2021 (FAO 2022) and it is the third largest source of vegetable oil worldwide (Statista 2023). It is primarily used to extract oil from its seeds, which can be used for a variety of purposes, from food to making lubricants and soap, while the seed residue after oil extraction can also be used as animal feed (Alford 2003). In some countries, it is also an important break crop in cereal-dominated crop rotations (Angus et al. 2015). Similar to expanding global production, oilseed rape cultivation in Serbia increased from 5000–6000 ha at the end of the last century to more than 29,000 ha in 2022 (Sekulić and Kereši 2007, RZS 2023).

There are six major insect pest species of oilseed rape in Europe which growers often need to control to protect seed yield. In addition, there are a number of minor pests that may be of greater importance in some countries and at certain times of the year (Alford 2003, Williams 2010) (Table 1). Control of these pests is usually achieved through conventional pesticide use, but also through integrated pest management (IPM), as naturally occurring biological control agents, such as predators, parasitoids and pathogens can provide biocontrol and reduce pesticide use (Williams 2010). Parasitoids play particularly important role in pest control because their development is closely linked to their host species and show positive density-dependent patterns, while being specialised to one or few host species (Gunton and Pöyry 2016, Dainese et al. 2017). Higher parasitism rates (30–40%) can successfully suppress pest populations (Hokkanen 2008), although even lower rates (around 15%), especially when complemented with ground-dwelling predators, can also negatively affect pest abundance (Dainese et al. 2017). Semi-natural habitats that remain in agricultural landscapes are important for the conservation of these natural enemies, as they provide nectar resources, alternative host or prey species and overwintering sites (Landis et al. 2000, Bianchi et al. 2006) and their positive effect on

oilseed rape pest suppression has been demonstrated (Thies and Tschardt 1999, Dainese et al. 2017). However, oilseed rape itself can be a rich and highly attractive source of pollen and nectar for a wide range of insect species (Stanley et al. 2013), including parasitoids that use nectar as a food source (Heimpel and Jervis 2005, Varennes et al. 2016). Therefore, some parasitoid species may be present in oilseed rape fields without direct trophic interactions with known oilseed rape pests.

Table 1.

List of major and minor oilseed rape pests (Alford 2003, Williams 2010)

	Latin name	Common name	Family
Major oilseed rape pests			
1	<i>Brassicogethes</i> Audisio & Cline, 2009	pollen beetles	Nitidulidae
2	<i>Psylliodes chrysocephala</i> Linnaeus, 1758	cabbage stem flea beetle	Chrysomelidae
3	<i>Ceutorhynchus obstrictus</i> (Marsham, 1802)	cabbage seed weevil	Curculionidae
4	<i>Ceutorhynchus pallidactylus</i> (Marsham, 1802)	cabbage stem weevil	Curculionidae
5	<i>Ceutorhynchus napi</i> Gyllenhal, 1837	rape stem weevil	Curculionidae
6	<i>Dasineura brassicae</i> (Winnertz, 1853)	cabbage pod midge	Cecidomyiidae
Minor oilseed rape pests			
1	<i>Ceutorhynchus picitarsis</i> Gyllenhal, 1837	rape winter stem weevil	Curculionidae
2	<i>Phyllotreta nemorum</i> (Linnaeus, 1758)	cabbage flea beetle	Chrysomelidae
3	<i>Athalia rosae</i> (Linnaeus, 1758)	turnip sawfly	Tenthredinidae
4	<i>Brevicoryne brassicae</i> (Linnaeus, 1758)	cabbage aphid	Aphididae
5	<i>Myzus persicae</i> (Sulzer, 1776)	peach-potato aphid	Aphididae

Host-parasitoid complexes for the aforementioned major and some minor pests have been extensively studied and recorded (Alford 2003, Williams 2010), but mainly based on data from two projects: BORIS (1997-2000, CT-96-1314) and MASTER (2001-2005, QLK5-CT-2001-01447), focusing mainly on western, central and northern Europe. Data from southern Europe and Serbia, in particular, have been largely absent. To the best of our knowledge, there has not been a comprehensive list of parasitoids and hyperparasitoids that occur in oilseed rape fields for this region. Todorov et al. (2022) provide a list of parasitoid species found in oilseed rape fields in Bulgaria with comments on perspectives for biological control, but only for the family *Pteromalidae*. Oilseed rape pests, on the other hand, are relatively well studied in Serbian fields from both taxonomical and traditional chemical management aspects (Kereši et al. 2007, Sekulić and Kereši 2007, Štrbac et al. 2007, Mitrović et al. 2008, Milovac et al. 2010, Graora et al. 2013, Milovanović et al. 2013, Graora et al. 2015, Sivčev et al. 2015, Sivčev et al. 2016, Milovac 2016, Marjanović and Prodanović 2021). Similarly, the parasitoid wasp fauna of other crops in Serbia, especially cereals and alfalfa, has also been well studied (Tomanović et al. 2003, Kavallieratos et al. 2004, Kavallieratos et al. 2005a, Kavallieratos et al. 2005b, Tomanović et al. 2008, Žikić et al. 2012, Tomanović et al. 2021), but currently, there is a gap in knowledge of the parasitoid

wasp fauna in oilseed rape. Therefore, we aim here to explore the entire parasitoid community in oilseed rape to better understand the trophic interactions and the wider potential for biological control in agricultural landscapes.

## Materials and methods

Sampling was conducted in two consecutive years with a similar sampling design. In 2018, we selected five oilseed rape fields in the southern part of the Bačka Region (Serbia) (Table 2, Fig. 1). A single sampling transect of 20 m length was established in each field. Two fields had adjacent herbaceous semi-natural habitats where an additional sampling transect was established. In 2019, four oilseed rape fields were selected in the northern part of the Bačka Region, each of which had an adjacent herbaceous or shrubland semi-natural habitat. Two sampling transects were established in each field, one near the field edge (2 m from the field edge) and the second deeper in the field (20 m from the field edge) and one transect was established in the adjacent semi-natural habitat. Two fields had a second nearby semi-natural habitat and one additional sampling transect was established in each of them.

Table 2.

Coordinates of the sampled oilseed rape fields.

	South Bačka region					North Bačka region			
	Čenej			Srbobran		Bajmok	Đurđin	Mišićevo	Pačir
<b>Latitude</b>	45.3241	45.3468	45.3296	45.5216	45.5501	45.9516	45.9631	45.9892	45.8936
<b>Longitude</b>	19.8481	19.8615	19.8313	19.7571	19.7371	19.4771	19.5129	19.5126	19.4088

Parasitoids and hyperparasitoids were collected multiple times using different methods during the period from mid-April to mid-June, i.e. from the beginning of the oilseed rape flowering to the pre-harvest stage of pod maturation. Along each transect, 20 sweeps were made at the top of the vegetation with a sweep net and the collected samples were immediately stored in 96% ethanol. Sweep-net samples were collected seven times in 2018 and five times in 2019. In addition, during oilseed rape flowering phase, three clusters of pan traps were placed along each transect, at the beginning, middle and at the end of the transect. Each cluster consisted of three coloured water-filled bowls (yellow, white and blue) that were placed on the ground when surrounding vegetation was low or mounted on a pole and raised to the top of the vegetation when vegetation was tall. Traps were left active for 72 hours in 2018 and 48 hours in 2019, after which samples were placed in 96% ethanol. In both years, pan-trap sampling was done three times, once in late April and twice in early May in 2018 and three times in mid- to late April in 2019. During each field visit, oilseed rape plants and other plants were systematically examined for the presence of aphids along each established transect, both in crop fields and in semi-natural habitats. If aphid colonies were found, they were placed in a mesh-top plastic box for aphid parasitoids rearing. After four weeks, emerged parasitoids were transferred to 96% ethanol. In addition, in 2019 during pod maturation phase, we collected up to 100 oilseed

rape pods per field transect and stored them in a mesh-top plastic box to allow the emergence of pod midge parasitoids, which were then transferred to 96% ethanol. All collected parasitoids were identified using morphological identification keys (Suppl. material 1) to the lowest possible taxonomic level (species, genus or family). In case of identification to higher taxonomic levels, taxa were characterised and organised into different morphospecies (e.g. *Eulophidae* sp. 1, *Eulophidae* sp. 2 etc.). Voucher specimens are deposited in the collection of the Institute of Zoology, Faculty of Biology, University of Belgrade, Serbia. To establish consistency of occurrence of parasitoids in oilseed rape fields, we selected taxa/genera, based on two combined criteria across two years of sampling: 1) present in at least three fields (out of nine total) and 2) present with at least 10 individuals. We constructed species accumulation curves to assess completeness of our fauna survey in oilseed rape fields. We defined each combination of locality and sampling period as a “sampling event” which consisted of combined data from all sampling methods. To estimate extrapolated species richness for our study, we used Chao, First order Jackknife (Jack1), Second order Jackknife (Jack2) and Bootstrap estimates from the R package *vegan* (v.2.6-4; Oksanen et al. (2022)) within R Statistical Software (v.4.3.1; R Core Team (2023)).



Figure 1. [doi](#)

Map of the study area.

## Checklist of parasitoid wasps found in oilseed rape fields and adjacent semi-natural habitats in Serbia

### Family Bethylidae

#### *Goniozus claripennis* (Förster, 1851)

##### Material

- a. country: Serbia; locality: Đurđin; samplingProtocol: Pan traps; eventDate: 22-24.04.2019; habitat: oilseed rape; individualCount: 1 male; behavior: primary parasitoids, larval; occurrenceID: 2100E2C9-88FF-50B0-96C1-065E02A65373

Parasite of: Tortricidae

Notes: oilseed rape pest host: unknown

#### *Plastanoxus westwoodi* (Kieffer, 1914)

##### Material

- a. country: Serbia; locality: Čenej; samplingProtocol: Pan traps; eventDate: 24-27.04.2018; habitat: oilseed rape; individualCount: 1 male; behavior: primary parasitoids, larval/pupal; occurrenceID: 69C5F48C-FA04-5E14-BD6A-6DCAFF0DB2BE

Parasite of: Cucujidae

Notes: oilseed rape pest host: unknown

### Family Braconidae

#### *Apanteles* sp. 1

##### Material

- a. country: Serbia; locality: Čenej, Mišičevo, Pačir; samplingProtocol: Sweep net; eventDate: 07.05.2018, 10.05.2018, 25.05.2018, 18.04.2019, 12.06.2019; habitat: oilseed rape, semi-natural habitat; individualCount: 8 males, 5 females; behavior: primary parasitoids, larval; occurrenceID: 670AD6EE-22E6-5A5C-8D72-C0625E726E99

Parasite of: Lepidoptera

Notes: oilseed rape pest host: unknown

***Aphidius ervi* Haliday, 1834****Material**

- a. country: Serbia; locality: Bajmok, Čenej, Đurđin, Srbobran; samplingProtocol: Sweep net, Pan traps, Aphid colony; eventDate: 24-27.04.2018, 04-07.05.2018, 07-10.05.2018, 25.05.2018, 22-24.04.2019, 23.05.2019; habitat: oilseed rape, semi-natural habitat; individualCount: 215 males, 60 females; behavior: primary parasitoids, larval; occurrenceID: C5B6BE40-91B1-5621-AC91-D69A68728B55

**Parasite of:** *Myzus persicae*

**Notes:** oilseed rape pest host: yes

***Aphidius matricariae* Haliday, 1834****Material**

- a. country: Serbia; locality: Čenej, Srbobran; samplingProtocol: Aphid colony; eventDate: 04.05.2018, 07.05.2018, 10.05.2018; habitat: oilseed rape; individualCount: 72 males, 40 females; behavior: primary parasitoids, larval; occurrenceID: 3950E1EC-D843-5AB9-B17F-937164B61976

**Parasite of:** *Myzus persicae*

**Notes:** oilseed rape pest host: yes

***Aphidius* sp. 1****Material**

- a. country: Serbia; locality: Đurđin; samplingProtocol: Aphid colony; eventDate: 05-23-19; habitat: oilseed rape; individualCount: 1 male, 1 female; behavior: primary parasitoids, larval; occurrenceID: 3E6F63DA-DB7D-5F1D-A55F-07289769670F

**Parasite of:** *Myzus persicae*

**Notes:** oilseed rape pest host: unknown, possible

***Binodoxys angelicae* (Haliday, 1833)****Material**

- a. country: Serbia; locality: Mišićevo; samplingProtocol: Aphid colony; eventDate: 05-23-19; habitat: oilseed rape; individualCount: 1 male; behavior: primary parasitoids, larval; occurrenceID: EBAA0EF7-3F19-5404-9F10-5282D552F36F

**Parasite of:** *Myzus persicae*

**Notes:** oilseed rape pest host: yes

***Blacus nigricornis* Haeselbarth, 1973****Material**

- a. country: Serbia; locality: Čenej, Srbobran; samplingProtocol: Sweep net, Pan traps; eventDate: 27.04.2018, 07-10.05.2018; habitat: oilseed rape, semi-natural habitat; individualCount: 2 males, 1 female; behavior: primary parasitoids, larval; occurrenceID: 92167F97-BDA3-57A5-9DB1-203EF3FAB78D

**Parasite of:** *Meligethes aeneus*

**Notes:** oilseed rape pest host: yes

***Bracon picticornis* (Wesmael, 1838)****Material**

- a. country: Serbia; locality: Bajmok; samplingProtocol: Pan traps; eventDate: 17-19.04.2019; habitat: oilseed rape; individualCount: 1 male; behavior: primary parasitoids, larval; occurrenceID: BD9DB36F-E474-5431-B537-EF01E6550C2E

**Parasite of:** *Meligethes aeneus*

**Notes:** oilseed rape pest host: yes

***Bracon variator* Nees, 1811****Material**

- a. country: Serbia; locality: Čenej; samplingProtocol: Sweep net, Pan traps; eventDate: 04-07.05.2018, 10.05.2018; habitat: oilseed rape; individualCount: 2 males; behavior: primary parasitoids, larval; occurrenceID: 76500E7C-7739-54B4-9FB6-1460E32CCBB5

**Parasite of:** *Ceutorhynchus assimilis*

**Notes:** oilseed rape pest host: yes

***Chelonus oculator* (Fabricius, 1775)****Material**

- a. country: Serbia; locality: Srbobran; samplingProtocol: Pan traps; eventDate: 04-07.05.2018; habitat: semi-natural habitat; individualCount: 1 male; behavior: primary parasitoids, egg/larval; occurrenceID: 981653E9-B91B-5795-8BE8-910E33C9FD2C

**Parasite of:** Lepidoptera, *Spodoptera exigua*

**Notes:** oilseed rape pest host: unknown



***Choeras parasitellae* (Bouché, 1834)****Material**

- a. country: Serbia; locality: Čenej; samplingProtocol: Sweep net; eventDate: 05-10-18; habitat: oilseed rape; individualCount: 1 female; behavior: primary parasitoids, larval; occurrenceID: A0968BCD-EB9E-5A04-A16E-734D2F88B3E8

**Parasite of:** Lepidoptera

**Notes:** oilseed rape pest host: unknown

***Cotesia glomerata* (Linnaeus, 1758)****Material**

- a. country: Serbia; locality: Srbobran; samplingProtocol: Sweep net, Pan traps; eventDate: 27.04.2018, 04-07.05.2018, 25.05.2018; habitat: oilseed rape; individualCount: 2 males, 1 female; behavior: primary parasitoids, larval; occurrenceID: 5DA837C7-FAF4-5E5D-BBAF-2679FD8A7656

**Parasite of:** *Pieris* spp.

**Notes:** oilseed rape pest host: yes

***Cotesia vestalis* (Haliday, 1834)****Material**

- a. country: Serbia; locality: Pačir; samplingProtocol: Sweep net; eventDate: 05-10-19; habitat: semi-natural habitat; individualCount: 1 female; behavior: primary parasitoids, larval; occurrenceID: D9CFE5F0-9B48-5FAB-887A-A19948176C6C

**Parasite of:** *Plutella xylostella*

**Notes:** oilseed rape pest host: yes

***Dacnusa* sp. 1****Material**

- a. country: Serbia; locality: Čenej; samplingProtocol: Pan traps; eventDate: 04-07.05.2018; habitat: oilseed rape; individualCount: 1 female; behavior: primary parasitoids, larval; occurrenceID: 9D21083D-9666-5D29-9B82-B599A478426B

**Parasite of:** *Phytomyza rufipes*?

**Notes:** oilseed rape pest host: unknown

***Diaeretiella rapae* (McIntosh, 1855)****Material**

- a. country: Serbia; locality: Bajmok, Čenej, Đurđin, Mišičevo, Pačir, Srbobran; samplingProtocol: Sweep net, Pan traps, Aphid colony; eventDate: 24-27.04.2018, 04-07.05.2018, 07-10.05.2018, 25.05.2018, 17-19.04.2019, 22-24.04.2019, 08.05.2019, 10.05.2019, 23.05.2019, 12.06.2019, 13.06.2019; habitat: oilseed rape, semi-natural habitat; individualCount: 816 males, 427 females; behavior: primary parasitoids, larval; occurrenceID: 9B75C3AA-8CF9-5C70-A0AD-910DE1A7E9EA

**Parasite of:** *Myzus persicae*

**Notes:** oilseed rape pest host: yes

***Diospilus capito* (Nees, 1834)****Material**

- a. country: Serbia; locality: Srbobran; samplingProtocol: Sweep net; eventDate: 04-27-18; habitat: semi-natural habitat; individualCount: 1 female; behavior: primary parasitoids, larval; occurrenceID: 38221A3B-0E37-569C-8E40-2F3A41611E99

**Parasite of:** *Meligethes aeneus*

**Notes:** oilseed rape pest host: yes

***Ephedrus persicae* Froggatt, 1904****Material**

- a. country: Serbia; locality: Čenej; samplingProtocol: Aphid colony; eventDate: 05-10-18; habitat: oilseed rape; individualCount: 1 male; behavior: primary parasitoids, larval; occurrenceID: 894A5022-E488-56DD-8F1B-563D4D8E8E23

**Parasite of:** *Myzus persicae*

**Notes:** oilseed rape pest host: yes

***Eubazus sigalphoides* (Marshall, 1889)****Material**

- a. country: Serbia; locality: Čenej, Srbobran; samplingProtocol: Sweep net, Pan traps, Aphid colony; eventDate: 24-27.04.2018, 04.05.2018, 07.05.2018, 10.05.2018, 25.05.2018, 27.04.2018; habitat: oilseed rape; individualCount: 44 males, 8 females; behavior: primary parasitoids, larval; occurrenceID: AD0C09B2-CCD1-5DFD-882F-0BE15DE7EDE2

**Parasite of:** *Meligethes aeneus*

Notes: oilseed rape pest host: yes

### ***Eubazus* sp. 1**

#### **Material**

- a. country: Serbia; locality: Srbobran; samplingProtocol: Sweep net; eventDate: 27.04.2018, 04.05.2018; habitat: oilseed rape, semi-natural habitat; individualCount: 1 male, 1 female; behavior: primary parasitoids, larval; occurrenceID: C0808E15-FC75-5133-9529-253FC39B21F3

Parasite of: Curculionidae, *Pissodes* spp.

Notes: oilseed rape pest host: unknown, possible

### ***Habrobracon hebetor* (Say, 1836)**

#### **Material**

- a. country: Serbia; locality: Čenej; samplingProtocol: Sweep net, Pan traps; eventDate: 27.04.2018, 04-07.05.2018, 07-10.05.2018; habitat: oilseed rape; individualCount: 3 males; behavior: primary parasitoids, larval; occurrenceID: D172A12C-83BF-5A6C-882D-B89A6B2A56B7

Parasite of: *Plutella xylostella*

Notes: oilseed rape pest host: yes

### ***Lysiphlebus fabarum* (Marshall, 1896)**

#### **Material**

- a. country: Serbia; locality: Pačir, Srbobran; samplingProtocol: Pan traps, Aphid colony; eventDate: 07-10.05.2018, 23.05.2019; habitat: semi-natural habitat; individualCount: 7 males, 8 females; behavior: primary parasitoids, larval; occurrenceID: 5E911AD9-61F7-5767-82D5-2C5F9C00A3CC

Parasite of: *Myzus persicae*

Notes: oilseed rape pest host: yes

### ***Microctonus* sp. 1**

#### **Material**

- a. country: Serbia; locality: Čenej; samplingProtocol: Sweep net; eventDate: 04-27-18; habitat: oilseed rape; individualCount: 1 female; behavior: primary parasitoids, adult; occurrenceID: 2C6284BF-9E5E-5350-9DC9-0302DA217FF7

Parasite of: *Psylliodes chrysocephala*

Notes: oilseed rape pest host: unknown, possible

### *Microctonus* sp. 2

#### Material

- a. country: Serbia; locality: Čenej; samplingProtocol: Sweep net; eventDate: 05-10-18; habitat: semi-natural habitat; individualCount: 1 male; behavior: primary parasitoids, adult; occurrenceID: 5867C306-6324-5480-92E0-45222448E6D9

Parasite of: *Psylliodes chrysocephala*

Notes: oilseed rape pest host: unknown, possible

### *Microplitis* sp. 1

#### Material

- a. country: Serbia; locality: Čenej; samplingProtocol: Sweep net; eventDate: 07.05.2018, 10.05.2018, 25.05.2018; habitat: oilseed rape, semi-natural habitat; individualCount: 4 males; behavior: primary parasitoids, larval; occurrenceID: DAB4FA8A-9140-5590-AC18-0A40980767C9

Parasite of: Lepidoptera

Notes: oilseed rape pest host: unknown

### *Peristenus* sp. 1

#### Material

- a. country: Serbia; locality: Mišićevo, Pačir; samplingProtocol: Sweep net; eventDate: 25.04.2019, 10.05.2019; habitat: semi-natural habitat; individualCount: 2 males; behavior: primary parasitoids, adult; occurrenceID: FF098CFC-54DE-59E1-8DEF-F0B13AA0C33C

Parasite of: Hemiptera, Miridae

Notes: oilseed rape pest host: unknown

### *Peristenus* sp. 2

#### Material

- a. country: Serbia; locality: Pačir; samplingProtocol: Sweep net; eventDate: 05-10-19; habitat: semi-natural habitat; individualCount: 1 male; behavior: primary parasitoids, adult; occurrenceID: 134DA297-FDA0-5920-8599-D9951B634070

Parasite of: Hemiptera, Miridae

Notes: oilseed rape pest host: unknown

***Praon volucre* (Haliday, 1933)****Material**

- a. country: Serbia; locality: Čenej, Srbobran; samplingProtocol: Sweep net, Pan traps, Aphid colony; eventDate: 27.04.2018, 04-07.05.2018, 10.05.2018; habitat: oilseed rape, semi-natural habitat; individualCount: 17 males, 12 females; behavior: primary parasitoids, larval; occurrenceID: 87BA73CA-4D26-5BD6-957F-D51999F207CC

**Parasite of:** *Myzus persicae*

**Notes:** oilseed rape pest host: yes

***Schizoprymnus obscurus* (Nees, 1816)****Material**

- a. country: Serbia; locality: Čenej, Mišičevo; samplingProtocol: Sweep net, Aphid colony; eventDate: 20.04.2018, 10.05.2018, 12.06.2019; habitat: oilseed rape, semi-natural habitat; individualCount: 6 males; behavior: primary parasitoids, larval; occurrenceID: E16619A8-405D-5DB5-865C-9A40D688C840

**Parasite of:** *Ceutorhynchus* spp.

**Notes:** oilseed rape pest host: yes

***Townesilitus bicolor* (Wesmael, 1835)****Material**

- a. country: Serbia; locality: Čenej, Pačir, Srbobran; samplingProtocol: Sweep net, Pan traps; eventDate: 24-27.04.2018, 07.05.2018, 10.05.2018, 25.05.2018, 10.05.2019; habitat: oilseed rape, semi-natural habitat; individualCount: 29 males, 4 females; behavior: primary parasitoids, adult; occurrenceID: 6C1B44DA-6B18-5F2A-AA95-117FA9C8DC30

**Parasite of:** *Phyllotreta* spp.

**Notes:** oilseed rape pest host: yes

***Triaspis thoracica* (Curtis, 1860)****Material**

- a. country: Serbia; locality: Čenej, Srbobran; samplingProtocol: Sweep net; eventDate: 27.04.2018, 07.05.2018, 10.05.2018, 25.05.2018; habitat: oilseed rape; individualCount: 17 males, 53 females; behavior: primary parasitoids, larval; occurrenceID: 31AF200E-A505-5FF0-BE16-9F8C0D660AAC

**Parasite of:** Chrysomelidae, *Bruchus* spp.

**Notes:** oilseed rape pest host: unknown

## Family Ceidae

### *Cea pulicaris* Walker, 1837

#### Material

- a. country: Serbia; locality: Srbobran; samplingProtocol: Pan traps; eventDate: 07-10.05.2018; individualCount: 1 male; behavior: primary parasitoids, larval; occurrenceID: 8C00D2C6-6FA5-5B26-9D99-ADDCC9E1D372

**Parasite of:** Agromyzidae, *Phytomyza* spp.

**Notes:** oilseed rape pest host: unknown

### *Spalangiopelta* sp. 1

#### Material

- a. country: Serbia; locality: Čenej, Đurđin, Mišićevo, Srbobran; samplingProtocol: Sweep net, Pan traps; eventDate: 27.04.2018, 04.05.2018, 07.05.2018, 25.05.2018, 22-24.04.2019, 25.04.2019, 12.06.2019; habitat: oilseed rape, semi-natural habitat; individualCount: 8 males, 4 females; behavior: primary parasitoids, larval; occurrenceID: E6ABC350-A923-5C28-BBBA-2465C9111481

**Parasite of:** Agromyzidae, Drosophilidae (miners)

**Notes:** oilseed rape pest host: unknown

## Family Ceraphronidae

### Ceraphronidae sp. 1

#### Material

- a. country: Serbia; locality: Čenej, Srbobran; samplingProtocol: Sweep net, Pan traps; eventDate: 24-27.04.2018, 07-10.05.2018, 25.05.2018; habitat: oilseed rape, semi-natural habitat; individualCount: 2 males, 1 female; behavior: primary parasitoids, larval; occurrenceID: 6977C9C7-9469-5D55-9EB8-7B56EF27769E

**Parasite of:** Cecidomyiidae, Hemiptera, Neuroptera, Thysanoptera

**Notes:** oilseed rape pest host: unknown

**Ceraphronidae sp. 2****Material**

- a. country: Serbia; locality: Đurđin; samplingProtocol: Pan traps; eventDate: 22-24.04.2019; habitat: semi-natural habitat; individualCount: 1 male; behavior: primary parasitoids, larval; occurrenceID: 3C03AC9C-9FF1-5178-9F37-501881047C03

**Parasite of:** Cecidomyiidae, Hemiptera, Neuroptera, Thysanoptera

**Notes:** oilseed rape pest host: unknown

***Aphanogmus abdominalis* (Thomson, 1858)****Material**

- a. country: Serbia; locality: Čenej, Srbobran; samplingProtocol: Sweep net, Pan traps; eventDate: 24-27.04.2018, 04-07.05.2018, 07-10.05.2018, 13.06.2018; habitat: oilseed rape, semi-natural habitat; individualCount: 9 males, 16 females; behavior: primary parasitoids, larval; occurrenceID: 67963D58-F8DD-5A8E-9A48-9100C67EC5F9

**Parasite of:** *Dasineura brassicae*

**Notes:** oilseed rape pest host: yes

***Ceraphron* sp. 1****Material**

- a. country: Serbia; locality: Srbobran; samplingProtocol: Pan traps; eventDate: 04-07.05.2018; habitat: semi-natural habitat; individualCount: 1 male; behavior: primary parasitoids, larval; occurrenceID: D3738836-AF98-56D5-8C8F-4906B24752AD

**Parasite of:** Cecidomyiidae, Hemiptera, Neuroptera, Thysanoptera

**Notes:** oilseed rape pest host: unknown

***Ceraphron* sp. 2****Material**

- a. country: Serbia; locality: Čenej, Đurđin; samplingProtocol: Pan traps; eventDate: 24-27.04.2018, 22-24.04.2019; habitat: oilseed rape; individualCount: 3 males; behavior: primary parasitoids, larval; occurrenceID: 4F1DC248-EF72-5FB1-9F45-07BCA5B0BBD8

**Parasite of:** Cecidomyiidae, Hemiptera, Neuroptera, Thysanoptera

**Notes:** oilseed rape pest host: unknown

**Family Chalcididae**

## ***Brachymeria tibialis*-group Steffan, 1958**

### **Material**

- a. country: Serbia; locality: Đurđin, Srbobran; samplingProtocol: Sweep net, Pan traps; eventDate: 24-27.04.2018, 24.04.2019; habitat: oilseed rape, semi-natural habitat; individualCount: 2 males; behavior: primary/secondary parasitoids, larval/pupal; occurrenceID: BB4DEADB-F72A-5BD6-B3F9-A220BDF54A02

**Parasite of:** Lepidoptera, Hymenoptera, Diprionidae, Diptera, Cecidomyiidae

**Notes:** oilseed rape pest host: unknown

## **Family Diapriidae**

### ***Lyteba* sp. 1**

#### **Material**

- a. country: Serbia; locality: Bajmok, Pačir, Srbobran; samplingProtocol: Sweep net, Pan traps; eventDate: 04-07.05.2018, 07-10.05.2018, 25.05.2018, 22-24.04.2019; habitat: oilseed rape, semi-natural habitat; individualCount: 10 males, 6 females; behavior: primary parasitoids, larval/pupal; occurrenceID: 7103B83E-8E05-5365-A4AE-3F10EDABCE1D

**Parasite of:** Diptera, Mycetophilidae, Sciaridae

**Notes:** oilseed rape pest host: unknown

### ***Trichopria* sp. 1**

#### **Material**

- a. country: Serbia; locality: Srbobran; samplingProtocol: Pan traps; eventDate: 24-27.04.2018; habitat: semi-natural habitat; individualCount: 1 male; behavior: primary parasitoids, larval/pupal; occurrenceID: 08A7A3D2-D256-5D3E-810C-1369C2BE9D0A

**Parasite of:** Drosophilidae, Sarcophagidae, Sepsidae, Muscidae, Calliphoridae

**Notes:** oilseed rape pest host: unknown



## Family Encyrtidae

### Encyrtidae sp. 1

#### Material

- a. country: Serbia; locality: Čenej; samplingProtocol: Pan traps; eventDate: 07-10.05.2018; habitat: oilseed rape; individualCount: 1 male; behavior: primary/secondary parasitoids, egg/larval; occurrenceID: 665B033C-6A98-58E7-84CD-7A1EAB4A40C9

**Parasite of:** Hemiptera, Homoptera, Coccoidea, Acarina

**Notes:** oilseed rape pest host: unknown

### Encyrtidae sp. 2

#### Material

- a. country: Serbia; locality: Đurđin; samplingProtocol: Pan traps; eventDate: 22-24.04.2019; habitat: semi-natural habitat; individualCount: 1 female; behavior: primary/secondary parasitoids, egg/larval; occurrenceID: B44CF365-920E-5918-B950-D845517334A2

**Parasite of:** Hemiptera, Homoptera, Coccoidea, Acarina

**Notes:** oilseed rape pest host: unknown

### Encyrtidae sp. 3

#### Material

- a. country: Serbia; locality: Mišićevo; samplingProtocol: Sweep net; eventDate: 04-25-19; habitat: oilseed rape, semi-natural habitat; individualCount: 2 females; behavior: primary/secondary parasitoids, egg/larval; occurrenceID: 329554B8-9214-5768-AC2A-A1C0CA3E1A72

**Parasite of:** Hemiptera, Homoptera, Coccoidea, Acarina

**Notes:** oilseed rape pest host: unknown

### *Anagyrus* sp. 1

#### Material

- a. country: Serbia; locality: Pačir; samplingProtocol: Pan traps; eventDate: 22-24.04.2019; habitat: semi-natural habitat; individualCount: 1 male, 2 females; behavior: primary parasitoids, egg; occurrenceID: EFEB1B0D-9452-5C73-960B-CEF3BB2D0609

**Parasite of:** Hemiptera, Pseudococcidae?

**Notes:** oilseed rape pest host: unknown

***Copidosoma bakeri* (Howard, 1898)****Material**

- a. country: Serbia; locality: Čenej, Pačir, Srbobran; samplingProtocol: Sweep net, Pan traps; eventDate: 04-07.05.2018, 07-10.05.2018, 22-24.04.2019; habitat: oilseed rape, semi-natural habitat; individualCount: 12 males, 1 female; behavior: primary parasitoids, egg/larval; occurrenceID: A39B9228-A151-5F73-AF4B-1EEB2059E5C4

**Parasite of:** Noctuidae (*Euxoa auxiliaris*)

**Notes:** oilseed rape pest host: unknown

***Eugahania fumipennis* (Ratzeburg, 1852)****Material**

- a. country: Serbia; locality: Bajmok; samplingProtocol: Pan traps; eventDate: 22-24.04.2019; habitat: semi-natural habitat; individualCount: 1 male; behavior: primary parasitoids, larval; occurrenceID: EAC5B5D2-1D84-55D3-916E-4B86FBA7F964

**Parasite of:** Cicadellidae, *Macropsis vicina*

**Notes:** oilseed rape pest host: unknown

***Metaphycus flavus* (Ashmead, 1901)****Material**

- a. country: Serbia; locality: Mišičevo, Srbobran; samplingProtocol: Sweep net; eventDate: 07.05.2018, 25.04.2019; habitat: oilseed rape; individualCount: 3 males; behavior: primary parasitoids, egg; occurrenceID: 762F7BCB-A08B-592F-943A-91081FB581A4

**Parasite of:** Hemiptera, Coccoidea

**Notes:** oilseed rape pest host: unknown

***Rhopus* sp. 1****Material**

- a. country: Serbia; locality: Đurđin; samplingProtocol: Pan traps; eventDate: 22-24.04.2019; habitat: semi-natural habitat; individualCount: 1 male; behavior: primary parasitoids, egg; occurrenceID: D3D8E152-A381-585A-BFB5-C1BC53F9DE00

**Parasite of:** Hemiptera, Pseudococcidae

**Notes:** oilseed rape pest host: unknown

## Family Eulophidae

### Eulophidae sp. 1

#### Material

- a. country: Serbia; locality: Mišičevo, Pačir; samplingProtocol: Sweep net; eventDate: 18.04.2019, 10.05.2019; habitat: oilseed rape, semi-natural habitat; individualCount: 2 males; behavior: primary/secondary parasitoids, egg/larval/pupal; occurrenceID: F9DCF374-157E-5B65-BD84-865CA67D29F5

**Parasite of:** Holometabolous insects

**Notes:** oilseed rape pest host: unknown

### Eulophidae sp. 2

#### Material

- a. country: Serbia; locality: Čenej; samplingProtocol: Sweep net, Pan traps; eventDate: 24-27.04.2018, 10.05.2018; habitat: oilseed rape, semi-natural habitat; individualCount: 1 male, 1 female; behavior: primary/secondary parasitoids, egg/larval/pupal; occurrenceID: A14BDA52-22D1-52AB-8ADD-D4687C566B92

**Parasite of:** Holometabolous insects

**Notes:** oilseed rape pest host: unknown

### Eulophidae sp. 3

#### Material

- a. country: Serbia; locality: Čenej, Mišičevo; samplingProtocol: Sweep net, Pan traps, Aphid colony; eventDate: 07-10.05.2018, 25.05.2018; habitat: oilseed rape, semi-natural habitat; individualCount: 7 males, 3 females; behavior: primary/secondary parasitoids, egg/larval/pupal; occurrenceID: CC0F692C-2F39-5777-B056-9BED5DFA170E

**Parasite of:** Holometabolous insects

**Notes:** oilseed rape pest host: unknown

### Eulophidae sp. 4

#### Material

- a. country: Serbia; locality: Mišičevo, Srbobran; samplingProtocol: Sweep net; eventDate: 07.05.2018, 12.06.2019; habitat: oilseed rape; individualCount: 2 males; behavior: primary/secondary parasitoids, egg/larval/pupal; occurrenceID: 33B3070D-A66C-5491-9238-C41CFD109A8F

**Parasite of:** Holometabolous insects

**Notes:** oilseed rape pest host: unknown

## **Eulophidae sp. 5**

### **Material**

- a. country: Serbia; locality: Bajmok; samplingProtocol: Sweep net; eventDate: 05-08-19; habitat: oilseed rape; individualCount: 1 male, 1 female; behavior: primary/secondary parasitoids, egg/larval/pupal; occurrenceID: 5B30C53D-6E6B-52CD-B264-53B03B03E44E

**Parasite of:** Holometabolous insects

**Notes:** oilseed rape pest host: unknown

## **Tetrastichinae sp. 1**

### **Material**

- a. country: Serbia; locality: Čenej, Srbobran; samplingProtocol: Sweep net, Pan traps; eventDate: 24-27.04.2018, 04-07.05.2018, 25.05.2018; habitat: oilseed rape, semi-natural habitat; individualCount: 15 males, 9 females; behavior: primary/secondary parasitoids, egg/larval/pupal; occurrenceID: 25F73A04-63DB-5BCC-8198-6D334C049E02

**Parasite of:** Holometabolous insects, spiders, mites, nematodes

**Notes:** oilseed rape pest host: unknown

## **Tetrastichinae sp. 2**

### **Material**

- a. country: Serbia; locality: Čenej, Srbobran; samplingProtocol: Sweep net, Pan traps; eventDate: 04-07.05.2018, 10.05.2018; habitat: oilseed rape; individualCount: 4 males; behavior: primary/secondary parasitoids, egg/larval/pupal; occurrenceID: 3B402071-358E-5202-9EE0-2FCB10998A76

**Parasite of:** Holometabolous insects, spiders, mites, nematodes

**Notes:** oilseed rape pest host: unknown

## ***Aprostocetus epicharmus* (Walker, 1839)**

### **Material**

- a. country: Serbia; locality: Mišićevo; samplingProtocol: Aphid colony; eventDate: 06-12-19; habitat: semi-natural habitat; individualCount: 2 males; behavior: primary parasitoids, larval; occurrenceID: DBE900AA-7874-5147-B369-EB8AE571ABD1

**Parasite of:** *Dasineura brassicae*

**Notes:** oilseed rape pest host: yes

### ***Aprostocetus* sp. 1**

#### **Material**

- a. country: Serbia; locality: Bajmok, Čenej, Mišičevo, Pačir, Srbobran; samplingProtocol: Sweep net, Aphid colony; eventDate: 07.05.2018, 10.05.2018, 25.05.2018, 25.04.2019, 08.05.2019; habitat: oilseed rape, semi-natural habitat; individualCount: 16 males, 1 female; behavior: primary parasitoids, larval; occurrenceID: 65C1DA0B-F6CE-5521-8C0E-697DB235729C

**Parasite of:** *Dasineura brassicae*

**Notes:** oilseed rape pest host: unknown, possible

### ***Diaulinopsis arenaria* (Erdős, 1951)**

#### **Material**

- a. country: Serbia; locality: Čenej, Srbobran; samplingProtocol: Sweep net, Pan traps; eventDate: 24-27.04.2018, 04-07.05.2018, 10.05.2018, 25.05.2018; habitat: oilseed rape, semi-natural habitat; individualCount: 4 males, 3 females; behavior: primary parasitoids, larval; occurrenceID: 31245DC2-09E3-5E20-AD85-28106E0475FC

**Parasite of:** *Liriomyza* spp.

**Notes:** oilseed rape pest host: unknown

### ***Diglyphus* aff. *isaea***

#### **Material**

- a. country: Serbia; locality: Srbobran; samplingProtocol: Pan traps; eventDate: 04-07.05.2018; habitat: oilseed rape; individualCount: 1 female; behavior: primary parasitoids, larval; occurrenceID: 77C33F6A-EF2C-5439-9474-63CAF51843D9

**Parasite of:** leaf miners

**Notes:** oilseed rape pest host: unknown

### ***Elasmus platyedrae* Ferrière, 1935**

#### **Material**

- a. country: Serbia; locality: Bajmok, Pačir; samplingProtocol: Pan traps; eventDate: 17-19.04.2019, 22-24.04.2019; habitat: semi-natural habitat; individualCount: 7 males;

behavior: primary/secondary parasitoids, larval; occurrenceID: 9EE8B34B-6E58-5C9F-A1CF-CAF99F0BE995

**Parasite of:** Gelechiidae

**Notes:** oilseed rape pest host: unknown

### ***Eulophus* sp. 1**

#### **Material**

- a. country: Serbia; locality: Bajmok, Čenej, Pačir, Srbobran; samplingProtocol: Sweep net; eventDate: 04.05.2018, 07.05.2018, 10.05.2018, 25.05.2018, 25.04.2019, 13.06.2019; habitat: oilseed rape, semi-natural habitat; individualCount: 8 males, 11 females; behavior: primary parasitoids, larval; occurrenceID: 67E92477-7F66-5525-8171-B91BF18A0BEC

**Parasite of:** Cabbage Seed Weevil, Lepidoptera

**Notes:** oilseed rape pest host: unknown, possible

### ***Necremnus* sp. 1**

#### **Material**

- a. country: Serbia; locality: Bajmok; samplingProtocol: Pan traps; eventDate: 17-19.04.2019; habitat: semi-natural habitat; individualCount: 1 female; behavior: primary parasitoids, larval; occurrenceID: 52828C98-25D5-5FAD-9603-BBEF1E42BD1D

**Parasite of:** Cabbage Seed Weevil, Lepidoptera

**Notes:** oilseed rape pest host: unknown, possible

### ***Omphale clypealis* (Thomson, 1878)**

#### **Material**

- a. country: Serbia; locality: Bajmok, Čenej, Pačir; samplingProtocol: Sweep net, Pan traps; eventDate: 24-27.04.2018, 24.04.2019, 25.04.2019; habitat: oilseed rape; individualCount: 3 males; behavior: primary parasitoids, larval; occurrenceID: 57F0A662-3BF1-5D85-B023-CCE77D6685A9

**Parasite of:** *Dasineura brassicae*

**Notes:** oilseed rape pest host: yes

## ***Pnigalio* sp. 1**

### **Material**

- a. country: Serbia; locality: Čenej, Pačir; samplingProtocol: Sweep net, Pan traps; eventDate: 24-27.04.2018, 25.04.2019; habitat: oilseed rape; individualCount: 1 male, 2 females; behavior: primary/secondary parasitoids, larval; occurrenceID: A73F7362-9DA0-512C-98E2-BA7EF36E0272

**Parasite of:** leaf miners: Lepidoptera, Diptera, Coleoptera, Hymenoptera

**Notes:** oilseed rape pest host: unknown, possible

## ***Tetrastichus* sp. 1**

### **Material**

- a. country: Serbia; locality: Čenej, Mišičevo, Srbobran; samplingProtocol: Sweep net, Pan traps, Aphid colony; eventDate: 24-27.04.2018, 04-07.05.2018, 07-10.05.2018, 25.05.2018, 25.04.2019; habitat: oilseed rape, semi-natural habitat; individualCount: 9 males, 8 females; behavior: primary parasitoids, larval/pupal; occurrenceID: E1CEB768-7D5D-5081-B997-CC5BD051F615

**Parasite of:** Buprestidae, Cerambycidae, Chrysomelidae, Curculionidae, Lepidoptera, Diptera, Hymenoptera

**Notes:** oilseed rape pest host: unknown, possible

## **Family Eurytomidae**

### ***Eurytoma* sp. 1**

#### **Material**

- a. country: Serbia; locality: Čenej, Mišičevo, Srbobran; samplingProtocol: Sweep net, Pan traps, Aphid colony; eventDate: 27.04.2018, 04-07.05.2018, 07-10.05.2018, 25.05.2018, 25.04.2019; habitat: oilseed rape, semi-natural habitat; individualCount: 4 males, 7 females; behavior: primary/secondary parasitoids, larval; occurrenceID: B8AA6924-7073-5B04-AFFD-EA1EA8F1323A

**Parasite of:** *Systole*, *Bruchophagus*, hyperparasitoid on *Tetramesa*

**Notes:** oilseed rape pest host: unknown, possible

## Family Figitidae

### Eucolinae sp. 1

#### Material

- a. country: Serbia; locality: Čenej, Đurđin, Srbobran; samplingProtocol: Sweep net, Pan traps; eventDate: 24-27.04.2018, 07-10.05.2018, 22-24.04.2019; habitat: oilseed rape, semi-natural habitat; individualCount: 8 males; behavior: primary parasitoids, larval/pupal; occurrenceID: 6E9D43A2-09E0-53E1-9353-764D0E4F36C8

**Parasite of:** Cyclorraphic dipterous larvae

**Notes:** oilseed rape pest host: unknown

### Eucolinae sp. 2

#### Material

- a. country: Serbia; locality: Čenej, Srbobran; samplingProtocol: Sweep net, Pan traps; eventDate: 24-27.04.2018, 07-10.05.2018; habitat: oilseed rape, semi-natural habitat; individualCount: 2 males, 3 females; behavior: primary parasitoids, larval/pupal; occurrenceID: D2EAF200-7A66-5FD2-B360-62432C85BE8A

**Parasite of:** Cyclorraphic dipterous larvae

**Notes:** oilseed rape pest host: unknown

### *Alloxysta* sp. 1

#### Material

- a. country: Serbia; locality: Srbobran; samplingProtocol: Aphid colony; eventDate: 04.05.2018, 07.05.2018; habitat: oilseed rape; individualCount: 2 males; behavior: secondary parasitoids, larval; occurrenceID: E59C956B-93C4-59D4-99D5-D8A84FE5ACD3

**Parasite of:** Aphidiinae, Aphelininae, *Encyrtidae*

**Notes:** oilseed rape pest host: unknown, possible

### *Rhoptomeris* sp. 1

#### Material

- a. country: Serbia; locality: Čenej, Srbobran; samplingProtocol: Pan traps; eventDate: 24-27.04.2018; habitat: oilseed rape; individualCount: 2 males, 1 female; behavior: primary parasitoids, larval/pupal; occurrenceID: A8E2E3BE-ED68-5E53-A541-274CD00A5354



**Parasite of:** Chloropidae, Diptera

**Notes:** oilseed rape pest host: unknown

## Family Ichneumonidae

### Ichneumonidae sp. 1

#### Material

- a. country: Serbia; locality: Srbobran; samplingProtocol: Sweep net; eventDate: 05-07-18; habitat: oilseed rape; individualCount: 1 female; behavior: primary/secondary parasitoids, egg/larval; occurrenceID: C67F9BF1-DD57-5167-ACC8-733B38E5777D

**Parasite of:** Holometabolous insects

**Notes:** oilseed rape pest host: unknown

### Phaeogenini sp. 1

#### Material

- a. country: Serbia; locality: Srbobran; samplingProtocol: Sweep net; eventDate: 05-25-18; habitat: oilseed rape; individualCount: 1 female; behavior: primary/secondary parasitoids, larval; occurrenceID: 064DA64F-7F98-5937-982C-A00C74CC4561

**Parasite of:** *Plutella xylostella*

**Notes:** oilseed rape pest host: yes

### Phygadeuontinae sp. 1

#### Material

- a. country: Serbia; locality: Đurđin; samplingProtocol: Pan traps; eventDate: 22-24.04.2019; habitat: semi-natural habitat; individualCount: 2 females; behavior: primary/secondary parasitoids, egg/larval; occurrenceID: B2778AA4-64F4-5F69-8FA4-DAD9C5EA9408

**Parasite of:** Holometabolous insects

**Notes:** oilseed rape pest host: unknown

### Phygadeuontini sp. 1

#### Material

- a. country: Serbia; locality: Srbobran; samplingProtocol: Sweep net; eventDate: 05-10-18; habitat: oilseed rape; individualCount: 1 female; behavior: primary/secondary parasitoids, larval; occurrenceID: 28B6A6C4-A371-5989-9208-88674A1797FE

**Parasite of:** Symphyta

**Notes:** oilseed rape pest host: unknown

### ***Aneuclis incidens* (Thomson, 1889)**

#### **Material**

- a. country: Serbia; locality: Čenej, Pačir, Srbobran; samplingProtocol: Sweep net, Pan traps; eventDate: 24-27.04.2018, 10.05.2019; habitat: oilseed rape, semi-natural habitat; individualCount: 1 male, 2 females; behavior: primary parasitoids, larval; occurrenceID: 183E1573-8527-5A4C-AD8B-C6D6D4B465BB

**Parasite of:** *Meligethes aeneus*

**Notes:** oilseed rape pest host: yes

### ***Aptesis flagitator* (Rossi, 1794)**

#### **Material**

- a. country: Serbia; locality: Bajmok; samplingProtocol: Sweep net; eventDate: 05-08-19; habitat: oilseed rape; individualCount: 1 male; behavior: primary parasitoids, larval; occurrenceID: C2101708-61BC-55AA-8D4F-478F8D7DDA85

**Parasite of:** *Agonopterix heracliana*, *Athalia spinarum*

**Notes:** oilseed rape pest host: yes

### ***Bathyplectes curculionis* (Thomson, 1887)**

#### **Material**

- a. country: Serbia; locality: Srbobran; samplingProtocol: Sweep net; eventDate: 04-27-18; habitat: semi-natural habitat; individualCount: 1 female; behavior: primary parasitoids, pupal; occurrenceID: AC0A6F81-CAE3-5C1C-BD6E-45A10A1B37D1

**Parasite of:** *Apion pisi*, *Hypera* spp.

**Notes:** oilseed rape pest host: unknown

### ***Collyria coxator* (Villers, 1789)**

#### **Material**

- a. country: Serbia; locality: Bajmok, Čenej, Đurđin, Pačir, Srbobran; samplingProtocol: Sweep net, Pan traps; eventDate: 04-07.05.2018, 10.05.2018, 25.05.2018, 22-24.04.2019, 08.05.2019; habitat: oilseed rape, semi-natural habitat; individualCount: 7 males, 14 females; behavior: primary parasitoids, larval; occurrenceID: CC1A9ECA-A7E3-52A2-9004-232735782E0D

**Parasite of:** *Cephus cinctus*, *Cephus pygmeus*

**Notes:** oilseed rape pest host: unknown

### ***Diadegma insulare* (Cresson, 1865)**

#### **Material**

- a. country: Serbia; locality: Mišičevo; samplingProtocol: Sweep net; eventDate: 06-12-19; habitat: oilseed rape; individualCount: 1 male; behavior: primary parasitoids, pupal; occurrenceID: CD57A189-7AE6-5C7A-B431-CBA52EAE2F73

**Parasite of:** *Plutella xylostella*

**Notes:** oilseed rape pest host: yes

### ***Diplazon laetatorius* (Fabricius, 1781)**

#### **Material**

- a. country: Serbia; locality: Čenej, Mišičevo; samplingProtocol: Sweep net, Aphid colony; eventDate: 27.04.2018, 12.06.2019; habitat: oilseed rape, semi-natural habitat; individualCount: 2 males; behavior: primary parasitoids, pupal; occurrenceID: 7FC941A6-E807-5486-B88F-9347CAAC84C4

**Parasite of:** Diptera, Syrphidae

**Notes:** oilseed rape pest host: unknown

### ***Dusona pugillator* (Linnaeus, 1758)**

#### **Material**

- a. country: Serbia; locality: Bajmok, Pačir; samplingProtocol: Sweep net, Pan traps; eventDate: 22-24.04.2019, 10.05.2019; habitat: semi-natural habitat; individualCount: 2 males; behavior: primary parasitoids, larval; occurrenceID: D3D73282-6F08-588D-B352-81F81E9CF17F

**Parasite of:** Lepidoptera

**Notes:** oilseed rape pest host: unknown

### ***Diphyus ochromelas* (Gmelin, 1790)**

#### **Material**

- a. country: Serbia; locality: Bajmok; samplingProtocol: Sweep net; eventDate: 05-08-19; habitat: oilseed rape; individualCount: 1 male; behavior: primary parasitoids, larval; occurrenceID: AD397A1A-B727-5C1F-BC52-DF4C511895AB

**Parasite of:** Lepidoptera

Notes: oilseed rape pest host: unknown

### **Mesochorus sp. 1**

#### **Material**

- a. country: Serbia; locality: Čenej; samplingProtocol: Sweep net; eventDate: 05-10-18; habitat: semi-natural habitat; individualCount: 1 male; behavior: secondary parasitoids, larval; occurrenceID: 747E1AF2-EA36-5BE5-B065-72D08031D9F9

Parasite of: *Cotesia* spp.

Notes: oilseed rape pest host: unknown, possible

### **Olesicampe sp. 1**

#### **Material**

- a. country: Serbia; locality: Srbobran; samplingProtocol: Pan traps; eventDate: 04-07.05.2018; habitat: oilseed rape; individualCount: 1 female; behavior: primary parasitoids, pupal; occurrenceID: 6489C6DD-884E-5CE6-9C8E-D4FA7EF45758

Parasite of: Tenthredinidae

Notes: oilseed rape pest host: unknown

### **Stibeutes curvispina (Thomson, 1884)**

#### **Material**

- a. country: Serbia; locality: Čenej, Đurđin, Srbobran; samplingProtocol: Sweep net, Pan traps; eventDate: 04-07.05.2018, 07-10.05.2018, 22-24.04.2019; habitat: oilseed rape; individualCount: 3 males, 2 females; behavior: primary parasitoids, larval; occurrenceID: 12C812FB-44D7-54FE-A5BC-CFEA81EDF1E1

Parasite of: *Ceutorhynchus pallidactylus*

Notes: oilseed rape pest host: yes

### **Syrphophilus bizonarius (Gravenhorst, 1829)**

#### **Material**

- a. country: Serbia; locality: Paćir; samplingProtocol: Sweep net; eventDate: 05-10-19; habitat: semi-natural habitat; individualCount: 1 female; behavior: primary parasitoids, larval; occurrenceID: C55EB950-248B-5402-B80F-E6B92AD77282

Parasite of: *Atherigona soccata*, *Delia radicum*, *Episyrphus balteatus*, *Eupeodes corollae*, *Eupeodes luniger*, *Emex spinosa*, *Loxostege sticticalis*, *Neocnemodon vitripennis*, *Sphaerophoria scripta*

Notes: oilseed rape pest host: unknown

### ***Tersilochus heterocerus* (Thomson, 1889)**

#### **Material**

- a. country: Serbia; locality: Čenej, Srbobran; samplingProtocol: Sweep net; eventDate: 04.05.2018, 07.05.2018, 10.05.2018, 27.04.2018; habitat: oilseed rape, semi-natural habitat; individualCount: 11 males, 7 females; behavior: primary parasitoids, larval; occurrenceID: FD6E7063-B95F-58FC-B7FB-D0FF05686E8E

Parasite of: *Meligethes aeneus*

Notes: oilseed rape pest host: yes

### ***Thrybius praedator* (Rossi, 1792)**

#### **Material**

- a. country: Serbia; locality: Srbobran; samplingProtocol: Sweep net; eventDate: 05-04-18; habitat: semi-natural habitat; individualCount: 1 female; behavior: primary parasitoids, larval; occurrenceID: E1618D03-CD78-5960-811F-FF74A0DC5B74

Parasite of: *Achnara spargani*, *Chilo phragmitellus*, *Oberea euphorbiae*

Notes: oilseed rape pest host: unknown

## **Family Megaspilidae**

### ***Conostigmus rufescens* Kieffer, 1907**

#### **Material**

- a. country: Serbia; locality: Srbobran; samplingProtocol: Sweep net; eventDate: 05-07-18; habitat: oilseed rape; individualCount: 1 male; behavior: primary parasitoids, egg/larval; occurrenceID: FEF6BD20-000B-5AB3-A033-AB1C56E92AE1

Parasite of: *Dasineura brassicae*

Notes: oilseed rape pest host: yes

### ***Lagynodes pallidus* (Boheman 1832)**

#### **Material**

- a. country: Serbia; locality: Čenej, Srbobran; samplingProtocol: Sweep net, Pan traps; eventDate: 24-27.04.2018, 04-07.05.2018, 07-10.05.2018; habitat: oilseed rape, semi-natural habitat; individualCount: 9 males, 6 females; behavior: secondary parasitoids, larval; occurrenceID: 6CEF44D2-BB26-55B3-8C3B-AAF783C929E2

**Parasite of:** *Cotesia* spp.

**Notes:** oilseed rape pest host: unknown

## Family Mymaridae

### Mymaridae sp. 1

#### Material

- a. country: Serbia; locality: Srbobran; samplingProtocol: Pan traps; eventDate: 24-27.04.2018; habitat: semi-natural habitat; individualCount: 1 male; behavior: primary parasitoids, egg; occurrenceID: 84DD4EAA-01BE-5B28-8A93-77EBB1BF35DF

**Parasite of:** Auchenorrhynchos, Hemiptera, Coleoptera, Psocoptera

**Notes:** oilseed rape pest host: unknown

### *Anagrus* sp. 1

#### Material

- a. country: Serbia; locality: Bajmok, Đurđin, Mišičevo; samplingProtocol: Sweep net, Pan traps; eventDate: 20-22.04.2019, 24.04.2019, 25.04.2019; habitat: oilseed rape, semi-natural habitat; individualCount: 3 males, 1 female; behavior: primary parasitoids, egg; occurrenceID: 0DAC860B-40C2-503B-BBBE-2E2E7A426D30

**Parasite of:** Cicadellidae

**Notes:** oilseed rape pest host: unknown

### *Anagrus* sp. 2

#### Material

- a. country: Serbia; locality: Čenej, Srbobran; samplingProtocol: Pan traps; eventDate: 04-07.05.2018; habitat: oilseed rape, semi-natural habitat; individualCount: 2 males; behavior: primary parasitoids, egg; occurrenceID: 97C28639-6481-501E-A8A1-08EE18C9F29A

**Parasite of:** Cicadellidae

**Notes:** oilseed rape pest host: unknown

### *Anagrus* sp. 3

#### Material

- a. country: Serbia; locality: Mišičevo; samplingProtocol: Sweep net; eventDate: 25.04.2019, 12.06.2019; habitat: oilseed rape; individualCount: 2 males; behavior: primary parasitoids, egg; occurrenceID: 2C0C5CF7-B1AD-502E-9B1F-61E21AE752D3

**Parasite of:** Cicadellidae

**Notes:** oilseed rape pest host: unknown

### *Anaphes* sp. 1

#### Material

- a. country: Serbia; locality: Bajmok, Čenej, Đurđin, Mišičevo, Pačir, Srbobran; samplingProtocol: Sweep net, Pan traps; eventDate: 24-27.04.2018, 04-07.05.2018, 10.05.2018, 22-24.04.2019, 25.04.2019; habitat: oilseed rape, semi-natural habitat; individualCount: 27 males, 6 females; behavior: primary parasitoids, egg; occurrenceID: 9047154B-F856-51C1-AB08-FEB894164FCC

**Parasite of:** Coleoptera, Curculionidae, Chrysomelidae, Hemiptera, Miridae

**Notes:** oilseed rape pest host: unknown, possible

### *Gonatocerus* sp. 1

#### Material

- a. country: Serbia; locality: Bajmok; samplingProtocol: Sweep net; eventDate: 05-08-19; habitat: oilseed rape; individualCount: 1 male, 1 female; behavior: primary parasitoids, egg; occurrenceID: B9F3DFBB-3D89-5BF3-BA8E-F5A92F59E93C

**Parasite of:** Cicadellidae

**Notes:** oilseed rape pest host: unknown

### *Litus cynipseus* Haliday, 1833

#### Material

- a. country: Serbia; locality: Čenej, Srbobran; samplingProtocol: Sweep net, Pan traps; eventDate: 04-07.05.2018, 10.05.2018; habitat: oilseed rape, semi-natural habitat; individualCount: 2 males; behavior: primary parasitoids, egg; occurrenceID: 5F95AF60-AC55-546E-A606-10F10185B721

**Parasite of:** Coleoptera, Staphylinidae

**Notes:** oilseed rape pest host: unknown

***Lymaenon* sp. 1****Material**

- a. country: Serbia; locality: Srbobran; samplingProtocol: Sweep net; eventDate: 05-04-18; habitat: oilseed rape; individualCount: 1 male, 1 female; behavior: primary parasitoids, egg; occurrenceID: 7E2F864C-48F5-5C7F-B559-119DB45F5F9E

**Parasite of:** Cicadellidae, Membracoidea

**Notes:** oilseed rape pest host: unknown

***Ooctonus* sp. 1****Material**

- a. country: Serbia; locality: Čenej, Srbobran; samplingProtocol: Sweep net, Pan traps; eventDate: 04-07.05.2018, 07-10.05.2018; habitat: oilseed rape, semi-natural habitat; individualCount: 1 male, 2 females; behavior: primary parasitoids, egg; occurrenceID: C1AD470E-FA4D-563F-AAAD-C752EA57D667

**Parasite of:** Cercopoidea, Cicadellidae

**Notes:** oilseed rape pest host: unknown

***Ooctonus vulgatus* Haliday, 1833****Material**

- a. country: Serbia; locality: Čenej; samplingProtocol: Pan traps; eventDate: 04-07.05.2018; habitat: oilseed rape; individualCount: 1 male; behavior: primary parasitoids, egg; occurrenceID: 3411EABA-B0DF-5C42-AF8D-86EE173E223D

**Parasite of:** *Philaenus leucophthalmus*, *Philaenus spumarius*

**Notes:** oilseed rape pest host: unknown

***Polynema* sp. 1****Material**

- a. country: Serbia; locality: Čenej; samplingProtocol: Pan traps; eventDate: 24-27.04.2018; habitat: oilseed rape; individualCount: 1 male; behavior: primary parasitoids, egg; occurrenceID: DF42C77F-441F-58DF-8209-E0D66D088FA6

**Parasite of:** Cicadellidae

**Notes:** oilseed rape pest host: unknown



## Family Perilampidae

### Perilampidae sp. 1

#### Material

- a. country: Serbia; locality: Bajmok; samplingProtocol: Sweep net; eventDate: 06-13-19; habitat: oilseed rape; individualCount: 1 male; behavior: primary/secondary parasitoids; occurrenceID: 7DAD5AEB-28D5-587B-949F-16730565D0F7

**Parasite of:** Hymenoptera, Diptera, Coleoptera, Lepidoptera, Neuroptera

**Notes:** oilseed rape pest host: unknown

### *Chrysolampus thenae* (Walker, 1848)

#### Material

- a. country: Serbia; locality: Čenej, Srbobran; samplingProtocol: Sweep net; eventDate: 27.04.2018, 04.05.2018, 07.05.2018, 10.05.2018; habitat: oilseed rape, semi-natural habitat; individualCount: 17 males, 12 females; behavior: primary parasitoids, larval/pupal; occurrenceID: F85D3D08-C1FE-52C7-B833-46B497D6904E

**Parasite of:** *Meligethes pedicularis*

**Notes:** oilseed rape pest host: unknown

### *Perilampus aeneus* (Rossius, 1790)

#### Material

- a. country: Serbia; locality: Bajmok; samplingProtocol: Sweep net; eventDate: 06-13-19; habitat: oilseed rape; individualCount: 1 female; behavior: primary parasitoids, larval; occurrenceID: BA6E5516-6F02-56B5-95EB-05FCDCE6A515

**Parasite of:** *Athalia rosae*

**Notes:** oilseed rape pest host: yes

## Family Pirenidae

### *Macroglenes* sp. 1

#### Material

- a. country: Serbia; locality: Čenej; samplingProtocol: Pan traps; eventDate: 04-07.05.2018; habitat: oilseed rape; individualCount: 1 female; behavior: primary parasitoids, egg/larval; occurrenceID: 64D88E4B-98DF-505A-8F73-991A3307575B

**Parasite of:** Cecidomyiidae

**Notes:** oilseed rape pest host: unknown

## Family Platygasteridae

### *Euxestonotus error* (Fitch, 1861)

#### Material

- a. country: Serbia; locality: Čenej, Mišičevo; samplingProtocol: Oilseed pods, Pan traps; eventDate: 04-07.05.2018, 12.06.2019; habitat: oilseed rape; individualCount: 2 males; behavior: primary parasitoids, larval; occurrenceID: D2051C6B-CCEF-5BD6-951B-4A4919461A19

**Parasite of:** *Sitodiplosis mosellana*, *Dasineura brassicae*?

**Notes:** oilseed rape pest host: unknown

### *Inostemma boscii* (Jurine, 1807)

#### Material

- a. country: Serbia; locality: Čenej, Mišičevo; samplingProtocol: Sweep net, Pan traps, Aphid colony; eventDate: 07-10.05.2018, 12.06.2019; habitat: oilseed rape, semi-natural habitat; individualCount: 3 males; behavior: primary parasitoids, egg/larval; occurrenceID: 679DEE51-0DD2-51E1-BD76-D94776FDB43E

**Parasite of:** *Dasineura brassicae*

**Notes:** oilseed rape pest host: yes

### *Platygaster* sp. 1

#### Material

- a. country: Serbia; locality: Mišičevo; samplingProtocol: Sweep net; eventDate: 04-25-19; habitat: semi-natural habitat; individualCount: 1 male; behavior: primary parasitoids, larval; occurrenceID: CE7D884E-5A2C-5411-BEAF-3B9BC5B0E1B2

**Parasite of:** Cecidomyiidae (*Dasineura brassicae*?)

**Notes:** oilseed rape pest host: unknown, possible

### *Platygaster* sp. 2

#### Material

- a. country: Serbia; locality: Bajmok, Đurđin, Srbobran; samplingProtocol: Sweep net, Pan traps; eventDate: 07.05.2018, 17-19.04.2019, 22-24.04.2019; habitat: oilseed rape, semi-

natural habitat; individualCount: 5 males, 1 female; behavior: primary parasitoids, larval;  
occurrenceID: 518A0586-0A1A-59F0-8C02-16CEDB1BED18

**Parasite of:** Cecidomyiidae (*Dasineura brassicae*?)

**Notes:** oilseed rape pest host: unknown, possible

## ***Platygaster subuliformis* Kieffer, 1926**

### **Material**

- a. country: Serbia; locality: Čenej; samplingProtocol: Pan traps; eventDate: 07-10.05.2018;  
habitat: oilseed rape; individualCount: 1 male; behavior: primary parasitoids, egg/larval;  
occurrenceID: 4D15DF4D-D4AF-522F-8D92-C9DCC1A223F2

**Parasite of:** *Dasineura brassicae*

**Notes:** oilseed rape pest host: yes

## ***Synopeas* sp. 1**

### **Material**

- a. country: Serbia; locality: Mišičevo; samplingProtocol: Sweep net; eventDate: 04-25-19;  
habitat: semi-natural habitat; individualCount: 1 male; behavior: primary parasitoids, egg/  
larval; occurrenceID: 290E0998-E75F-52A9-9155-ADFAB09FA382

**Parasite of:** Cecidomyiidae (*Dasineura brassicae*?)

**Notes:** oilseed rape pest host: unknown, possible

## ***Telenomus* sp. 1**

### **Material**

- a. country: Serbia; locality: Bajmok, Čenej, Đurđin, Mišičevo, Srbobran; samplingProtocol:  
Sweep net, Pan traps; eventDate: 24-27.04.2018, 04-07.05.2018, 07-10.05.2018,  
22-24.04.2019, 25.04.2019, 12.06.2019, 13.06.2019; habitat: oilseed rape, semi-natural  
habitat; individualCount: 25 males, 6 females; behavior: primary parasitoids, egg;  
occurrenceID: 8A98FE66-1FF3-5068-B91F-7CE2629032F3

**Parasite of:** Lepidoptera, Heteroptera, Diptera, Neuroptera

**Notes:** oilseed rape pest host: unknown

## ***Telenomus* sp. 2**

### **Material**

- a. country: Serbia; locality: Đurđin, Srbobran; samplingProtocol: Pan traps; eventDate:  
24-27.04.2018, 04-07.05.2018, 07-10.05.2018, 22-24.04.2019; habitat: oilseed rape,

semi-natural habitat; individualCount: 5 males, 3 females; behavior: primary parasitoids, egg; occurrenceID: 3FF5D53B-3C0F-5687-8C81-91E1203DE58E

**Parasite of:** Lepidoptera, Heteroptera, Diptera, Neuroptera

**Notes:** oilseed rape pest host: unknown

### *Telenomus* sp. 3

#### Material

- a. country: Serbia; locality: Bajmok, Čenej, Đurđin, Mišićevo, Pačir, Srbobran; samplingProtocol: Sweep net, Pan traps; eventDate: 24-27.04.2018, 04-07.05.2018, 07-10.05.2018, 22-24.04.2019, 25.04.2019, 13.06.2019; habitat: oilseed rape, semi-natural habitat; individualCount: 17 males, 2 females; behavior: primary parasitoids, egg; occurrenceID: F2466748-82B0-547A-B168-23D648548A6E

**Parasite of:** Lepidoptera, Heteroptera, Diptera, Neuroptera

**Notes:** oilseed rape pest host: unknown

### *Telenomus* sp. 4

#### Material

- a. country: Serbia; locality: Bajmok, Čenej, Srbobran; samplingProtocol: Sweep net, Pan traps; eventDate: 24-27.04.2018, 04-07.05.2018, 07-10.05.2018, 13.06.2019; habitat: oilseed rape, semi-natural habitat; individualCount: 15 males, 6 females; behavior: primary parasitoids, egg; occurrenceID: 85A0CC52-0238-5F96-B65D-D7F0887B1630

**Parasite of:** Lepidoptera, Heteroptera, Diptera, Neuroptera

**Notes:** oilseed rape pest host: unknown

## Family Pteromalidae

### Pteromalidae sp. 1

#### Material

- a. country: Serbia; locality: Čenej, Srbobran; samplingProtocol: Sweep net, Pan traps; eventDate: 27.04.2018, 04-07.05.2018; habitat: oilseed rape; individualCount: 4 males; behavior: primary/secondary parasitoids, larval/pupal; occurrenceID: 3AFD2F76-5B82-5461-9D51-BB9993A1B4AF

**Parasite of:** Lepidoptera, Coleoptera, Diptera

**Notes:** oilseed rape pest host: unknown

## Pteromalidae sp. 2

### Material

- a. country: Serbia; locality: Čenej; samplingProtocol: Pan traps; eventDate: 24-27.04.2018; habitat: oilseed rape; individualCount: 1 male; behavior: primary/secondary parasitoids, larval/pupal; occurrenceID: 9C1CFA2D-48B0-5510-9CE6-46D7A0C71457

**Parasite of:** Lepidoptera, Coleoptera, Diptera

**Notes:** oilseed rape pest host: unknown

## Pteromalidae sp. 3

### Material

- a. country: Serbia; locality: Srbobran; samplingProtocol: Pan traps; eventDate: 04-07.05.2018; habitat: semi-natural habitat; individualCount: 1 male; behavior: primary/secondary parasitoids, larval/pupal; occurrenceID: 6FF49984-DB97-5383-8383-99B291993177

**Parasite of:** Lepidoptera, Coleoptera, Diptera

**Notes:** oilseed rape pest host: unknown

## Pteromalidae sp. 4

### Material

- a. country: Serbia; locality: Čenej, Srbobran; samplingProtocol: Sweep net; eventDate: 04.05.2018, 07.05.2018; habitat: oilseed rape; individualCount: 2 males; behavior: primary/secondary parasitoids, larval/pupal; occurrenceID: 0179DAD3-C720-5A57-AEAC-723A1C9ADB93

**Parasite of:** Lepidoptera, Coleoptera, Diptera

**Notes:** oilseed rape pest host: unknown

## Pteromalidae sp. 5

### Material

- a. country: Serbia; locality: Čenej, Đurđin, Pačir, Srbobran; samplingProtocol: Sweep net, Pan traps; eventDate: 04-07.05.2018, 25.04.2019, 08.05.2019; habitat: oilseed rape; individualCount: 8 males; behavior: primary/secondary parasitoids, larval/pupal; occurrenceID: 7B2133A9-66DC-583C-B802-5C239B0A4033

**Parasite of:** Lepidoptera, Coleoptera, Diptera

**Notes:** oilseed rape pest host: unknown

**Pteromalidae sp. 6****Material**

- a. country: Serbia; locality: Čenej; samplingProtocol: Sweep net; eventDate: 27.04.2018, 04.05.2018, 07.05.2018; habitat: oilseed rape; individualCount: 3 males; behavior: primary/secondary parasitoids, larval/pupal; occurrenceID: F3CE52EF-8D40-5F57-8755-722D2E3E1A4C

**Parasite of:** Lepidoptera, Coleoptera, Diptera

**Notes:** oilseed rape pest host: unknown

**Pteromalidae sp. 7****Material**

- a. country: Serbia; locality: Čenej, Mišićevo, Srbobran; samplingProtocol: Sweep net; eventDate: 27.04.2018, 10.05.2018, 25.05.2018; habitat: oilseed rape, semi-natural habitat; individualCount: 5 males, 2 females; behavior: primary/secondary parasitoids, larval/pupal; occurrenceID: D1A62987-9E64-53EE-A8CC-F58803407DFD

**Parasite of:** Lepidoptera, Coleoptera, Diptera

**Notes:** oilseed rape pest host: unknown

**Pteromalidae sp. 8****Material**

- a. country: Serbia; locality: Bajmok, Čenej, Srbobran; samplingProtocol: Sweep net; eventDate: 04.05.2018, 10.05.2018, 24.04.2019; habitat: oilseed rape; individualCount: 4 males, 2 females; behavior: primary/secondary parasitoids, larval/pupal; occurrenceID: 7E4CFEEE-8C69-5CBC-B5B5-63E2603EB967

**Parasite of:** Lepidoptera, Coleoptera, Diptera

**Notes:** oilseed rape pest host: unknown

**Pteromalidae sp. 9****Material**

- a. country: Serbia; locality: Čenej; samplingProtocol: Sweep net; eventDate: 27.04.2018, 10.05.2018; habitat: oilseed rape, semi-natural habitat; individualCount: 1 male, 1 female; behavior: primary/secondary parasitoids, larval/pupal; occurrenceID: 3F827E96-D1E1-5159-AB7A-167D27E787FE

**Parasite of:** Lepidoptera, Coleoptera, Diptera

**Notes:** oilseed rape pest host: unknown

**Pteromalidae sp. 10****Material**

- a. country: Serbia; locality: Srbobran; samplingProtocol: Aphid colony; eventDate: 05-07-18; habitat: oilseed rape; individualCount: 1 male; behavior: primary/secondary parasitoids, larval/pupal; occurrenceID: DE482C0B-356E-597F-9158-22C2E8DA290F

**Parasite of:** Lepidoptera, Coleoptera, Diptera

**Notes:** oilseed rape pest host: unknown

**Pteromalidae sp. 11****Material**

- a. country: Serbia; locality: Bajmok; samplingProtocol: Pan traps; eventDate: 17-19.04.2019; habitat: semi-natural habitat; individualCount: 1 female; behavior: primary/secondary parasitoids, larval/pupal; occurrenceID: 4FBAA759-55F7-51FB-AA60-7D18FFC729CC

**Parasite of:** Lepidoptera, Coleoptera, Diptera

**Notes:** oilseed rape pest host: unknown

***Dibrachys microgastri* (Bouché, 1834)****Material**

- a. country: Serbia; locality: Srbobran; samplingProtocol: Pan traps; eventDate: 24-27.04.2018; habitat: oilseed rape; individualCount: 1 male; behavior: secondary parasitoids, pupal; occurrenceID: C2BE8652-2DBC-5A8D-9A41-B2A6C789A542

**Parasite of:** *Cotesia* spp.

**Notes:** oilseed rape pest host: unknown

***Mesopolobus incultus* (Walker, 1834)****Material**

- a. country: Serbia; locality: Bajmok, Mišičevo, Pačir; samplingProtocol: Sweep net; eventDate: 25.04.2019, 12.06.2019, 13.06.2019; habitat: oilseed rape; individualCount: 4 males, 1 female; behavior: primary parasitoids, larval; occurrenceID: 4FB3965B-1340-5F9B-BB35-111CAE2B6B29

**Parasite of:** Curculionidae, *Gymnetron* sp., *Gymnetron pascuorum*, *Mecinus* sp., Scolytidae, *Polygraphus poligraphus*, Agromyzidae, *Phytobia humeralis*, Cecidomyiidae, *Kaltenbachiola strobi*

**Notes:** oilseed rape pest host: unknown

***Mesopolobus morys* (Walker, 1848)****Material**

- a. country: Serbia; locality: Bajmok, Čenej, Đurđin, Mišičevo, Srbobran; samplingProtocol: Sweep net, Oilseed pods; eventDate: 27.04.2018, 10.05.2018, 25.05.2018, 12.06.2019, 13.06.2019; habitat: oilseed rape, semi-natural habitat; individualCount: 23 males, 5 females; behavior: primary parasitoids, larval; occurrenceID: 009636D4-DFBA-522C-90B6-8DBA0B11E1EB

**Parasite of:** *Ceutorhynchus assimilis*

**Notes:** oilseed rape pest host: yes

***Mesopolobus* sp. 1****Material**

- a. country: Serbia; locality: Čenej; samplingProtocol: Pan traps; eventDate: 04-07.05.2018, 07-10.05.2018; habitat: oilseed rape; individualCount: 10 males; behavior: primary parasitoids, larval; occurrenceID: 33B2176A-BC4D-5619-822B-22D79B58F442

**Parasite of:** *Ceutorhynchus* spp.

**Notes:** oilseed rape pest host: unknown, possible

***Mesopolobus* sp. 2****Material**

- a. country: Serbia; locality: Čenej, Srbobran; samplingProtocol: Sweep net, Pan traps; eventDate: 24-27.04.2018, 10.05.2018; habitat: oilseed rape; individualCount: 3 males; behavior: primary parasitoids, larval; occurrenceID: 3D0EEB7D-A975-53E6-B51B-B046F124045D

**Parasite of:** *Ceutorhynchus* spp.

**Notes:** oilseed rape pest host: unknown, possible

***Mesopolobus* sp. 3****Material**

- a. country: Serbia; locality: Čenej, Srbobran; samplingProtocol: Sweep net, Pan traps; eventDate: 04-07.05.2018, 10.05.2018; habitat: oilseed rape, semi-natural habitat; individualCount: 2 males, 1 female; behavior: primary parasitoids, larval; occurrenceID: E5605028-C316-573A-90BC-AA00987B6E8D

**Parasite of:** *Ceutorhynchus* spp.

**Notes:** oilseed rape pest host: unknown, possible



### ***Pteromalus* sp. 1**

#### **Material**

- a. country: Serbia; locality: Čenej, Srbobran; samplingProtocol: Sweep net; eventDate: 27.04.2018, 04.05.2018, 25.05.2018; habitat: oilseed rape; individualCount: 2 males, 3 females; behavior: primary parasitoids, pupal; occurrenceID: 963C73AA-A9C4-5BA1-9EA7-1B189335F65C

**Parasite of:** Lepidoptera, Tenthredinidae

**Notes:** oilseed rape pest host: unknown

### ***Pteromalus* sp. 2**

#### **Material**

- a. country: Serbia; locality: Mišićevo; samplingProtocol: Sweep net; eventDate: 04-25-19; habitat: semi-natural habitat; individualCount: 1 male; behavior: primary parasitoids, pupal; occurrenceID: 9F2959EB-40AA-5EBB-8492-CEC29BDA2C93

**Parasite of:** Lepidoptera, Tenthredinidae

**Notes:** oilseed rape pest host: unknown

### ***Pteromalus* sp. 3**

#### **Material**

- a. country: Serbia; locality: Mišićevo; samplingProtocol: Sweep net; eventDate: 25.04.2019, 10.05.2019; habitat: semi-natural habitat; individualCount: 2 males; behavior: primary parasitoids, pupal; occurrenceID: E8377C37-16CC-5F4F-ADF7-46E4C90C4146

**Parasite of:** Lepidoptera, Tenthredinidae

**Notes:** oilseed rape pest host: unknown

### ***Trichomalus lucidus* (Walker, 1835)**

#### **Material**

- a. country: Serbia; locality: Bajmok, Čenej, Đurđin, Pačir, Srbobran; samplingProtocol: Sweep net, Pan traps, Aphid colony; eventDate: 27.04.2018, 04-07.05.2018, 07-10.05.2018, 25.05.2018, 22-24.04.2019, 08.05.2019, 13.06.2019; habitat: oilseed rape, semi-natural habitat; individualCount: 244 males, 150 females; behavior: primary parasitoids, larval; occurrenceID: 9E42182F-CD4C-58E3-A006-500646839E1B

**Parasite of:** *Ceutorhynchus* spp., *Psylliodes chrysocephala*

**Notes:** oilseed rape pest host: yes

***Trichomalus* sp. 1****Material**

- a. country: Serbia; locality: Čenej, Srbobran; samplingProtocol: Pan traps; eventDate: 04-07.05.2018, 07-10.05.2018; habitat: oilseed rape, semi-natural habitat; individualCount: 3 males, 1 female; behavior: primary parasitoids, larval; occurrenceID: C0DFE327-FD39-5199-A973-1F370CF44F89

**Parasite of:** *Ceutorhynchus* spp.

**Notes:** oilseed rape pest host: unknown, possible

***Trichomalus* sp. 2****Material**

- a. country: Serbia; locality: Čenej, Srbobran; samplingProtocol: Pan traps; eventDate: 04-07.05.2018, 07-10.05.2018; habitat: oilseed rape; individualCount: 2 males, 4 females; behavior: primary parasitoids, larval; occurrenceID: 5B64E486-9913-590C-9EA2-3C7FEAF2EAD7

**Parasite of:** *Ceutorhynchus* spp.

**Notes:** oilseed rape pest host: unknown, possible

***Trichomalus* sp. 3****Material**

- a. country: Serbia; locality: Čenej; samplingProtocol: Sweep net; eventDate: 05-10-18; habitat: oilseed rape; individualCount: 1 male; behavior: primary parasitoids, larval; occurrenceID: 08BC1E84-4526-5A64-A66F-11A613C156A9

**Parasite of:** *Ceutorhynchus* spp.

**Notes:** oilseed rape pest host: unknown, possible

***Trichomalus* sp. 4****Material**

- a. country: Serbia; locality: Srbobran; samplingProtocol: Pan traps; eventDate: 04-07.05.2018; habitat: semi-natural habitat; individualCount: 1 male; behavior: primary parasitoids, larval; occurrenceID: 635BACFB-24DC-51D4-B87A-E94BBACEAD18

**Parasite of:** *Ceutorhynchus* spp.

**Notes:** oilseed rape pest host: unknown, possible

## Family Scelionidae

### Scelionidae sp. 1

#### Material

- a. country: Serbia; locality: Srbobran; samplingProtocol: Sweep net; eventDate: 05-07-18; habitat: oilseed rape; individualCount: 1 female; behavior: primary parasitoids; occurrenceID: 155A11AF-C04E-5ACF-B427-5945A80564DA

**Parasite of:** Insects, arachnids

**Notes:** oilseed rape pest host: unknown

### *Eumicrosoma* sp. 1

#### Material

- a. country: Serbia; locality: Srbobran; samplingProtocol: Sweep net; eventDate: 05-07-18; habitat: oilseed rape; individualCount: 1 male; behavior: primary parasitoids, egg; occurrenceID: 80F2947C-A65A-5166-98C3-25C427411C13

**Parasite of:** Heteroptera, Pentatomidae, Lygaeidae

**Notes:** oilseed rape pest host: unknown

### *Gryon* sp. 1

#### Material

- a. country: Serbia; locality: Srbobran; samplingProtocol: Sweep net, Pan traps; eventDate: 24-27.04.2018, 07.05.2018, 10.05.2018; habitat: oilseed rape; individualCount: 1 male, 2 females; behavior: primary parasitoids, egg; occurrenceID: 8D48E981-9573-5984-A23B-2C7D628D0D7B

**Parasite of:** Hemiptera, Coreidae

**Notes:** oilseed rape pest host: unknown

### *Trimorus* sp. 1

#### Material

- a. country: Serbia; locality: Bajmok, Srbobran; samplingProtocol: Sweep net; eventDate: 07.05.2018, 24.04.2019; habitat: oilseed rape; individualCount: 1 male, 1 female; behavior: primary parasitoids, egg; occurrenceID: 02F2A15C-4302-5C1A-908C-8F0612805AEE

**Parasite of:** Carabidae

**Notes:** oilseed rape pest host: unknown

### ***Trissolcus basalis* (Wollaston, 1858)**

#### **Material**

- a. country: Serbia; locality: Bajmok, Čenej, Srbobran; samplingProtocol: Sweep net, Pan traps; eventDate: 24-27.04.2018, 04-07.05.2018, 07-10.05.2018, 25.05.2018, 13.06.2019; habitat: oilseed rape, semi-natural habitat; individualCount: 14 males, 5 females; behavior: primary parasitoids, egg; occurrenceID: F640CCB8-EB59-5547-8493-53E50E3F9240

**Parasite of:** Heteroptera, *Nezara viridula*

**Notes:** oilseed rape pest host: unknown

### **Family Spalangiidae**

### ***Spalangia nigra* Latreille, 1805**

#### **Material**

- a. country: Serbia; locality: Čenej, Pačir; samplingProtocol: Pan traps; eventDate: 07-10.05.2018, 22-24.04.2019; habitat: oilseed rape, semi-natural habitat; individualCount: 3 males; behavior: primary parasitoids, pupal; occurrenceID: E0BC2BE7-A9F1-5930-B13D-8B4980F0D92D

**Parasite of:** Diptera puparia

**Notes:** oilseed rape pest host: unknown

### **Family Systasidae**

### ***Asaphes vulgaris* Walker, 1834**

#### **Material**

- a. country: Serbia; locality: Čenej, Srbobran; samplingProtocol: Sweep net; eventDate: 10.05.2018, 25.05.2018; habitat: oilseed rape; individualCount: 2 males; behavior: secondary parasitoids, larval; occurrenceID: 0354FF52-0A62-5246-8340-A904585A61BD

**Parasite of:** aphid parasitoids

**Notes:** oilseed rape pest host: yes

## Family Torymidae

### *Podagrion pachymerum* (Walker, 1833)

#### Material

- a. country: Serbia; locality: Mišićevo; samplingProtocol: Sweep net; eventDate: 04-25-19; habitat: semi-natural habitat; individualCount: 1 male; behavior: primary parasitoids, egg; occurrenceID: D3EA9F47-C288-59B0-8F97-02BF3560AC92

**Parasite of:** Mantodea, Mantidae

**Notes:** oilseed rape pest host: unknown

### *Pseudotorymus napi* (Amerling & Kirchner, 1860)

#### Material

- a. country: Serbia; locality: Čenej, Mišićevo, Srbobran; samplingProtocol: Sweep net, Pan traps, Aphid colony, Oilseed pods; eventDate: 24-27.04.2018, 04-07.05.2018, 07-10.05.2018, 25.05.2018, 25.05.2019; habitat: oilseed rape, semi-natural habitat; individualCount: 144 males, 74 females; behavior: primary parasitoids, larval; occurrenceID: A46E8930-EDC8-58B9-9BBA-D60067B74506

**Parasite of:** *Dasineura brassicae*

**Notes:** oilseed rape pest host: yes

### *Torymus* sp. 1

#### Material

- a. country: Serbia; locality: Pačir; samplingProtocol: Sweep net; eventDate: 04-25-19; habitat: oilseed rape; individualCount: 1 male; behavior: primary parasitoids, larval; occurrenceID: 59EE6B5E-7094-559B-BD13-2305E7E082C7

**Parasite of:** Ectoparasitoids of gall-forming insects (Cecidomyiidae, Cynipidae)

**Notes:** oilseed rape pest host: unknown

## Family Trichogrammatidae

### Trichogrammatidae sp. 1

#### Material

- a. country: Serbia; locality: Bajmok; samplingProtocol: Sweep net; eventDate: 04-24-19; habitat: oilseed rape; individualCount: 1 male; behavior: primary parasitoids, egg; occurrenceID: 844E2307-1DB9-578E-9CE2-2A8A79EE2C32

**Parasite of:** Lepidoptera, Coleoptera, Neuroptera, Diptera, Hymenoptera

**Notes:** oilseed rape pest host: unknown

### ***Trichogramma evanescens* Westwood, 1833**

#### **Material**

- a. country: Serbia; locality: Srbobran; samplingProtocol: Pan traps; eventDate: 24-27.04.2018; habitat: semi-natural habitat; individualCount: 1 male; behavior: primary parasitoids, egg; occurrenceID: 1E041ED8-6884-5503-B24E-7115C06B7382

**Parasite of:** Lepidoptera, Chrysomelidae

**Notes:** oilseed rape pest host: unknown

### **Analysis**

During this two-year study, we found a total of 3135 specimens of primary or secondary parasitoids, of which 2855 were found in oilseed rape fields and 280 in semi-natural habitats. In the 2018 season, we found 2926 individuals, while in 2019, we found only 209 individuals. The temporal dynamics of parasitoid abundance were also different in the two years. The highest abundance was in May in 2018 (2694) and in April in 2019 (107). Most individuals were found when sampling with sweep nets (1715), followed by rearing from aphid colonies (1001), pan traps (404) and collecting oilseed pods (15).

We found 153 different taxa, of which 119 were found in oilseed rape fields and 87 in semi-natural habitats. Not all specimens could be identified to species level, but only to genus or family level. We found a total of 92 different genera, 73 in oilseed rape fields and 59 in semi-natural habitats and 22 families, all of which occurred in oilseed rape and 19 in semi-natural habitats. The species accumulation curve shows that the number of detected species in oilseed rape fields has not reached a plateau (Fig. 2), while extrapolated species richness differed somewhat between different estimators (Chao =  $173.2 \pm 21.5$ , Jack1 =  $165.4 \pm 12.3$ , Jack2 = 193.2, Bootstrap =  $138.1 \pm 6.8$ ). The number of singletons and doubletons in oilseed rape fields was 45 and 19, respectively.

The four most abundant families were Braconidae (1880), *Pteromalidae* (500), Torymidae (220) and *Eulophidae* (123) with more than 85% of all specimens captured and the four most numerous genera were *Diaeretiella* Starý, 1960 (1243), *Trichomalus* Thomson, 1878 (406), *Aphidius* Nees, 1818 (389) and *Pseudotorymus* Masi, 1921 (218) with more than 70%. The number of different taxa found in each family, sorted by number of taxa is shown in Table 3.

Of all the taxa we found, 119 are described in literature as primary parasitoids and five as hyperparasitoids, with another 29 that can be either primary or secondary parasitoids. Based on the literature examined, in both habitats, we found 31 genera (33 species) described as parasitoids of oilseed rape pests and 54 genera (97 species) parasitising

non-pest species. We also identified 10 genera (23 species) as possible parasitoids of oilseed rape pests. These taxa could only be identified to the genus level and belong to genera that have at least one other species of oilseed rape pest parasitoids. In oilseed rape fields, we found 28 genera (29 species) of oilseed rape pest parasitoids, 40 genera (73 species) parasitising non-pest species and eight genera (17 species) of possible oilseed rape pest parasitoids.

Table 3.

The number of parasitoid taxa found in each family in two habitats.

Total		Oilseed rape fields		Semi-natural habitats	
Family	No. of taxa	Family	No. of taxa	Family	No. of taxa
Braconidae	29	Braconidae	22	Braconidae	16
<i>Pteromalidae</i>	25	<i>Pteromalidae</i>	20	<i>Eulophidae</i>	11
<i>Ichneumonidae</i>	18	<i>Eulophidae</i>	14	<i>Pteromalidae</i>	11
<i>Eulophidae</i>	17	<i>Ichneumonidae</i>	12	<i>Ichneumonidae</i>	10
<i>Mymaridae</i>	11	<i>Mymaridae</i>	10	Platygastridae	8
Platygastridae	10	Platygastridae	8	<i>Encyrtidae</i>	6
<i>Encyrtidae</i>	8	<i>Scelionidae</i>	5	<i>Mymaridae</i>	6
<i>Ceraphronidae</i>	5	<i>Encyrtidae</i>	4	<i>Ceraphronidae</i>	4
<i>Scelionidae</i>	5	Figitidae	4	Ceidae	2
Figitidae	4	<i>Ceraphronidae</i>	3	Diapriidae	2
<i>Perilampidae</i>	3	<i>Perilampidae</i>	3	Figitidae	2
Torymidae	3	Bethylidae	2	Torymidae	2
Bethylidae	2	Megaspilidae	2	Chalcididae	1
Ceidae	2	Torymidae	2	Eurytomidae	1
Diapriidae	2	Ceidae	1	Megaspilidae	1
Megaspilidae	2	Chalcididae	1	<i>Perilampidae</i>	1
<i>Trichogrammatidae</i>	2	Diapriidae	1	<i>Scelionidae</i>	1
Chalcididae	1	Eurytomidae	1	Spalangiidae	1
Eurytomidae	1	Pirenidae	1	<i>Trichogrammatidae</i>	1
Pirenidae	1	Spalangiidae	1		
Spalangiidae	1	Systasidae	1		
Systasidae	1	<i>Trichogrammatidae</i>	1		

Using two criteria for consistency of occurrence, we selected 25 taxa (22 genera) that regularly occur in our oilseed rape fields, of which 13 taxa (11 genera) were not reported as parasitoids of oilseed rape pests, although four taxa are identified as possible parasitoids (Table 4).




Figure 2. [doi](#)

Species accumulation curve for oilseed rape fields. Light blue area represents 95% confidence intervals.

Table 4.

List of taxa consistently found in oilseed rape fields.

Taxon	Oilseed rape pest host	No. of specimens	No. of fields
<i>Diaeretiella rapae</i> (McIntosh, 1855)	<i>Myzus persicae</i> , <i>Brevicoryne brassicae</i>	1197	9
<i>Trichomalus lucidus</i> (Walker, 1835)	<i>Psylliodes chrysocephala</i> , <i>Ceutorhynchus pallidactylus</i>	392	6
<i>Aphidius ervi</i> Haliday, 1834	<i>Myzus persicae</i> , <i>Brevicoryne brassicae</i>	270	6
<i>Pseudotorymus napi</i> (Amerling & Kirchner, 1860)	<i>Dasineura brassicae</i>	217	5
<i>Aphidius matricariae</i> Haliday, 1834	<i>Myzus persicae</i> , <i>Brevicoryne brassicae</i>	112	4
<i>Triaspis thoracica</i> (Curtis, 1860)	unknown	70	4
<i>Eubazus sigalphoides</i> (Marshall, 1889)	<i>Brassicogethes aeneus</i> (Fabricius, 1775)	52	5
<i>Mesopolobus morys</i> (Walker, 1848)	<i>Ceutorhynchus obstrictus</i>	27	6
<i>Praon volucre</i> (Haliday, 1833)	<i>Myzus persicae</i> , <i>Brevicoryne brassicae</i>	27	5
<i>Anaphes</i> sp. 1	unknown/possible	24	7



Taxon	Oilseed rape pest host	No. of specimens	No. of fields
<i>Aphanogmus abdominalis</i> (Thomson, 1858)	<i>Dasineura brassicae</i>	23	3
<i>Townesilitus bicolor</i> (Wesmael, 1835)	<i>Phyllotreta nemorum</i>	20	3
<i>Eulophus</i> sp. 1	unknown/possible	18	6
<i>Collyria coxator</i> (Villers, 1789)	unknown	18	3
<i>Trissolcus basalis</i> (Wollaston, 1858)	<i>Nezara viridula</i> (Linnaeus, 1758)	17	4
<i>Tersilochus heterocerus</i> (Thomson, 1889)	<i>Brassicogethes aeneus</i>	17	4
<i>Aprostocetus</i> sp. 1	unknown/possible	16	7
<i>Telenomus</i> sp. 3	unknown	15	7
<i>Chrysolampus thenae</i> (Walker, 1848)	unknown	15	3
<i>Telenomus</i> sp. 4	unknown	14	5
<i>Telenomus</i> sp. 1	unknown	13	5
<i>Lagynodes pallidus</i> (Boheman, 1832)	unknown	12	4
<i>Tetrastichus</i> sp. 1	unknown/possible	12	3
<i>Spalangiopecta</i> sp. 1	unknown	11	5
<i>Apanteles</i> sp. 1	unknown	10	3

Detailed information on locations, collection dates, habitat, sampling methods, host, parasitism type, oilseed rape pest host status and number of specimens for each taxon can be found in Suppl. material 2 and Suppl. material 3 and additional data on host stage and references in Suppl. material 4.

## Discussion

Oilseed rape fields are known to harbour a variety of parasitoids, which are natural enemies of oilseed rape pests (Alford 2003, Williams 2010). However, we have found that the diversity of parasitoids found in our oilseed rape fields goes beyond those associated with crop pests. In fact, only slightly less than 25% of the parasitoid taxa identified in oilseed rape fields are linked to crop pest species. This suggests that oilseed rape fields serve multiple functions for parasitoids, including providing host species that are not pests and floral resources that attract parasitoids. Although the aim of this study was not to conduct a detailed analysis of differences between oilseed rape fields and nearby semi-natural habitats, we observed a noticeably higher abundance and diversity of parasitoids in oilseed rape fields compared to nearby semi-natural habitats. The presence of a diverse parasitoid community in fields, including those not typically associated with oilseed rape pests, may contribute to the natural pest control of other crops and promote biodiversity in the surrounding environment. However, it is worth noting that some parasitoid species

might also provide disservice to biological control by parasitising on other beneficial insects, like *Diplazon laetatorius* that can parasitise hoverfly pupae (Townes 1971), *Asaphes vulgaris* which is hyperparasitoid of several of the aphid parasitoids (Žikić et al. 2012) and *Podagrion* species that specialise on praying mantis eggs (Thompson 1958).

There are several taxa that were collected consistently and in greater numbers in our oilseed rape fields, but are not known to parasitise oilseed rape pest species. *Triaspis thoracica* was the most numerous of them and parasitises species of the genus *Bruchus* (Chrysomelidae), which are important pests in various types of bean crops, such as peas, lentils and vetches (Yu et al. 2011). In 2018, it was found in four fields, but only in one field in large numbers (65 out of a total of 70 individuals). Although this cannot be verified, it is possible that this particular field had a high understorey of some Fabaceae species that provided resources for the *Bruchus* population and consequently for *T. thoracica*. Similarly, *Trissolcus basalus* is an egg parasitoid of invasive pentatomid species, such as *Nezara viridula* and *Halyomorpha halys* (Stål, 1855), which are pests in many legumes, fruits and vegetables (Powell and Shepard 1982, Schaefer and Panizzi 2000, Zapponi et al. 2021). The sporadic presence of other plant species in the oilseed rape fields, such as wild grasses, may also possibly explain the presence of *Collyria coxator*, a specialised parasitoid of the European wheat stem sawfly (*Cephus pygmaeus* (Linnaeus, 1767)), which is a pest of cereal crops, but also feeds on various grass species. Likewise, *Telenomus* species parasitise various Heteroptera, but also Lepidoptera, which can be readily found as non-pests in oilseed rape fields. On the other hand, *Chrysolampus thenae* is known to parasitise *Lamiogethes pedicularis* (Gyllenhal, 1808) which are found on Lamiaceae plants (Noyes 2019). It is possible that this parasitoid also attacks other species from the subfamily Meligethinae, such as *B. aeneus* or that host plants were present in the undergrowth. Given the significant presence of these parasitoids not associated with pest species, it would be valuable for future studies to note the previous season's crop as this may help explain the presence of volunteer plants and associated pests and parasitoids.

There are also some parasitoid species that are widely distributed in oilseed rape throughout Europe, but are absent or present only in small numbers in our study. According to Nilsson (2003), the most important pollen beetle parasitoid species are *Phradis interstitialis* (Thomson, 1889), *P. morionellus* (Holmgren, 1860) and *Tersilochus heterocerus*, with *Diospilus capito*, *Blacus nigricornis* and *Brachyserphus parvulus* (Nees, 1834) also being quite common. Only *T. heterocerus* was found in our study in noteworthy numbers to be considered as an important biological control agent. Two other species were found in very low numbers, *D. capito* only in semi-natural habitats and *B. nigricornis* in both habitats, while other species were not found. Interestingly, the most abundant pollen beetle parasitoid in our study was *Eubazus sigalphoides*, previously reported only from France and Poland (Nilsson 2003, Ulber et al. 2010). Likewise, for the cabbage seed weevil (*C. obstrictus*), only one species (*Mesopolobus morys*) of the three most common larval ectoparasitoids (*Stenomalina gracilis* (Walker, 1834), *Trichomalus perfectus* (Walker, 1835)) was detected in our study. We also found the two most common parasitoids of the brassica pod midge in Europe, *Platygaster subuliformis* and *Omphale clypealis*, but both in

low abundance, while the dominant species in our study were *Pseudotorymus napi* and *Aphanogmus abdominalis*. The second most abundant parasitoid in our study was the pteromalid *Trichomalus lucidus*, which parasitises the cabbage stem flea beetle (*P. chrysocephala*) and the cabbage stem weevil (*C. pallidactylus*). This species is one of eight recorded parasitoids of the cabbage stem flea beetle and of several parasitoids of the cabbage stem weevil (Ulber 2003). Other parasitoid species of these two pests were either not observed or observed in very low numbers. Todorov et al. (2022) recently presented a checklist of the pteromalid fauna in Bulgarian oilseed rape fields and recorded 26 taxa, but, interestingly, did not find any *Trichomalus* species. Parasitoids of the cabbage flea beetle (*P. nemorum*) are represented only by *Townesilitus bicolor*, while other known species are absent or could not be determined at the species level, but only at the genus level. The most numerous group of parasitoids we found were aphid parasitoids, although their dominance was partly influenced by sampling methods, as about half of the specimens were from aphid rearing. *Diaeretiella rapae* and *Praon volucre* are known parasitoids of both cabbage aphid (*B. brassicae*) and peach-potato aphid (*M. persicae*), while *Aphidus ervi* and *A. matricariae* parasitise *M. persicae* (Tomanović et al. 2021).

It is unclear why some of the parasitoid species, which are widespread and numerous in various previously studied parts of Europe, do not occur at all or in greater numbers in Serbia. Nilsson (2003), while noting that there are no records from most of the Balkans, states that the parasitoid wasp fauna of this region should largely resemble that of Central Europe and that all Central European species should probably be present here. This is a reasonable expectation since, for example, the three most important pollen beetle parasitoids (*P. morionellus*, *P. interstitialis* and *T. heterocerus*) occur in neighbouring Hungary (all three species) and Bulgaria (*T. heterocerus*) (van Achterberg and Zwakhals 2023). Furthermore, since the Balkans is a hotspot of biodiversity in Europe, the parasitoid wasp fauna of Serbia can be expected to be richer than that of Central Europe (Griffiths et al. 2004). Petrović (2022) compared the parasitoid wasp fauna of the subfamily Aphidiinae from different European countries and found that only the Czech fauna had greater diversity (about 135 species) than the Serbian fauna (121 species), while other countries, such as Germany (109), the United Kingdom (96) and Norway (26), had fewer species, although the Czech fauna may simply be much better studied. To date, only Todorov et al. (2022) have published a partial list of oilseed parasitoids for one Balkan country, Bulgaria and only for the family Pteromalidae. However, only two species (*Mesopolobus incultus* and *M. morys*) are recorded in both Bulgaria and Serbia, with the possible existence of several other taxa of the genera *Mesopolobus* and *Pteromalus* that are difficult to identify to species level.

We obtained the most specimens by two sampling methods, sweep-net sampling (about 1700) and rearing from aphid colonies (about 1000). However, it should be noted that sampling with sweep nets and pan traps yielded more diverse samples (106 and 88 taxa, respectively), although there was no way to directly associate the collected specimens with host pests or plants. On the other hand, rearing from oilseed pods or from aphid colonies provided more targeted samples of either seed pod pests (such as *D. brassicae* and *C. obstrictus*) or aphid species (*B. brassicae* and *M. persicae*). Most of the detected parasitoid

families, with meaningfully large number of individuals, tended to be collected with sweep-net sampling, with the exception of *Platygastridae* and *Mymaridae*, which tended to occur more in pan traps. Therefore, the choice of sampling method can considerably limit the range of species detected and should be adapted to the research questions. Based on the examination of the species accumulation curve for samples from oilseed rape fields, it is clear that observed species richness obtained through our combined sampling methodology did not reach a plateau and that actual species richness is even higher. This is further corroborated by different species richness estimators which show a range of values, from 138 species as lowest estimate to 193 species as highest. We also found a seemingly large proportion of rare species (singletons and doubletons) in the oilseed rape fields in our study. About 38% were found only with one specimen and about 19% with two individuals. However, Scharff et al. (2003) give an overview of undersampling bias in terrestrial arthropod surveys and argue that, even in the “state of the art” arthropod surveys, about 50–70% of species are known from just one or two individuals.

## Conclusions

Our study of the parasitoid community associated with oilseed rape fields revealed great diversity of parasitoid species. Interestingly, the majority of these parasitoids were not known to parasitise oilseed rape pest species, suggesting the existence of alternative hosts or other ecological interactions within oilseed rape fields. Additional factors should be carefully measured, such as the previous season's crop in each field, the presence of volunteer plants and weed species and associated herbivores, to gain a better insight into the various interactions that appear to exist in oilseed rape fields. We also observed notable differences in the occurrence patterns of different parasitoid species in Serbia and other parts of Europe, indicating the need for further studies of the ecological dynamics and host associations of parasitoid species in oilseed rape fields. Understanding the factors affecting the presence, abundance and distribution of parasitoids will contribute to more effective regional and local pest management strategies.

## Acknowledgements

This work was supported by the European Union's Horizon 2020 Research and Innovation Programme, under grant agreement no. 773554 (EcoStack) and by the Serbian Ministry of Science and Education (451-03-47/2023-01/ 200178) and Serbian Academy of Sciences and Arts (grant F131). We thank Alex Gumovsky (Schmalhausen Institute of Zoology, Kiev, Ukraine) for help in identification of Eulophids. We also thank colleagues from the Institute for Field and Vegetable Crops, Novi Sad and Florijan Farkaš for providing contacts with the farmers.

## References

- Alford DV (2003) Biocontrol of oilseed rape pests. Blackwell Science Ltd. [ISBN 9780632054275] <https://doi.org/10.1002/9780470750988>
- Angus JF, Kirkegaard JA, Hunt JR, Ryan MH, Ohlander L, Peoples MB (2015) Break crops and rotations for wheat. *Crop and Pasture Science* 66 (6). <https://doi.org/10.1071/cp14252>
- Bianchi FJ, Booij CJ, Tscharntke T (2006) Sustainable pest regulation in agricultural landscapes: a review on landscape composition, biodiversity and natural pest control. *Proceedings of the Royal Society B: Biological Sciences* 273 (1595): 1715-1727. <https://doi.org/10.1098/rspb.2006.3530>
- Dainese M, Schneider G, Krauss J, Steffan-Dewenter I (2017) Complementarity among natural enemies enhances pest suppression. *Scientific Reports* 7 (1). <https://doi.org/10.1038/s41598-017-08316-z>
- FAO (2022) Crops and livestock products. URL: <https://www.fao.org/faostat/en/#data/QCL>
- Graora D, Sivčev I, Sivčev L, Buchs W (2013) Brojnost i štetnost *Dasineura brassicae* (Winn.) u različitim sistemima gajenja uljane repice. *Zaštita bilja* 64 (3): 144-153. [In Serbian].
- Graora D, Sivcev I, Sivcev L, Buchs W, Tomic V, Dudic B, Gotlin-Culjak T (2015) Biology and harmfulness of Brassica pod midge (*Dasineura brassicae* Winn.) in winter oilseed rape. *Pesticidi i fitomedicina* 30 (2): 85-90. <https://doi.org/10.2298/pif1502085g>
- Griffiths HI, Kryštufek B, Reed JM (Eds) (2004) *Balkan biodiversity*. Springer Netherlands, 358 pp. <https://doi.org/10.1007/978-1-4020-2854-0>
- Gunton R, Pöyry J (2016) Scale-specific spatial density dependence in parasitoids: a multi-factor meta-analysis. *Functional Ecology* 30 (9): 1501-1510. <https://doi.org/10.1111/1365-2435.12627>
- Heimpel GE, Jervis MA (2005) Does floral nectar improve biological control by parasitoids? In: Wäckers FL, van Rijn PC, Bruin J (Eds) *Plant-provided food for voracious insects: a protective mutualism and its applications*. Cambridge University Press <https://doi.org/10.1017/cbo9780511542220.010>
- Hokkanen HMT (2008) Biological control methods of pest insects in oilseed rape. *EPPO Bulletin* 38 (1): 104-109. <https://doi.org/10.1111/j.1365-2338.2008.01191.x>
- Kavallieratos N, Tomanović Ž, Starý P, Athanassiou C, Sarlis G, Petrović O, Niketić M, Veroniki MA (2004) A survey of aphid parasitoids (Hymenoptera: Braconidae: Aphidiinae) of Southeastern Europe and their aphid-plant associations. *Applied Entomology and Zoology* 39 (3): 527-563. <https://doi.org/10.1303/aez.2004.527>
- Kavallieratos N, Tomanović Ž, Athanassiou C, Starý P, Žikić V, Sarlis G, Fasseas C (2005a) Aphid parasitoids infesting cotton, citrus, tobacco, and cereal crops in southeastern Europe: aphid-plant associations and keys. *The Canadian Entomologist* 137 (5): 516-531. <https://doi.org/10.4039/n05-020>
- Kavallieratos N, Tomanović Ž, Starý P, Athanassiou C, Fasseas C, Petrović O, Stanisavljević L, Veroniki MA (2005b) *Praon* Haliday (Hymenoptera: Braconidae: Aphidiinae) of Southeastern Europe: key, host range and phylogenetic relationships. *Zoologischer Anzeiger - A Journal of Comparative Zoology* 243 (3): 181-209. <https://doi.org/10.1016/j.jcz.2004.11.001>

- Kereši T, Sekulić R, Štrbac P (2007) Ostale važne štetočine uljane repice. Biljni lekar XXXV (4). [In Serbian].
- Landis D, Wratten S, Gurr G (2000) Habitat Management to Conserve Natural Enemies of Arthropod Pests in Agriculture. Annual Review of Entomology 45 (1): 175-201. <https://doi.org/10.1146/annurev.ento.45.1.175>
- Marjanović J, Prodanović S (2021) Good agricultural practices and technologies to mitigate the impacts of natural disasters in oilseed rape production in Serbia. FAO. URL: <https://www.fao.org/documents/card/en/c/CB5193EN>
- Milovac Ž, Pešić S, Kereši T, Marinković R (2010) Weevils (Coleoptera: Curculionoidea) – important members of rapeseed entomofauna in vicinity of Novi Sad. Kragujevac J. Sci. 32: 141-148.
- Milovac Ž (2016) Bionomija i mogućnosti suzbijanja repičinih pipa *Ceutorhynchus pallidactylus* (Marsham) i *Ceutorhynchus napi* Gyllenhal (Coleoptera: Curculionidae). University of Belgrade, Belgrade, 120 pp. [In Serbian].
- Milovanović P, Kljajić P, Andrić G, Prazić-Golić M, Popović T (2013) Efficacy of different insecticides in controlling pollen beetle (*Meligethes aeneus* F.) in rapeseed crop. Pesticidi i fitomedicina 28 (4): 255-263. <https://doi.org/10.2298/pif1304255m>
- Mitrović P, Milovac Ž, Marinković R (2008) Aktuelni problemi u zaštiti ozime uljane repice (*Brassica napus* var. *napus* L.). Zaštita bilja 60 (2): 127-144. [In Serbian].
- Nilsson C (2003) Parasitoids of pollen beetles. In: Alford DV (Ed.) Biocontrol of oilseed rape pests. Blackwell Science Ltd <https://doi.org/10.1002/9780470750988.ch4>
- Noyes JS (2019) Universal chalcidoidea database. URL: <http://www.nhm.ac.uk/chalcidoidea>
- Oksanen J, Simpson G, Blanchet F, Kindt R, Legendre P, Minchin P, O'Hara R, Solymos P, Stevens M, Szoecs E, Wagner H, Barbour M, Bedward M, Bolker B, Borcard D, Carvalho G, Chirico M, De Caceres M, Durand S, Evangelista H, FitzJohn R, Friendly M, Furneaux B, Hannigan G, Hill M, Lahti L, McGlenn D, Ouellette M, Ribeiro Cunha E, Smith T, Stier A, Ter Braak C, Weedon J (2022) `vegan`: Community Ecology Package. R package version 2.6-4. URL: <https://CRAN.R-project.org/package=vegan>
- Petrović A (2022) tSizing the knowledge gap in axonomy: The last dozen years of Aphidiinae research. Insects 13 (2). <https://doi.org/10.3390/insects13020170>
- Powell JE, Shepard M (1982) Biology of Australian and United States strains of *Trissolcus basalıs*, a parasitoid of the green vegetable bug, *Nezara viridula*. Austral Ecology 7 (2): 181-186. <https://doi.org/10.1111/j.1442-9993.1982.tb01591.x>
- R Core Team (2023) R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing. URL: <https://www.R-project.org/>
- RZS (2023) Republički zavod za statistiku STAT baza.
- Schaefer CW, Panizzi AR (Eds) (2000) Heteroptera of economic importance. CRC Press, Boca Raton, 856 pp. <https://doi.org/10.1201/9781420041859>
- Scharff N, Coddington J, Griswold C, Hormiga G, Bjørn PdP (2003) When to quit? Estimating spider species richness in a northern European deciduous forest. Journal of Arachnology 31 (2): 246-273. [https://doi.org/10.1636/0161-8202\(2003\)031\[0246:wqess\]2.0.co;2](https://doi.org/10.1636/0161-8202(2003)031[0246:wqess]2.0.co;2)
- Sekulić R, Kereši T (2007) Repičin sjajnik *Meligethes aeneus* najvažnija stetočina ozime uljane repice. Biljni lekar 25.

- Sivčev I, Sivčev L, Pešić S, Graora D, Tomić V, Dudić B (2015) Weevils of the genus *Ceutorhynchus* Germ associated with oilseed rape in northern Serbia. *Pesticidi i fitomedicina* 30 (3): 155-159. <https://doi.org/10.2298/pif1503155s>
- Sivčev L, Graora D, Sivčev I, Tomić V, Dudić B (2016) Phenology of cabbage stem flea beetle (*Psylliodes chrysocephala* L.) in oilseed rape. *Pesticidi i fitomedicina* 31: 139-144. <https://doi.org/10.2298/pif1604139s>
- Stanley D, Gunning D, Stout J (2013) Pollinators and pollination of oilseed rape crops (*Brassica napus* L.) in Ireland: ecological and economic incentives for pollinator conservation. *Journal of Insect Conservation* 17 (6): 1181-1189. <https://doi.org/10.1007/s10841-013-9599-z>
- Statista (2023) Production of major vegetable oils worldwide from 2012/13 to 2022/2023, by type. URL: <https://www.statista.com/statistics/263933/production-of-vegetable-oils-worldwide-since-2000/>
- Štrbac P, Kereši T, Sekulić R (2007) Zaštita uljane repice of repičine ose (*Athalia rosae*). *Biljni lekar* XXXV (4). [In Serbian].
- Thies C, Tscharntke T (1999) Landscape Structure and Biological Control in Agroecosystems. *Science* 285 (5429): 893-895. <https://doi.org/10.1126/science.285.5429.893>
- Thompson WR (1958) A catalogue of the parasites and predators of insect pests. Section 2. Host parasite catalogue, Part 5. Commonwealth Agricultural Bureaux, Commonwealth Institute of Biological Control, Ottawa, Ontario, Canada.
- Todorov I, Ljubomirov T, Peneva V (2022) Pteromalid fauna (Hymenoptera, Pteromalidae) in oilseed rape (*Brassica napus* L.) fields in Bulgaria – species composition and perspectives for biological control. *BioRisk* 17: 329-342. <https://doi.org/10.3897/biorisk.17.77454>
- Tomanović Ž, Kavallieratos NG, Starý P, Athanassiou CG, Žikić V, Petrović-Obradović O, Sarlis GP (2003) *Aphidius* Nees aphid parasitoids (Hymenoptera, Braconidae, Aphidiinae) in Serbia and Montenegro: tritrophic associations and key. *Acta Entomologica Serbica* 8 (1/2): 15-39.
- Tomanović Ž, Kavallieratos NG, Starý P, Petrović-Obradović O, Athanassiou CG, Stanisavljević L (2008) Cereal aphids (Hemiptera: Aphidoidea) in Serbia: Seasonal dynamics and natural enemies. *Eur. J. Entomol.* 105 (3): 495-501. <https://doi.org/10.14411/eje.2008.064>
- Tomanović Ž, Žikić V, Petrović A (2021) Fauna parazoidnih osa (Hymenoptera, Braconidae, Aphidiinae) Srbije. *Srpska Akademija Nauka i umetnosti*
- Townes H (1971) The Genera of Ichneumonidae, Part 4 (Vol. 17). *Memoirs of the American Entomological Institute*
- Ulber B (2003) Parasitoids of Ceutorhynchid Stem Weevils. In: Alford DV (Ed.) *Biocontrol of oilseed rape pests*. Blackwell Science Ltd, 45–76 pp. <https://doi.org/10.1002/9780470750988.ch5>
- Ulber B, Williams IH, Klukowski Z, Luik A, Nilsson C (2010) Parasitoids of oilseed rape pests in Europe: key species for conservation biocontrol. In: Williams I (Ed.) *Biocontrol-based integrated management of oilseed rape pests*. Springer, Dordrecht. [https://doi.org/10.1007/978-90-481-3983-5\\_2](https://doi.org/10.1007/978-90-481-3983-5_2)
- van Achterberg K, Zwakhals K (2023) Fauna Europaea: Ichneumonidae. *Fauna Europaea* version 2023.05.

- Varennes YD, Gonzalez Chang M, Boyer S, Wratten SD (2016) Nectar feeding increases exploratory behaviour in the aphid parasitoid *Diaeretiella rapae* (McIntosh). *Journal of Applied Entomology* 140 (6): 479-483. <https://doi.org/10.1111/jen.12281>
- Williams I (2010) Biocontrol-based integrated management of oilseed rape pests. Springer, Dordrecht. <https://doi.org/10.1007/978-90-481-3983-5>
- Yu DS, van Achterberg C, Horstmann K (2011) *World Ichneumonidea 2011. Taxonomy, biology, morphology and distribution.*
- Zapponi L, Tortorici F, Anfora G, Bardella S, Bariselli M, Benvenuto L, Bernardinelli I, Butturini A, Caruso S, Colla R, Costi E, Culatti P, Di Bella E, Falagiarda M, Giovannini L, Haye T, Maistrello L, Malossini G, Marazzi C, Marianelli L, Mele A, Michelon L, Moraglio ST, Pozzebon A, Preti M, Salvetti M, Scaccini D, Schmidt S, Szalatnay D, Roversi PF, Tavella L, Tommasini MG, Vaccari G, Zandigiaco P, Sabbatini-Peverieri G (2021) Assessing the distribution of exotic egg parasitoids of *Halyomorpha halys* in Europe with a large-scale monitoring program. *Insects* 12 (4). <https://doi.org/10.3390/insects12040316>
- Žikić V, Ilić-Milošević M, Stanković S, Petrović A, Petrović-Obrovčić O, Kavallieratos NG, Starý P, Tomanović Ž (2012) Aphidiinae (Hymenoptera: Braconidae) of Serbia and Montenegro – tritrophic interactions. *Acta Entomologica Serbica* 17 (1/2): 83-105.

## Supplementary materials

### Suppl. material 1: Supplements Table 1. List of identification keys used. [doi](#)

**Authors:** Milan Plečaš, Vladimir Žikić, Korana Kocić, Jelisaveta Črkrić, Anđeljko Petrović, Željko Tomanović

**Data type:** List of references

[Download file](#) (17.87 kb)

### Suppl. material 2: Supplement Table 2a. Checklist of parasitoids found in oilseed rape fields in Serbia [doi](#)

**Authors:** Milan Plečaš, Vladimir Žikić, Korana Kocić, Jelisaveta Črkrić, Anđeljko Petrović, Željko Tomanović

**Data type:** Checklist with additional data

**Brief description:** Detailed information on locations, collection dates, habitat, sampling methods, parasitism type, oilseed rape pest host status and number of specimens for each taxon.

[Download file](#) (68.73 kb)



**Suppl. material 3: Supplement Table 2b. Checklist of parasitoids found in oilseed rape fields in Serbia in csv format** [doi](#)

**Authors:** Milan Plečaš, Vladimir Žikić, Korana Kocić, Jelisaveta Črkić, Anđeljko Petrović, Željko Tomanović

**Data type:** Checklist with additional data

**Brief description:** Detailed information on locations, collection dates, habitat, sampling methods, parasitism type, oilseed rape pest host status and number of specimens for each taxon provided in cvs format.

[Download file](#) (38.57 kb)

**Suppl. material 4: Supplement Table 3. Checklist of parasitoids found in oilseed rape fields in Serbia with additional host information.** [doi](#)

**Authors:** Milan Plečaš, Vladimir Žikić, Korana Kocić, Jelisaveta Črkić, Anđeljko Petrović, Željko Tomanović

**Data type:** Checklist with additional data

**Brief description:** Detailed information on OSR pest host status, parasitism type, host species and host stage and references.

[Download file](#) (70.02 kb)