



# First report of *Trichopoda pictipennis* (Diptera, Tachinidae) for the Canary Islands

Pablo Caloca<sup>‡</sup>, Daniel Suárez<sup>‡</sup>, Gustavo Peña<sup>‡</sup>, Carlos Ruiz<sup>‡</sup>

<sup>‡</sup> Departamento de Biología Animal, Edafología y Geología, Facultad de Ciencias, Universidad de La Laguna, San Cristóbal de La Laguna, Spain

Corresponding author: Daniel Suárez ([danielsura94@gmail.com](mailto:danielsura94@gmail.com))

Academic editor: Pierfilippo Cerretti

Received: 25 Sep 2024 | Accepted: 13 Dec 2024 | Published: 05 Feb 2025

Citation: Caloca P, Suárez D, Peña G, Ruiz C (2025) First report of *Trichopoda pictipennis* (Diptera, Tachinidae) for the Canary Islands. Biodiversity Data Journal 13: e137821.

<https://doi.org/10.3897/BDJ.12.e137821>

## Abstract

## Background

The genus *Trichopoda* Berthold, 1827 is distributed in the Neotropical and Nearctic Regions and some species are very important for biological control. During the last decades, the species *Trichopoda pictipennis* Bigot, 1876 has received much attention. It is of Neotropical origin, but it has been introduced throughout the western Palaearctic, probably through exchanges that transported its main host, the 'southern green stink bug' *Nezara viridula*.

## New information

*Trichopoda pictipennis* is reported for the first time from the Canary Islands. To date, this introduced species has been detected throughout the island of Tenerife and in a single locality on the island of La Gomera. This finding confirms that this species of Neotropical origin is expanding its range across the Palaearctic realm of this species of Neotropical origin. Parasitised *Nezara viridula* were collected and reared under laboratory conditions

to document the complete life cycle of *T. pictipennis*. Its potential effects on its main host, as well as on the Canary fauna, are discussed.

## Keywords

new records, parasitoid, faunistics, pest control, Macaronesia, introduced species, Phasiinae, Pentatomidae, *Nezara viridula*

## Introduction

The family Tachinidae (Calyptatae, Oestroidea) comprises ca. 8,500 valid species worldwide (O'Hara et al. 2020). This family is especially interesting, as all species are endoparasitoids of arthropods (Stireman et al. 2006) and several species visit flowers for feeding (Tooker et al. 2006). The genus *Trichopoda* Berthold, 1827, which includes the species commonly called 'feather-legged flies', is distributed in the Neotropical and Nearctic realms and some species are very important for biological control (Dios and Nihei 2020). During the last decades, the species *Trichopoda pictipennis* Bigot, 1876 has received much attention. It is of Neotropical origin (Dios et al. 2021), but was unintentionally introduced in Italy on the 1980s, probably through exchanges that transported its only known main host, the 'southern green stink bug' *Nezara viridula* (Linnaeus, 1758) (Colazza et al. 1996). Since then, it has been spreading throughout the western Palaearctic. The second record of the species was in north-eastern Spain in 1995 (Peris 1998). During the last two decades, it has been detected in the following countries: France (Tschorasnig et al. 2000), Slovenia (de Groot et al. 2007), The Netherlands (Zeegers 2010), Italy (Cargnus et al. 2011), Israel (Freidberg et al. 2011), Croatia (Bystrowski 2012), Switzerland (Obrecht 2015), Portugal (Almeida et al. 2017), Malta (Darmanin and Cerretti 2019), Egypt (El-Hawagry et al. 2020), Albania, Cyprus, Greece, Turkey, Russia (Kazilas et al. 2020), Algeria (Meriem et al. 2021), Germany (Dios et al. 2021) and Morocco (Kettani et al. 2022). It is still unknown whether this expansion is the result of long-distance dispersal from Italy or if multiple introduction events occurred (Bystrowski 2012). A distributional study of *T. pictipennis* in Spain and Portugal, which revealed a strong affinity for coastal Mediterranean areas, suggested that the main way of entrance for this species might be port areas, probably associated with parasitised *N. viridula* individuals on fresh fruit and vegetables (Ricarte et al. 2020).

The Canary Islands, an oceanic archipelago within the Macaronesian Region, stands out for its high endemicity rate in arthropods (Oromí et al. 2015, Florencio et al. 2021). Diptera are relevant, as they constitute the second most species-rich insect order of the archipelago, with ca. 1,200 species, almost a third of them being endemics (Gobierno de Canarias 2024). In the Canary Islands, the tachinid assemblage has not received much attention historically, with only few contributions focusing specifically on this family (Báez et al. 1986, Tschorasnig et al. 2007, Suárez et al. 2020). In the present paper, *Trichopoda pictipennis* is reported for the first time from the Canary Islands, more specifically from the islands of Tenerife and La Gomera.

## Materials and methods

The collected individuals were pinned and examined under a Zeiss Stemi 2000 stereomicroscope and individuals were identified to species level by using the key of Dios and Nihei (2020). For study of the female genitalia, the last abdominal segments of one specimen were dissected and placed in 10% potassium hydroxide (KOH). Once the genitalia were visible, they were placed in glycerine and examined under a stereomicroscope. Illustrations of the terminalia were made using the vector graphics editor Inkscape, based on photos taken with a Canon EOS 750D camera. The material is deposited in the collection of the Department of Animal Biology of the University of La Laguna, Tenerife (DZUL). Additionally, in order to document the complete life cycle of *T. pictipennis*, a sampling of its main host *Nezara viridula* was conducted. Live specimens that had eggs on the integument, indicating that they were potentially parasitised, were collected and reared in the laboratory.

## Taxon treatment

### *Trichopoda pictipennis* Bigot, 1876

#### Materials

- a. scientificName: *Trichopoda pictipennis* Bigot, 1876; taxonID: <https://www.gbif.org/es/species/5059806>; order: Diptera; family: Tachinidae; genus: *Trichopoda*; island: Tenerife; country: Spain; stateProvince: Santa Cruz de Tenerife; municipality: Fasnia; locality: Fasnia; decimalLatitude: 28.245555; decimalLongitude: -16.435555; georeferenceProtocol: GPS; eventDate: 19/01/2022; eventRemarks: On flowers of *Argyranthemum frutescens* (Asteraceae); individualCount: 1; lifeStage: adult; recordedBy: Gustavo Peña; identifiedBy: Gustavo Peña; dateIdentified: 2023; basisOfRecord: HumanObservation; occurrenceID: CB27062D-1AF5-59F2-955F-4A40C2482BA8
- b. scientificName: *Trichopoda pictipennis* Bigot, 1876; taxonID: <https://www.gbif.org/es/species/5059806>; order: Diptera; family: Tachinidae; genus: *Trichopoda*; island: Tenerife; country: Spain; stateProvince: Santa Cruz de Tenerife; municipality: Fasnia; locality: Fasnia; decimalLatitude: 28.245555; decimalLongitude: -16.435555; georeferenceProtocol: GPS; eventDate: 20/06/2022; eventRemarks: On flowers of *Zygophyllum fontanesii* (Zygophyllaceae); individualCount: 1; lifeStage: adult; recordedBy: Gustavo Peña; identifiedBy: Gustavo Peña; dateIdentified: 2023; basisOfRecord: HumanObservation; occurrenceID: D28506C1-567C-5AAA-B53A-D442F905DC51
- c. scientificName: *Trichopoda pictipennis* Bigot, 1876; taxonID: <https://www.gbif.org/es/species/5059806>; order: Diptera; family: Tachinidae; genus: *Trichopoda*; island: Tenerife; country: Spain; stateProvince: Santa Cruz de Tenerife; municipality: Puerto de la Cruz; locality: Puerto de la Cruz; decimalLatitude: 28.402587; decimalLongitude: -16.556857; georeferenceProtocol: GPS; eventDate: 01/03/2022; eventRemarks: On leaves of *Bosea yervamora* (Amaranthaceae); individualCount: 1; sex: female; lifeStage: adult; catalogNumber: DZUL-36962; recordedBy: Pablo Caloca; identifiedBy: Daniel Suárez; dateIdentified: 2023; basisOfRecord: PreservedSpecimen; occurrenceID: CA2F2897-64C6-5DB7-BE32-193571A1A877

- d. scientificName: *Trichopoda pictipennis* Bigot, 1876; taxonID: <https://www.gbif.org/es/species/5059806>; order: Diptera; family: Tachinidae; genus: *Trichopoda*; island: Tenerife; country: Spain; stateProvince: Santa Cruz de Tenerife; municipality: Puerto de la Cruz; locality: Puerto de la Cruz; decimalLatitude: 28.402587; decimalLongitude: -16.556857; georeferenceProtocol: GPS; eventDate: 01/03/2022; eventRemarks: On leaves of *Bosea yervamora* (Amaranthaceae); individualCount: 1; sex: male; lifeStage: adult; catalogNumber: DZUL-36963; recordedBy: Pablo Caloca; identifiedBy: Daniel Suárez; dateIdentified: 2023; basisOfRecord: PreservedSpecimen; occurrenceID: 08A9DD48-C7BB-564B-A415-D59575239239
- e. scientificName: *Trichopoda pictipennis* Bigot, 1876; taxonID: <https://www.gbif.org/es/species/5059806>; order: Diptera; family: Tachinidae; genus: *Trichopoda*; island: Tenerife; country: Spain; stateProvince: Santa Cruz de Tenerife; municipality: San Cristóbal de La Laguna; locality: San Cristóbal de La Laguna; decimalLatitude: 28.403688; decimalLongitude: -16.335111; georeferenceProtocol: GPS; eventDate: 28/02/2022; individualCount: 1; sex: male; lifeStage: adult; recordedBy: María de Fuentes; identifiedBy: Pablo Caloca; dateIdentified: 2023; basisOfRecord: HumanObservation; occurrenceID: ED223ECF-8281-5D7C-BAC5-01F6BB577CC3
- f. scientificName: *Trichopoda pictipennis* Bigot, 1876; taxonID: <https://www.gbif.org/es/species/5059806>; order: Diptera; family: Tachinidae; genus: *Trichopoda*; island: Tenerife; country: Spain; stateProvince: Santa Cruz de Tenerife; municipality: Los Realejos; locality: Los Realejos; decimalLatitude: 28.403688; decimalLongitude: -16.575571; georeferenceProtocol: GPS; eventDate: 16/01/2023; eventRemarks: On leaves of *Bosea yervamora* (Amaranthaceae); individualCount: 1; sex: male; lifeStage: adult; catalogNumber: DZUL-36964; recordedBy: Pablo Caloca; identifiedBy: Pablo Caloca; dateIdentified: 2023; basisOfRecord: PreservedSpecimen; occurrenceID: DEEF9D8F-D991-55FF-87CB-07D8A95011E0
- g. scientificName: *Trichopoda pictipennis* Bigot, 1876; taxonID: <https://www.gbif.org/es/species/5059806>; order: Diptera; family: Tachinidae; genus: *Trichopoda*; island: Tenerife; country: Spain; stateProvince: Santa Cruz de Tenerife; municipality: Tegueste; locality: Tegueste; decimalLatitude: 28.544774; decimalLongitude: -16.318082; georeferenceProtocol: GPS; eventDate: 16/01/2023; eventRemarks: On flowers of *Foeniculum vulgare* (Apiaceae); individualCount: 1; sex: male; lifeStage: adult; recordedBy: Carlos Ruiz; identifiedBy: Carlos Ruiz; dateIdentified: 2023; basisOfRecord: HumanObservation; occurrenceID: 226F34DD-88B0-5189-AD16-9EB4D917469D
- h. scientificName: *Trichopoda pictipennis* Bigot, 1876; taxonID: <https://www.gbif.org/es/species/5059806>; order: Diptera; family: Tachinidae; genus: *Trichopoda*; island: Tenerife; country: Spain; stateProvince: Santa Cruz de Tenerife; municipality: San Cristóbal de La Laguna; locality: Punta del Hidalgo; decimalLatitude: 28.544774; decimalLongitude: -16.318082; georeferenceProtocol: GPS; eventDate: 02/02/2023; eventRemarks: On flowers of *Crithmum maritimum* (Apiaceae); individualCount: 1; lifeStage: adult; recordedBy: Eduardo Jiménez; identifiedBy: Eduardo Jiménez; dateIdentified: 2023; basisOfRecord: HumanObservation; occurrenceID: 8A6D3CD8-1C58-5568-9C19-6901CA0988B7
- i. scientificName: *Trichopoda pictipennis* Bigot, 1876; taxonID: <https://www.gbif.org/es/species/5059806>; order: Diptera; family: Tachinidae; genus: *Trichopoda*; island: Tenerife; country: Spain; stateProvince: Santa Cruz de Tenerife; municipality: San Miguel de Abona; locality: San Miguel de Abona; decimalLatitude: 28.030278; decimalLongitude: -16.627952; georeferenceProtocol: GPS; eventDate: 05/02/2023; individualCount: 1; lifeStage: adult; recordedBy: Carmen Delía González; identifiedBy: Carlos Ruiz;

dateIdentified: 2023; basisOfRecord: HumanObservation; occurrenceID: 2FD96D2F-7FC1-528F-B196-8CFCBF2B23001  
j. scientificName: *Trichopoda pictipennis* Bigot, 1876; taxonID: <https://www.gbif.org/es/species/5059806>; order: Diptera; family: Tachinidae; genus: *Trichopoda*; island: La Gomera; country: Spain; stateProvince: Santa Cruz de Tenerife; municipality: San Sebastián; locality: San Sebastián; decimalLatitude: 28.092127; decimalLongitude: -17.114348; georeferenceProtocol: GPS; eventDate: 29/01/2023; individualCount: 2; lifeStage: adult; recordedBy: Uquén Fernández; identifiedBy: Pablo Caloca; dateIdentified: 2023; basisOfRecord: HumanObservation; occurrenceID: 93AA6364-7D8B-5233-BD62-D09B0112CBAC

### Diagnosis

The collected individuals were identified as *Trichopoda pictipennis* by the following combination of diagnostic characters. Males: yellow to orange abdomen, yellow scutellum, black wing with a yellow marking, lower calypter orange (Fig. 1A). Females: black abdomen and scutellum, black wing with a yellow marking (Fig. 1B). Female genitalia: sternite 7 as a trapezoidal plate, distal margin slightly rounded; sternite 8 lingulate, apically round; cercus subquadrate (Fig. 1C).




Figure 1. [doi](#)

Habitus, genitalia and life cycle (partial) of *Trichopoda pictipennis*. **A** Male of *Trichopoda pictipennis*, photo by Pablo Caloca; **B** Female of *Trichopoda pictipennis*, photo by Pablo Caloca; **C** Female genitalia of *Trichopoda pictipennis*; **D** Parasitised individuals of *Nezara viridula* at different life stages, photo by Pablo Caloca.

### Biology

Parasitised individuals of *Nezara viridula* were collected from leaves of *Bosea yervamora* (Amaranthaceae) in the localities of Puerto de La Cruz and Los Realejos

(north of Tenerife) on 17 December 2022 (Fig. 1D). *Trichopoda pictipennis* full-grown larva emerged on 22 December 2022 and puparated outside its host. The adult emerged on 8 January 2023.

## Discussion

The present record of *Trichopoda pictipennis* is not only the first report of the species for the Canary Islands, but also for the Macaronesian Region (Arechavaleta et al. 2005, Borges et al. 2008, Borges et al. 2022, Gobierno de Canarias 2024). The species can be easily distinguished from any other tachinid species currently known from Macaronesia by the presence of feather-like setae on the hind tibia. *T. pictipennis* had previously been misidentified as *T. pennipes* Fabricius, 1781 throughout the Palaearctic realm until Dios et al. (2021). Its correct identification for further use in biological control is crucial to avoid harmful impacts on Pentatomidae native communities (Dios and Nihei (2020)). The distribution of *T. pictipennis* in the Canary Islands partially fits (a third of the records) with the commercial ports present at Tenerife and La Gomera (Fig. 2), possibly confirming the hypothesis of Ricarte et al. (2020) for mainland Spain. Two-thirds of the records were found far from commercial ports, possibly indicating expansions throughout the islands. This mode of entrance has also been proposed for other introduced species on the Canary Islands (Martínez and Barone 2006, López-dos-Santos et al. 2013, Ruiz et al. 2020, Lugo et al. 2022). However, this hypothesis should be tested with a phylogeographic study comprising material from across its distribution range, both native and introduced.




Figure 2. [doi](#)

Map showing the localities where *T. pictipennis* has been reported (red dots). Commercial ports in Tenerife and La Gomera have been marked with a ship symbol.

The main host of *T. pictipennis* is *Nezara viridula* (Cerretti and Tschorasnig 2010, Tschorasnig 2017), which is present on all the major islands of the Canary Archipelago (Gobierno de Canarias 2024), implying the potential for *T. pictipennis* to spread to any island. This globally introduced species is known to feed on native plants in natural and agricultural ecosystems, thus causing economic damage to crops where it has been introduced, especially in subtropical and tropical regions (García et al. 1992, Esquivel et al. 2019). The recent arrival and establishment of *T. pictipennis* may have a regulatory impact on *N. viridula*. Although data collected in Italy suggests a moderate effect on parasitism rate (Salerno et al. 2002), differences in climatic conditions may lead to different rates on the Archipelago. Potentially, *T. pictipennis* could also shift to other Pentatomidae species. Outside its native area, there is only a single record of a host other than *N. viridula*, the Mediterranean species *Graphosoma lineatum* (Linneaus, 1758) (Colazza et al. 1996, Cerretti and Tschorasnig 2010, Tschorasnig 2017). However, under laboratory conditions, *T. pictipennis* was able to parasitise additional three Australian native species belonging to different genera of Pentatomidae (Sands and Coombs 1999). The fact that there is just one known record of a host different from *N. viridula* suggests that *T. pictipennis* is a specialist. To date, we have only detected parasitism on *N. viridula*, with adults emerging from pupae 18 days after larvae emerged from its host. However, the idea of a potential niche shift should not be discarded. Insular biota exhibit several geographic, demographic and genetic characteristics that make them particularly vulnerable to invasive species (Leclerc et al. 2018, Russell and Kaiser-Bunbury 2019, Fernández-Palacios et al. 2021). In the Canary Islands, there are 38 native species of Pentatomidae, including seven endemic species. Therefore, it is imperative to study whether *T. pictipennis* can parasitise other species in the Canary Islands, with the aim of mitigating potential adverse impacts on the native Pentatomidae fauna.

## Acknowledgements

The authors wish to thank María de Fuentes, Eduardo Jiménez, Carmen Delia González and Uquèn Fernández for letting us use their observations in the present article. We also want to thank Rodrigo Dios for his help with the identification of this species. We are very grateful to Daniel Whitmore, Hans-Peter Tschorasnig and Pierfilippo Cerretti for their useful comments and suggestions on a previous version of the manuscript.

## References

- Almeida JM, Andrade R, Gonçalves AR, Jacinto V, Raper C, Tschorasnig HP, Zeegers T (2017) An annotated checklist of the Tachinidae (Diptera) of mainland Portugal. Boletín de la S.E.A. 60: 62-76.
- Arechavaleta M, Zurita N, Marrero MC, Martín JL (2005) Lista preliminar de especies silvestres de Cabo Verde (hongos, plantas y animales terrestres). Consejería de Medio Ambiente y Ordenación Territorial, Gobierno de Canarias, Santa Cruz de Tenerife.

- Báez M, Herting B, Tschorsnig HP (1986) The Tachinidae (Diptera) of the Canary Islands. Stuttgarter Beiträge Zur Naturkunde 394: 1-15.
- Borges P, Abreu C, Aguiar A, Carvalho P, Jardim R, Melo I, Oliveira P, Sérgio C, Serrano A, Vieira P (2008) A list of the terrestrial fungi, flora and fauna of Madeira and Selvagens archipelagos. Direcção Regional do Ambiente da Madeira and Universidade dos Açores, Funchal and Angra do Heroísmo.
- Borges P, Lamelas-Lopez L, Andrade R, Lhoumeau S, Vieira V, Soares A, Borges I, Boieiro M, Cardoso P, Crespo LC, Karsholt O, Schülke M, Serrano A, Quartau JA, Assing V (2022) An updated checklist of Azorean arthropods (Arthropoda). Biodiversity Data Journal 10 <https://doi.org/10.3897/bdj.10.e97682>
- Bystrowski C (2012) Thoughts on how *Trichopoda pennipes* (F.) reached Israel. The Tachinid Times 25: 13-14.
- Cargnus E, Buian FM, Zandigiacomo P (2011) Presenza di *Trichopoda pennipes* (Diptera, Tachinidae) nell'Italia nord-orientale. Bollettino Società Naturalisti "Silvia Zenari" 35: 123-130.
- Cerretti P, Tschorsnig HP (2010) Annotated host catalogue for the Tachinidae (Diptera) of Italy. Stuttgarter Beiträge Zur Naturkunde 3: 305-340.
- Colazza S, Giangiuliani G, Bin F (1996) Fortuitous introduction and successful establishment of *Trichopoda pennipes* F.: Adult parasitoid of *Nezara viridula* (L.). Biological Control 6 (3): 409-411. <https://doi.org/10.1006/bcon.1996.0053>
- Darmanin A, Cerretti P (2019) *Trichopoda pennipes* (Fabricius, 1781) - A new tachinid fly (Diptera) for Malta. Bulletin of the Entomological Society of Malta 10: 103-105.
- de Groot M, Virant-Doberlet M, Zunic A (2007) *Trichopoda pennipes* F. Diptera, Tachinidae): a new natural enemy of *Nezara viridula* (L.) in Slovenia. Agricultura 5: 25-26.
- Dios RDVP, Nihei SS (2020) Taxonomic revision of the genus *Trichopoda* Berthold, 1827 (Diptera: Tachinidae: Phasiinae), with emphasis on the Neotropical fauna. Zootaxa 4870 (1). <https://doi.org/10.11646/zootaxa.4870.1.1>
- Dios RP, Ziegler J, Zeegers T (2021) The American genus *Trichopoda* (Diptera: Tachinidae) in Europe – Decades of a misidentified invasive species. Beiträge zur Entomologie 71 (2): 221-225. <https://doi.org/10.21248/contrib.entomol.71.2.221-225>
- El-Hawagry MSA, Ebrahim AME, Nada MSE (2020) First detection of the Nearctic parasitoid species *Trichopoda pennipes* (Fabricius) (Diptera: Tachinidae) in Egypt. Egyptian Journal of Biological Pest Control 30 (1). <https://doi.org/10.1186/s41938-020-0211-z>
- Esquivel J, Musolin D, Jones W, Rabitsch W, Greene J, Toes M, Schwertener C, Grazia J, McPherson R (2019) *Nezara viridula* (L.). In: McPherson J (Ed.) Invasive stink bugs and related species (Pentatomidae). Taylor & Francis, Boca Ratón, 351-425 pp.
- Fernández-Palacios JM, Kretz H, Irl SH, Norder S, Ah-Peng C, Borges PV, Burns K, de Nascimento L, Meyer J, Montes E, Drake D (2021) Scientists' warning – The outstanding biodiversity of islands is in peril. Global Ecology and Conservation 31 <https://doi.org/10.1016/j.gecco.2021.e01847>
- Florencio M, Patiño J, Nogué S, Traveset A, Borges P, Schaefer H, Amorim I, Arnedo M, Ávila S, Cardoso P, de Nascimento L, Fernández-Palacios J, Gabriel S, Gil A, Gonçalves V, Haroun R, López-Darias M, Martínez A, Martins G, Neto A, Nogales M, Oromí P, Rando J, Raposeiro P, Rigal F, Romerias M, Silva L, Valido A, Vanderpoorten A, Vasconcelos R, Santos A (2021) Macaronesia as a fruitful arena for ecology, evolution, and conservation biology. Frontiers in Ecology and Evolution 9: 1-19.

- Freidberg A, Morgulis E, Cerretti P (2011) The invasive species, *Trichopoda pennipes* (F.) (Diptera: Tachinidae), found in Israel. The Tachinid Times 24: 14-15.
- García R, Ortega G, Pérez J (1992) Insectos de Canarias. Ediciones del Cabildo Insular de Gran Canaria, Las Palmas de Gran Canaria.
- Gobierno de Canarias (2024) Banco de Datos de Biodiversidad de Canarias. <http://www.biodiversidadcanarias.es/biota/>. Accessed on: 2024-3-01.
- Kazilas C, Demetriou J, Kalaantzis K (2020) Filling the gaps in the distribution of an alien species: The case of the feather-legged fly *Trichopoda pennipes* (Diptera: Tachinidae) in the Western Palearctic. Entomologia Hellenica 29: 8-16. <https://doi.org/10.12681/eh.21774>
- Kettani K, Ebejer M, Ackland D, Bächli G, Barracough D, Barták M, Carles-Tolrá M, Černý M, Cerretti P, Chandler P, Dakki M, Daugeron C, De Jong H, Dils J, Disney H, Droz B, Evenhuis N, Gatt P, Graciolli G, Grichanov I, Haenni J, Hauser M, Himmi O, MacGowan I, Mathieu B, Mouna M, Munari L, Nartshuk E, Negrobov O, Oosterbroek P, Pape T, Pont A, Popov G, Rognes K, Skuhravá M, Skuhravý V, Speight M, Tomasovic G, Trari B, Tschorasnig HP, Vala J, von Tschirnhaus M, Wagner R, Whitmore D, Woźnica A, Zatwarnicki T, Zwick P (2022) Catalogue of the Diptera (Insecta) of Morocco—an annotated checklist, with distributions and a bibliography. ZooKeys 1094: 1-466. <https://doi.org/10.3897/zookeys.1094.62644>
- Leclerc C, Courchamp F, Bellard C (2018) Insular threat associations within taxa worldwide. Scientific Reports 8 (1). <https://doi.org/10.1038/s41598-018-24733-0>
- López-dos-Santos N, López-Jurado L, Hernández-Peña A, Mateo J (2013) Una nueva población de lagartija de Madeira (*Teira dugesii*) en Las Palmas de Gran Canaria. Boletín la Asociación Herpetológica Española 24: 102-103.
- Lugo D, Peña G, De la Rúa P, Ruiz C (2022) Presence of exotic species of the wild bee genus *Hylaeus* (Hymenoptera: Colletidae) in the Canary Islands revealed by molecular and citizen science. Journal of Apicultural Research 61 (4): 460-468. <https://doi.org/10.1080/00218839.2022.2053034>
- Martínez J, Barone R (2006) Observation d'un Corbeau pie *Corvus albus* en haute mer près des îles Canaries et du Sahara Occidental. Alauda 74: 275-276.
- Meriem A, Doumanddj-Mitiche B, Chebli A (2021) First record of the feather-legged fly *Trichopoda pennipes* (Fabricius, 1781) (Diptera Tachinidae) a parasitoid of the southern green stink bug *Nezara viridula* (Linnaeus, 1758) (Hemiptera Pentatomidae) in Algeria. Biodiversity Journal 12 (3): 561-566. <https://doi.org/10.31396/biodiv.jour.2021.12.3.561.566>
- Obrecht E (2015) Erstfunde von *Trichopoda pennipes* (Fabricius, 1781) (Diptera, Tachinidae) in der Schweiz, und eine Würdigung einer Amateurentomologin. Contributions to Natural History: Scientific Papers from the Natural History Museum Bern 25: 71-79.
- O'Hara J, Henderson S, Wood D (2020) Preliminary checklist of the Tachinidae (Diptera) of the world. [www.nadsdiptera.org/Tach/WorldTachs/Checklist/Worldchecklist.html](http://www.nadsdiptera.org/Tach/WorldTachs/Checklist/Worldchecklist.html). Accessed on: 2024-3-01.
- Oromí P, Zurita N, Morales E, López H (2015) Diversidad de artrópodos terrestres en las Islas Canarias. Revista IDE@-SEA 4: 1-14.
- Peris S (1998) Un Trichopodini (Diptera, Tachinidae, Phasiinae) en España. Boletín de la Sociedad Española de Historia Natural 94: 163-164.

- Ricarte A, Álvarez-Fidalgo P, Álvarez-Fidalgo M (2020) Distribución actualizada y fenología de *Trichopoda pennipes* (Fabricius, 1781) (Diptera: Tachinidae) en la Península Ibérica. Graellsia 76 (2). <https://doi.org/10.3989/graeellsia.2020.v76.271>
- Ruiz C, Suárez D, Naranjo M, De la Rúa P (2020) First record of the carpenter bee *Xylocopa pubescens* (Hymenoptera, Apidae) in the Canary Islands confirmed by DNA barcoding. Journal of Hymenoptera Research 80: 169-175. <https://doi.org/10.3897/jhr.80.59649>
- Russell J, Kaiser-Bunbury C (2019) Consequences of multispecies introductions on island ecosystems. Annual Review of Ecology, Evolution, and Systematics 50 (1): 169-190. <https://doi.org/10.1146/annurev-ecolsys-110218-024942>
- Salerno G, Colazza S, Bin F (2002) *Nezara viridula* parasitism by the tachinid fly *Trichopoda pennipes* ten years after its accidental introduction into Italy from the New World. BioControl 47: 617-624. <https://doi.org/10.1023/A:1020521607677>
- Sands DP, Coombs MT (1999) Evaluation of the argentinian parasitoid, *Trichopoda giacomellii* (Diptera: Tachinidae), for biological control of *Nezara viridula* (Hemiptera: Pentatomidae) in Australia. Biological Control 15 (1): 19-24. <https://doi.org/10.1006/bcon.1999.0695>
- Stireman J, O'Hara J, Wood DM (2006) Tachinidae: Evolution, behavior, and ecology. Annual Review of Entomology 51 (1): 525-555. <https://doi.org/10.1146/annurev.ento.51.110104.151133>
- Suárez D, García J, Santos I, Ruiz C, Peña G, Pérez AJ, Lugo D, García R, Báez M (2020) Annotated check-list of the family Tachinidae (Diptera: Calyptratae: Oestroidea) in the Canary Islands. Annales de la Société entomologique de France (N.S.) 56 (2): 135-152. <https://doi.org/10.1080/00379271.2020.1753574>
- Tooker J, Hauser M, Hanks L (2006) Floral host plants of Syrphidae and Tachinidae (Diptera) of Central Illinois. Annals of the Entomological Society of America 99 (1): 96-112. [https://doi.org/10.1603/0013-8746\(2006\)099\[0096:fhpso\]2.0.co;2](https://doi.org/10.1603/0013-8746(2006)099[0096:fhpso]2.0.co;2)
- Tschorasnig HP, Zeegers T, Holstein J (2000) Further records of the introduced parasitoid *Trichopoda pennipes* (Fabricius, 1781) (Diptera, Tachinidae) from northeastern Spain and southern France. Boletín de la Sociedad Española de Historia Natural 96: 215-216.
- Tschorasnig HP, Domingo-Quero T, Alonso-Zarazaga M (2007) Tachinidae (Diptera) from the Caldera de Taburiente National Park, La Palma (Canary Islands). Graellsia 63: 43-51. <https://doi.org/10.3989/graeellsia.2007.v63.i1.79>
- Tschorasnig HP (2017) Preliminary host catalogue of Palaearctic Tachinidae (Diptera). <http://www.nadsdiptera.org/Tach/WorldTachs/CatPalHosts/Home.html>. Accessed on: 2024-12-03.
- Zeegers T (2010) Tweede aanvulling op de naamlijst van Nederlandse Sluipvliegen (Diptera: Tachinidae). Nederlandse Faunistische Mededelingen 34: 55-66.