



# Taxonomic review of Korean *Siphonaria* species (Mollusca, Gastropoda, Siphonariidae)

Yukyung Kim<sup>‡</sup>, Jina Park<sup>‡</sup>, Ui Wook Hwang<sup>§</sup>, Joong-Ki Park<sup>‡</sup>

<sup>‡</sup> Division of EcoScience, Ewha Womans University, Seoul, Republic of Korea

<sup>§</sup> Department of Biology Education, Teachers College and Institute for Phylogenomics and Evolution, Kyungpook National University, Daegu, Republic of Korea

Corresponding author: Ui Wook Hwang ([uwhwang@knu.ac.kr](mailto:uwhwang@knu.ac.kr)), Joong-Ki Park ([jkpark@ewha.ac.kr](mailto:jkpark@ewha.ac.kr))

Academic editor: Andrew Davinack

Received: 15 Oct 2024 | Accepted: 10 Feb 2025 | Published: 18 Feb 2025

Citation: Kim Y, Park J, Hwang UW, Park J-K (2025) Taxonomic review of Korean *Siphonaria* species (Mollusca, Gastropoda, Siphonariidae). Biodiversity Data Journal 13: e139388.

<https://doi.org/10.3897/BDJ.13.e139388>

## Abstract

### Background

Many molluscan species exhibit a high degree of shell morphological plasticity in their shape (including sculptures), size and colour patterns, which can vary significantly depending on environmental conditions. These shell morphological variations make it challenging to differentiate species, based on morphology alone, often resulting in various taxonomic errors, such as misidentifications, overlooking cryptic species diversity or a plethora of nominal species. The genus *Siphonaria* constitutes a significant component of the macrobenthic invertebrate fauna in intertidal habitats across temperate to tropical regions. Given the limited attention to shell variation in previous taxonomic studies on the Korean *Siphonaria* species, the extensive range of ecophenotypic shell variations documented in this group raises questions about the taxonomic validity of previously reported *Siphonaria* species in Korea.

## New information

The present study provides a comprehensive taxonomic review of Korean *Siphonaria* species using a combination of shell morphology, radula structure and phylogenetic analysis of the mtDNA *cox1* sequences. This integrative analysis confirmed the validity of *S. acmaeoides*, *S. japonica* and *S. sirius* in Korea, highlighting differences in shell and siphonal groove morphology amongst these species. Detailed descriptions of shell and radula characteristics, along with mtDNA *cox1* sequences as DNA barcodes, are also provided, which are very useful for the accurate identification of *Siphonaria* species. Unlike these three *Siphonaria* species, the taxonomic validity of the four other species (*S. coreensis*, *S. javanica*, *S. laciniosa* and *S. rucuana*) previously reported from Korean waters is questionable, given their documented geographic distribution ranges and the potential misidentification of shell variants in Korean malacofaunal studies.

## Keywords

Panpulmonata, Korean *Siphonaria*, taxonomic review, shell morphology, radula, mtDNA *cox1*

## Introduction

The genus *Siphonaria* G. B. Sowerby I, 1823, also known as the ‘false limpet’ genus, is a marine panpulmonate group consisting of 106 species worldwide (MolluscaBase eds 2024). *Siphonaria* species constitute a significant component of the macrobenthic community in intertidal/subtidal habitats across temperate to tropical regions, but some are found in cold temperature environments, such as sub-Antarctic shores (Vermeij 1971, Hodgson 1999, Dayrat et al. 2014, González-Wevar et al. 2018). They are typically inhabit intertidal rocky shores, but are occasionally found at the bottom of rocks or amongst seagrass in the subtidal zone (Collin 2000, Toyohara et al. 2001, Dayrat et al. 2014, Okutani 2000, Okutani 2017). Taxonomically, Sowerby (1823) first introduced the genus name *Siphonaria* with the type species *S. siphonaria* Sowerby, 1823 (now considered as a junior synonym of *S. javanica* (Lamarck, 1819), distinguishing it from the ‘true limpet’ genera *Patella* Linnaeus, 1758 and *Emarginula* Lamarck, 1801 by its posteriorly curved apex and a prominent siphonal groove on the lateral side. The siphonal groove allows animals to perform air-breathing through the mantle cavity, while the secondary gill supports aquatic respiration. These characteristics distinguish them from Patellostropoda (i.e. true limpets), which also have a similar conical or cap-like shell, but lack the siphonal groove, relying solely on gills for their respiration.

*Siphonaria* species exhibit remarkably high ecophenotypic shell variation, leading to many taxonomic complications such as species misidentification and the potential for unrecognised cryptic species (White and Dayrat 2012, Dayrat et al. 2014, De Coito 2021). To date, seven species have been recorded in Korea (National Institute of

Biological Resources 2023, Jenkins and Köhler 2024): *S. acmaeoides* Pilsbry, 1894, *S. coreensis* A. Adams and Reeve, 1848, *S. japonica* (Donovan, 1824), *S. javanica* (Lamarck, 1819), *S. laciniosa* (Linnaeus, 1758), *S. sirius* Pilsbry 1894, and *S. rucuana* Pilsbry, 1904. Of these, *S. coreensis* is designated as a "*taxon inquirendum*" in the World Register of Marine Species (MolluscaBase eds 2025), reflecting the view of several authorities who have suggested that it may represent a misidentification of *S. atra* Quoy & Gaimard, 1833 (Lischke 1871, Shikama 1964, Cernohorsky 1972, Jenkins and Köhler 2024). Furthermore, its occurrence in Korea, as well as northern Pacific Region, has not been substantiated since its initial report. In addition, most previous Korean *Siphonaria* records relied primarily on species checklists or taxonomic illustrations, lacking detailed descriptions of diagnostic characters and overlooking individual shell variation in their species identification. Given the limited attention to shell variation in previous studies, the extensive ecophenotypic morphological plasticity observed in this group raises questions regarding the taxonomic validity of previously reported Korean *Siphonaria* species. Insufficient and/or erroneous taxonomic information has continued to pose challenges not only in taxonomy, but also in other scientific fields that rely on accurate species identification.

In this study, we re-examined the taxonomic validity of six *Siphonaria* species previously reported in Korea by conducting a comprehensive comparison of morphological characters described in previous records against the original descriptions, based on an in-depth analysis of shell and radula morphologies, combined with mtDNA *cox1* sequence data. Based on the results of this taxonomic re-assessment, we provide detailed morphological descriptions of the shell morphology and radula microstructures of three *Siphonaria* species (*S. acmaeoides*, *S. japonica* and *S. sirius*) confirmed to occur along the Korean coast using scanning electron microscopy (SEM), as well as a phylogenetic analysis of the mtDNA *cox1* sequences.

## Materials and methods

Specimens were collected from intertidal rocky shores in Korea (Fig. 1) and preserved in 95% ethanol solution. For species identification and morphological descriptions, shells were observed using a stereoscopic microscope (Leica M205C, Wetzlar, Germany) and radula characters were examined using a scanning electron microscope (SEM). To prepare the radula for SEM examination, the radula sac was extracted from the buccal mass and residual tissue was dissolved in a mixture of 180  $\mu$ l ATL buffer and 20  $\mu$ l Proteinase K at 56°C for about 1 hour. Then the radula was rinsed with distilled water and ethanol, mounted on nickel tape attached to an SEM stub, air-dried and coated with platinum. Radula images were captured using an SEM (Zeiss Ultra Plus, Germany) and a FE-SEM (Jeol JSM-7800F, Japan). The examined specimens are deposited at the National Institute of Biological Resources (NIBR) in Incheon, Korea and the Animal Phylogenomics Laboratory at Ewha Womans University, Seoul, Korea.

For sequencing of the mtDNA *cox1* gene fragments, genomic DNA was extracted from foot tissue by using a DNeasy Blood and Tissue kit (QIAGEN, Germany) following the

manufacturer's protocol. The partial mitochondrial *cox1* gene sequence was amplified by polymerase chain reaction (PCR) using the primer pair LCO1490 and HCO2198 (Folmer et al. 1994). Amplifications were performed in a 50 µl total volume, containing 3 µl of template DNA, 35.75 µl of distilled water, 5 µl of 10x Ex Taq buffer, 1 µl of each primer, 4 µl of dNTP and 0.25 µl of TaKaRa Ex Taq DNA polymerase (TaKaRa Bio, Japan) under the following conditions: an initial denaturation at 95°C for 5 min, 40 cycles of denaturation at 94°C for 40 s, annealing at 48°C for 1 min, elongation at 72°C for 1 min and a final extension at 72°C for 10 min. The PCR-amplified target fragment was purified using a Qiaquick gel extraction kit (Qiagen Valencia, USA), sequenced using an ABI PRISM 3730xl DNA analyser (Applied Biosystems, USA) and analysed using Geneious Prime v. 2022.2.2 (Biomatters, Auckland, New Zealand).




Figure 1. [doi](#)

Map showing the sampling localities for Korean *Siphonaria* examined in this study. **1** Daejin-ri, Hyeonnae-myeon, Goseong-gun, Gangwon-do; **2** Jeonjin-ri, Ganghyeon-myeon, Yangyang-gun, Gangwon-do; **3** Namae-ri, Hyeonnam-myeon, Yangyang-gun, Gangwon-do; **4** Namyang-ri, Seo-myeon, Ulleung-gun, Gyeongsangbuk-do; **5** Dokdo-ri, Ulleung-eup, Ulleung-gun, Gyeongsangbuk-do; **6** Jukbyeon-ri, Jukbyeon-myeon, Ulin-gun, Gyeongsangbuk-do; **7** Guryongpo-eup, Nam-gu, Pohang-si, Gyeongsangbuk-do; **8** Gujora-ri, Irun-myeo, Geoje-si, Gyeongsangnam-do; **9** Mendehae-an-gil, Tongyeong-si, Gyeongsangnam-do; **10** Gahak-ri, Jisan-myeon, Jindo-gun, Jeollanam-do; **11** Seopo-ri, Deokjeok-myeon, Ongjin-gun, Incheon; **12** Jeju-si Samyang 1(il)-dong, Jeju-do; **13** Seongsan-eup, Seogwipo-si, Jeju-do; **14** Andeok-myeon, Seogwipo-si, Jeju-do. Circles with multiple colours represent localities where the corresponding species were found together.

The newly-determined *cox1* sequences of *S. acmaeoides*, *S. japonica* and *S. sirius* were deposited in GenBank (accession numbers in Table 1). The *cox1* sequences from the

three Korean *Siphonaria* species and the homologous gene sequences from NCBI database (Table 1) were used for phylogenetic analysis with two outgroup species; *Trimusculus afer* (Gmelin, 1791) and *T. reticulatus* (G. B. Sowerby I, 1835). The sequences were aligned using the default settings of MUSCLE (Edgar 2004) in the Geneious software and trimmed to a length of 630 bp. MEGA X was used to calculate genetic distances within and between species, applying uncorrected *p*-distance (Collins et al. 2012; Srivathsan and Meier 2012). To select the best-fit nucleotide substitution model, ModelFinder (Kalyaanamoorthy et al. 2017), implemented in IQ-TREE v.1.6.12. Nguyen et al. (2015) was applied with the corrected Akaike Information Criterion (AICc). Phylogenetic analysis was performed using Maximum Likelihood (ML) estimation with IQ-TREE software, with the K3Pu+F+I+G4 substitution model and 1,000 standard bootstrap pseudoreplicates (Felsenstein 1985) to assess branch support values.

Table 1.

Sampling localities and GenBank accession numbers of mtDNA *cox1* sequences of *Siphonaria* species used for molecular analysis.

Species	Locality	GenBank accession nos.	References
<i>S. acmaeoides</i> _01	Guryongpo-eup, Nam-gu, Pohang-si, Gyeongsangbuk-do	<a href="#">PQ422946</a>	This study
<i>S. acmaeoides</i> _02	Sagye-ro, Andeok-myeon, Seogwipo-si, Jeju-do	<a href="#">PQ422947</a>	This study
<i>S. acmaeoides</i> _03	Sagye-ro, Andeok-myeon, Seogwipo-si, Jeju-do	<a href="#">PQ422948</a>	This study
<i>S. acmaeoides</i> _04	Hyeongjehae-an-ro, Andeok-myeon, Seogwipo-si, Jeju-do	<a href="#">PQ422949</a>	This study
<i>S. acmaeoides</i> _05	Sagye-ro, Andeok-myeon, Seogwipo-si, Jeju-do	<a href="#">PQ422950</a>	This study
<i>S. atra</i>	China	<a href="#">OQ735434</a>	Unpublished
<i>S. japonica</i>	Japan	<a href="#">KF000756</a>	Dayrat et al. (2014)
<i>S. japonica</i>	Taiwan	<a href="#">KF000759</a>	Dayrat et al. (2014)
<i>S. japonica</i>	Taiwan	<a href="#">KF000760</a>	Dayrat et al. (2014)
<i>S. japonica</i>	Taiwan	<a href="#">KF000762</a>	Dayrat et al. (2014)
<i>S. japonica</i>	Japan	<a href="#">KF000835</a>	Dayrat et al. (2014)
<i>S. japonica</i> _01	Onpyeong-ro, Seongsan-eup, Seogwipo-si, Jeju-do	<a href="#">PQ422951</a>	This study
<i>S. japonica</i> _02	Onpyeong-ro, Seongsan-eup, Seogwipo-si, Jeju-do	<a href="#">PQ422952</a>	This study
<i>S. japonica</i> _03	Daejin-ri, Hyeonnae-myeon, Goseong-gun, Gangwon-do	<a href="#">PQ422953</a>	This study
<i>S. javanica</i>	Singapore	<a href="#">KF690503</a>	Ip et al. (2019)

Species	Locality	GenBank accession nos.	References
<i>S. laciniosa</i>	Japan	<a href="#">KF000727</a>	Dayrat et al. (2014)
<i>S. laciniosa</i>	Japan	<a href="#">KF000728</a>	Dayrat et al. (2014)
<i>S. laciniosa</i>	Japan	<a href="#">KF000758</a>	Dayrat et al. (2014)
<i>S. sirius</i>	Japan	<a href="#">KF000832</a>	Dayrat et al. (2014)
<i>S. sirius</i>	Japan	<a href="#">KF000833</a>	Dayrat et al. (2014)
<i>S. sirius_01</i>	Mendehaean-gil, Tongyeong-si, Gyeongsangnam-do	<a href="#">PQ422954</a>	This study
<i>S. sirius_02</i>	Nambu-myeon, Geoje-si, Gyeongsangnam-do	<a href="#">PQ422955</a>	This study
<i>S. sirius_03</i>	Mendehaean-gil, Tongyeong-si, Gyeongsangnam-do	<a href="#">PQ422956</a>	This study
<i>S. subatra</i>	Japan	<a href="#">KF000743</a>	Dayrat et al. (2014)
<i>S. subatra</i>	Japan	<a href="#">KF000754</a>	Dayrat et al. (2014)
<i>S. subatra</i>	Japan	<a href="#">KF000755</a>	Dayrat et al. (2014)
<i>S. subatra</i>	Japan	<a href="#">KF000757</a>	Dayrat et al. (2014)
<i>S. zelandica</i>	Australia	<a href="#">JX680967</a>	Colgan and Da Costa (2013)
<i>S. zelandica</i>	Australia	<a href="#">KF000725</a>	Dayrat et al. (2014)
<i>S. zelandica</i>	Australia	<a href="#">KF000791</a>	Dayrat et al. (2014)
<i>S. zelandica</i>	Australia	<a href="#">KF000793</a>	Dayrat et al. (2014)
<i>T. afer</i>	-	<a href="#">EF489388</a>	Klussmann-Kolb et al. (2008)
<i>T. reticulatus</i>	-	<a href="#">JN632509</a>	White et al. (2011)

## Taxon treatments

### *Siphonaria acmaeoides* Pilsbry, 1894

- GBIF <https://www.gbif.org/species/6787904>
- WoRMs <https://www.marinespecies.org/aphia.php?p=taxdetails&id=599431>

### Nomenclature

*Siphonaria acmaeoides* Pilsbry, 1894: *Pilsbry 1894*: 16; Pilsbry 1895: 6, pl. 6, figs. 19–22; Lee 1956: 79; Hirase 1941: 94, pl. 121, fig. 15; Kuroda and Habe 1952: 86 (cited from Jenkins and Köhler 2024); Azuma 1960: 62 (cited from Jenkins and Köhler (2024)); Baker 1964: 159 (cited from Jenkins and Köhler (2024)); Christiaens 1980b: 466; White and Dayrat 2012: 60 [checklist]; Jenkins and Köhler 2024: 114, 115, figs. 43A–D, M, N, 44A–C.

*Siphonaria (Patellopsis) acmaeoides*: Hubendick 1945: 70, fig. 19 (cited from Jenkins and Köhler (2024)); Hubendick 1946: 30, pl. 6, figs. 12–15; Habe and Kikuchi 1960: 64 (cited from Jenkins and Köhler (2024)); Kira 1962: 201, pl. 69, fig. 9a, b (misspelled as '*Patellops*'; cited from Jenkins and Köhler (2024)).

*Siphonaria zebra*: Kuroda and Habe 1952: 86 (non *Siphonaria zebra* Reeve, 1856) (cited from Jenkins and Köhler (2024)).

*Planesiphon acmaeoides*: Kuroda et al. 1971: 484, pl. 64, fig. 8; Choe 1992: 749, pl. 126, fig. 224.

*Siphonaria (Mouretus) acmaeoides*: Christiaens 1980a: 79, 80.

*Siphonaria (Planesiphon) acmaeoides*: Inaba 1983: 149 (cited from Jenkins and Köhler (2024)); Je 1989: 29 [checklist]; Higo et al. 1999: 402, G4976 [checklist]; Okutani 2000: 815, pl. 405, fig. 7; Cheng et al. 2005: 8, fig. 1e–g; Noseworthy et al. 2007: 90 [checklist]; Okutani 2017: 1101, pl. 403, fig. 7.

## Materials

- a. scientificName: *Siphonaria acmaeoides*; kingdom: Animalia; phylum: Mollusca; class: Gastropoda; order: Siphonariida; family: Siphonariidae; genus: *Siphonaria*; specificEpithet: *acmaeoides*; scientificNameAuthorship: Pilsbry, 1894; country: Korea; locality: Hyeongjehae-ro, Andeok-myeon, Seogwipo-si, Jeju-do; verbatimCoordinates: 33°13'37.13"N 126°18'30.92"E; eventDate: 2012-01-07; individualCount: 1; catalogNumber: NIBRIV0000307707; occurrenceID: 83FCD050-F70C-5128-B42F-DE6C1A9BA020
- b. scientificName: *Siphonaria acmaeoides*; kingdom: Animalia; phylum: Mollusca; class: Gastropoda; order: Siphonariida; family: Siphonariidae; genus: *Siphonaria*; specificEpithet: *acmaeoides*; scientificNameAuthorship: Pilsbry, 1894; country: Korea; locality: Sagye-ro, Andeok-myeon, Seogwipo-si, Jeju-do; verbatimCoordinates: 33°13'13.2"N 126°17'42.0"E; eventDate: 2022-04-21; individualCount: 3; occurrenceID: D7664B8A-483D-5DD2-9933-CECE40648E65
- c. scientificName: *Siphonaria acmaeoides*; kingdom: Animalia; phylum: Mollusca; class: Gastropoda; order: Siphonariida; family: Siphonariidae; genus: *Siphonaria*; specificEpithet: *acmaeoides*; scientificNameAuthorship: Pilsbry, 1894; country: Korea; locality: Donghaean-ro, Guryongpo-eup, Nam-gu, Pohang-si, Gyeongsangbuk-do; verbatimCoordinates: 35°57'05.7"N 129°33'04.4"E; eventDate: 2023-07-23; individualCount: 3; occurrenceID: A1D5EA8B-4851-5AD4-A0A1-2FCC3E1B0328
- d. scientificName: *Siphonaria acmaeoides*; kingdom: Animalia; phylum: Mollusca; class: Gastropoda; order: Siphonariida; family: Siphonariidae; genus: *Siphonaria*; specificEpithet: *acmaeoides*; scientificNameAuthorship: Pilsbry, 1894; country: Korea; locality: Irun-myeon, Geoje-si, Gyeongsangnam-do; verbatimCoordinates: 34°58'34.9"N 128°41'18.18"E; eventDate: 2023-04-19; individualCount: 1; occurrenceID: 84F9DDD5-5BD5-5645-8D86-F3D843A9ACD2

## Description

Measurements: Shell length [SL] 11.32–17.58 mm, Shell width [SW] 9.42–14.87 mm, Shell height [SH] 2.63–6.73 mm.

Shell (Fig. 2a, b) solid, oval, small to medium in size (in examined specimens, SL 11.32–17.58 mm), medium-high in height (about 2/7 on its length). Exterior colour generally greyish-brown or yellowish-brown with dark brown irregular maculations. Shell surface with 14–17 wide, light-cream or white-coloured primary ribs, unevenly spaced, sometimes discontinuous from apex to outer shell margin in juvenile. Interspaces between primary ribs filled with 1–7 thin secondary ribs. Apex white, glossy, spiral in counterclockwise direction (Fig. 2bC), often eroded due to shell growth, positioned in posterior one-third of shell, biased towards left. Shell margin weakly undulated. Anterior slope slightly convex, posterior slope short, nearly straight. Siphonal groove indistinct on outer shell surface. Interior colour dark brown, reddish-brown, with light cream bands near margin, reddish-brown or pale at centre.




Figure 2.

Shell morphology of three Korean *Siphonaria* species. Red arrowhead: *S. acmaeoides*; Yellow arrowhead: *S. japonica*, Blue arrowhead: *S. sirius*. Scale bars = 5 mm.

**a:** **A** *S. acmaeoides* on rocky substrate of Korea, Jeju; **B** Dorsal, ventral and lateral view (left) of *S. acmaeoides* from Jeju. [doi](#)

**b:** **C** Apex of *S. acmaeoides* juvenile from Jeju; **D** Dorsal, ventral, and lateral view (left) of *S. acmaeoides* juvenile from Jeju. [doi](#)

**c:** **E** *S. japonica* on rocky substrate of Korea, Jeju; **F** Dorsal, ventral and lateral view (left) of *S. japonica* from Goseong. [doi](#)

**d:** **G** *S. sirius* on rocky substrate of Korea, Jeju; **H** Dorsal, ventral and lateral view of *S. sirius* from Geoje. [doi](#)

Radula (Fig. 3a, b) dentition formulae 27:1:27 (in specimens measuring SL 16.23 mm, SW 11.88 mm, SH 4.80 mm). Each transverse row with narrow central rachidian tooth flanked by symmetrical half rows with lateral teeth, decreasing in size outwards. Rachidian tooth (Fig. 3a) short, approximately one-third length of lateral tooth, with sharply-pointed cusp. Innermost lateral teeth (first 7 lateral teeth), each tooth consisting of mesocone without ectocone; mesocone with bicuspid (Fig. 3a). Middle lateral teeth (following 5 lateral teeth), each tooth consisting of mesocone and ectocone; mesocone with bicuspid to blunt towards the outer margins; ectocone sharply pointed. Outermost lateral teeth (remaining 12 lateral teeth) consisting of endocone, ectocone and mesocone; mesocone subquadrate-shaped; endocone and ectocone short, sharply pointed (Fig. 3b).






Figure 3.

Radula morphology of three Korean *Siphonaria* species. Abbreviations: m, mesocone; ec, ectocone. Scale bars = 20  $\mu$ m.

a: *S. acmaeoides* - Details of rachidian teeth (arrowhead) and innermost lateral teeth. [doi](#)

b: *S. acmaeoides* - Details of outermost lateral teeth. [doi](#)

c: *S. japonica* - Details of rachidian teeth (arrowhead) and innermost lateral teeth. [doi](#)

d: *S. japonica* - Details of outermost lateral teeth. [doi](#)

e: *S. sirius* - Details of rachidian teeth (arrowhead) and innermost lateral teeth. [doi](#)

f: *S. sirius* - Details of outermost lateral teeth. [doi](#)

## Distribution

Korea, Japan and Taiwan.

**Type locality:** Japan; Boshu island (Boso Peninsula).

**Habitat:** On rocky substrate in high intertidal zones, typically in shallow pools (Fig. 2a A).

## Taxon discussion

The Korean *S. acmaeoides* corresponds well with Pilsbry (1894)'s original description. The external shell morphology is characterised by wide, low primary ribs with fine secondary ribs in the interspaces, a glossy spiral apex coiling counterclockwise and a weakly undulated shell margin. Christiaens (1980a) described a new subspecies, *S. acmaeoides paulae* Christiaens, 1980, from Hong Kong, distinguishing it by its finer, thinner, more elliptical, lighter-coloured shell. This subspecies was also noted to lack a marked central area and exhibit a more bulging siphon compared to *S. acmaeoides* reported from Japan. However, Jenkins and Köhler (2024) later synonymised *S. a. paulae* with the *S. acmaeoides*, citing the morphological similarity of the shell. Based on our examination of the Korean specimens (Fig. 2b), we did not observe the light-coloured shell characteristics described in *S. a. paulae* by Christiaens (1980a). Further morphological and molecular analyses for *S. a. paulae* sampled from Hong Kong are necessary to confirm the taxonomic validity of this subspecies. In addition, *S. acmaeoides* has repeatedly been noted for its similarity in shell and radula morphology to *S. zelandica* Quoy and Gaimard, 1833 which is distributed in Australia (Hubendick 1946, Dayrat et al. 2014, Jenkins and Köhler 2024). Both species share the following morphological characteristics: a homostrophic apex, broad primary ribs with fine secondary ribs and an innermost lateral tooth in the radula with a mesocone that lacks ectocone and endocone (Quoy and Gaimard 1833, Pilsbry 1894, Jenkins 1983, Jenkins and Köhler 2024). Hubendick (1946) reported the radula formula of *S. acmaeoides* as 26:1:26, noting the absence of an ectocone on the lateral teeth. In this study, we observed a similar pattern of radula characters, with a radula formula of 27:1:27, also lacking an ectocone on the lateral teeth. This can be generally included within the documented range of intraspecific variation for the genus *Siphonaria* (e.g. in *S. lessonii* Blainville, 1827, the radula formula varies from 13:1:13 to 76:1:76; Hubendick (1946), Güller et al. (2016)). In shell morphology, *S. acmaeoides* is distinguished by having its less prominent, almost flat radial ribs and a more indistinct siphonal groove, whereas *S. zelandica* is characterised by relatively finer and dual radial ribs within the siphonal groove. In our phylogenetic tree, the two species form a sister clade with 95% bootstrap support (Fig. 4). The uncorrected *p*-distances between *S. acmaeoides* and *S. zelandica* range from 5.71 to 6.51%, while *S. sirius*, the next closest related species shows a significantly higher genetic divergence (27.94%) (Table 2). Thus, a taxonomic re-assessment of the two species is necessary to confirm their taxonomic validity.

Table 2.

Uncorrected *p*-distance (%) for the mtDNA *cox1* sequences amongst *Siphonaria* species. The species which include newly-determined species in this study are denoted by asterisks (\*).

	<i>S. japonica</i> *	<i>S. zelandica</i>	<i>S. acmaeoides</i> *	<i>S. javanica</i>	<i>S. lacinosia</i>	<i>S. sirius</i> *	<i>S. atra</i>	<i>S. subatra</i>
<i>S. japonica</i> *	0.16–1.75							
<i>S. zelandica</i>	28.89–29.84	0.16–1.90						
<i>S. acmaeoides</i> *	29.37–30.95	5.71–6.51	0.16–3.33					
<i>S. javanica</i>	29.52–30.32	29.05–30.00	29.37–30.16	-				
<i>S. lacinosia</i>	29.05–30.14	28.57–29.52	28.57–29.68	15.56–16.03	0.16–0.79			
<i>S. sirius</i> *	30.79–31.90	28.25–29.68	27.94–29.21	30.63–30.79	29.52–30.48	0.16–0.79		
<i>S. atra</i>	31.27–31.90	27.78–28.25	28.25–28.82	27.46	27.14–27.30	22.38–22.86	-	
<i>S. subatra</i>	31.90–33.81	29.68–30.63	29.37–30.63	29.68–30.79	29.21–30.48	25.24–25.87	15.87–16.51	2.06–3.33

## *Siphonaria japonica* (Donovan, 1824)

- GBIF <https://www.gbif.org/species/5859486>
- WoRMS <https://www.marinespecies.org/aphia.php?p=taxdetails&id=740941>

### Nomenclature

*Patella japonica* Donovan 1824: pl. 79; Kuroda et al. 1971: 483, pl. 64, fig. 7; White and Dayrat 2012: 64 [checklist].

? *Siphonaria radiata*: Adams and Reeve 1850: 69, pl. 13, fig. 2.

? *Siphonaria radians*: Adams and Adams 1855: 271; Hubendick 1946: 43.

*Siphonaria cochleariformis*: Reeve 1856: pl. 6, fig. 28; Hubendick 1946: 43, pl. 2, figs. 33–35; Kuroda et al. 1971: 483, pl. 64, fig. 7.

*Siphonaria japonica*: Hanley 1858: 152 [checklist]; Pilsbry 1920: 141; Abe 1940: 59 (cited from Jenkins and Köhler (2024)); Hirase 1941: 94, pl. 121, fig. 12; Kuroda 1941: 137; Hubendick 1945: 27, fig. 34, 36, 39, 41 (cited from Jenkins and Köhler (2024)); Kuroda and Habe 1952: 86 (cited from Jenkins and Köhler (2024)); Lee 1956: 79; Kuroda 1960: 43 (cited from Jenkins and Köhler (2024)); Habe and Kikuchi 1960: 64 (cited from Jenkins and Köhler 2024); Shikama 1964: 6 (cited from Jenkins and

Köhler (2024)); Berry 1977: 197, fig. 19 (cited from Jenkins and Köhler (2024)); Lee and Chao 2003: 35, pl. 6, fig. 173; Qi 2004: 198, pl. 108, fig. E; Dayrat et al. 2014: 266, fig. 3B; Jenkins and Köhler 2024: 31–34, figs. 13A–L, Q–S, 15B–E.




Figure 4. [doi](#)

Phylogenetic relationships amongst some selected NWP *Siphonaria* species inferred from the Maximum Likelihood method using the mtDNA *cox1* sequences. The bootstrap supporting values ( $\geq 50\%$ ) are indicated on the branches. Asterisks (\*) represents the mtDNA *cox 1* sequences determined in this study.

*Siphonaria alterniplicata*: Grabau and King 1928: 237, pl. 11, fig. 117; Hubendick 1946: 62; White and Dayrat 2012: 60 [checklist]; Coan et al. 2015: 221, fig. 38A, B.

*Siphonaria (Sacculosiphonaria) japonica*: Hubendick 1946: 42, pl. 2, figs. 29–32; Kira 1962: 201, pl. 69, fig. 8a, b (cited from Jenkins and Köhler (2024)); Habe 1971: 15, pl. 4, fig. 16 (cited from Jenkins and Köhler (2024)); Yoo 1976: 89, pl. 19, figs. 1–4; Christiaens 1980a: 79; Christiaens 1980b: 466, 467; Je 1989: 89 [checklist]; Fukuda

et al. 1992: 76, pl. 23, fig. 360a, b (cited from Jenkins and Köhler (2024)); Kwon et al. 1993: 334, fig. 61-1; Okutani 2000: 815, pl. 405, fig. 6; Kwon 2001: 191, fig. 723; Hamamura 2004: 115; Min et al. 2004: 335, fig. 1056; Noseworthy et al. 2007: 90 [checklist]; Okutani 2017: 1101, pl. 403, fig. 6.

*Sacculosiphonaria japonica*: Kuroda et al. 1971: 483, pl. 64, fig. 7; Trew 1983: 9 (cited from Jenkins and Köhler (2024)); Choe 1992: 747, 748, pl. 125, fig. 222.

*Siphonaria (Mastosiphon) sirius*: Yoo 1976: 89, pl. 19, fig. 5 (non *Siphonaria sirius* Pilsbry 1894).

*Siphonaria acmaeoides*: Kwon et al. 1993: 335, fig. 61-3 (non-*Siphonaria acmaeoides* Pilsbry 1894).

*Siphonaria (Planesiphon) acmaeoides*: Kwon 2001: 191, fig. 724; Min et al. 2004: 337, fig. 1057 (non-*Siphonaria acmaeoides* in Pilsbry (1894)).

*Siphonaria japonica* tall form: Yokogawa et al. 2010: fig. 1A.

## Materials

- a. scientificName: *Siphonaria japonica*; kingdom: Animalia; phylum: Mollusca; class: Gastropoda; order: Siphonariida; family: Siphonariidae; genus: *Siphonaria*; specificEpithet: *japonica*; scientificNameAuthorship: (Donovan, 1824); country: Korea; locality: Namyang-ri, Seo-myeon, Ulleung-gun, Gyeongsangbuk-do; verbatimCoordinates: 37°27'34.5"N, 130°51'27.4"E; eventDate: 2011-11-13; individualCount: 20; catalogNumber: NIBRIV0000894428–NIBRIV0000894447; occurrenceID: 8CB2FFA8-7177-5D58-9987-D944894409D8
- b. scientificName: *Siphonaria japonica*; kingdom: Animalia; phylum: Mollusca; class: Gastropoda; order: Siphonariida; family: Siphonariidae; genus: *Siphonaria*; specificEpithet: *japonica*; scientificNameAuthorship: (Donovan, 1824); country: Korea; locality: Jeonjin-ri, Ganghyeon-myeon, Yangyang-gun, Gangwon-do; verbatimCoordinates: 38°07'31.5"N, 128°37'52.3"E; eventDate: 2014-04-09; individualCount: 10; catalogNumber: NIBRIV0000894458–NIBRIV0000894467; occurrenceID: E2FA26B9-DF79-5C3B-BAA0-2A7982854C38
- c. scientificName: *Siphonaria japonica*; kingdom: Animalia; phylum: Mollusca; class: Gastropoda; order: Siphonariida; family: Siphonariidae; genus: *Siphonaria*; specificEpithet: *japonica*; scientificNameAuthorship: (Donovan, 1824); country: Korea; locality: Jukbyeon-ri, Jukbyeon-myeon, Uljin-gun, Gyeongsangbuk-do; verbatimCoordinates: 37°3'32.50"N, 129°25'43.92"E; eventDate: 2014-04-09; individualCount: 20; catalogNumber: NIBRIV0000894468–NIBRIV0000894487; occurrenceID: 59D7504B-74E8-514D-BCCA-3C6F3557C300
- d. scientificName: *Siphonaria japonica*; kingdom: Animalia; phylum: Mollusca; class: Gastropoda; order: Siphonariida; family: Siphonariidae; genus: *Siphonaria*; specificEpithet: *japonica*; scientificNameAuthorship: (Donovan, 1824); country: Korea; locality: Namae-ri, Hyeonnam-myeon, Yangyang-gun, Gangwon-do; verbatimCoordinates: 37°56'33.7"N, 128°47'17.8"E; eventDate: 2014-08-24; individualCount: 3; catalogNumber: NIBRIV0000894448NIBRIV0000894450; occurrenceID: 8451E1F2-DB21-501A-A3C9-52209A958584

- e. scientificName: *Siphonaria japonica*; kingdom: Animalia; phylum: Mollusca; class: Gastropoda; order: Siphonariida; family: Siphonariidae; genus: *Siphonaria*; specificEpithet: *japonica*; scientificNameAuthorship: (Donovan, 1824); country: Korea; locality: Samyang 1(ii)-dong Jeju-si Jeju-do; verbatimCoordinates: 33°31'34.6"N, 126°35'09.0"E; eventDate: 2015-12-21; individualCount: 15; catalogNumber: NIBRIV0000894548-NIBRIV0000894562; occurrenceID: 7E7E0FC6-5FF9-5A57-A3BD-6F006D828021
- f. scientificName: *Siphonaria japonica*; kingdom: Animalia; phylum: Mollusca; class: Gastropoda; order: Siphonariida; family: Siphonariidae; genus: *Siphonaria*; specificEpithet: *japonica*; scientificNameAuthorship: (Donovan, 1824); country: Korea; locality: Namae-ri, Hyeonnam-myeon, Yangyang-gun, Gangwon-do; verbatimCoordinates: 37°56'33.7"N, 128°47'17.8"E; eventDate: 2016-03-02; individualCount: 7; catalogNumber: NIBRIV0000894451-NIBRIV0000894457; occurrenceID: CC219B8C-1E0B-5581-B327-D492B0178063
- g. scientificName: *Siphonaria japonica*; kingdom: Animalia; phylum: Mollusca; class: Gastropoda; order: Siphonariida; family: Siphonariidae; genus: *Siphonaria*; specificEpithet: *japonica*; scientificNameAuthorship: (Donovan, 1824); country: Korea; locality: Dokdo-ri, Ulleung-eup, Ulleung-gun, Gyeongsangbuk-do; verbatimCoordinates: 37°14'20.12"N, 131°52'05.74"E; eventDate: 2016-06-22; individualCount: 17; catalogNumber: NIBRIV0000894408-NIBRIV0000894424; occurrenceID: 2CD256A6-3489-5322-9FA4-8F75AF0BCF39
- h. scientificName: *Siphonaria japonica*; kingdom: Animalia; phylum: Mollusca; class: Gastropoda; order: Siphonariida; family: Siphonariidae; genus: *Siphonaria*; specificEpithet: *japonica*; scientificNameAuthorship: (Donovan, 1824); country: Korea; locality: Gahak-ri, Jisan-myeon, Jindo-gun, Jeollanam-do; verbatimCoordinates: 34°25'48.1"N, 126°05'55.6"E; eventDate: 2016-07-08; individualCount: 20; catalogNumber: NIBRIV0000894508-NIBRIV0000894527; occurrenceID: 0D774AF9-C5DA-5C87-B0AE-26873ED2A1F7
- i. scientificName: *Siphonaria japonica*; kingdom: Animalia; phylum: Mollusca; class: Gastropoda; order: Siphonariida; family: Siphonariidae; genus: *Siphonaria*; specificEpithet: *japonica*; scientificNameAuthorship: (Donovan, 1824); country: Korea; locality: Ojo-ri, Seongsan-eup, Seogwipo-si, Jeju-do; verbatimCoordinates: 33°28'14.8"N, 126°55'21.9"E; eventDate: 2017-06-27; individualCount: 2; catalogNumber: NIBRIV0000894563-NIBRIV0000894564; occurrenceID: EE49E3D2-2BA0-5D88-8772-0303B4B71649
- j. scientificName: *Siphonaria japonica*; kingdom: Animalia; phylum: Mollusca; class: Gastropoda; order: Siphonariida; family: Siphonariidae; genus: *Siphonaria*; specificEpithet: *japonica*; scientificNameAuthorship: (Donovan, 1824); country: Korea; locality: Hwasun-ri, Andeok-myeon, Seogwipo-si, Jeju-do; verbatimCoordinates: 33°14'20.7"N, 126°20'01.6"E; eventDate: 2017-06-28; individualCount: 3; catalogNumber: NIBRIV0000894565-NIBRIV0000894567; occurrenceID: 0CBFE4F7-A55D-52DF-AAC8-16500149F3AD
- k. scientificName: *Siphonaria japonica*; kingdom: Animalia; phylum: Mollusca; class: Gastropoda; order: Siphonariida; family: Siphonariidae; genus: *Siphonaria*; specificEpithet: *japonica*; scientificNameAuthorship: (Donovan, 1824); country: Korea; locality: Seopo-ri, Deokjeok-myeon, Ongjin-gun, Incheon; verbatimCoordinates: 37°13'03.16"N, 126°06'55.67"E; eventDate: 2018-04-17; individualCount: 20; catalogNumber: NIBRIV0000894488-NIBRIV0000894507; occurrenceID: BEF4F2B8-098D-5566-8920-C4545FAEE777

- l. scientificName: *Siphonaria japonica*; kingdom: Animalia; phylum: Mollusca; class: Gastropoda; order: Siphonariida; family: Siphonariidae; genus: *Siphonaria*; specificEpithet: *japonica*; scientificNameAuthorship: (Donovan, 1824); country: Korea; locality: Gujora-ri, Irun-myeon, Geoje-si, Gyeongsangnam-do; verbatimCoordinates: 34°48'31.93"N, 128°41'25.97"E; eventDate: 2018-06-12; individualCount: 20; catalogNumber: NIBRIV0000894528-NIBRIV0000894547; occurrenceID: B872D531-A994-59F3-A5E9-E1FDD9EAB14F
- m. scientificName: *Siphonaria japonica*; kingdom: Animalia; phylum: Mollusca; class: Gastropoda; order: Siphonariida; family: Siphonariidae; genus: *Siphonaria*; specificEpithet: *japonica*; scientificNameAuthorship: (Donovan, 1824); country: Korea; locality: Dokdo-ri, Ulleung-eup, Ulleung-gun, Gyeongsangbuk-do; verbatimCoordinates: 37°14'20.12"N, 131°52'05.74"E; eventDate: 2021-06-02; individualCount: 3; catalogNumber: NIBRIV0000894425-NIBRIV0000894427; occurrenceID: 3B25F340-A40E-5225-BBC0-AC7D16CA5F33
- n. scientificName: *Siphonaria japonica*; kingdom: Animalia; phylum: Mollusca; class: Gastropoda; order: Siphonariida; family: Siphonariidae; genus: *Siphonaria*; specificEpithet: *japonica*; scientificNameAuthorship: (Donovan, 1824); country: Korea; locality: Guryongpo-eup, Nam-gu, Pohang-si, Gyeongsangbuk-do; verbatimCoordinates: 35°57'11.5"N, 129°32'50.4"E; eventDate: 2023-07-12; individualCount: 1; occurrenceID: FB198DA4-1645-5901-BDFA-506AE0CDD712
- o. scientificName: *Siphonaria japonica*; kingdom: Animalia; phylum: Mollusca; class: Gastropoda; order: Siphonariida; family: Siphonariidae; genus: *Siphonaria*; specificEpithet: *japonica*; scientificNameAuthorship: (Donovan, 1824); country: Korea; locality: Daejin-ri, Hyeonae-myeon, Goseong-gun, Gangwon-do; verbatimCoordinates: 38°29'57.1"N, 128°25'38.7"E; eventDate: 2024-03-20; individualCount: 1; occurrenceID: 1B68A9AE-75A2-505F-8F9E-E9AE8E3C17F9
- p. scientificName: *Siphonaria japonica*; kingdom: Animalia; phylum: Mollusca; class: Gastropoda; order: Siphonariida; family: Siphonariidae; genus: *Siphonaria*; specificEpithet: *japonica*; scientificNameAuthorship: (Donovan, 1824); country: Korea; locality: Onpyeong-ro, Seongsan-eup, Seogwipo-si, Jeju-do; verbatimCoordinates: 33°24'03.7"N, 126°54'17.7"E; eventDate: 2024-07-28; individualCount: 3; occurrenceID: C26236C2-BD2D-5ED5-94A0-A09287579017
- q. scientificName: *Siphonaria japonica*; kingdom: Animalia; phylum: Mollusca; class: Gastropoda; order: Siphonariida; family: Siphonariidae; genus: *Siphonaria*; specificEpithet: *japonica*; scientificNameAuthorship: (Donovan, 1824); country: Korea; locality: Sagye-ro, Andeok-myeon, Seogwipo-si, Jeju-do; verbatimCoordinates: 33°13'13.2"N 126°17'42.0"E; eventDate: 2022-04-21; individualCount: 2; occurrenceID: DE709C43-9BAD-5511-BB20-64DF8FE5144F

## Description

Measurements: SL 10.5–15.45 mm, SW 7.5–12.24 mm, SH 2.81–4.7 mm.

Shell (Fig. 2c) thin, oval, small size (in examined specimens, SL 10.5–15.45 mm), medium-high in height (2/7 of its length). Exterior colour greyish-brown. Shell surface with 17–23 thin, yellowish white-coloured primary radial ribs. Interspaces between radial ribs sometimes with 1–2 secondary ribs. Apex eroded, located centrally in posterior one-third of shell. Shell margin weakly undulated. Anterior slope slightly convex, posterior slope shorter, nearly straight. Siphonal groove weakly prominent with two radial ribs on outer shell surface. Interior colour reddish-brown.

Radula (Fig. 3c, d) dentition formulae 29:1:29 (in specimens measuring SL 10.54 mm, SW 7.47 mm, SH 2.81 mm). Each transverse row with narrow central rachidian tooth flanked by symmetrical half rows with lateral teeth, decreasing in size outwards. Rachidian tooth (Fig. 3c) short, -approximately one-third length of lateral tooth, with sharply pointed cusp. Innermost lateral teeth (first 17 lateral teeth), each tooth consisting of mesocone and ectocone; mesocone with bicuspid or unicuspid (Fig. 3c); ectocone shorter, pointed. Outermost lateral teeth (remaining 12 lateral teeth) consisting of endocone, ectocone and mesocone; mesocone paddle-shaped, not angled; endocone and ectocone short, sharply pointed (Fig. 3d).

### Distribution

Korea, China, Japan and Taiwan.

**Type locality:** Japan.

**Habitat:** On rocky substrate in low to high intertidal zones (Fig. 2cE).

### Taxon discussion

Jenkins and Köhler (2024) designated a neotype for *S. japonica* due to the absence of the original type specimens and provided a detailed description of the shell morphology, reproductive system and morphological features of spermatophore. The shell morphology of the Korean *S. japonica* specimens match es well both by Donovan (1824)'s original description and the observations by Jenkins and Köhler (2024). Nevertheless, moderate variations in the number of ribs were observed in this study, with the number of primary ribs ranging from 17 to 23 and secondary ribs ranging from 1 to 2. These ranges differ from those reported by Jenkins and Köhler (2024), which included up to 20 primary ribs and 2 to 3 secondary ribs. In addition, Jenkins and Köhler (2024) considered the shell images of Figs. 1 and 4 of *S. japonica* described by Yoo (1976) to likely represent misidentification, without providing detailed information on their interpretation. Upon re-examination of the Figs. 1 and 4, we observed that the specimens show characteristics consistent with *S. japonica*, including 20–25 primary ribs with secondary rib interstices (though the exact number is ambiguous) and two primary ribs on the siphonal groove. Based on our integrated analyses of the morphology (including the number of radial ribs on the shell surface) and molecular sequence data, we have concluded that all the shell images in Yoo (1976), including Figs. 1 and 4, represent within the normal range of shell variations typically observed in *S. japonica*. In radula morphology, Hubendick (1946) provided a brief sketch of radula characters, based on only two specimens in his description of *S. japonica*. He provided a radula formula ranging from 32:1:32 to 40:1:40 and noted the presence of an ectocone on the innermost lateral teeth. Our examination of the Korean specimens revealed a radula formula of 29:1:29, which differs from those reported in Hubendick (1946), while the presence of an ectocone on the innermost lateral teeth was consistent with his findings. These observations suggest that the radula formula can vary within the same species.



*S. japonica* is commonly found in intertidal zones along the Korean coast, often occurring alongside *S. acmaeoides* and *S. sirius* (Fig. 2aA, Fig. 2dG). Their wide range of distribution and variations in shell morphology can lead to confusion in identification, resulting in past misidentifications in Korean illustrations as *S. sirius* (Yoo 1976) and *S. acmaeoides* (Kwon et al. 1993, Kwon 2001, Min et al. 2004). However, this species can be distinguished from other sympatric *Siphonaria* species by its thinner, regularly spaced primary ribs, siphonal groove with two radial ribs, undulated margin and dark brown-coloured interior (Fig. 2cF). Yokogawa et al. (2010) described two types from *S. japonica*, based on the combination of morphological, molecular, and ecological data, as the “tall form” and the “short form”. The tall form is characterised by weak radial ribs over the shell margin and alternating sized (thick and thin) radial ribs, while the short form has projected radial ribs over the shell margin and generally thick radial ribs. The Korean *S. japonica* morphologically matches the “tall form”. A comprehensive analysis of shell morphological data, coupled with molecular sequence information from broader sampling of this species, is required to confirm whether the “short form” of *S. japonica* is also distributed along the Korean sea coast, as previously reported in Japan (Yokogawa et al. 2010). Our phylogenetic tree indicates that *S. japonica* is positioned basal amongst selected Siphonariidae species in the north-western Pacific, with 99% bootstrap support value (Fig. 4). This species exhibits relatively high sequence divergence from other *Siphonaria* species, ranging from 28.89% (*S. zelandica*) to 33.81% (*S. subatra*) (Table 2).

### ***Siphonaria javanica* (Lamarck, 1819)**

- GBIF <https://www.gbif.org/species/5859486>
- WoRMS <https://www.marinespecies.org/aphia.php?p=taxdetails&id=740945>

#### **Nomenclature**

*Siphonaria* (*Siphonaria*) *javanica*: Lee and Min 2002: 146 [checklist]; Min et al. 2004: 334, fig. 1054.

*Siphonaria javanica*: Noseworthy et al. 2007: 90 [checklist].

#### **Distribution**

Japan (Okinawa island), Malaysia, Palau, Indonesia and northeast Australia.

**Type locality:** Indonesia; Java Island.

#### **Taxon discussion**

This species was initially included by Lee and Min (2002) in their checklist. Later, Min et al. (2004) described the shell characters of this species as having a large shell size (SL 20 mm, SW 18 mm, SH 9 mm) and white primary ribs with 2–3 thin secondary ribs

in the interspaces. However, the shell characters in their description do not align with the original description of *S. javanica* provided by Lamarck (1819) or the subsequent studies by Morrison (1972) and Jenkins and Köhler (2024). The original description of *S. javanica* is brief, noting that the species possesses ‘white radiating ribs and crenated margin’ (translation from Latin). Morrison (1972) further noted that *S. javanica* was readily distinguished from other congeneric species by its high conical shell with buttress-like strong white radial ribs. Jenkins and Köhler (2024) provided a further detailed description of *S. javanica* by designating a lectotype and refined its taxonomy by clarifying its diagnostic characteristics. Their study also emphasised that *S. javanica* possesses well-developed primary ribs that extend beyond the shell margin, resulting in a coarsely undulated shell margin. This contrasts with the shell image provided by Min et al. (2004), which depicts the species with a smooth shell margin. Given the restricted distribution range of *S. javanica* to Indonesia and Timor-Leste, as clarified by the recent revision in Jenkins and Köhler (2024), Morrison (1972) and the discordance between the shell description of Korean “*S. javanica*” samples and both the original and subsequent descriptions, it is likely that the previously reported Korean specimens of “*S. javanica*” are likely misidentifications of *S. japonica*. Further studies with broader taxon sampling are necessary to resolve this taxonomic issue.

### ***Siphonaria laciniosa* (Linnaeus, 1758)**

- GBIF <https://www.gbif.org/species/5189835>
- WoRMS <https://www.marinespecies.org/aphia.php?p=taxdetails&id=215290>

#### **Nomenclature**

*Siphonaria laciniosa*: Kwon et al. 1993: 335, fig. 61-4.

*Siphonaria* (*Siphonaria*) *laciniosa*: Kwon 2001: 191, fig. 721.

*Siphonaria* (*Mestosiphon*) *laciniosa*: Lee and Min 2002 [checklist]: 146, fig. 268; Min et al. 2004: 336, fig. 1059; Noseworthy et al. 2007: 90 [checklist].

#### **Distribution**

Persian Gulf, Red Sea, Samoan Islands, Tonga, Fiji (Morrison 1972).

**Type locality:** India.

#### **Taxon discussion**

Kwon et al. (1993) first reported the occurrence of this species in Korea, noting its high shell height (SL 20 mm, SW 18 mm, SH 9.4 mm), with approximately 50 radial ribs and a pointed apex. Subsequent literature described this species as having less pronounced radial ribs, with white spots along the inner shell margin (Kwon 2001, Min et al. 2004), which are shell features also commonly observed in some

individuals of *S. japonica*. Jenkins and Köhler (2024) recently designated *S. laciniosa* as a “*nomen dubium*”, highlighting the “utter confusion” in its original description and subsequent literature. Jenkins and Köhler (2024) noted that many type specimens attributed to *S. laciniosa* are either lost or represent mixed lots. Given the substantial taxonomic confusion regarding the validity of *S. laciniosa* and the morphological overlap between Korean specimens previously identified as *S. laciniosa* (Kwon et al. 1993, Kwon 2001, Min et al. 2004) and *S. japonica* (as observed in this study), it is likely that the earlier records of *S. laciniosa* in Korea represent misidentifications of *S. japonica*. As a result, we excluded *S. laciniosa* from the list of Korean *Siphonaria* species.

## ***Siphonaria rucuana* Pilsbry, 1904**

- GBIF <https://www.gbif.org/species/6126792>
- WoRMS <https://www.marinespecies.org/aphia.php?p=taxdetails&id=599440>

### **Nomenclature**

*Siphonaria rucuana*: Kwon et al. 1993: 335, fig. 61-5.

*Siphonaria (Siphonaria) rucuana*: Kwon 2001: 191, fig. 722; Lee and Min 2002: 146 [checklist]; Min et al. 2004: 335, fig. 1055; Noseworthy et al. 2007: 90 [checklist].

### **Distribution**

Japan (Okinawa Island).

**Type locality**: Japan; Ryukyu island (Okinawa Island).

### **Taxon discussion**

Kwon et al. (1993) first reported this species in Korea, based on shell morphology, describing it as having a flatly convex shell with a brown apex curving posteriorly and brown intervals with secondary ribs. However, in contrast to this description, Pilsbry (1904) described *S. rucuana* as having a steeply conic shell with a glossy brown-coloured apex curving posteriorly and dusky-coloured intervals without secondary ribs. These characteristics are consistent with the recent revision of Jenkins and Köhler (2024). Yokogawa et al. (2010) divided *S. japonica* into two shell forms—“tall form” and “short form”—based on an integrative analysis of their morphology, habitat ecology and allelic frequencies. Compared to these two shell forms of *S. japonica*, the specimens previously described as *S. rucuana* in Korean waters (Kwon et al. 1993, Kwon 2001, Min et al. 2004) do not accord with the Pilsbry (1904)’s original description and morphological characters described in the recent revisional study (Jenkins and Köhler 2024). A re-evaluation of prior records of *S. rucuana* in Korea suggests that the specimens described in Kwon et al. (1993) correspond to the “tall form” of *S. japonica*, representing the original *S. japonica*. On the other hand, the

other two reports (Kwon 2001, Min et al. 2004) are assumed to correspond to the “short form” of *S. japonica* discovered first by Yokogawa et al. (2010) in Japan. In addition to these morphological aspects, *S. rucuana* is known to have a limited geographic distribution, being reported exclusively from Okinawa Island, a subtropical region of Japan (Pilsbry 1904, Oyama et al. 1954, Okutani 2000, Okutani 2017). Given this restricted geographic distribution range and the morphological discrepancies observed in the Korean nominal records of “*S. rucuana*”, it is likely that these previous records in the Korean malacofauna result from misidentifications.

## ***Siphonaria sirius* Pilsbry, 1894**

- GBIF <https://www.gbif.org/species/6126791>
- WoRMS <https://www.marinespecies.org/aphia.php?p=taxdetails&id=740940>

### **Nomenclature**

*Siphonaria sirius* Pilsbry 1894: 9, 10; Pilsbry 1895: 5, 6, pl. 6, figs. 23–28; Hirase 1941: 94, pl. 121, fig. 16; Hubendick 1945: 29 (cited from Jenkins and Köhler (2024)); Kuroda and Habe 1952: 86 (cited from Jenkins and Köhler (2024)); Azuma 1960: 62 (cited from Jenkins and Köhler (2024)); Kuroda 1960: 43 (cited from Jenkins and Köhler (2024)); Baker 1964: 159 (cited from Jenkins and Köhler (2024)); Shikama 1964: 6 (cited from Jenkins and Köhler (2024)); Habe and Igarashi 1967: 28 (cited from Jenkins and Köhler (2024)); Inaba 1983: 145 (cited from Jenkins and Köhler (2024)); Trew 1983: 7 (cited from Jenkins and Köhler (2024)); Je 1989: 29 [checklist]; Morton and Morton 1983: 298, pl. 1K (cited from Jenkins and Köhler (2024)); Hylleberg and Kilburn 2003, 2003: 133; White and Dayrat 2012: 68 [checklist]; Dayrat et al. 2014: 269, fig. 5H; Jenkins and Köhler 2024: 115–118, figs. 43E, F, O, P, 44D, E.

*Siphonaria (Siphonaria) sirius*: Hubendick 1946: 50, 51, pl. 3, figs. 24–27.

*Siphonaria (Mestosiphon) sirius*: Habe and Kikuchi 1960: 60 (cited from Jenkins and Köhler (2024)); Kira 1962: 200, 201, text-fig, pl. 69, fig. 12a, b (cited from Jenkins and Köhler (2024)); Habe 1971: 15, pl. 4, fig. 12 (cited from Jenkins and Köhler (2024)).

*Anthosiphonaria sirius*: Kuroda et al. 1971: 483, 484, fig. 9; Inaba 1983: 145 (cited from ); Je 1989: 29 [checklist]; Choe 1992: 748, 749, pl. 126, fig. 223; Fukuda et al. 1992: 76, pl. 23, fig. 361a, b (cited from Jenkins and Köhler (2024)).

*Siphonaria laciniosa* forma *sirius*: Christiaens 1980a: 79.

*Siphonaria laciniosa*: Springsteen and Leobrera 1986: 285, pl. 81, fig. 19 (non *Siphonaria laciniosa* (Linnaeus, 1758); cited from Jenkins and Köhler (2024)).

*Siphonaria subatra*: Je 1989: 29 [checklist].

*Siphonacmea oblongata*: Je 1989: 29 [checklist].

*Siphonaria* (*Anthosiphonaria*) *sirius*: Kwon et al. 1993: 334, fig. 61-2; Higo et al. 1999: 402, G4977 [checklist]; Kwon 2001: 191, fig. 725; Min et al. 2004: 336, fig. 1058; Okutani 2000: 815, pl. 405, fig. 5; Noseworthy et al. 2007: 90 [checklist]; Okutani 2017: 1101, pl. 403, fig. 5.

## Materials

- a. scientificName: *Siphonaria sirius*; kingdom: Animalia; phylum: Mollusca; class: Gastropoda; order: Siphonariida; family: Siphonariidae; genus: *Siphonaria*; specificEpithet: *sirius*; scientificNameAuthorship: Pilsbry, 1894; country: Korea; locality: Dokdo-ri, Ulleung-eup, Ulleung-gun, Gyeongsangbuk-do; verbatimCoordinates: 37°14'20.12"N 131°52'05.74"E; eventDate: 2016-06-22; individualCount: 1; occurrenceID: D54B007E-530E-5A64-9E37-238C909DFC8B
- b. scientificName: *Siphonaria sirius*; kingdom: Animalia; phylum: Mollusca; class: Gastropoda; order: Siphonariida; family: Siphonariidae; genus: *Siphonaria*; specificEpithet: *sirius*; scientificNameAuthorship: Pilsbry, 1894; country: Korea; locality: Sagye-ro, Andeok-myeon, Seogwipo-si, Jeju-do; verbatimCoordinates: 33°13'58.7"N 126°22'23.8"E; verbatimCoordinateSystem: 33°13'13.2"N 126°17'42.0"E; eventDate: 2022-04-21; individualCount: 1; occurrenceID: C0A81431-17F4-52C1-891F-62FF72C4302D
- c. scientificName: *Siphonaria sirius*; kingdom: Animalia; phylum: Mollusca; class: Gastropoda; order: Siphonariida; family: Siphonariidae; genus: *Siphonaria*; specificEpithet: *sirius*; scientificNameAuthorship: Pilsbry, 1894; country: Korea; locality: Irun-myeon., Geoje-si, Gyeongsangnam-do; verbatimCoordinates: 34°58'34.9"N 128°41'18.18"E; eventDate: 2023-04-19; individualCount: 2; occurrenceID: BEE8EC41-E8D8-5759-BB91-328F2C2C0A3E
- d. scientificName: *Siphonaria sirius*; kingdom: Animalia; phylum: Mollusca; class: Gastropoda; order: Siphonariida; family: Siphonariidae; genus: *Siphonaria*; specificEpithet: *sirius*; scientificNameAuthorship: Pilsbry, 1894; country: Korea; locality: Namyang-ri, Seo-myeon, Ulleung-gun, Gyeongsangbuk-do; verbatimCoordinates: 37°27'35.6"N 130°51'27.0"E; eventDate: 2023-05-23; individualCount: 3; occurrenceID: 2213E690-E902-5F33-99A8-212A4C10304D
- e. scientificName: *Siphonaria sirius*; kingdom: Animalia; phylum: Mollusca; class: Gastropoda; order: Siphonariida; family: Siphonariidae; genus: *Siphonaria*; specificEpithet: *sirius*; scientificNameAuthorship: Pilsbry, 1894; country: Korea; locality: Mendehaeon-gil, Tongyeong-si, Gyeongsangnam-do; verbatimCoordinates: 34°50'37.8"N 128°26'36.0"E; eventDate: 2023-06-21; individualCount: 2; occurrenceID: 37D006D8-B765-5917-A845-CD6DD8473CD9
- f. scientificName: *Siphonaria sirius*; kingdom: Animalia; phylum: Mollusca; class: Gastropoda; order: Siphonariida; family: Siphonariidae; genus: *Siphonaria*; specificEpithet: *sirius*; scientificNameAuthorship: Pilsbry, 1894; country: Korea; locality: Guryongpo-eup, Nam-gu, Pohang-si, Gyeongsangbuk-do; verbatimCoordinates: 35°57'05.7"N 129°33'04.4"E; eventDate: 2023-07-23; individualCount: 1; occurrenceID: 2A53BAB4-3A3B-59CA-8E8A-716A5BC80FD6

## Description

Measurements: SL 11.23–19.63 mm, SW 9.09–15.62 mm, SH 2.87–4.93 mm.

Shell (Fig. 2d) solid, oval, small to medium in size (in examined specimens, SL 11.23–19.63 mm), low in height (about 1/4 of its length). Exterior colour greyish-brown or blackish-brown. Shell surface with 5–11 strong, white-coloured primary radial ribs. Interspaces between the primary radial ribs with 1–8 thinner secondary ribs, coloured white, sometimes same as shell colour. Apex often eroded, located central of the shell. Shell margin prominently undulated by primary ribs. Anterior slope slightly convex, posterior slope nearly straight. Siphonal groove protruded with one radial rib on outer shell surface. Interior colour reddish-brown to black with white bands, sometimes white or pale-coloured at centre.

Radula (Fig. 3e, f) dentition formulae 34:1:34 (in specimens measuring SL 14.69 mm, SW 11.80 mm, SH 3.89 mm). Each transverse row with narrow central rachidian tooth flanked by symmetrical half rows with lateral teeth, decreasing in size outwards. Rachidian tooth (Fig. 3e) short, approximately one-third length of lateral tooth, bud-shaped, middle widest, tapering gently towards the top with sharply-pointed cusp. Innermost lateral teeth (first 15 lateral teeth), each tooth consisting of mesocone and ectocone; mesocone bicuspid, with U-shaped cleft; ectocone shorter, sharp (Fig. 3e). Outermost lateral teeth (remaining 19 lateral teeth) with short and sharply-pointed endocone, ectocone and mesocone; mesocone subquadrate (Fig. 3f).

### Distribution

Korea, China, Japan, Philippines, Singapore, Indonesia (Sumatra Island) and Vietnam.

**Type locality:** Japan; Sagami, Kashiurazaki, Boshu (Boso Peninsula).

**Habitat:** On rocky substrate in middle to low intertidal zones (Fig. 2dG).

### Taxon discussion

This species is clearly distinguished from other Korean sympatric species (*S. acmaeoides* and *S. japonica*) by its flat shell, remarkably extended white radial ribs, siphonal groove with a single radial rib and pale-coloured inner centre (Fig. 2d). It closely resembles *S. atra* Quoy and Gaimard, 1833, by having a flat shell and strongly stretched radial ribs. Christiaens (1980a) noted that *S. sirius* represents one of the three forms of *S. laciniosa* (Linnaeus, 1758); (*S. laciniosa* forma *sirius*, *S. laciniosa* forma *atra* (representing *S. atra* Quoy and Gaimard, 1833) and *S. laciniosa* forma *subatra* (representing *S. subatra* Pilsbry, 1904). He reported no differences in niche selection between *S. atra* and *S. sirius*, both of which inhabit highly exposed rocky substrates in the low tidal zone at Ping Chau, Hong Kong. However, the two species differ in colouration and the number of ribs in the siphonal groove. While *S. atra* has a dusky brown shell with radial ribs of the same colour (sometimes paler than the interspaces) and a siphonal groove with two radial ribs, *S. sirius* is characterised by a black or dark brown-coloured shell with solid white radial ribs and a siphonal groove with only one radial rib. In the radula formula, as mentioned in the cases of *S.*

*acmaeoides* and *S. japonica*, *Siphonaria* species exhibits high levels of intraspecific variation. Similarly, the radula formulae of *S. sirius* reported in previous studies showed considerable variation depending on the authority, ranging from 40:1:40 (Hubendick 1946) to 34:1:34 (Christiaens 1980a). The radula formula (34:1:34) observed in this study is consistent with the finding of Christiaens (1980a). The phylogenetic tree shows that *S. sirius* forms a sister group with *S. atra* and *S. subatra* receiving 99% bootstrap value (Fig. 4). The *p*-distances within *S. sirius* species range from 0.16% to 0.79%, while this species shows high sequence divergence from other congeneric species ranging from 22.38% (*S. atra*) to 31.90% (*S. japonica*) (Table 2).

## Discussion

Previously, a total of seven *Siphonaria* species have been sparsely reported in Korea, based solely on external shell morphologies that has often resulted in taxonomic complications, including misidentification of species due to their remarkably high ecophenotypic variations. From a comprehensive analysis of shell morphology, radula structure and molecular analysis of mtDNA *cox1* sequences in this study, three species (*S. acmaeoides*, *S. japonica* and *S. sirius*) have been confirmed to occur in Korea. The three Korean *Siphonaria* species commonly share the presence of secondary ribs and often exhibit an eroded apex as they grow (Fig. 2). However, they are distinguished from each other by the number of primary ribs: 14–17 (in *S. acmaeoides*), 17–23 (in *S. japonica*) and 5–11 (in *S. sirius*), respectively. Furthermore, their primary ribs vary in thickness, elevation and spacing: *S. acmaeoides* has wide, low, unevenly spaced ribs; *S. japonica* has thin, moderately elevated, evenly spaced ribs; and *S. sirius* has wide, moderately elevated, unevenly spaced ribs. The morphology of the siphonal groove also varies amongst the species: *S. acmaeoides* has an indistinct groove with uncountable radial ribs, *S. japonica* has a weakly projected groove with two radial ribs and *S. sirius* has a strongly projected groove with a single radial rib. Very recently, Jenkins and Köhler (2024) compared the three species (*S. acmaeoides*, *S. japonica* and *S. sirius*) in their shell morphology and reproductive system, including the morphology of the spermatophore. Their shell morphology comparison focused on a set of characters such as the solidity and crenulation of the shell margin, which are less informative for distinguishing sympatric species due to substantial intraspecific variation that complicates precise species identification. Nevertheless, they provided anatomical comparisons, noting that *S. acmaeoides* differs from *S. japonica* by having a smaller accessory organ and bursa copulatrix, a shorter flagellum and a short drop-like spermatophore. In contrast, *S. sirius* has a larger accessory organ and a spermatophore lacking barbs. In radula morphology, *S. acmaeoides* has only a mesocone in the innermost lateral teeth, whereas both *S. japonica* and *S. sirius* possess a mesocone with sharply-pointed ectocone in the inner lateral teeth (Fig. 3). Despite these distinct morphological features, the radula formulae described in previous studies, including our observations, showed considerable individual variations (Hubendick 1946, Christiaens 1980a).

Our phylogenetic analysis using mtDNA *cox1* sequences of these three Korean *Siphonaria* species and some other congeneric species confirmed their morphology-based species identification (Fig. 4). The resulting tree indicated that the mtDNA *cox1* sequences determined from *S. acmaeoides*, *S. japonica* and *S. sirius* respectively formed monophyletic groups, corroborating their identification based on morphological characteristics. A significantly high sequence gap was observed between the lowest intraspecific distance (3.33%) and the highest interspecific distance (27.94%) amongst these three species. Interestingly, *S. acmaeoides* was found as closely related to *S. zelandica*, showing a relatively low genetic divergence (5.71–6.51%; Table 2). *S. acmaeoides* is known to be distributed in Japan, Korea (Kuroda et al. 1971, Choe 1992, Okutani 2000, Cheng et al. 2005, Yokogawa et al. 2010, Okutani 2017), whereas *S. zelandica* is found along the southern coasts of Australia (Quoy and Gaimard 1833, Jenkins 1983). Dayrat et al. (2014) proposed the possibility of an extensive distribution of *S. zelandica* ranging from Japan to Australia and suggested that *S. acmaeoides* could be a junior synonym of *S. zelandica*. However, Jenkins and Köhler (2024) confirmed that *S. acmaeoides* and *S. zelandica* are distinct species, based on comparative analyses of shell morphology, soft-body anatomy and mitochondrial phylogenetics. Unlike the three *Siphonaria* species confirmed to occur in Korea, the taxonomic validity of the other four species (*S. coreensis*, *S. javanica*, *S. laciniosa* and *S. rucuana*), previously reported from Korean waters is questionable. Despite extensive follow-up efforts over nearly the last two decades, these species have rarely been found. In particular, the shell images and morphological features described in the previous Korean literature do not closely match the original descriptions, but instead appear very similar to *S. japonica*, one of the most widely and abundantly found *Siphonaria* species in the north-western Pacific, including the Korean coastline. Moreover, none of these studies provided detailed descriptions of diagnostic characters and their species identifications were based solely on shell characteristics, raising questions about the taxonomic validity of their conclusions.

In conclusion, many molluscan species, including *Siphonaria* species, exhibit a high degree of morphological plasticity in shell shape (including sculptures), size and colour, which can vary significantly depending on environmental conditions. These variations in shell morphology make it challenging to differentiate species based on morphology alone, often resulting in various taxonomic errors, such as misidentifications, overlooking cryptic species diversity or a plethora of nominal species. Our taxonomic review incorporating both morphological and molecular data revealed that Korean *Siphonaria* species also display a wide range of shell morphological variations within species (e.g. shell colour, number of radial ribs and the slope of the anterior and posterior shell). Moreover, in many cases — including those observed in the present study — some species exhibiting a wide range of shell morphological variations are often found co-occurring in the same intertidal habitats. A comprehensive analysis of integrated morphological and molecular data, obtained through extensive taxon sampling along the Korean coastline can provide better resolution to address taxonomic ambiguities amongst *Siphonaria* species.



## Acknowledgements

This work was supported by a grant from the National Institute of Biological Resources (NIBR), funded by the Ministry of Environment (MOE) of the Republic of Korea (NIBR202231206) and the National Research Foundation of Korea (NRF) grant funded by the Korea government (MSIT) (2020R1A2C2005393) to JKP and by Global-Learning & Academic research institution for Masters'-PhD students and Postdocs (LAMP) Program of the National Research Foundation of Korea (NRF) grant funded by the Ministry of Education (No. RS-2023-00301914) to UWH.

## References

- Abe N (1940) The homing, spawning and other habits of a limpet, *Siphonaria japonica* Donovan. Scientific Reports of the Tohoku Imperial University 15 (1): 59-95.
- Adams A, Reeve L (1850) The Zoology of the voyage of H.M.S. Samarang, under the command of Captain Sir Edward Belcher, C.B., F.R.A.S., F.G.S., during the years 1843-1846 (Mollusca). Reeve and Benham, London.
- Adams H, Adams A (1855) The genera of recent Mollusca. Vol. 2. John Van Voorst, London.
- Azuma M (1960) A catalogue of the shell-bearing Mollusca of Okinoshima, Kasiwajima and the adjacent area (Tosa Province) Shikoku, Japan. ConchBooks, Harxheim, 102 pp.
- Baker B (1964) Type land snails in the Academy of Natural Sciences of Philadelphia Part III. Limnophile and Thalassophile Pulmonata Part IV. Land and fresh-water Prosobranchia. Proceedings of the Academy of Natural Sciences of Philadelphia 116 (4): 149-193.
- Berry AJ (1977) Chapter 3. Gastropoda: Pulmonata. In: Giese AC, Pearse JS (Eds) Reproduction of Marine Invertebrates. 4. Academic Press, USA, 181-226 pp. <https://doi.org/10.1016/B978-0-12-282504-0.50010-X>
- Cernohorsky WO (1972) Marine shells of the Pacific. 2. Pacific, Sydney, 411 pp.
- Cheng YT, Yeh TY, Chang CY, Hsueh PW (2005) New records of marine gastropods from Taiwan. Collections and Research of the National Museum of Natural Science 18: 7-9.
- Choe BL (1992) Illustrated Encyclopedia of Fauna and Flora of Korea. Vol. 33, Mollusca (II). Ministry of Education, Seoul, 859 pp.
- Christiaens J (1980a) The limpets of Hong Kong with descriptions of seven new species and subspecies. In: Morton B (Ed.) The Malacofauna of Hong Kong and Southern China. Hong Kong University Press, Hong Kong.
- Christiaens J (1980b) Supplementary notes on Hong Kong limpets. In: Morton BS, Tseng CK (Eds) Proceedings from the First International Marine Biological Workshop: The Marine Flora and Fauna of Hong Kong and Southern China, Hong Kong. Hong Kong University Press, Hong Kong, 459-468 pp.
- Coan EV, Lutaenko KA, Zhang J, Sun Q (2015) The molluscan taxa of AW Grabau & SG King (1928) and their types. Malacologia 58: 179-224. <https://doi.org/10.4002/040.058.0206>

- Colgan DJ, Da Costa P (2013) Possible drivers of biodiversity generation in the *Siphonaria* of southeastern Australia. *Marine Biodiversity* 43: 73-85. <https://doi.org/10.1007/s12526-012-0127-2>
- Collin R (2000) The development of three heterobranch mollusks from California, USA. *Nautilus* 114 (3): 117-119.
- Collins RA, Boykin LM, Cruickshank RH, Armstrong KF (2012) Barcoding's next top model: an evaluation of nucleotide substitution models for specimen identification. *Methods in Ecology and Evolution* 3 (3): 457-465. <https://doi.org/10.1111/j.2041-210X.2011.00176.x>
- Dayrat B, Goulding TC, White TR (2014) Diversity of Indo-West Pacific *Siphonaria* (Mollusca: Gastropoda: Euthyneura). *Zootaxa* 3779 (2): 246-276. <https://doi.org/10.11646/zootaxa.3779.2.7>
- De Coito PM (2021) Phylogeny and biogeography of southern African limpets in the genus *Siphonaria* in the context of a global phylogeny. University of Cape Town, Cape Town, 187 pp.
- Donovan E (1824) *Naturalist's repository, or monthly miscellany of exotic natural history*. W. Simpkin and R. Marshall, London.
- Edgar RC (2004) MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research* 32 (5): 1792-1797. <https://doi.org/10.1093/nar/gkh340>
- Felsenstein J (1985) Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* 39 (4): 783-791. <https://doi.org/10.2307/2408678>
- Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R (1994) DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology* 5: 294-299.
- Fukuda H, Mashino K, Sugimura T (1992) A review of the molluscan fauna of Yamaguchi Prefecture, western Japan. Yamaguchi Museum, Yamaguchi.
- González-Wevar CA, Segovia NI, Rosenfeld S, Ojeda J, Hüne M, Naretto J, Saucedo T, Brickle P, Morley S, Féral J (2018) Unexpected absence of island endemics: Long-distance dispersal in higher latitude sub-Antarctic *Siphonaria* (Gastropoda: Euthyneura) species. *Journal of Biogeography* 45 (4): 874-884. <https://doi.org/10.1111/jbi.13174>
- Grabau AW, King SG (1928) *Shells of Peitaiho*. 2nd Edition. Peking Society of Natural History, Beijing, 279 pp.
- Güller M, Zelaya DG, Ituarte C (2016) How many *Siphonaria* species (Gastropoda: Euthyneura) live in southern South America? *Journal of Molluscan Studies* 82 (1): 80-96.
- Habe T, Kikuchi T (1960) *Fauna and Flora of the Sea around the Amakusa Marine Biological Laboratory, Part I Mollusca*. Tomioka, Reihoku-cho, Amakusa, Kumamoto-ken, Japan.
- Habe T, Igarashi T (1967) A list of marine molluscan shells in the Fisheries Museum, Faculty of Fisheries, Hokkaido University (Kawasaki collection and specimens collected by the Marine Zoological Laboratory). 6. *Bulletin of the Faculty of Fisheries Hokkaido University, Hokkaido*, 56 pp.
- Habe T (1971) *Shells of Japan*. Hoikusha, Japan, 139 pp.
- Hamamura Y (2004) *An Illustrated Guide to Marine Shells of Geinan Region*. Ranto Bunka Shinko Zaidan, Hiroshima, 220 pp.
- Hanley S (1858) On *Siphonaria*. *Proceedings of the Zoological Society of London* 26: 151-153.

- Higo S, Callomon P, Goto Y (1999) Catalogue and bibliography of the marine shell-bearing Mollusca of Japan. Elle Scientific Publications
- Hirase S (1941) A collection of Japanese shells with illustrations in natural colours. 7. Matsumura Sanshodo, Tokyo, 1-217 pp.
- Hodgson AN (1999) The biology of siphonariid limpets (Gastropoda: Pulmonata). In: Margaret Barnes ADR (Ed.) Oceanography and marine biology. 37. CRC Press, The Dunstaffnage Marine Laboratory, Oban, Argyll, Scotland, 69 pp.
- Hubendick B (1945) Phylogenie und Tiergeographie der Siphonariidae: Zur Kenntnis der Phylogenie in der Ordnung Basommatophora und des Ursprungs der Pulmonatengruppe. Almquist och Wiksells boktryck, Upsala, 1-216 pp.
- Hubendick B (1946) Systematic monograph of the Patelliformia. Kungliga Svenska Vetenskapsakademiens Handlingar Ser. 3 23 (5): 1-93.
- Hylleberg J, Kilburn RN (2003) Marine molluscs of Vietnam. Phuket Marine Biological Center, Special Publication 28: 1-300.
- Inaba A (1983) Fauna and flora of the Seto Inland Sea. 2, 1. Mukaishima Marine Biological Station, Hiroshima University
- Ip YCA, Ywee CT, Su XG, Hui PA, Karenne T, Loke MC, Danwei H, Rudolf M (2019) From marine park to future genomic observatory? Enhancing marine biodiversity assessments using a biocode approach. Biodiversity Data Journal 7.
- Je JG (1989) Korean names of molluscs in Korea. 1. The Korean Journal of Malacology, Seoul, 90 pp.
- Jenkins B (1983) Redescriptions and relationship of *Siphonaria zelandica* Quoy and Gaimard to *S. australis* Quoy and Gaimard with a description of *S. propria* sp. nov. (Mollusca: Pulmonata: Siphonariidae). Journal of the Malacological Society of Australia 6 (1-2): 1-35. <https://doi.org/10.1080/00852988.1983.10673952>
- Jenkins B, Köhler F (2024) Hidden in plain sight: Systematic review of Indo-West Pacific Siphonariidae uncovers extensive cryptic diversity based on comparative morphology and mitochondrial phylogenetics (Mollusca, Gastropoda). Megataxa 13 (1): 1-217. <https://doi.org/10.11646/megataxa.13.1.1>
- Kalyaanamoorthy S, Minh BQ, Wong TK, Von Haeseler A, Jermini LS (2017) ModelFinder: fast model selection for accurate phylogenetic estimates. Nature Methods 14 (6): 587-589. <https://doi.org/10.1038/nmeth.4285>
- Kira T (1962) Shells of the western Pacific in color. Hoikusha, Japan, 1-244 pp.
- Klussmann-Kolb A, Dinapoli A, Kuhn K, Streit B, Albrecht C (2008) From sea to land and beyond—new insights into the evolution of euthyneuran Gastropoda (Mollusca). BMC Evolutionary Biology 8: 1-16. <https://doi.org/10.1186/1471-2148-8-57>
- Kuroda T (1941) A catalogue of molluscan shells from Taiwan (Formosa), with descriptions of new species. Memoirs of the Faculty of Science and Agriculture 22 (2): 65-216.
- Kuroda T, Habe T (1952) Check list and bibliography of the recent marine Mollusca of Japan. Hosokawa, Japan, 1-210 pp.
- Kuroda T (1960) A Catalogue of Molluscan fauna of the Okinawa Islands. Ryukyu University Kyomubu Fukyuka, Tokyo, 1-106 pp.
- Kuroda T, Habe T, Oyama K (1971) The sea shells of Sagami Bay: collected by His Majesty the Emperor of Japan. Maruzen, Tokyo, 1280 pp.
- Kwon OK, Park GM, Lee JS (1993) Coloured Shells of Korea. Academy Publishing Company, Seoul, 445 pp.

- Kwon OK (2001) Korean mollusks with color illustration. Shell House, Busan, 332 pp.
- Lamarck JdMd (1819) Histoire naturelle des animaux sans vertèbres. 6. Verdière, Paris, 343 pp.
- Lee BD (1956) The catalogue of molluscan shells of Korea. Bulletin of Pusan Fisheries College, 48 pp.
- Lee JS, Min DK (2002) A catalogue of molluscan fauna in Korea. The Korean Journal of Malacology 18 (2): 93-217.
- Lee SC, Chao SM (2003) Shallow-water marine shells from northeastern Taiwan. Collection and Research 16: 29-59.
- Lischke VE (1871) Japanische Meeres-Conchyliden. Ein Beitrag zur Kenntnis der Mollusken Japans, mit besonderer Rücksicht auf die geographische Verbreitung derselben. 2. Theodor Fischer, Cassel, 184 pp.
- Min DK, Lee JS, Koh DB, Je JG (2004) Mollusks in Korea (revised supplementary edition). Min Molluscan Research Institute, Seoul, 566 pp.
- MolluscaBase eds (2024) MolluscaBase. <https://www.molluscabase.org>. Accessed on: 2024-8-01.
- MolluscaBase eds (2025) MolluscaBase. *Siphonaria coreensis* A. Adams & Reeve, 1848. <https://www.marinespecies.org/aphia.php?p=taxdetails&id=883559>. Accessed on: 2025-2-13.
- Morrison J (1972) Mediterranean *Siphonaria*: west and east—old and new. Argamon 3: 51-62.
- Morton B, Morton J (1983) The sea shore ecology of Hong Kong. Hong Kong University Press, Hong Kong, 1-350 pp. <https://doi.org/10.1163/9789004630673>
- National Institute of Biological Resources (2023) National List of Korea. <http://kbr.go.kr/>. Accessed on: 2024-1-30.
- Nguyen L, Schmidt HA, Von Haeseler A, Minh BQ (2015) IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. Molecular Biology and Evolution 32 (1): 268-274. <https://doi.org/10.1093/molbev/msu300>
- Noseworthy RG, Lim N, Choi K (2007) A catalogue of the mollusks of Jeju Island, South Korea. Korean Journal of Malacology 23 (1): 65-104.
- Okutani T (2000) Marine mollusks in Japan. Tokai University Press, Tokyo, 1173 pp.
- Okutani T (2017) Marine mollusks in Japan. 2nd Edition. Tokai University Press, Hiratsuka, 1375 pp.
- Oyama K, Yamamoto T, Tokioka T (1954) Invertebrate fauna of the intertidal zone of the Tokara Islands. Publications of the Seto Marine Biological Laboratory 4 (1): 1-15. <https://doi.org/10.5134/174506>
- Pilsbry HA (1894) Notices of new Japanese mollusks, II. The Nautilus 8 (1): 9-10.
- Pilsbry HA (1895) Catalogue of the marine mollusks of Japan, with descriptions of new species and notes on others collected by Frederick Stearns. The Nautilus 8: 9-10.
- Pilsbry HA (1904) New Japanese marine mollusca: Gastropoda. Vol. 56. Proceedings of the Academy of Natural Sciences of Philadelphia, 895 pp.
- Pilsbry HA (1920) *Siphonaria japonica* Donovan an earlier name for *S. cochleariformis* Reeve. Nautilus 34: 141.
- Qi Z (2004) Seashells of China. China Ocean Press, Beijing, 418 pp.
- Quoy J, Gaimard J (1833) Zoophytes. Voyage de découvertes de l'Astrolabe exécuté par ordre du Roi, pendant les années 2: 1-686.

- Reeve LA (1856) *Conchologia iconica, or, Illustrations of the shells of molluscous animals*. Vol. 9. Benham & Reeve, London.
- Shikama T (1964) *Selected shells of the world: Illustrated in colours (II)*. Japan Fisheries Resource Conservation Association, Hokuryu-Kan, Tokyo, 1-212 pp.
- Sowerby GB (1823) *Siphonaria*. In: Sowerby GB (Ed.) *The genera of Recent and fossil shells*. London.
- Springsteen FJ, Leobrera FM (1986) *Shells of the Philippines*. Carfel Seashell Museum, Manila, 1-377 pp.
- Srivathsan A, Meier R (2012) On the inappropriate use of Kimura-2-parameter (K2P) divergences in the DNA-barcoding literature. *Cladistics* 28 (2): 190-194. <https://doi.org/10.1111/j.1096-0031.2011.00370.x>
- Toyohara T, Nakaoka M, Tsuchida E (2001) Population dynamics and life history traits of *Siphonacmea oblongata* (Yokoyama) on seagrass leaves in Otsuchi Bay, north-eastern Japan (Siphonariidae, Pulmonata). *Venus* 60 (1-2): 27-36.
- Trew A (1983) Part 16, Siphonariacea. National Museum of Wales, Wales. UK.
- Vermeij G (1971) Temperature relationships of some tropical Pacific intertidal gastropods. *Marine Biology* 10: 308-314. <https://doi.org/10.1007/BF00368090>
- White TR, Conrad M, Tseng R, Balayan S, Golding R, Martins AMdF, Dayrat B (2011) Ten new complete mitochondrial genomes of pulmonates (Mollusca: Gastropoda) and their impact on phylogenetic relationships. *BMC Evolutionary Biology* 11: 1-15. <https://doi.org/10.1186/1471-2148-11-295>
- White TR, Dayrat B (2012) Checklist of genus-and species-group names of the false limpets *Siphonaria* (Mollusca: Gastropoda: Euthyneura). *Zootaxa* 3538 (1): 78-54.
- Yokogawa K, Ishikawa H, Hamamura Y (2010) Genetic divergence between two forms of the pulmonate limpet *Siphonaria japonica* (Gastropoda: Siphonariidae). *Venus* 68 (3): 139-149.
- Yoo J (1976) *Korean Shells in Colour*. Iljisa, Seoul, 196 pp.