



ISSN: 2617-1295

Effect of organic and inorganic nutrients on rice

S. Manimaran*, V. Prakash

Department of Agronomy, Faculty of Agriculture, Annamalai University, Annamalai Nagar 608 002, Tamil Nadu, India

ABSTRACT

Field experiment was conducted to study the influence of organic and inorganic sources of nutrients on yield parameters of rice (ADT 36). The experiment consists of twelve treatments with three replications viz., T₁-100 % Recommended Dose (RD) of NPK kg ha⁻¹ (120:38:38), T₂ - RD with soil application of Beema green granules 25 Kg ha⁻¹ at 20 DAT, T₃ - 75 % RD of NPK kg ha⁻¹ (90:38:38) with soil application of Beema green granules 25 Kg ha⁻¹ at 20 DAT, T₄ - RD with foliar spray of panchagavya 3 % at 30 and 45 DAT, T₅ - RD with soil application of Beema green granules 25 Kg ha⁻¹ 20 DAT with foliar spray of panchagavya 3 % at 30 and 45 DAT, T₆ - 75 % RD of NPK kg ha⁻¹ (90:38:38) with soil application of Beema green granules 25 Kg ha⁻¹ 20 DAT with foliar spray of panchagavya 3 % at 30 and 45 DAT, T₇ - RD with foliar spray of Nitrobenzene 750ml ha⁻¹ at 20 DAT, T₈ - RD with soil application of Beema green granules 25 Kg ha⁻¹ at 20 DAT with foliar spray of Nitrobenzene 750ml ha⁻¹ at 30 DAT, T₉ - 75 % RD of NPK kg ha⁻¹ (90:38:38) with soil application of Beema green granules 25 Kg ha⁻¹ at 20 DAT with foliar spray of Nitrobenzene 750ml ha⁻¹ at 30 DAT, T₁₀ - RD with foliar spray of panchagavya 3 % at 30 DAT and foliar spray of Nitrobenzene 750ml ha⁻¹ at 45 DAT, T₁₁ - RD with soil application of Beema green granules 25 Kg ha⁻¹ at 20 DAT and foliar spray of panchagavya 3 % at 30 DAT and foliar spray of Nitrobenzene 750ml ha⁻¹ at 45 DAT, T₁₂ - 75 % RD of NPK kg ha⁻¹ (90:38:38) with soil application of Beema green granules 25 Kg ha⁻¹ at 20 DAT and foliar spray panchagavya 3 % at 30 DAT and foliar spray of Nitrobenzene 750ml ha⁻¹ at 45 DAT. The growth parameters of rice viz., plant height, number of tillers per hill, LAI and DMP was recorded higher in 100 % recommended dose of NPK kg ha⁻¹ (120:38:38) with soil application of Beema green granules 25 Kg ha⁻¹ at 20 DAT and foliar spray of panchagavya 3 % at 30 DAT and foliar spray of Nitrobenzene 750ml ha⁻¹ at 45 DAT (T₁₁) compared to other treatments.

Received: February 18, 2018

Accepted: March 27, 2018

Published: March 31, 2018

*Corresponding Author:

S. Manimaran

Email:sambamaran@gmail.com

KEYWORDS: Rice, beema green, panchagavya, nitrobenzene, foliar application

INTRODUCTION

Rice (*Oryza sativa*) is important staple food crop that is being extensively cultivated in India [1]. Even though rice cultivation is common in India and provides a stable income to people, the yield is still lower, when compared to the average productivity of rice producing countries such as Japan, China, Egypt and Israel [2,3].

If the inorganic fertilizers are used without any proper care and application practices, it can lead soil degradation [4]. Now a day the sources of organic matter are scarce, due to shortage and non-availability of labour to rear animals. Humic acid, a decomposed product of organic matter, influences plant growth by modifying the physiology of plant and by improving the physical, chemical and biological properties of soil and ultimately crop yield [5]. Organics like humic substances will enhance the plant and soil properties [6]. Panchagavya, an organic source of nutrition, is an indigenous material which is used widely for agricultural and horticultural crops. Panchagavya had positive influence on

beneficial microorganisms present in the soil and influence the crop growth and yield [7]. Panchagavya is now gaining attention as an efficient organic growth promotor [8,9]. Keeping in view of sustainable agronomic practices to increase the yield of rice, the field investigation was taken up.

MATERIALS AND METHODS

The field experiment was conducted at the Experimental Farm, Department of Agronomy, Annamalai University, Tamil Nadu, during the period of February to May, 2014, to study the influence of organic and inorganic sources of nutrients on growth parameters of rice. The cropping period the mean maximum temperature ranges between 29.2°C and 37.3°C, while the minimum temperature ranges from 20.9°C to 26.6°C. The RH ranges from 76 to 96 percent. The soil was clay loam in texture with low in available nitrogen, medium in available phosphorus and high in available potassium. A short duration rice variety ADT 36 was chosen with a duration of 110 days.

Copyright: © 2018 The authors. This article is open access and licensed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>) which permits unrestricted, use, distribution and reproduction in any medium, or format for any purpose, even commercially provided the work is properly cited. Attribution — You must give appropriate credit, provide a link to the license, and indicate if changes were made.

The experiment was laid out in Randomized Block Design (RBD) with three replication with twelve treatments *viz.*, T₁-100 % Recommended Dose (RD) of NPK kg ha⁻¹ (120:38:38), T₂-T₁ with soil application of Beema green granules 25 Kg ha⁻¹ at 20 DAT, T₃-75 % RD of NPK kg ha⁻¹ (90:38:38) with soil application of Beema green granules 25 Kg ha⁻¹ at 20 DAT, T₄-T₁ with foliar spray of panchagavya 3 % at 30 and 45 DAT, T₅-T₁ with soil application of Beema green granules 25 Kg ha⁻¹ 20 DAT with foliar spray of panchagavya 3 % at 30 and 45 DAT, T₆-75 % RD of NPK kg ha⁻¹ (90:38:38) with soil application of Beema green granules 25 Kg ha⁻¹ 20 DAT and foliar spray of panchagavya 3 % at 30 and 45 DAT, T₇-T₁ with foliar spray of Nitrobenzene 750ml ha⁻¹ at 20 DAT, T₈-T₁ with soil application of Beema green granules 25 Kg ha⁻¹ at 20 DAT and foliar spray of Nitrobenzene 750ml ha⁻¹ at 30 DAT, T₉-75 % RD of NPK kg ha⁻¹ (90:38:38) with soil application of Beema green granules 25 Kg ha⁻¹ at 20 DAT and foliar spray of Nitrobenzene 750ml ha⁻¹ at 30 DAT, T₁₀-T₁ with foliar spray of panchagavya 3 % at 30 DAT and foliar spray of Nitrobenzene 750ml ha⁻¹ at 45 DAT, T₁₁-T₁ with soil application of Beema green granules 25 Kg ha⁻¹ at 20 DAT and foliar spray of panchagavya 3 % at 30 DAT and foliar spray of Nitrobenzene 750ml ha⁻¹ at 45 DAT, T₁₂-75 % RD of NPK kg ha⁻¹ (90:38:38) with soil application of Beema green granules 25 Kg ha⁻¹ at 20 DAT and foliar spray panchagavya 3 % at 30 DAT and foliar spray of Nitrobenzene 750ml ha⁻¹ at 45 DAT.

Beema green granules is an organic humic granules and it contain humic acid, fulvic acid, amino acid and micronutrients. Beema green granules @ 25 kg ha⁻¹ was applied at 20 DAS evenly over respective plots. The panchagavya was applied through foliar @ 3 % at different growth stages of rice. Nitrobenzene plant growth promoter was used as foliar spray @ 750 ml ha⁻¹. The solution was uniformly sprayed using hand sprayer (knap sack) at 30 and 45 DAT as per the treatment schedule. The 26 days old paddy seedlings were transplanted in lines @ 2 seedlings hill⁻¹ in a depth of 2-3cm. A spacing of 12.5 was adopted.

RESULTS AND DISCUSSION

Plant Height at Harvest

The plant height observed at harvest is presented in Table 1. Among the different treatments, the highest plant height at harvest was observed under T₁₁-100% recommended dose of NPK kg ha⁻¹ (120:38:38) with soil application of Beema green granules 25 kg ha⁻¹ and foliar spray of Panchagavya 3% and foliar spray of Nitrobenzene 750 ml ha⁻¹. This treatment was found to be superior and recorded the plant height of 98.95 cm at harvest. It was on par with the treatment T₁₂-75% recommended dose of NPK kg ha⁻¹ (90:38:38) with soil application of Beema green granules 25 kg ha⁻¹ and foliar spray of Panchagavya 3% and foliar spray of Nitrobenzene 750 ml ha⁻¹. The least plant height (60.42 cm) was recorded in T₃-75 % recommended dose of NPK kg ha⁻¹ (90:38:38) with soil application of Beema green granules 25kg ha⁻¹. The enhancement in growth with increase in nutrient availability was owing to rapid conversion of synthesized photosynthates into protein to form more

protoplasm, this increased the number and size of the cell which might have increased the plant height. Similar finding was reported by Sridevi [10].

Number of Tillers Hill⁻¹

Among the treatments tried, highest values on number of tillers hill⁻¹ (Table 1) was observed under T₁₁-100% recommended dose of NPK kg ha⁻¹ (120:38:38) with soil application of Beema green granules 25 kg ha⁻¹ and foliar spray of Panchagavya 3 % and foliar spray of Nitrobenzene 750 ml ha⁻¹ with a value of 7.00. It was on par with the treatment T₁₂-75% recommended dose of NPK kg ha⁻¹ (90:38:38) with soil application of Beema green granules 25 kg ha⁻¹ and foliar spray of Panchagavya 3% and foliar spray of Nitrobenzene 750ml ha⁻¹. The least number of tillers were recorded in T₃-75% recommended dose of NPK kg ha⁻¹ (90:38:38) with soil application of Beema green granules 25 kg ha⁻¹ of 5.02. The increase in number of tillers hill⁻¹ may be due to cumulative effect of fertilizer under increasing trend of nutrients. Similar result was reported by Nageswari and Balasubramaniyan [11].

Leaf Area Index (LAI)

The leaf area index at flowering are presented in Table 1. Among the various treatments, T₁₁-100 % Recommended dose of NPK kg ha⁻¹ (120:38:38) with soil application of Beema green granules 25 kg ha⁻¹ and foliar spray of Panchagavya 3% and foliar spray of Nitrobenzene 750 ml ha⁻¹ recorded a highest LAI of 6.48. It was on par with the treatment T₁₂-75 % recommended dose of NPK kg ha⁻¹ (90:38:38) with soil application of Beema green granules 25 kg ha⁻¹ + foliar spray of Panchagavya 3% and foliar spray of Nitrobenzene 750 ml ha⁻¹. The lowest LAI was observed under T₃-75 % Recommended dose of NPK kg ha⁻¹ (90:38:38) with soil application of Beema green granules 25 kg ha⁻¹ with a value of 4.29. Adequate supply of the essential nutrients i.e. N, P and K might have exerted a positive influence on the tillers and leaves leading to higher leaf area index. The results are in conformity with the findings of Manivannan [12].

Dry Matter Production

Among the different treatments, the highest dry matter production (Table 1) at harvest was recorded under T₁₁-100 % recommended dose of NPK kg ha⁻¹ (120:38:38) with soil application of Beema green granules 25 Kg ha⁻¹ and foliar spray of Panchagavya 3% and foliar spray of Nitrobenzene 750 ml ha⁻¹ with a value of 12034 kg ha⁻¹. The T₁₁ treatment on par with T₁₂-75 % Recommended dose of NPK kg ha⁻¹ (90:38:38) with soil application of Beema green granules 25 kg ha⁻¹ and foliar spray of Panchagavya 3% and foliar spray of Nitrobenzene 750 ml ha⁻¹. The lowest dry matter production was recorded under T₃-75 % Recommended dose of NPK kg ha⁻¹ (90:38:38) with soil application of Beema green granules 25 kg ha⁻¹ which recorded a value of 6670 kg ha⁻¹. Increased dry matter accumulation observed under higher NPK level due higher portion of green leaves with increased chlorophyll content enabling the plants to have increase the photosynthetic

Table 1: Influence of organic and inorganic sources of nutrients on growth parameters of rice

Treatment	Plant height at harvest	No. of tillers hill ⁻¹ at tillering	LAI at flowering	DMP at harvest (kg ha ⁻¹)
T ₁ – 100% RDF (120:38:38 NPK kg ha ⁻¹)	64.80	5.39	4.69	7705
T ₂ – 100% RDF (120:38:38 NPK kg ha ⁻¹) + Beema green granules 25 kg ha ⁻¹	68.29	5.20	4.86	7886
T ₃ – 75% RDF (90:38:38 NPK kg ha ⁻¹) + Beema green granules 25 kg ha ⁻¹	60.42	5.02	4.29	6670
T ₄ – 100% RDF (120:38:38 NPK kg ha ⁻¹) + Panchagavya 3%	71.68	5.57	5.02	8845
T ₅ – 100% RDF (120:38:38 NPK kg ha ⁻¹) + Beema green granules 25 kg ha ⁻¹ + Panchagavya 3%	88.78	6.57	5.68	9817
T ₆ – 75% RDF (90:38:38 NPK kg ha ⁻¹) + Beema green granules 25 kg ha ⁻¹ + Panchagavya 3%	78.56	5.94	5.29	9340
T ₇ – 100% RDF (120:38:38 NPK kg ha ⁻¹) + Nitrobenzene 750ml ha ⁻¹	75.11	5.76	5.14	8948
T ₈ – 100% RDF (120:38:38 NPK kg ha ⁻¹) + Beema green granules 25 kg ha ⁻¹ + Nitrobenzene 750ml ha ⁻¹	85.39	6.37	5.56	9648
T ₉ – 75% RDF (90:38:38 NPK kg ha ⁻¹) + Beema green granules 25 kg ha ⁻¹ + Nitrobenzene 750ml ha ⁻¹	81.98	6.16	5.43	9381
T ₁₀ – 100% RDF (120:38:38 NPK kg ha ⁻¹) + Panchagavya 3% + Nitrobenzene 750ml ha ⁻¹	92.21	6.76	6.26	11176
T ₁₁ – 100% RDF (120:38:38 NPK kg ha ⁻¹) + Beema green granules 25 kg ha ⁻¹ + Panchagavya 3% + Nitrobenzene 750ml ha ⁻¹	98.95	7.00	6.48	12034
T ₁₂ – 75% RDF (90:38:38 NPK kg ha ⁻¹) + Beema green granules 25 kg ha ⁻¹ + Panchagavya 3% + Nitrobenzene 750ml ha ⁻¹	97.62	6.95	6.45	11990
SEd	1.56	0.07	0.05	84
CD (P = 0.05)	3.38	0.17	0.11	180

efficiency and resulted in vigorous crop growth coupled with more tillers. The above finding was in conformity with the result of Latt *et al.* [13].

REFERENCES

- Gokulakrishnan J. Heterosis for yield and yield attributed traits in rice (*Oryza sativa* L.). Horticultural Biotechnology Research. 2018; 4: 26-29.
- Bruinsma J. The resource outlook to 2050: by how much do land, water and crop yields need to increase by 2050. In Expert meeting on how to feed the world in 2009 (Vol. 2050, pp. 24-26).
- Dhakar R, Yadav RK, Paudel DP, Deosarkar DB. Studies on genetic divergence and reaction to iron chlorosis in aerobic rice (*Oryza sativa* L.) on vertisols in Maharashtra. Journal of Plant Stress Physiology. 2017; 3: 12-21.
- Tilman D, Cassman KG, Matson PA, Naylor R, Polasky S. Agricultural sustainability and intensive production practices. Nature. 2002;418(6898):671.
- Bot A, Benites J. The importance of soil organic matter: Key to drought-resistant soil and sustained food production. Food & Agriculture Org.; 2005.
- Schmidt MW, Torn MS, Abiven S, Dittmar T, Guggenberger G, Janssens IA, Kleber M, Kögel-Knabner I, Lehmann J, Manning DA, Nannipieri P. Persistence of soil organic matter as an ecosystem property. Nature. 2011;478(7367):49.
- Mehdizadeh M, Darbandi EI, Naseri-Rad H, Tobeh A. Growth and yield of tomato (*Lycopersicon esculentum* Mill.) as influenced by different organic fertilizers. International journal of Agronomy and plant production. 2013;4(4):734-8.
- Nene YL. A critical discussion on the methods currently recommended to support organic crop farming in India. Asian Agri-History. 2017;21(4):265-83.
- Nagaraj N, Sreenivasa MN. Influence of bacteria isolated from panchagavya on seed germination and seed vigour in wheat. Karnataka Journal of Agricultural Sciences. 2009;22(1):231-2.
- Sridevi, V. Studies on crop establishment techniques on weed and nutrient management in lowland rice. 2011; Ph.D., Thesis, TNAU, Coimbatore
- Nageswari R, Balasubramaniyan P. Influence of delayed basal dressing and split application of nitrogen in wet-seeded rice (*Oryza sativa*). Indian Journal of Agronomy (India). 2004.
- Manivannan, K. Studies on the effect of different levels of nitrogen and potassium on the growth and yield of rice hybrids. DRRH-1 and KRH2. 2003; M.Sc.(Ag.) Thesis, Annamalai Univ., Annamalai Nagar
- Latt YK, Myint AK, Yamakawa T, Ogata K. The effects of green manure (*Sesbania rostrata*) on the growth and yield of rice. Journal of Faculty of Agriculture of Kyushu University. 2009;54:313-9.