

***ATRACTOSTEUS SPATULA* (ACTINOPTERYGII: LEPISTOSTEIFORMES: LEPISTOSTEIDAE): A DELIBERATE AQUARIUM TRADE INTRODUCTION INCIDENCE IN THE SHATT AL-ARAB RIVER, BASRAH, IRAQ**

Falah MUTLAK¹, Laith JAWAD², and Abbas AL-FAISAL¹

¹ Marine Science Centre, University of Basrah, Basrah, Iraq

² Flat Bush, Manukau, Auckland, New Zealand

Mutlak F., Jawad L., Al-Faisal A. 2017. *Atractosteus spatula* (Actinopterygii: Lepisosteiformes: Lepisosteidae): A deliberate aquarium trade introduction incidence in the Shatt al-Arab River, Basrah, Iraq. *Acta Ichthyol. Piscat.* 47 (2): 205–207.

Abstract. The first record of a single specimen of alligator gar, *Atractosteus spatula* (Lacepède, 1803), native to North America, is reported from the inland waters of Iraq. The specimen was caught from southern reaches of the Shatt al-Arab River at Om-al-Rasas Island, Basrah, Iraq during an ichthyologic survey in September 2016 using gillnets with mesh size of 40 mm. The capture of a sub-adult specimen, 900 mm in total length, marks the first record of a member of the family Lepisosteidae in the freshwater system of Iraq. Aquarium trade pathway is the suspected factor for the presence of this species in the Iraqi freshwater area.

Keywords: biodiversity, new locality, new occurrence, aquarium trade, invasion

The introduction of exotic species, is widely considered to be the second most important cause of species extinction after habitat destruction (Simberloff 2003). Acosta and Gupta (2005) suggested that exotic fishes have been deliberately introduced primarily to enhance fisheries and aquaculture production. Fishes may be also introduced to support recreational fishery activities in developed countries. Other main reasons include pest control, research activities, and aquarium trade (Cohen et al. 2007, Chang et al. 2009).

Basrah Province, Iraq has many aquatic resources including coastal waters, marshes, and rivers, which are potentially a suitable environment for exotic fish species capable of surviving the local environmental conditions. With the absence of regulations on importing fish species and deficiencies in the knowledge on the biology of the fish species being imported by the aquarium trade in Iraq, such events give the chance for unfavourable fish species to be included to the freshwater system of Iraq.

The only exotic fish introduction via aquarium trade reported from the marshes of central Iraq is *Pangasianodon hypophthalmus* (Sauvage, 1878) (see Khamees et al. 2013), *Mollienesia latipinna* (Lesueur, 1821) (see Al-Faisal and Mutlak 2014). The same species were also mentioned by Al-Lamy et al. (2012) and Mohamed et al. (2008).

A single specimen of alligator gar, *Atractosteus spatula* (Lacepède, 1803), native to North America, was captured from southern reaches of the Shatt al-

Arab River at Om-al-Rasas Island, Basrah, Iraq (30°30'04.56"N; 47°51'57.18"E) during a fish survey in the Shatt al-Arab River basin. The aim of this study was to report the presence of *A. spatula* in the inland waters of Iraq and discuss the possible impact of the introduction of this exotic species on the native fish fauna of the area. The fish specimen was captured in September 2016 using gillnets with mesh size of 40 mm. The specimen was identified according to Page and Burr (2011) and deposited in the fish collection at the Marine Science Centre, University of Basrah, Iraq (MSC 00132). The nomenclature follows Eschmeyer et al. (2017). A series of morphometric measurements were made with a ruler to the nearest 1 mm following the methods of Wiley (1976) (Table 1). Gonads were examined using laboratory microscope.

Atractosteus spatula can be distinguished from the other members of the family Lepisosteidae in having a large, round body, with an elongated, but relatively broad snout (Figs. 1 and 2). The fish examined was an immature male, with gonad weight of 85.9 g.

The total length of our specimen (900 mm) is less than the common total length given for this species (2000 mm) (Wiley 1978).

The alligator gar can be harmful if it becomes a successful invasive species and it is capable of creating a sustainable population. Alligator gar feeds mainly on fishes and it is a successful predator in its native environment. The eggs of alligator gar are poisonous,

* Correspondence: Laith A. Jawad, 4 Tintum Place, Flat Bush, Manukau, Auckland, New Zealand, phone +64 9 2715165, e-mail: (LAJ) laith_jawad@hotmail.com, (FM) falahmutlak@yahoo.com, (AAF) abbasjsm71@yahoo.com.

but their consumption by people is unlikely (Boschung and Mayden 2004, Goddard 2009). Additionally, this fish could damage fishing nets due to the shape of its head and the sharp teeth.

There has been no published information about established population alligator gar outside its native America. The nearest locality where alligator gar has been reported is the Caspian Sea coasts of Turkmenistan (Salinkov 2010). Considering the substantial distance the possible migration of the fish to Iraq seems to be unlikely. It is difficult to tell precisely how this specimen was introduced and how it penetrated into the Shatt al-Arab River of Iraq, but it is known that these fish are kept in aquaria all over the world. Recently, the aquarium

trade in Iraq has become very popular with the easing of regulations importing fishes into Iraq, and increase of aquarium shops (personal observation). Alligator gars are sold in these shops, mainly in large cities like Baghdad and Basrah. Aquarists frequently get rid of unwanted fish by releasing them into open water bodies. It is also likely that alligator gars are released, because of their substantial annual length increment of 250–300 mm (Boschung and Mayden 2004, Roberts and Harrel 2009).

The presently reported finding represents the third Iraqi case of the exotic species introduction, blamed on aquarium trade.

REFERENCES

- Acosta B.O., Gupta M.V.** 2005. The status of introduced carp species in Asia. Pp. 121–1230. *In*: Penman D.J., Gupta M.V. (eds.) *Carp genetic resources for aquaculture in Asia*, World Fish Centre, Penang, Malaysia.
- Al-Faisal A.J., Mutlak F.M.** 2014. First record of the Nile tilapia *Oreochromis niloticus* (Linnaeus, 1758), from the Shatt al-Arab River, southern Iraq. *Mesopotamian Journal of Marine Science* **29** (1): 45–50.
- Al-Lamy J.H., Resen A.K., Taher M.M., Muhammed S.J.** 2012. Specialization and diet overlapped between two alien species (*Carassius auratus* and *Tilapia zillii*) and native species (*Barbus luteus*) in East Hammar Marsh, southern Iraq. *Journal of King Abdulaziz University Marine Sciences* **23** (2): 25–37. DOI: [10.4197/Mar.23-2.2](https://doi.org/10.4197/Mar.23-2.2)
- Boschung H.T., Mayden R.L.** 2004. *Fishes of Alabama*. Smithsonian Institution Press, Washington DC, USA.
- Chang A.L., Grossman J.D., Spezio T.S., Weiskel H.W., Blum J.C., Burt J.W., Muir A.A., Piovio-Scott J., Veblen K.E., Grosholz E.D.** 2009. Tackling aquatic invasions: Risks and opportunities for the aquarium fish industry. *Biological Invasions* **11** (4): 773–785. DOI: [10.1007/s10530-008-9292-4](https://doi.org/10.1007/s10530-008-9292-4)
- Cohen J., Mirotchnick N., Leung B.** 2007. Thousands introduced annually: the aquarium pathway for non-indigenous plants to the St Lawrence Seaway. *Frontiers in the Ecology and the Environment* **5** (10): 528–532. DOI: [10.1890/060137](https://doi.org/10.1890/060137)
- Eschmeyer W.N., Fricke R., van der Laan R.** (eds.) 2017. *Catalog of fishes: Genera, species, references*. California Academy of Sciences, San Francisco, USA. <http://researcharchive.calacademy.org/research/ichthyology/catalog/fishcatmain.asp> [Accessed on 4 March 2017.]
- Goddard N.** 2009. Alligator gar. *Ichthyology at the Florida Museum of Natural History*. <https://www.floridamuseum.ufl.edu/fish/discover/species-profiles/alligator-gar>
- Khamees N.R., Ali A.H., Abed J.M., Adday T.K.** 2013. First record of striped catfish *Pangasianodon hypophthalmus* (Sauvage, 1878) (Pisces: Pangasiidae) from inland waters of Iraq. *Basrah Journal of Agricultural Sciences* **26** (Special issue 1): 178–183.
- Mohamed A.-R.M., Hussain N.A., Al-Noor S.S., Mutlak F.M., Sudani I.M., Mojer A.M., Toman**

Table 1

Biometric characters of *Atractosteus spatula* collected from Shatt al-Arab River, Basrah, Iraq

Character	Absolute value	Relative value
Weight	5252 g	
Total length (TL)	900 mm	
Standard length (SL)	780 mm	86.7% in TL
Body depth	130 mm	16.7% in SL
Head length (HL)	245 mm	31.4% in SL
Head depth	74.8 mm	30.5% in HL
Head width	82.1 mm	33.5% in HL
Snout length	128 mm	52.2% in HL
Eye diameter	19.9 mm	8.1% in HL
Interorbital distance	66.6 mm	27.2% in HL
Dorsal fin length	46.8 mm	6.1% in SL
Pectoral fin length	75.3 mm	9.7% in SL
Pelvic fin length	78.8 mm	10.1% in SL
Anal fin length	47.3 mm	6.1% in SL
Predorsal fin length	670 mm	85.9% in SL
Postdorsal fin length	735 mm	94.2% in SL
Dorsal fin rays	7	
Anal fin rays	7	
Pectoral fin rays	14	
Pelvic fin rays	7	
Lateral line scales	60	



Fig. 1. *Atractosteus spatula*, 900 mm TL, collected from Shatt al-Arab River, Basrah, Iraq, mouth view



Fig. 2. *Atractosteus spatula*, 900 mm TL collected from Shatt al-Arab River, Basrah, Iraq; lateral view

- A.J., Abdad M.A.** 2008 Fish assemblage of restored Al-Hawizeh marsh, southern Iraq. *Ecology and Hydrobiology* **8** (2–4): 375–384.
DOI: [10.2478/v10104-009-0029-5](https://doi.org/10.2478/v10104-009-0029-5)
- Page L.M., Burr B.M.** 1991. *Freshwater fishes: A field guide to freshwater fishes of North America north of Mexico*. Peterson Field Guide Series. Houghton Mifflin, Boston MA, USA.
- Roberts D., Harrel S.** 2006. *Atractosteus spatula* gemfish. Animal Diversity Web, University of Michigan, Museum of Zoology. http://animaldiversity.org/accounts/Atractosteus_spatula
- Salnikov V.B.** 2010. First finding of gar *Atractosteus* sp. (Actinopterygii, Lepisosteiformes, Lepisosteidae) in the Caspian Sea near the coast of Turkmenistan. *Russian Journal of Biological Invasion* **1** (1): 17–20.
DOI: [10.1134/S2075111710010042](https://doi.org/10.1134/S2075111710010042)
- Simberloff D.** 2003. Confronting introduced species: A form of xenophobia? *Biological Invasions* **5** (3): 179–192.
DOI: [10.1023/A:1026164419010](https://doi.org/10.1023/A:1026164419010)
- Wiley E.O.** 1976. The phylogeny and biogeography of fossil and recent gars (Actinopterygii: Lepisosteidae). University of Kansas Museum of Natural History, Miscellaneous Publication **64**: 1–111.
- Wiley E.O.** 1978. Lepisosteidae. In: Fischer W. (ed.) *FAO species identification sheets for fishery purposes. Western Central Atlantic (Fishing Area 31)*. Vol. 3. FAO, Rome.

Received: 19 December 2016

Accepted: 17 March 2017

Published electronically: 30 June 2017