















PREPRINT

Author-formatted, not peer-reviewed document posted on 09/06/2025

DOI: <https://doi.org/10.3897/arhpreprints.e161414>

The first national assessment of established non-native species in Türkiye

 Ali Serhan Tarkan,  Sercan Yapıcı,  Irmak Kurtul,  Dagmara Błońska,  Lorenzo Vilizzi, 
Hayrunisa Bas,  Shahid Farooq,  Ceray Aldemir, Tuğba Ucma Uysal,  Daniela Giannetto, 
Gokcen Bilge, Mülkibar Çiftçioğlu,  Elnaz Najafi-Majd, Lale Aktay-Sözüer, Cüneyt Kaya,  Esra
Bayçelebi, Ilker Aydın,  Phillip Haubrock,  Elizabeta Briski, Ismael Soto

1 The first national assessment of established non-native species 2 in Türkiye

3
4 Ali Serhan Tarkan^{1,2*}, Sercan Yapıcı², Irmak Kurtul^{3,4}, Dagmara Błońska^{1,4}, Lorenzo Vilizzi¹,
5 Hayrunisa Baş Sermenli⁵, Shahid Farooq⁶, Ceray Aldemir⁷, Tuğba Uçma Uysal⁸, Daniela
6 Giannetto⁵, Gökçen Bilge², Mülkibar Çiftçioğlu², Elnaz Najafi-Majd^{9,10}, Lale Aktay-Sözüer^{11,12},
7 Cüneyt Kaya^{13,14}, Esra Bayçelebi¹³, İlker Aydın^{15*}, Phillip J. Haubrock^{4,16,17}, Elizabeta Briski¹⁸,
8 Ismael Soto¹⁶

9
10 ¹University of Lodz, Faculty of Biology and Environmental Protection, Department of Ecology and Vertebrate
11 Zoology, Lodz, Poland

12 ²Department of Basic Sciences, Faculty of Fisheries, Muğla Sıtkı Koçman University, Muğla, Türkiye

13 ³Marine and Inland Waters Sciences and Technology Department, Faculty of Fisheries, Ege University, İzmir, Türkiye

14 ⁴Department of Life and Environmental Sciences, Faculty of Science and Technology, Bournemouth University, Poole,
15 Dorset, UK

16 ⁵Department of Biology, Faculty of Sciences, Muğla Sıtkı Koçman University, Muğla, Türkiye

17 ⁶Department of Plant Protection, Faculty of Agriculture, Harran University, Şanlıurfa, Türkiye

18 ⁷Department of Public Administration, Faculty of Economics and Administrative Sciences, Muğla Sıtkı Koçman
19 University, Muğla, Türkiye

20 ⁸Department of International Trade and Finance, Faculty of Economics and Administrative Sciences, Muğla Sıtkı
21 Koçman University, Muğla, Türkiye

22 ⁹Department of Biology, Faculty of Science, Ege University, İzmir, Türkiye

23 ¹⁰Department of Microbiology and Immunology, University of Rochester Medical Center, MRBX 2-11001.
24 Rochester, NY, 14642, USA

25 ¹¹Department of Biology, Institute of Science, Akdeniz University, Antalya, Türkiye

26 ¹²Raptor Research and Conservation Working Group of Türkiye, Antalya, Türkiye

27 ¹³Faculty of Fisheries, Recep Tayyip Erdogan University, 53100, Rize, Türkiye

28 ¹⁴Department of Ecology, Faculty of Science, Charles University, Prague 2, Czech Republic

29 ¹⁵Ege University Faculty of Fisheries Department of Fishing Technology, 35100, İzmir, Türkiye

30 ¹⁶University of South Bohemia in České Budějovice, Faculty of Fisheries and Protection of Waters, South Bohemian
31 Research Centre of Aquaculture and Biodiversity of Hydrocenoses, Zátíší 728/II, 389 25 Vodňany, Czech Republic

32 ¹⁷Center for Applied Mathematics and Bioinformatics, Department of Mathematics and Natural Sciences, Gulf
33 University for Science and Technology, Hawally, Kuwait

34 ¹⁸GEOMAR Helmholtz-Zentrum für Ozeanforschung Kiel, Wischhofstraße 1-3, 24148 Kiel, Germany
35

36 *Corresponding Author(s) e-mail: serhantarkan@gmail.com, ilker.aydin@ege.edu.tr

37

38

39

40

41

42

43

44

45

46

47 **ABSTRACT**

48

49 The ecological and socio-economic impacts of non-native species underscore the importance of
50 maintaining accurate national lists to support biosecurity and management strategies. Türkiye's
51 most recent national list highlights ongoing challenges in maintaining up-to-date records. To
52 advance Türkiye's national assessment of non-native species, we present the first comprehensive
53 list of established non-native species in the country, documenting 268 species from 157 families
54 across diverse taxonomic groups. Marine species ($n = 199$) dominated the list, with notable
55 contributions from Chordata, Mollusca, and Arthropoda, followed by Terrestrial (41), Freshwater
56 (24), Terrestrial/Freshwater (3) and Freshwater/Marine (1). Our spatial analysis revealed
57 considerable regional variation, with urban and coastal areas exhibiting higher rates of species
58 establishment likely facilitated by human-mediated pathways. We also identify Türkiye's role as a
59 crucial hub for the spread of non-native species across the country's borders. Further, we show
60 that 17% of non-native species have known ecological or economic impacts, emphasizing the
61 need for targeted impact assessments, monitoring, and proactive management. By providing a
62 detailed and up-to-date inventory of established non-native species and their impacts, this study
63 offers a critical foundation for strengthening biosecurity policies, guiding targeted monitoring
64 efforts, and fostering coordinated management responses at local, national, and international
65 scales to safeguard Türkiye's biodiversity.

66 **Key-words:** Biodiversity, Impacts, Species management, Human-mediated pathways, Trade
67 networks, Regional patterns

68

69

70 Introduction

71
72 Non-native species pose significant ecological threats to biodiversity, and their socio-economic
73 impacts represent not only a burden on national economies but are also detrimental to human
74 well-being (Cucherousset and Olden 2011, Diagne et al. 2020). Despite the population-level
75 differences of invasion impacts (Haubrock et al. 2024), identifying and assessing non-native
76 species at the national level is essential for their effective management in view of legislation
77 measures. National lists of non-native species therefore serve multiple purposes: (i) they compile
78 species seen as relevant in terms of their potential or existing threats, (ii) they help prioritize
79 species for management actions, and (iii) they inform biosecurity measures by identifying those
80 species that may threaten neighbouring countries through trade and travel (Hulme 2009).
81 However, these lists, whenever available, are frequently inconsistent, reflecting the subjective
82 biases of assessors and differences in research fields (e.g. aquatic vs terrestrial, animals vs
83 plants), the status of non-native species (i.e. whether introduced, established, or invasive: *sensu*
84 Soto et al., 2024), and the criteria for determining a species as 'invasive' (i.e. spread vs impact:
85 (Cuthbert et al. 2022, Vilizzi et al. 2022a, Haubrock et al. 2024, Soto et al. 2024b). This can result
86 in misclassification of non-native species, with some being listed as established without clear
87 evidence, and some established being flagged as invasive without proper justification—although
88 we acknowledge that drawing a line to distinguish the different statuses of non-native species can
89 be obscure.

90
91 Several national lists of non-native species have emerged over the past two decades, providing
92 essential tools for guiding non-native species management, with most European Union countries
93 having compiled such lists even before formal regulations (Genovesi and Monaco 2013, Cerri et
94 al. 2022). Türkiye, however, exemplifies the challenges of maintaining accurate and updated
95 national lists. Given the country's significant economic position (ranked 18th globally: IMF, 2025),
96 coupled with its rich biodiversity (Şekercioğlu et al. 2011) and strategic location at the intersection
97 of the Middle East and Europe (Kahraman et al. 2012), such a national list is urgently needed to
98 mitigate the undesirable impacts of biological invasions. This is because non-native species in
99 Türkiye have been shown to pose substantial ecological and economic threats, adversely
100 affecting various sectors including agriculture, fisheries, tourism, and human health (Tarkan et al.
101 2012, 2024b). Their economic impact, while significant, is often underrepresented in national
102 financial assessments, despite the costs they impose on food production, biodiversity, and public
103 infrastructure (Shackelford et al. 2013, Bălăcenoiu et al. 2023). Incorporating these costs into

104 environmental policy and fiscal planning—such as through ‘green budgeting’—could improve
105 resource allocation and support more proactive management responses. Additionally, given
106 Türkiye's trade routes and the high volume of international travel, the risk of new non-native
107 species introductions in the country remains high (Kahraman and Altun 2016).

108
109 Although the first and only national list of non-native species in Türkiye dates back to 2020 and
110 includes 872 species (Uludag et al. 2020), questions remain about its accuracy and utility, as the
111 list includes all recorded occurrences of non-native species regardless of their establishment
112 status rather than focusing solely on those non-native species with self-sustaining populations,
113 i.e. established non-native species (hereafter, ‘established species’: see Soto et al., 2024). It is
114 therefore crucial to differentiate between those non-native species that are established, hence
115 likely to pose an immediate threat, and those that might be introduced, thereby presenting a
116 potential danger.

117
118 Updating existing national lists to guide future monitoring is essential for developing effective
119 early-warning systems and intervention strategies. In this respect, Türkiye exemplifies the
120 challenges of maintaining accurate and updated national lists, and the need for such updates is
121 critical in addressing the ecological and economic threats posed by non-native species. To fill this
122 gap, this study aims to provide a detailed overview of the established species in Türkiye by
123 compiling a new and up-to-date database. By analyzing the composition of these species, we
124 seek to understand the patterns and implications of their establishment in Türkiye. We
125 hypothesize that (1) the composition of established species will be dominated by terrestrial plants
126 and marine invertebrates, reflecting patterns observed in comparable regions, and (2) regions
127 within Türkiye with higher human activity, particularly those with dense populations, significant
128 trade and transportation infrastructure, and intensive land use, will exhibit higher rates of non-
129 native species establishment. In contrast, we hypothesize that (3) regions with less human activity
130 or more remote locations, such as certain inland or mountainous areas, will show lower rates of
131 invasion due to reduced opportunities for species introduction and less disturbance to native
132 ecosystems. We also anticipate that (4) the known ecological and economic impacts of these
133 established species are highly biased towards well-known invaders and taxonomic groups.
134 Finally, this study aims to assess known invasion impacts and, whenever possible, the level of
135 risk posed by the established species in Türkiye so as to provide targeted recommendations for
136 management and control.

137

138
 139
 140
 141
 142
 143
 144
 145
 146
 147
 148
 149
 150
 151
 152
 153
 154
 155
 156
 157
 158

Material and methods

Data compilation

To compile a comprehensive and up-to-date database of established species in Türkiye, we used the recent global database of occurrences by Briski et al. (2024). This database gathers information from several sources, including the online database Standardising and Integrating Alien Species (SInAS: Seebens, 2021) and non-native species lists. We used the SInAS database as the base file for our study (see also Briski et al., 2024; Haubrock et al., 2025). We restricted usage to the spatial scale of species assigned only as CASUAL and ABSENT in the columns ‘degreeOfEstablishment’ and ‘occurrenceStatus’, respectively, which were removed due to their undetermined non-native establishment status (Groom et al. 2019). We emphasize that our dataset also contains taxonomically accepted subspecies as per the Global Biodiversity Information Facility (GBIF: <https://www.gbif.org/>). When a species was not found in GBIF, a manual search was done to verify its scientific name by consulting databases such as the World Register of Marine Species (WoRMS: <https://www.marinespecies.org/>), coupled with a general internet search in September 2024 (Briski et al. 2024). This process yielded a total of 1,092 established species in Türkiye. After a comprehensive review of each major species group by Turkish experts—based on the most up-to-date literature to filter out native, cryptogenic, and non-established species—a total of 268 species (≈ 25%) remained classified as established in Türkiye.

Table 1. Breakdown of non-native organism habitat classifications (after Haubrock et al., 2025).

Habitat	Description
Freshwater	Species classified as freshwater inhabit environments such as rivers, lakes, and streams, with salinity < 0.5 g kg ⁻¹ .
Marine	Species classified as marine occupy environments such as oceans, seas, coastal areas, and estuaries with salinity ≥ 0.5 g kg ⁻¹ .
Terrestrial	Species classified as terrestrial are found on land, including habitats such as forests, grasslands, deserts, and urban areas.
Semi-aquatic	Species associated with both terrestrial and aquatic habitats (e.g. freshwater-terrestrial), depend on both aquatic and terrestrial habitats to complete their life cycle or daily activities.

159
 160

Habitat and native range

161 For every species reported in the dataset, we identified the habitat it predominantly occupies (i.e.
162 terrestrial, freshwater, marine, or a mixture thereof) by consulting the
163 Step2_StandardTerms_GRIIS file (Seebens 2021) with habitat data originating from the Global
164 Register of Introduced and Invasive Species (GRIIS: (Pagad et al. 2018). We then assigned one
165 or more habitats (i.e. a combination of them, such as freshwater-marine) to each species. Where
166 habitat information was not available, we conducted an additional manual search of WoRMS and
167 a general internet search between July and September 2024. Of note, brackish habitats were
168 defined as marine based on the Venice System (1958)

169 We extracted the native range of each established species in Türkiye in two ways: (i) by employing
170 the web-scraping technique on several online databases, including Plants of the World Online
171 (<https://powo.science.kew.org/>) and GBIF, and (ii) by consulting targeted publications. We
172 identified the native range at the biogeographic realm (i.e. Nearctic, Neotropical, Palearctic,
173 Afrotropical, Indo-Malayan, Australasian, and Oceanian). Due to the low number of species from
174 the Oceanian realm, these were combined with the Australasian. Species associated with two
175 realms of equal weight were assigned to each realm (i.e. the taxon was distributed in both realms).
176 Species distribution across two or more realms (i.e. cosmopolitan) and species with unclear native
177 range were not considered in the analyses.

178 *Distribution of established species in Türkiye*

179 To unveil the spatial pattern of the established species, we extracted GBIF occurrences for each
180 of them in Türkiye. We restricted downloads to only those occurrences categorized as present,
181 with coordinates without geospatial issues and removal of fossil records. The download was done
182 by using the *occ_download* function from the *rgbif* R package (Chamberlain et al. 2017,
183 GBIF.org 2025). To identify the number of established species in each administrative area, we
184 overlapped the established non-native richness within the boundaries of each administrative area,
185 including marine areas. We downloaded the terrestrial administrative areas from the *geodata* R
186 package (Hijmans et al. 2024), whereas the marine areas (Exclusive Economic Zones) were
187 extracted from Marine Regions (<https://marineregions.org/>: Flanders Marine Institute, 2023).

188

189 *Compilation of impact and risk*

190 For each established species, we extracted all available information on its reported impacts (i.e.
191 environmental, economic, or social) at two levels: (i) reported in Türkiye and (ii) globally from

192 GRIIS. In the case of established species with a reported impact in Türkiye, we checked the
 193 Centre for Agriculture and Biosciences International (CABI: <https://www.cabi.org/>) compendium
 194 and the InvaCost database (v4.1: Diagne et al., 2020) for information on the type of impact. If the
 195 data on the impact were not available in the CABI or InvaCost database, we used Google Scholar
 196 to search for the information using the following search string: “species scientific name” and
 197 “impact” or “invasive” in order to broaden the search, especially for less-studied species.
 198 Comparison between regional (Türkiye) and global scale allowed us to identify potential gaps in
 199 impact reporting.

200 For those established non-native species in the list for which published information was available
 201 on the level of risk posed in risk assessment areas across Türkiye (including the country as a
 202 whole), we included screening data from studies using the Aquatic Species Invasiveness
 203 Screening Kit (AS-ISK: (Copp et al. 2016, Vilizzi et al. 2025). The AS-ISK (available in its v2.4.1
 204 at <https://tinyurl.com/ISK-toolkits>) is a taxon-generic, multilingual decision-support tool that
 205 complies with minimum standards for assessing non-native species (Copp et al. 2016). It consists
 206 of 55 questions, 49 of which are for Basic Risk Assessment (BRA) and six are for Climate Change
 207 Assessment (CCA). For each screened species, two scores are obtained: BRA, under current
 208 climate conditions, and BRA+CCA, under predicted climate (change) conditions. Scores < 1
 209 suggest a ‘low risk’ of the species being or becoming invasive in the risk assessment area, scores
 210 ≥ 1 indicate a ‘medium risk’ or a ‘high risk’, which are distinguished using a calibrated threshold
 211 (Vilizzi et al. 2022a). For those species classified as high risk, we also distinguished a ‘very high
 212 risk’ category using an *ad hoc* threshold ≥ 45 based on the modular approach described by Britton
 213 et al. (2011). Identification of very high-risk species helps to prioritize allocation of resources for
 214 comprehensive (follow-up) risk assessment (Vilizzi et al. 2022a). In total, there were six AS-ISK
 215 applications of relevance to this study (i.e. Bilge et al., 2019; Killi et al., 2020; Tarkan et al., 2021,
 216 2017; Yapici, 2021), noting that the screenings by Tarkan et al. (2017) updated the previous ones
 217 by Tarkan et al. (2014) using a predecessor toolkit to the AS-ISK.


218

219 **Results**

220

221 In the final dataset listing the 268 established species in Türkiye, we identified a total of 156
 222 families from 24 classes and 10 phyla (Fig. 1; Supplementary Material 1). The families with the
 223 highest number of established species were Formicidae ($n = 14$), followed by Pyramidellidae (9),
 224 Penaeidae (8), and Tetraodontidae (7), with each of the remaining families containing fewer than
 225 five species. In terms of classes, the most species rich was Actinopterygii (98), followed by

226 Gastropoda (48) and Malacostraca (26). Regarding phyla, Chordata was the most represented
227 (109 species), followed by Mollusca (67) and Arthropoda (52). The majority of species were strictly
228 marine (199), followed by terrestrial (41) and by those classified as strictly freshwater (24). Only
229 a few species were classified as freshwater-terrestrial (3) and freshwater-marine (1) (Fig. 2).




230

231 **Figure 1.** Breakdown of Phyla and Classes with the highest number of established species in Türkiye.

232

233




234

235 **Figure 2.** Barplot of established species in Türkiye broken down by habitat.


236

237 From GBIF, we gathered a total of 16,160 records of the identified 268 established species in
 238 Türkiye. Most of the records came from three bird species, namely the rose-ringed parakeet
 239 *Psittacula krameri* ($n = 8,037$; 49.1%), the Alexandrine parakeet *Psittacula eupatria* (3,424;
 240 21.2%), and the common myna *Acridotheres tristis* (1,300; 8.0%). These species together
 241 constituted a total of 12,761 occurrences (78.3%). We observed a latitude gradient, with most of
 242 the species concentrated in the western part of the country and decreasing eastwards. The areas
 243 or zones with the highest numbers of reported species occurrences were Istanbul ($n = 184$),
 244 followed by Antalya (167) and Muğla (139) (Fig. 3).



245
 246 **Figure 3.** Number of established species in each administrative area of Türkiye based on GBIF records.
 247 Black areas represent administrative areas without non-native species based on GBIF. Marine areas
 248 represent the Exclusive Economic Zone.

249
 250 We identified the native range for 256 established species in Türkiye (84% of the total). The
 251 biogeographic realm where most established species originate was Indo-Malayan, with more than
 252 two times the number of species in Australasia (Fig. 4). Other realms with a lower number of
 253 established species were Palearctic, Afrotropical, Nearctic, and Neotropical.



254

255 **Figure 4.** Biogeographic origin of established non-native species in Türkiye (highlighted in black).


256 Numbers represent how many of the established species are native to their regions of origin.

257

258 The majority of established species in Türkiye had no reported impacts ($n = 222$; 83%), although
 259 this number decreased to 157 (59%) at the global scale. Conversely, the 46 species with reported
 260 impacts in Türkiye (17%) increased to 111 (41%) at the global scale. The proportion of established
 261 species with documented impacts was generally lower than those without reported impacts,
 262 highlighting a considerable gap in impact assessment and reporting across taxa (Fig. 5). Among
 263 species with known impacts, ecological and environmental were the most frequently cited ($n =$
 264 89; 33%), followed by economic (43; 16%) and socio-cultural (28; 11%). In monetary terms, 87
 265 species (33%) recorded some form of economic costs in Türkiye, increasing to 133 (50%) with
 266 monetary terms at the global scale. Regarding ecological impact mechanisms (Fig. 6),
 267 competition was the most commonly reported (64), followed by predation (32) and hybridisation
 268 (13). Other recorded mechanisms included habitat alteration, disease transmission, and, among
 269 others, resource use, though these were reported less frequently.


270

271



272
273
274
275
276
277


Figure 5. Doughnut plot indicating established species with (red) and without (grey) known from Türkiye (a) and (b) anywhere else



278
279
280
281
282

Figure 6. (a) impacts and (b) impact mechanisms by species.

283 The six AS-ISK applications of relevance to this study provided screenings for 62 established
 284 species, representing $\approx 23\%$ of the 268 species in total from the list (Supplementary Table 1). The
 285 risk assessment areas consisted of the inland waters of Türkiye (Tarkan et al. 2017, 2021, Aksu
 286 et al. 2024) and of the marine waters of the south-western coasts of Anatolia (Bilge et al. 2019)
 287 and of the Mediterranean Sea (Killi et al. 2020, Tarkan et al. 2021, Yapici 2021). Of the 62
 288 established species in total: (i) under current climate conditions (BRA scores), 3 (4.8%) posed a
 289 very high risk of invasiveness, 25 (40.3%) a high risk, 33 (53.2%) a medium risk, and 1 (1,6%) a
 290 low risk; (ii) under predicted climate (change) conditions (BRA+CCA scores), 13 (21.0%) posed
 291 a very high risk of invasiveness, 25 (40.3%) a high risk, 23 (37.1%) a medium risk, and 1 (1,6%)
 292 a low risk (Fig. 7). Overall, the number of very high-risk species substantially increased under
 293 predictions of climate change conditions for Türkiye and the risk assessment areas therein.



296 **Figure 7.** Risk outcome scores for 62 established species in Türkiye screened with the Aquatic Species
297 Invasiveness Screening Kit. (a) Basic Risk Assessment (BRA) scores for the first 31 species; (b) BRA +
298 Climate Change Assessment (BRA+CCA) scores for the first 31 species; (c) BRA scores for the last 31
299 species; (d) BRA+CCA scores for the last 31 species. Red bars = very high-risk (VH) species (score ≥ 45);
300 black bars = high-risk (H) species; grey bars = medium-risk (M) species; white bars = low-risk (L) species.
301 Species-specific risk levels are based on the thresholds provided in Table S1.

302
303

304 **Discussion**

305 Türkiye has a long-standing tradition of research in the natural sciences, yet national efforts to
306 address the threats posed by biological invasions remain limited (Şekercioğlu et al. 2011). This
307 study has offered the first comprehensive national assessment of established species in Türkiye,
308 identifying 268 species that present substantial threats to the country's biodiversity and social
309 well-being. Our compilation points to clear biases, with the majority of the identified species
310 occupying marine environments in contrast to the less-represented freshwater species (Butchart
311 et al. 2010), with the same being true for certain taxonomic groups such as insects and fishes.
312 Even more notable is the lack of information on these species' impacts, raising concerns about
313 biological invasions being a silently growing threat, especially in the case of understudied
314 taxonomic groups and regions. In relation to our hypotheses, we did not find support for (1) the
315 expectation that the composition of established species would be dominated by terrestrial plants
316 but with marine fishes being more prevalent, and (2) confirmed that regions with greater human
317 activity tend to host more established species, while (3) less accessible or remote areas
318 experience lower invasion rates. However, (4) the anticipated regional variation in impact severity
319 was only partially supported, suggesting that impact may not scale directly with establishment
320 density.

321 Compared to the 268 established species we identified here, the national list of non-native species
322 published by Uludag et al. (2020) reported 872 species—a number much higher than in the
323 present study. The main reason for this difference is that the latter list included all non-native
324 species ever recorded in Türkiye, without confirming whether they are actually established or still
325 present today. While our focus was on species that are confirmed to be established, the 2020 list
326 still represents an important resource. It provides a broader picture of potential non-native
327 species, which could help identify future threats. In that sense, even though the two lists serve

328 different purposes, both are valuable for understanding and managing biological invasions in the
329 country.

330 *Taxonomic composition and biases*

331 The taxonomic profile of Türkiye's established species is shaped by a mix of ecological, economic,
332 and research-related factors. Our results show that Chordata, Mollusca, and Arthropoda are the
333 most represented Phyla, with a clear dominance of marine species. This is not surprising given
334 Türkiye's extensive coastline along three major seas, namely Black, Aegean, and Mediterranean,
335 which provide a broad range of habitats. These areas are also under intense pressure from
336 maritime traffic, aquaculture, and fisheries, all of which are known pathways for species
337 introductions (Çinar et al. 2005). Such conditions help explain the high number of marine species
338 in the national list.

339 The comparatively low numbers of terrestrial and freshwater species reflect a combination of
340 factors. In the case of terrestrial plants, the lower figures may be somewhat misleading. Türkiye
341 has a long history of plant introductions, especially for agriculture and horticulture, but not all
342 introduced species establish successfully or are monitored closely (Uludag et al. 2017). For
343 instance, the relatively small number of Tracheophyta in the dataset likely reflects gaps in
344 research and reporting rather than true absence. Therefore, there is still a need for more detailed
345 studies to fully understand the scale of non-native plant establishments.

346 Freshwater species were even less represented, which may be explained by both ecological and
347 research-related reasons. Many freshwater habitats in Türkiye, particularly those in regions of
348 high human impact, have been degraded by pollution, habitat destruction, and hydrological
349 changes (Pinarlik 2023). Such conditions can limit the success of new invasions (Rahel 2002). At
350 the same time, freshwater environments in more remote areas might be under-surveyed,
351 reflecting a lack of funding, institutional support, and research infrastructure in these regions.
352 Therefore, both real ecological constraints and a lack of data have likely contributed to the
353 observed pattern.

354 Regarding the role played by economic and industrial activities, marine ecosystems are heavily
355 exploited for fisheries and aquaculture, which increases the risk of species introductions and
356 establishment (Nunes et al. 2014). On the other hand, certain terrestrial sectors, such as
357 agriculture, may have stricter biosecurity protocols in place, which could explain the lower
358 representation of some groups (Pimentel et al. 2005). However, these patterns may also reflect

359 broader priorities, with more research and monitoring focused on economically important areas,
360 leaving others underexplored.

361 Overall, these taxonomic and environmental biases underline the need for a more balanced
362 research agenda. Without comprehensive assessments across all groups and environments,
363 gaps in knowledge will persist, increasing the risk of unforeseen ecological and economic
364 consequences. It is essential to strengthen research efforts, particularly in underrepresented
365 areas like freshwater systems and terrestrial flora, to ensure that management and policy
366 responses are based on a complete understanding of the country's non-native species landscape.

367 *Spatial heterogeneity and regional variability*

368 The distribution of non-native species across Türkiye displays marked spatial differences, with
369 the highest numbers recorded in urban and especially coastal regions. This pattern largely reflects
370 the role of human-mediated pathways, as areas with dense human populations, strong
371 economies, and well-developed trade networks—such as Istanbul, Antalya, and Muğla—offer
372 multiple routes for species introductions and establishment (Aikio et al. 2010). Key vectors in
373 these regions include maritime transport, aquaculture, the ornamental plant and pet trades, and
374 ballast water discharge, all of which are intensified by high volumes of trade and human activity.
375 In contrast, areas with lower population densities and limited economic activity, such as more
376 remote inland or mountainous regions, generally exhibit fewer invasions, which aligns with
377 patterns reported globally (Essl et al. 2015).

378 The underlying reasons for these differences are multifaceted. Urban and coastal centres are
379 hubs of international and domestic trade, dense infrastructure, and economic growth, which all
380 contribute to the introduction and establishment of non-native species (Hulme 2009). Increasingly,
381 global trade, tourism, and financial networks are recognized as key human-mediated pathways
382 for biological invasions, particularly in metropolitan settings (Seebens et al. 2021). In Türkiye,
383 rapid urbanisation and economic development have intensified these pathways, highlighting the
384 critical role of customs controls, ballast water regulations, and cross-border enforcement in
385 limiting new introductions. However, lax biosecurity enforcement, profit-driven trade in exotic
386 species, and limited institutional capacity continue to undermine prevention efforts (Tarkan et al.
387 2024b). Financial incentives often favour short-term economic gains over long-term ecological
388 health, placing additional pressure on regulatory systems that may lack political will or resources
389 to implement stringent controls (Early et al. 2016). In many cases, regions with fewer resources

390 and less institutional capacity also lack the funding, personnel, and infrastructure needed to
391 monitor and manage non-native species effectively (Pyšek et al. 2020). Local communities, often
392 burdened by more immediate socio-economic concerns, may have little opportunity to engage in
393 biodiversity monitoring or early detection programs. This disparity likely contributes not only to
394 lower actual introduction rates in remote regions but also to significant gaps in detection and
395 reporting (Pyšek et al. 2010).

396 Importantly, areas where non-native species have already established may also serve as stepping
397 stones for further spread within Türkiye. Once a species gains a foothold in a region, intentional
398 or accidental translocations—often linked to activities such as recreational angling, aquaculture,
399 and efforts to enhance local fisheries—can facilitate its movement to new areas (Tarkan et al.
400 2024a). These secondary introductions are often motivated by economic considerations or by the
401 aim of improving local food security, but they can have far-reaching ecological consequences if
402 left unchecked (Gaygusuz et al. 2015).

403 These observations highlight the need for management strategies that are tailored to the specific
404 risks and realities of different regions. Urban and coastal areas with high introduction pressure
405 require intensive monitoring and stricter biosecurity protocols, while less-affected regions would
406 benefit from investments in local capacity, training, and public awareness. Furthermore, policy
407 frameworks must account not only for initial introductions but also for the increasing risk of
408 secondary spread within the country, which remains a significant challenge for invasive species
409 management. Coordinated national efforts will be essential to mitigate the long-term impacts on
410 Türkiye's biodiversity and ecosystems (Early et al. 2016).

411 *Current and unknown impacts*

412 A clear pattern emerging from our analysis is the high proportion of non-native species for which
413 no impacts have been reported. This likely reflects two main issues: first, many species have not
414 yet been subject to formal impact assessment; second, some species—particularly those with
415 restricted distributions or narrow ecological niches—may indeed have limited or negligible
416 impacts. Initial impacts of biological invasions might seem benign or even positive, and while
417 being catastrophic in the long term (Soto et al. 2024a), this might create a lag in the reporting of
418 impacts. However, our results also indicate biases in which environments and taxonomic groups
419 are reported to have impacts. For instance, while marine species make up the largest portion of
420 the dataset, the ratio of reported impacts to non-impacts is around 19%, whereas freshwater

421 species show a notably higher ratio of 38%. Across taxonomic groups, Chordata has 14%,
422 Arthropoda 23%, and Mollusca 18%, with Annelida (20%) and Tracheophyta (28%) also showing
423 some level of recorded impacts. Many other groups have no reported impacts at all, underscoring
424 a significant gap in current knowledge.

425 This pattern likely stems from a focus on species already recognized as problematic—such as
426 large-bodied fishes or economically relevant invertebrates—while less conspicuous or poorly
427 studied taxa remain overlooked. Such gaps are concerning because they can result in delayed
428 responses to emerging threats, ultimately increasing biodiversity loss and the economic costs of
429 management. For Türkiye, with its rich biodiversity and reliance on ecosystems that are
430 vulnerable to invasive species (Tarkan et al. 2024b), the absence of robust impact data represents
431 a serious blind spot. Closing this gap requires systematic and ongoing assessments, particularly
432 targeting under-studied groups and habitats. Expanding monitoring and research to reflect the full
433 spectrum of Türkiye’s ecosystems will be crucial for strengthening biosecurity and ensuring timely,
434 evidence-based management actions.

435 *Invasiveness risk*

436 Having found more than 50% of established species in Türkiye—among those for which screening
437 data are available—to pose a high to very high risk of invasiveness, which increases to over 60%
438 under predicted future climate (change) conditions, is concerning. As a representative subset of
439 the whole list of 268 established species in Türkiye, a similar outcome may be expected should
440 the remaining species be eventually screened for their invasiveness. The limited number of
441 species screened so far is likely due to the unavailability of data for terrestrial species, as the
442 relevant tools for animals (Vilizzi et al. 2022b) and plants (Vilizzi et al. 2024)—have only recently
443 been developed (see also Vilizzi et al., 2025). Given the availability of our up-to-date list of
444 established species for Türkiye, future studies should therefore consider screening not only the
445 other aquatic species but also the terrestrial animals and plants in the list. This would provide a
446 comprehensive database of information based on which local legislation for the monitoring,
447 control, containment, and, if at all feasible, eradication of any established species in Türkiye could
448 be gauged accordingly. Moreover, given that nearly two-thirds of the screened species fall into
449 the high to very high risk categories under future climate conditions indicates the need to establish
450 a prioritization framework for their management. This could help allocate limited conservation and
451 management resources more effectively by targeting species and regions at the greatest risk.
452 Such a framework should consider both the risk scores (BRA and BRA+CCA) and ecological or

453 economic impact potential, especially for species expected to escalate in risk due to climate
454 change. Prioritization is also critical to guide early detection and rapid response measures.

455 **Conclusion**

456 Türkiye's strategic location at the intersection of Europe, Asia, and the Middle East makes it a
457 central hub for the spread of non-native species. Our findings show that Türkiye shares a
458 significant number of non-native species with neighbouring European countries (Güneralp et al.
459 2013, Uludag et al. 2020, Henry et al. 2023). Our findings also suggest that it not only receives
460 species from other regions but may also act as a source of invasions to nearby countries (Galil
461 2008). The observed spatial heterogeneity and regional variability highlight the need for tailored
462 management strategies that consider both local and international contexts. Moving forward,
463 addressing the identified biases, expanding research into underrepresented groups, and
464 strengthening biosecurity measures will be critical to preventing future invasions. A coordinated
465 approach, built on the insights from this comprehensive list, is essential for protecting Türkiye's
466 biodiversity and ensuring the sustainability of its ecosystems.

467 **Acknowledgements**

468 PJH was supported by the Marie Skłodowska-Curie Postdoctoral Fellowship HORIZON-MSCA-
469 2022-PF-01 (Project DIRECT; Grant No. 101203662) within the European Union's Horizon 2022
470 research and innovation programme. DB was supported by Marie Curie Individual Fellowship
471 HORIZON-MSCA-2022-PF-01 (project 101105250 - PROSPER) within the European Union's
472 Horizon 2022 research and innovation programme, funded by UKRI.

473

474 **References**

475

- 476 Aikio S, Duncan RP, Hulme PE (2010) Lag-phases in alien plant invasions: separating the facts
477 from the artefacts. *Oikos* 119: 370–378. <https://doi.org/10.1111/j.1600-0706.2009.17963.x>
- 478 Aksu S, Mercan D, Arslan N, Emiroğlu Ö, Haubrock PJ, Soto I, Tarkan AS (2024) Determining
479 environmental drivers of global mud snail invasions using climate and hydroclimate models.
480 *Hydrobiologia* 851: 3991–4006. <https://doi.org/10.1007/s10750-024-05554-x>
- 481 Bălăcenoiu F, Japelj A, Bernardinelli I, Castagneyrol B, Csóka G, Glavendekić M, Hoch G,
482 Hrasovec B, Krajter Ostoić S, Paulin M, Williams D, Witters J, de Groot M (2023) Ascertaining
483 the knowledge of the general public and stakeholders in the forestry sector to invasive alien
484 species—A Pan-European study. *Land* 12: 642. <https://doi.org/10.3390/land12030642>
- 485 Bilge G, Filiz H, Yapıcı S, Tarkan A, Vilizzi L (2019) A risk screening study on the potential
486 invasiveness of Lessepsian fishes in the South-Eastern coasts of Anatolia. *Acta*
487 *Ichthyologica et Piscatoria* 49: 23–31. <https://doi.org/10.3750/AIEP/02422>

- 488 Briski E, Kotronaki SG, Cuthbert RN, Bortolus A, Campbell ML, Dick JTA, Fofonoff P, Galil BS,
489 Hewitt CL, Lockwood JL, Maclsaac HJ, Ricciardi A, Ruiz G, Schwindt E, Sommer U, Zhan
490 A, Carlton JT (2024) Does non-native diversity mirror Earth's biodiversity? *Global Ecology*
491 *and Biogeography* 33: 48–62. <https://doi.org/10.1111/geb.13781>
- 492 Britton JR, Copp GH, Brazier M, Davies GD (2011) A modular assessment tool for managing
493 introduced fishes according to risks of species and their populations, and impacts of
494 management actions. *Biological Invasions* 13: 2847–2860. [https://doi.org/10.1007/s10530-](https://doi.org/10.1007/s10530-011-9967-0)
495 [011-9967-0](https://doi.org/10.1007/s10530-011-9967-0)
- 496 Butchart SHM, Walpole M, Collen B, van Strien A, Scharlemann JPW, Almond REA, Baillie JEM,
497 Bomhard B, Brown C, Bruno J, Carpenter KE, Carr GM, Chanson J, Chenery AM, Csirke J,
498 Davidson NC, Dentener F, Foster M, Galli A, Galloway JN, Genovesi P, Gregory RD,
499 Hockings M, Kapos V, Lamarque J-F, Leverington F, Loh J, McGeoch MA, McRae L,
500 Minasyan A, Morcillo MH, Oldfield TEE, Pauly D, Quader S, Revenga C, Sauer JR, Skolnik
501 B, Spear D, Stanwell-Smith D, Stuart SN, Symes A, Tierney M, Tyrrell TD, Vié J-C, Watson
502 R (2010) Global biodiversity: Indicators of recent declines. *Science* 328: 1164–1168.
503 <https://doi.org/10.1126/science.1187512>
- 504 Cerri J, Carnevali L, Monaco A, Genovesi P, Bertolino S (2022) Blacklists do not necessarily make
505 people curious about invasive alien species. A case study with Bayesian structural time
506 series and Wikipedia searches about invasive mammals in Italy. *NeoBiota* 71: 113–128.
- 507 Chamberlain S, Barve V, Mcglinn D, Oldoni D, Desmet P, Geffert L, Ram K (2017) *rgbif*: Interface
508 to the Global Biodiversity Information Facility API. R package version 0.9.9. [https://CRAN.R-](https://CRAN.R-project.org/package=rgbif)
509 [project.org/package=rgbif](https://CRAN.R-project.org/package=rgbif).
- 510 Çinar ME, Bilecenoglu M, Öztürk B, Katagan T, Aysel V (2005) Alien species on the coasts of
511 Turkey. *Mediterranean Marine Science* 6: 119–146. <https://doi.org/10.12681/mms.187>
- 512 Copp G, Vilizzi L, Tidbury H, Stebbing P, Tarkan AS, Miossec L, Gouletquer P (2016)
513 Development of a generic decision-support tool for identifying potentially invasive aquatic
514 taxa: AS-ISK. *Management of Biological Invasions* 7: 343–350.
515 <https://doi.org/10.3391/mbi.2016.7.4.04>
- 516 Cucherousset J, Olden JD (2011) Ecological impacts of non-native freshwater fishes. *Fisheries*
517 36: 215–230. <https://doi.org/10.1080/03632415.2011.574578>
- 518 Cuthbert RN, Diagne C, Haubrock PJ, Turbelin AJ, Courchamp F (2022) Are the “100 of the
519 world's worst” invasive species also the costliest? *Biological Invasions* 24: 1895–1904.
520 <https://doi.org/10.1007/s10530-021-02568-7>
- 521 Diagne C, Leroy B, Gozlan RE, Vaissière A-C, Assailly C, Nuninger L, Roiz D, Jourdain F, Jarić
522 I, Courchamp F (2020) InvaCost, a public database of the economic costs of biological
523 invasions worldwide. *Scientific Data* 7: 277. <https://doi.org/10.1038/s41597-020-00586-z>
- 524 Early R, Bradley BA, Dukes JS, Lawler JJ, Olden JD, Blumenthal DM, Gonzalez P, Grosholz ED,
525 Ibañez I, Miller LP, Sorte CJB, Tatem AJ (2016) Global threats from invasive alien species
526 in the twenty-first century and national response capacities. *Nature Communications* 7:
527 12485. <https://doi.org/10.1038/ncomms12485>
- 528 Essl F, Dullinger S, Rabitsch W, Hulme PE, Pyšek P, Wilson JRU, Richardson DM (2015)
529 Historical legacies accumulate to shape future biodiversity in an era of rapid global change.
530 *Diversity and Distributions* 21: 534–547. <https://doi.org/10.1111/ddi.12312>

- 531 Galil BS (2008) Alien species in the Mediterranean Sea—which, when, where, why?
532 Hydrobiologia 606: 105–116. <https://doi.org/10.1007/s10750-008-9342-z>
- 533 Gaygusuz O, Tarkan AS, Aydin H, Dorak Z, Top N, Karakus U, Vilizzi L (2015) Stocking of
534 common carp (*Cyprinus carpio*) into some newly-established reservoirs of North-West
535 Anatolia may enhance the spread of non-native fish. Turkish Journal of Fisheries and Aquatic
536 Sciences 15. https://doi.org/10.4194/1303-2712-v15_4_06
- 537 GBIF.org (2025) GBIF Occurrence Download. <https://doi.org/10.15468/dl.b8fdm>.
- 538 Genovesi P, Monaco A (2013) Guidelines for Addressing Invasive Species in Protected Areas.
539 In: Foxcroft LC et al. (Ed.), Plant Invasions in Protected Areas: Patterns, Problems and
540 Challenges. Springer, 487–506.
- 541 Groom Q, Desmet P, Reyserhove L, Adriaens T, Oldoni D, Vanderhoeven S, Baskauf SJ,
542 Chapman A, McGeoch M, Walls R, Wieczorek J, Wilson J, Zermoglio PF, Simpson A (2019)
543 Improving Darwin Core for research and management of alien species. Biodiversity
544 Information Science and Standards 3: e38084. <https://doi.org/10.3897/biss.3.38084>
- 545 Güneralp B, Tezer A, Albayrak İ (2013) Local assessment of İstanbul: Biodiversity and ecosystem
546 services. In: Urbanization, Biodiversity and Ecosystem Services: Challenges and
547 Opportunities. Springer Netherlands, Dordrecht, 291–311. https://doi.org/10.1007/978-94-007-7088-1_16
- 549 Haubrock PJ, Soto I, Cano-Barbacil C, Theissing K, Rios-Henriquez C, Parker B, Weithoff G,
550 Briski E (2025) Germany’s established non-native species: a comprehensive breakdown.
551 Environmental Sciences Europe 37: 56. <https://doi.org/10.1186/s12302-025-01094-w>
- 552 Haubrock PJ, Soto I, Ahmed DA, Ansari AR, Tarkan AS, Kurtul I, Macêdo RL, Lázaro-Lobo A,
553 Toutain M, Parker B, Błońska D, Guareschi S, Cano-Barbacil C, Dominguez Almela V,
554 Andreou D, Moyano J, Akalın S, Kaya C, Bayçelebi E, Yoğurtçuoğlu B, Briski E, Aksu S,
555 Emiroğlu Ö, Mammola S, De Santis V, Kourantidou M, Pincheira-Donoso D, Britton JR,
556 Kouba A, Dolan EJ, Kirichenko NI, García-Berthou E, Renault D, Fernandez RD, Yapıcı S,
557 Giannetto D, Nuñez MA, Hudgins EJ, Pergl J, Milardi M, Musolin DL, Cuthbert RN (2024)
558 Biological invasions are a population-level rather than a species-level phenomenon. Global
559 Change Biology 30. <https://doi.org/10.1111/gcb.17312>
- 560 Henry M, Leung B, Cuthbert RN, Bodey TW, Ahmed DA, Angulo E, Balzani P, Briski E,
561 Courchamp F, Hulme PE, Kouba A, Kourantidou M, Liu C, Macêdo RL, Oficialdegui FJ,
562 Renault D, Soto I, Tarkan AS, Turbelin AJ, Bradshaw CJA, Haubrock PJ (2023) Unveiling
563 the hidden economic toll of biological invasions in the European Union. Environmental
564 Sciences Europe 35: 43. <https://doi.org/10.1186/s12302-023-00750-3>
- 565 Hijmans RJ, Barbosa M, Ghosh A, Mandel A, Hijmans MRJ (2024) Package ‘geodata’. CRAN.
566 Geodata.
- 567 Hulme PE (2009) Trade, transport and trouble: managing invasive species pathways in an era of
568 globalization. Journal of Applied Ecology 46: 10–18. <https://doi.org/10.1111/j.1365-2664.2008.01600.x>
- 570 Kahraman A, Altun Y (2016) The role of international trade and travel in the introduction of non-
571 native species in Turkey. Turkish Journal of Environmental and Biological Science 5: 203–
572 211.

- 573 Kahraman A, Onder M, Ceyhan E (2012) The importance of bioconservation and biodiversity in
574 Turkey. *International Journal of Bioscience, Biochemistry and Bioinformatics* 2: 95–99.
575 <https://doi.org/10.7763/IJBBB.2012.V2.79>
- 576 Killi N, Tarkan AS, Kozic S, Copp GH, Davison PI, Vilizzi L (2020) Risk screening of the potential
577 invasiveness of non-native jellyfishes in the Mediterranean Sea. *Marine Pollution Bulletin*
578 150: 110728. <https://doi.org/10.1016/j.marpolbul.2019.110728>
- 579 Nunes AL, Katsanevakis S, Zenetos A, Cardoso AC (2014) Gateways to alien invasions in the
580 European seas. *Aquatic Invasions* 9: 133–144. <https://doi.org/10.3391/ai.2014.9.2.02>
- 581 Pagad S, Genovesi P, Carnevali L, Schigel D, McGeoch MA (2018) Introducing the Global
582 Register of Introduced and Invasive Species. *Scientific Data* 5: 170202.
583 <https://doi.org/10.1038/sdata.2017.202>
- 584 Pimentel D, Zuniga R, Morrison D (2005) Update on the environmental and economic costs
585 associated with alien-invasive species in the United States. *Ecological Economics* 52: 273–
586 288. <https://doi.org/10.1016/j.ecolecon.2004.10.002>
- 587 Pinarlik M (2023) The effects of climate change on water resources in Turkey. In: Sogut MZ,
588 Karakoc TH, Secgin O, Dalkiran A (Eds), *Proceedings of the 2022 International Symposium*
589 *on Energy Management and Sustainability*. Springer, Cham, 595–603.
590 https://doi.org/10.1007/978-3-031-30171-1_64
- 591 Pyšek P, Hulme PE, Simberloff D, Bacher S, Blackburn TM, Carlton JT, Dawson W, Essl F,
592 Foxcroft LC, Genovesi P, Jeschke JM, Kühn I, Liebhold AM, Mandrak NE, Meyerson LA,
593 Pauchard A, Pergl J, Roy HE, Seebens H, van Kleunen M, Vilà M, Wingfield MJ, Richardson
594 DM (2020) Scientists' warning on invasive alien species. *Biological Reviews* 95: 1511–1534.
595 <https://doi.org/10.1111/brv.12627>
- 596 Pyšek P, Jarošík V, Hulme PE, Kühn I, Wild J, Arianoutsou M, Bacher S, Chiron F, Didžiulis V,
597 Essl F, Genovesi P, Gherardi F, Hejda M, Kark S, Lambdon PW, Desprez-Loustau M-L,
598 Nentwig W, Pergl J, Pobljšaj K, Rabitsch W, Roques A, Roy DB, Shirley S, Solarz W, Vilà
599 M, Winter M (2010) Disentangling the role of environmental and human pressures on
600 biological invasions across Europe. *Proceedings of the National Academy of Sciences* 107:
601 12157–12162. <https://doi.org/10.1073/pnas.1002314107>
- 602 Rahel FJ (2002) Homogenization of Freshwater Faunas. *Annual Review of Ecology and*
603 *Systematics* 33: 291–315. <https://doi.org/10.1146/annurev.ecolsys.33.010802.150429>
- 604 Seebens H (2021) SInAS database of alien species occurrences (2.4.1) [Data set]. Zenodo.
- 605 Seebens H, Bacher S, Blackburn TM, Capinha C, Dawson W, Dullinger S, Genovesi P, Hulme
606 PE, van Kleunen M, Kühn I, Jeschke JM, Lenzner B, Liebhold AM, Pattison Z, Pergl J, Pyšek
607 P, Winter M, Essl F (2021) Projecting the continental accumulation of alien species through
608 to 2050. *Global Change Biology* 27: 970–982. <https://doi.org/10.1111/gcb.15333>
- 609 Şekercioğlu ÇH, Anderson S, Akçay E, Bilgin R, Can ÖE, Semiz G, Tavşanoğlu Ç, Yokeş MB,
610 Soyumert A, İpekdal K, Sağlam İK, Yücel M, Nüzhet Dalfes H (2011) Turkey's globally
611 important biodiversity in crisis. *Biological Conservation* 144: 2752–2769.
612 <https://doi.org/10.1016/j.biocon.2011.06.025>
- 613 Shackelford N, Renton M, Perring MP, Hobbs RJ (2013) Modeling disturbance-based native
614 invasive species control and its implications for management. *Ecological Applications* 23:
615 1331–1344. <https://doi.org/10.1890/12-1881.1>

- 616 Soto I, Macêdo RL, Carneiro L, Briski E, Kouba A, Cuthbert RN, Haubrock PJ (2024a) Divergent
617 temporal responses of native macroinvertebrate communities to biological invasions. *Global*
618 *Change Biology* 30. <https://doi.org/10.1111/gcb.17521>
- 619 Soto I, Balzani P, Carneiro L, Cuthbert RN, Macêdo R, Serhan Tarkan A, Ahmed DA, Bang A,
620 Bacela-Spychalska K, Bailey SA, Baudry T, Ballesteros-Mejia L, Bortolus A, Briski E, Britton
621 JR, Buřič M, Camacho-Cervantes M, Cano-Barbacil C, Copilaş-Ciocianu D, Coughlan NE,
622 Courtois P, Csabai Z, Dalu T, De Santis V, Dickey JWE, Dimarco RD, Falk-Andersson J,
623 Fernandez RD, Florencio M, Franco ACS, García-Berthou E, Giannetto D, Glavendekic MM,
624 Grabowski M, Heringer G, Herrera I, Huang W, Kamelamela KL, Kirichenko NI, Kouba A,
625 Kourantidou M, Kurtul I, Laufer G, Lipták B, Liu C, López-López E, Lozano V, Mammola S,
626 Marchini A, Meshkova V, Milardi M, Musolin DL, Nuñez MA, Oficialdegui FJ, Patoka J,
627 Pattison Z, Pincheira-Donoso D, Piria M, Probert AF, Rasmussen JJ, Renault D, Ribeiro F,
628 Rilov G, Robinson TB, Sanchez AE, Schwindt E, South J, Stoett P, Verreycken H, Vilizzi L,
629 Wang Y, Watari Y, Wehi PM, Weiperth A, Wiberg-Larsen P, Yapıcı S, Yoğurtçuoğlu B, Zenni
630 RD, Galil BS, Dick JTA, Russell JC, Ricciardi A, Simberloff D, Bradshaw CJA, Haubrock PJ
631 (2024b) Taming the terminological tempest in invasion science. *Biological Reviews* 99:
632 1357–1390. <https://doi.org/10.1111/brv.13071>
- 633 Tarkan AS, Güler Ekmekçi F, Vilizzi L, Copp GH (2014) Risk screening of non-native freshwater
634 fishes at the frontier between Asia and Europe: first application in Turkey of the fish
635 invasiveness screening kit. *Journal of Applied Ichthyology* 30: 392–398.
636 <https://doi.org/10.1111/jai.12389>
- 637 Tarkan AS, Gaygusuz Ö, Gürsoy Gaygusuz Ç, Saç G, Copp GH (2012) Circumstantial evidence
638 of gibel carp, *Carassius gibelio*, reproductive competition exerted on native fish species in a
639 mesotrophic reservoir. *Fisheries Management and Ecology* 19: 167–177.
640 <https://doi.org/10.1111/j.1365-2400.2011.00839.x>
- 641 Tarkan AS, Kurtul I, Błońska D, Britton JR, Haubrock PJ (2024a) Resolving the issues of
642 translocated species in freshwater invasions. *NeoBiota* 93: 177–186.
643 <https://doi.org/10.3897/neobiota.93.122837>
- 644 Tarkan AS, Vilizzi L, Top N, Ekmekçi FG, Stebbing PD, Copp GH (2017) Identification of
645 potentially invasive freshwater fishes, including translocated species, in Turkey using the
646 Aquatic Species Invasiveness Screening Kit (AS-ISK). *International Review of Hydrobiology*
647 102: 47–56. <https://doi.org/10.1002/iroh.201601877>
- 648 Tarkan AS, Tricarico E, Vilizzi L, Bılge G, Ekmekçi FG, Filiz H, Giannetto D, İlhan A, Kılıç N,
649 Kirankaya ŞG, Koutsikos N, Kozic S, Kurtul I, Lazzaro L, Marchini A, Occhipinti-Ambrogi A,
650 Perdikaris C, Piria M, Pompei L, Sari H, Smeti E, Stasolla G, Top N, Tsiamis K, Vardakas L,
651 Yapıcı S, Yoğurtçuoğlu B, Copp GH (2021) Risk of invasiveness of non-native aquatic
652 species in the eastern Mediterranean region under current and projected climate conditions.
653 *The European Zoological Journal* 88: 1130–1143.
654 <https://doi.org/10.1080/24750263.2021.1980624>
- 655 Tarkan AS, Bayçelebi E, Giannetto D, Özden ED, Yazlık A, Emiroğlu Ö, Aksu S, Uludağ A, Aksoy
656 N, Baytaşoğlu H, Kaya C, Mutlu T, Kirankaya ŞG, Ergüden D, Per E, Üremiş İ, Candan O,
657 Kekillioğlu A, Yoğurtçuoğlu B, Ekmekçi FG, Başak E, Özkan H, Kurtul I, Innal D, Killi N,
658 Yapıcı S, Ayaz D, Çiçek K, Mol O, Çınar E, Yeğen V, Angulo E, Cuthbert RN, Soto I,
659 Courchamp F, Haubrock PJ (2024b) Economic costs of non-native species in Türkiye: A first

- 660 national synthesis. *Journal of Environmental Management* 358: 120779.
661 <https://doi.org/10.1016/j.jenvman.2024.120779>
- 662 Uludag A, Aksoy N, Uremis I, Yazlık A, Arslan ZF, Demir Ozden E, Kirankaya SG, Ekmekçi FG,
663 Wong LJ, Pagad S (2020) Global register of introduced and invasive species - Turkey.
664 Invasive Species Specialist Group ISSG Version 1.2.
- 665 Uludag A, Aksoy N, Yazlık A, Arslan ZF, Yazmış E, Uremis I, Cossu TA, Groom Q, Pergl J, Pyšek
666 P, Brundu G (2017) Alien flora of Turkey: checklist, taxonomic composition and ecological
667 attributes. *NeoBiota* 35: 61–85. <https://doi.org/10.3897/neobiota.35.12460>
- 668 Venice System (1958) Symposium on the classification of brackish waters. *Oikos* 9: 311–312.
- 669 Vilizzi L, Hill JE, Piria M, Copp GH (2022a) A protocol for screening potentially invasive non-native
670 species using Weed Risk Assessment-type decision-support tools. *Science of The Total*
671 *Environment* 832: 154966. <https://doi.org/10.1016/j.scitotenv.2022.154966>
- 672 Vilizzi L, Piria M, Herczeg G, Almeida D, Al-Wazzan Z, Bakiu R, Boggero A, Chaichana R,
673 Dashinov D, De Zoysa M, Gilles A, Gouletquer P, Interesova E, Kopecký O, Koutsikos N,
674 Koyama A, Kristan P, Li S, Lukas J, Moghaddas S, Monteiro J, Mumladze L, Oh C, Olsson
675 K, Pavia R, Perdikaris C, Pickholtz R, Preda C, Ristovska M, Švolíková K, Števo B, Ta K,
676 Uzunova E, Vardakas L, Verreycken H, Wei H, Yoğurtçuoğlu B, Giannetto D, Pietraszewski
677 D (2025) Questionnaire improvements in second-generation, multilingual decision support
678 tools for invasion risk screening of non-native taxa. *Management of Biological Invasions* 16:
679 33–44. <https://doi.org/10.3391/mbi.2025.16.1.03>
- 680 Vilizzi L, Piria M, Pietraszewski D, Kopecký O, Špelić I, Radočaj T, Šprem N, Ta KAT, Tarkan AS,
681 Weiperth A, Yoğurtçuoğlu B, Candan O, Herczeg G, Killi N, Lemić D, Szajbert B, Almeida D,
682 Al-Wazzan Z, Atique U, Bakiu R, Chaichana R, Dashinov D, Ferincz Á, Flieller G, Gilles Jr
683 AS, Gouletquer P, Interesova E, Iqbal S, Koyama A, Kristan P, Li S, Lukas J, Moghaddas
684 SD, Monteiro JG, Mumladze L, Olsson KH, Paganelli D, Perdikaris C, Pickholtz R, Preda C,
685 Ristovska M, Švolíková KS, Števo B, Uzunova E, Vardakas L, Verreycken H, Wei H, Zięba
686 G (2022b) Development and application of a multilingual electronic decision-support tool for
687 risk screening non-native terrestrial animals under current and future climate conditions.
688 *NeoBiota* 76: 211–236. <https://doi.org/10.3897/neobiota.76.84268>
- 689 Vilizzi L, Piria M, Pietraszewski D, Giannetto D, Flory SL, Herczeg G, Sermenli HB, Britvec M,
690 Jukoniene I, Petrulaitis L, Vitasović-Kosić I, Almeida D, Al-Wazzan Z, Bakiu R, Boggero A,
691 Chaichana R, Dashinov D, De Zoysa M, Gilles AS, Gouletquer P, Interesova E, Kopecký O,
692 Koutsikos N, Koyama A, Kristan P, Li S, Lukas J, Moghaddas SD, Monteiro JG, Mumladze
693 L, Oh C, Olsson KH, Pavia RT, Perdikaris C, Pickholtz R, Preda C, Ristovska M, Švolíková
694 KS, Števo B, Ta KAT, Uzunova E, Vardakas L, Verreycken H, Wei H, Yoğurtçuoğlu B,
695 Ferincz Á, Kirkendall LR, Marszał L, Paganelli D, Stojchevska C, Tarkan AS, Yazlık A (2024)
696 Development and application of a second-generation multilingual tool for invasion risk
697 screening of non-native terrestrial plants. *Science of The Total Environment* 917: 170475.
698 <https://doi.org/10.1016/j.scitotenv.2024.170475>
- 699 Yapici S (2021) A risk screening of potential invasiveness of alien and neontative marine fishes in
700 the Mediterranean Sea: Implications for sustainable management. *Sustainability* 13: 13765.
701 <https://doi.org/10.3390/su132413765>
- 702