

PREPRINT

Author-formatted, not peer-reviewed document posted on 01/09/2022

DOI: <https://doi.org/10.3897/arphapreprints.e94201>

***Phaeoisaria laianensis* (Pleurotheciales, Pleurotheciaceae), a new species from freshwater habitats in China**

Yu Liu, Gui-Ping Xu, Xin-Yi Yan, Min-Hui Chen, Yang Gao,  Dian-Ming Hu, Zhi-Jun Zhai

Disclaimer on biological nomenclature and use of preprints

The preprints are preliminary versions of works accessible electronically in advance of publication of the final version. They are not issued for purposes of botanical, mycological or zoological nomenclature and **are not effectively/validly published in the meaning of the Codes**. Therefore, nomenclatural novelties (new names) or other nomenclatural acts (designations of type, choices of priority between names, choices between orthographic variants, or choices of gender of names) **should NOT be posted in preprints**. The following provisions in the Codes of Nomenclature define their status:

International Code of Nomenclature for algae, fungi, and plants (ICNafp)

Article 30.2: "An electronic publication is not effectively published if there is evidence within or associated with the publication that its content is merely preliminary and was, or is to be, replaced by content that the publisher considers final, in which case only the version with that final content is effectively published." In order to be validly published, a nomenclatural novelty must be effectively published (Art. 32.1(a)); in order to take effect, other nomenclatural acts must be effectively published (Art. 7.10, 11.5, 53.5, 61.3, and 62.3).

International Code of Zoological Nomenclature (ICZN)

Article: 21.8.3: "Some works are accessible online in preliminary versions before the publication date of the final version. Such advance electronic access does not advance the date of publication of a work, as preliminary versions are not published (Article 9.9)".

Phaeoisaria laianensis (Pleurotheciales, Pleurotheciaceae), a new species from freshwater habitats in China

Yu Liu[‡], Gui-Ping Xu[‡], Xin-Yi Yan[‡], Min-Hui Chen^{‡,§}, Yang Gao^{‡,§}, Dian-Ming Hu^{‡,§}, Zhi-Jun Zhai^{‡,§}

[‡] College of Bioscience and Bioengineering, Jiangxi Agricultural University, Nanchang, China

[§] Bioengineering and Technological Research Centre for Edible and Medicinal Fungi, Jiangxi Agricultural University, Nanchang, China

Corresponding author: Dian-Ming Hu (hudianming1@163.com), Zhi-Jun Zhai (zhjzh002@163.com)

Abstract

Background

Freshwater fungi play an indispensable role in the ecosystem and have great research value. Based on morphological and phylogenetic analyses of a concatenated dataset of ITS, LSU and SSU sequences, a new species, *Phaeoisaria laianensis*, was introduced as a freshwater hyphomycete from Anhui Province, China.

New information

Phaeoisaria laianensis was morphologically described as erect, rigid, dark brown to black, velvety synnemata which has macronematous, septate, branched, brown to dark brown, parallelly adressed conidiophores with polyblastic, integrated, terminal, hyaline to pale brown, smooth, denticulate, sympodial conidiogenous cells and ellipsoidal to obovoid, rounded at the apex, obtuse and tapering towards base, septate, guttulate conidia. Based on molecular and morphological characteristics, it is confirmed to be a new species, as all illustrations and descriptions had been provided.

Keywords

Ascomycota *Phaeoisaria*, morphology, phylogenetic anaysis, taxonomy

Introduction

Phaeoisaria (Pleurotheciales) was established by Höhnelt (1909) to accommodate *Phaeoisaria bambusae* as the type species, a hyphomycetous taxon isolated from a bamboo substrate. This species was later treated as a synonym of *P. clematidis* (Fuckel) S. Hughes (Deighton 1974). This genus is characterized by indeterminate synnemata with parallelly addressed conidiophores with numerous sympodially extending denticulate conidiogenous cells and aseptate or septate ellipsoidal, obovoidal, fusiform-cylindrical to falcate, hyaline conidia (Höhnelt 1909, Réblová et al. 2016, Hyde et al. 2018, Luo et al. 2018, Boonmee et al. 2021). Nevertheless, some species in this genus were not observed to have indeterminate synnemata, such as *P. curvata* (de Hoog and Papendorf 1976), *P. glauca* (de Hoog and Papendorf 1976), *P. loranthacearum* (Crous et al. 2015), *P. fasciculata* (Réblová et al. 2016), *P. annesophieae* (Crous et al. 2017) and *P. dalbergiae* (Crous et al. 2021).

In the past decades, an increasing number of new species was assigned to *Phaeoisaria* by having distinguishing characters (Crous et al. 2017, Hyde et al. 2018, Hyde et al. 2019, Luo et al. 2019, Boonmee et al. 2021, Crous et al. 2021). Until now, 28 species has been accepted in the genus *Phaeoisaria*. These species are relatively common and have a worldwide distribution, while only four of which have been recorded in China. Moreover, there are presently only 15 species having the molecular data in *Phaeoisaria*. In this study, we depicted a new species, *Phaeoisaria laianensis*, from submerged wood in Anhui Province of China, with both morphological examination and molecular phylogenetic analysis.

Materials and methods

Samples collection, specimen examination and isolation

Submerged rotting wood samples were gathered from Laian County, Anhui Province, China and were brought back to the laboratory to be incubated in plastic boxes at room temperature. Fungi on the host surface were observed with a Nikon SMZ-1270 microscope (Nikon Corporation, Japan) and morphologically photographed with a Nikon ECLIPSE Ni-U compound microscope (Nikon Corporation, Japan), which was outfitted with a Nikon DS-Fi3 camera. The structure of fungi was determined by PhotoRuler 1.1.3.0 (The Genus Inocybe, Hyogo, Japan) and figures were processed by Adobe Photoshop 2020 (Adobe Systems, USA). According to the method of Li et al. (2021), single spore isolation and pure culture were carried out. Fungal specimens were deposited in the Fungus Herbarium, Jiangxi Agricultural University, Nanchang, China.

DNA extraction, PCR amplification and sequencing

By using the improved CTAB method (Doyle and Doyle 1987), fungal total genomic DNA was extracted from fresh mycelium. Four gene regions (ITS, LSU and SSU), were respectively amplified by polymerase chain reaction (PCR) using the primers of ITS1/ITS4 (White et al. 1990), LROR/LR7 (Hopple and Vilgalys 1999) and NS1/NS4 (White et al. 1990), with 25 µl of the final volume including 9.5 µl ddH₂O, 12.5 µl 2 × Taq PCR MasterMix (Qingke, Changsha, China), 1 µl of DNA template and 1 µl of each primer (10 µM). Then amplifications were conducted under the PCR conditions described by Zhai et al. 2022. The PCR products were purified and the sequencing reactions were commercially conducted with the corresponding forward and reverse primers by QingKe Biotechnology Co. (Changsha, China). All sequences were edited with SeqMan v. 7.1.0 (DNASTAR, Inc, Madison, WI) and were deposited in the NCBI GenBank database.

Phylogenetic analyses

The sequences of 69 strains were retrieved from recent articles (Luo et al. 2018, Hyde et al. 2019, Boonmee et al. 2021) and downloaded from GenBank (Table 1). Each matrix of ITS, LSU and SSU was aligned using the online service of MAFFT v.7 (<http://mafft.cbrc.jp/alignment/server/large.html>, Katoh et al. 2019) and then the sequences of three regions were concatenated by PhyloSuite v.1.2.2 (Zhang et al. 2020). By using RAxML v.7.2.6 (Stamatakis and Alachiotis 2010), Maximum Likelihood (ML) analysis was performed, which used a GTRGAMMA substitution model with 1000 bootstrap replicates. Markov chain Monte Carlo (MCMC) methods in MrBayes was used to estimate the posterior probabilities (PP) (Zhaxybayeva and Gogarten 2002), and it was set as 4 chains (2 hot chains and 2 cold chains) running 2,000,000 generations synchronously, resulting 40002 trees totally. Based on the initial 25% of sampled data were cut off as burn-in, PhyloSuite v.1.2.2 (Zhang et al. 2020) was used to infer Bayesian inference phylogeny under JC+I+G+F model of the concatenation of ITS, LSU and SSU. With visualised by FigTree v. 1.4.4 (Rambaut 2018), phylogenetic trees were edited and beautified using Adobe Illustrator 2020 (Adobe Systems Inc., the United States). The aligned matrices and trees were submitted to TreeBASE (<http://purl.org/phylo/treebase/phyloids/study/TB2:S29465>).

Taxon treatment

***Phaeoisaria laianensis* Y. Liu, G.P. Xu, X.Y. Yan, D.M. Hu & Z.J. Zhai, sp. nov.**

- MycoBank [844773](https://www.ncbi.nlm.nih.gov/genbank/844773)

Material

Holotype:

- a. scientificName: *Phaeoisaria laianensis*; acceptedNameUsage: *Phaeoisaria laianensis* Y. Liu, D.M. Hu & Z.J. Zhai; kingdom: Fungi; phylum: Ascomycota; class: Sordariomycetes;

order: Pleurotheciales; family: Pleurotheciaceae; taxonRank: species;
verbatimTaxonRank: species; genus: *Phaeoisaria*; specificEpithet: *laianensis*;
scientificNameAuthorship: Y. Liu, D.M. Hu & Z.J. Zhai; continent: Asia; country: China;
stateProvince: Anhui; county: Laian; locality: Wawuzhuang; verbatimElevation: 35;
locationRemarks: Label transliteration; verbatimCoordinates: 32.66 N, 118.65 E;
verbatimLatitude: 32.66; verbatimLongitude: 118.65; samplingProtocol: collecting;
eventDate: 2021-05-06; year: 2021; month: 5; day: 6; habitat: Freshwater; recordedBy:
Yu Liu; identifiedBy: Yu Liu and Zhi-jun Zhai; dateIdentified: 2021; type: PhysicalObject;
language: en; rightsHolder: Dian-Ming Hu and Zhi-jun Zhai; institutionID: HFJAU10040;
collectionID: LKJ17; institutionCode:
the Herbarium of Fungi, Jiangxi Agricultural University (HFJAU); collectionCode: Fungi;
ownerInstitutionCode: HFJAU

Description

Saprobic on decaying wood submerged in freshwater habitats. **Sexual morph:** Undetermined. **Asexual morph:** Colonies effuse, solitary, scattered, dark brown to black, hairy, covered by white conidial mass. Mycelium partly superficial, partly immersed. Synnemata 290–848 × 9.3–30.7 μm (\bar{x} = 532 × 18.6, SD = 159×5, n = 20), erect, rigid, dark brown to black, velvety, smooth, composed of compactly and parallel adressed conidiophores. Conidiophores 116.2–491.1 × 2–3.2 μm (\bar{x} = 276.1 × 2.4, SD = 96.7×0.5, n = 10), macronematous, synnematous, septate, branched, brown to dark brown, smooth. Conidiogenous cells 8.3–27.5 × 2.3–3.8 μm (\bar{x} = 17.1 × 2.7, n = 10), polyblastic, integrated, terminal, hyaline to pale brown, smooth, denticulate, sympodial, each with several denticulate conidiogenous loci, 0.8–1.6 × 0.4–0.8 μm (\bar{x} = 1.3 × 0.7, n = 10). Conidia 5–7.2 × 1.7–2.9 μm (\bar{x} = 5.9 × 1.7, SD = 0.5 × 0.3, n = 50), ellipsoidal to obovoid, straight, rounded at the apex, obtuse and tapering towards base, hyaline, aseptate, guttulate, smooth-walled. (Fig. 1).

Culture characteristics: Conidia germinated within 24h which germ tubes were produced from both ends or sides at 28°C on PDA. The colony on PDA grows up slowly and reaches 24.5 mm in 26 days, periphery grey, surface folded, middle gray-green to black, raised with mycelium in the centre, covered with lots of white conidia, powdery, reverse gray to black.

Material examined: China, Anhui Province, alt. 35 m, near 32.66°N, 118.65°E, on decaying wood submerged in a freshwater stream, 6 May 2021, Y. Liu, G.P. Xu and Z.J. Zhai, LKJ17 (HFJAU10040, holotype), ex-type living culture, JAUCC 4967 = JAUCC 4968.

Etymology

Name reflect the district of this fungus.

Notes

Phylogenetic analyses show that *Phaeoisaria laianensis* a phylogenetically-distinct species, most closely related to *P. dalbergiae* and then to *P. clematidis* (Fig. 2).

However, *P. laianensis* is easily distinguished from *P. dalbergiae* by its ellipsoidal to obovoid, rounded at the apex, and tapering towards base conidia and by its existence of synnemata, which is absent in the latter species, and in *P. curvata*, *P. glauca* (de Hoog and Papendorf 1976), *P. loranthacearum* (Crous et al. 2015), *P. annesophieae*, *P. fasciculata* (Réblová et al. 2016) as well (Table 2). The new species is similar to *P. clematidis* in having resembling synnemata or conidia (Hughes 1958, Luo et al. 2018), while the former has shorter synnemata (290–848 µm vs. 1000–1500 µm) and smaller conidia (5–7.2 µm wide vs. 4–10 µm wide) than *P. clematidis* (Table 2). Likewise, *P. laianensis* has longer synnemata than *P. siamensis* (290–848 µm vs. 330–380 µm), smaller conidiophores than *P. guttulata* (Hyde et al. 2018) and *P. aquatica* (116.2–491.1 × 2–3 µm vs. 480–700 × 2–5 µm and 1028–1262 × 3.5–4.5 µm) (Luo et al. 2018), and smaller conidia (5–7.2 × 1.7–2.9 µm) than *P. annesophieae* (4.5–9 × 2–3.5 µm) (Crous et al. 2017), *P. synnemata* (4–11 µm long) (Boonmee et al. 2021), *P. siamensis* (3–4 µm wide) (Table 2). In addition, it can be differentiated from *P. filiformis* by the indeterminate asexual morph of the latter species (Luo et al. 2019). All in all, *P. laianensis* is morphologically-distinct not only from its sister species (*P. dalbergiae* and *P. clematidis*) but also other species in *Phaeoisaria*.

Analysis

Phylogenetic analyses

The aligned matrix for the combined analysis, ITS+LSU+SSU had 3105bp, including ITS 509 bp, LSU 1172 bp and SSU 1424 bp. No topological conflict exists between the tree generated by ML analyses and the Bayesian tree. The Bayesian tree are shown with BS and PP in Fig. 2. All 15 *Phaeoisaria* species in our analyses form a monophyletic group (BS/PP = 59/1.00). Most importantly, the two collections of *Phaeoisaria laianensis* form an independent lineage with strong support (BS/PP = 100/1.00). This lineage groups with *P. dalbergiae* into a highly supported clade (BS/PP = 98/1.00), which is sister to *P. clematidis* (BS/PP = 54/1.00). Searching of NCBI's GenBank nucleotide database based on a megablast, the ITS sequence of *P. laianensis* shares 97.46% similarity with *P. dalbergiae* (CPC 39540), and 96.35% similarity with *P. clematidis* (DAOM 226789). In addition, the sequence has 9 different loci from that of *P. dalbergiae*, and 15 different loci from that of *P. clematidis*.

Discussion

In our molecular phylogenetic tree, *Phaeoisaria* consists of 15 species and is supported as a monophyletic group (BS/PP = 59/1.00, Fig. 2). The low ML bootstrap might be due to a large number of unavailable sequences for 13 species in *Phaeoisaria*. However, the independent lineage of *P. laianensis* (BS/PP = 100/1.00, Fig. 2) are established and groups with *P. dalbergiae* into a highly supported clade (BS/PP = 98/1.00, Fig. 2). This clade is sister to the four collections of *P. clematidis* although with lower support (BS/PP = 54/1.00, Fig. 2). In addition, the morphological characters of *P. laianensis* can be effortlessly

distinguished from *P. dalbergiae* and *P. clematidis* and other species in *Phaeoisaria* (Tables 2, 3). Notably, our results favor *P. laianensis* as a new species in the genus. However, molecular data for *Phaeoisaria* species are needed to be enriched to clarify more species relationships in the genus.

Phaeoisaria predominantly occurs on leaves, barks, decaying wood and twigs of plants from the freshwater or terrestrial habitats (Table 3), while some is isolated from surface marine sediment (e.g., *P. sedimenticola*, Cheng et al. 2014), some from soil (e.g., *P. annesophieae*, Crous et al. 2017), and some from saprobic decaying fruits (e.g., *P. siamensis*, Hyde et al. 2019). Consequently, the habitats of *Phaeoisaria* are various. In this research, we introduce another lignicolous freshwater fungus, *P. laianensis*, discovered in China, other than that the freshwater it exists has been somewhat polluted. *Phaeoisaria* is thought to play an important role in nutrient and carbon cycling, biological diversity and ecosystem functioning of freshwater ecosystems, for their ability to decompose lignocellulose in woody litter, softening the wood and releasing nutrients (Bucher et al. 2004, Vijaykrishna et al. 2005, Hyde et al. 2016, Luo et al. 2018). Nonetheless, some *Phaeoisaria* species is pathogenic to humans, for example, it has been reported that *P. clematidis* and *Phaeoisaria* sp. can cause corneal inflammation of the eye (keratitis) (Guarro et al. 2000, Chew et al. 2010), and the former species is saprotrophic, which is similar to *P. laianensis*. What is the role of *P. laianensis* in ecosystem functioning? Is this species also pathogenic to humans? Such questions are waiting to be investigated by researchers.

Acknowledgements

We are grateful to Deng-Mei Fan (Agricultural college, Jiangxi Agricultural University) for very helpful comments on earlier drafts of this manuscript. This study was supported by the National Natural Science Foundation of China (NSFC 32070023 and NSFC 32060014), the Natural Science Foundation of Jiangxi Province (20151BAB214002) and Science and Technology Plan Project of Jiangxi Province (GJJ160417).

References

- Boonmee S, Wanasinghe DN, Calabon MS, Huanraluek N, Chandrasiri SKU, Jones GEB, Rossi W, Leonardi M, Singh SK, Rana S, Singh PN, Maurya DK, Lagashetti AC, Choudhary D, Dai YC, Zhao CL, Mu YH, Yuan HS, He SH, Phookamsak R, Jiang HB, Martín MP, Dueñas M, Telleria MT, Kalucka IL, Jagodziński AM, Liimatainen K, Pereira DS, Phillips AJL, Suwannarach N, Kumla J, Khuna S, Lumyong S, Potter TB, Shivas RG, Sparks AH, Vaghefi N, Abdel-Wahab MA, Abdel-Aziz FA, Li GJ, Lin WF, Singh U, Bhatt RP, Lee HB, Nguyen TTT, Kirk PM, Dutta AK, Acharya K, Sarma VV, Niranjana M, Rajeshkumar KC, Ashtekar N, Lad S, Wijayawardene NN, Bhat DJ, Xu RJ, Wijesinghe SN, Shen HW, Luo ZL, Zhang ZY, Sysouphanthong P, Thongklang N, Bao DF, Aluthmuhandiram JVS, Abdollahzadeh J, Javadi A, Dovana F, Usman M, Khalid AN, Dissanayake AJ, Telagathoti A, Probst M, Peintner U, Garrido-Benavent I, Bóna L,

- Merényi Z, Boros L, Zoltán B, Stielow JB, Jiang N, Tian CM, Shams E, Dehghanizadeh F, Pordel A, Javan-Nikkhah M, Denchev TT, Denchev CM, Kemler M, Begerow D, Deng CY, Harrower E, Bozorov T, Kholmuradova T, Gafforov Y, Abdurazakov A, Xu JC, Mortimer PE, Ren GC, Jeewon R, Maharachchikumbura SSN, Phukhamsakda C, Mapook A, Hyde KD (2021) Fungal diversity notes 1387-1511: taxonomic and phylogenetic contributions on genera and species of fungal taxa. *Fungal Diversity* 111: 1-335. <https://doi.org/10.1007/s13225-021-00489-3>
- Bucher VVC, Pointing SB, Hyde KD, Reddy CA (2004) Production of wood decay enzymes, loss of mass, and lignin solubilization in wood by diverse tropical freshwater fungi. *Microbial Ecology* 48: 331-337. <https://doi.org/10.1007/s00248-003-0132-x>
 - Castañeda Ruiz RF, Velasquez S, Cano J, Saikawa M, Guarro J (2002) *Phaeoisaria aguilerae* anam. sp. nov. from submerged wood in Cuba with notes and reflections in the genus *Phaeoisaria*. *Cryptogamie Mycologie* 23 (1): 9-18.
 - Cheng X, Li W, Zhang T (2014) A new species of *Phaeoisaria* from intertidal marine sediment collected in Weihai, China. *Mycotaxon* 127 (1): 17-24. <https://doi.org/10.5248/127.17>
 - Chew HF, Jungkind DL, Mah DY, Raber IM, Toll AD, MJ T, Cohen EJ (2010) Post-traumatic fungal keratitis caused by *Carpoligna* sp. *Cornea* 29 (4): 449-452. <https://doi.org/10.1097/ICO.0b013e3181af3954>
 - Crous PW, Schumacher RK, Wingfield MJ, Lombard L, Giraldo A, Christensen M, Gardiennet A, Nakashima C, Pereira OL, AJ S, Groenewald JZ (2015) Fungal systematics and evolution: FUSE 1. *Sydowia* 67: 81-118. <https://doi.org/10.12905/0380.sydowia67-2015-0081>
 - Crous PW, Wingfield MJ, Burgess TI, Carnegie AJ, Hardy GE, Smith D, Summerell BA, Cano-Lira JF, Guarro J, Houbraken J, Lombard L, Martín MP, Sandoval-Denis M, Alexandrova AV, Barnes CW, Baseia IG, Bezerra JD, Guarnaccia V, May TW, Hernández-Restrepo M, Stchigel AM, Miller AN, Ordoñez ME, Abreu VP, Accioly T, Agnello C, Agustíncolmn A, Albuquerque CC, Alfredo DS, Alvarado P, Araújo-Magalhães GR, Arauzo S, Atkinson T, Barili A, Barreto RW, Bezerra JL, Cabral TS, Camello RF, Cruz RH, Daniëls PP, Da Silva BD, De Almeida DA, De Carvalho Júnior AA, Decock CA, Delgat L, Denman S, Dimitrov RA, Edwards J, Fedosova AG, Ferreira RJ, Firmino AL, Flores JA, García D, Gené J, Giraldo A, Góis JS, Gomes AA, Gonçalves CM, Gouliamova DE, Groenewald M, Guéorguiev BV, Guevara-Suarez M, Gusmão LF, Hosaka K, Hubka V, Huhndorf SM, Jadan M, Jurjević Ž, B K, V K, TKA K, I L, Lamlertthon S, Lisboa WS, Loizides M, Luangsa-Ard JJ, Lysková P, McCormack WP, Macedo DM, Machado AR, Malysheva EF, Marinho P, Matočec N, Meijer M, Mešič A, Mongkolsamrit S, Moreira KA, Morozova OV, Nair KU, Nakamura N, Noisripoom W, Olariaga I, Oliveira RJ, Paiva LM, Pawar P, Pereira OL, Peterson SW, Prieto M, Rodríguez-Andrade E, Rojodeblas C, Roy M, Santos ES, Sharma R, Silva GA, Souza-Motta CM, Takeuchi-Kaneko Y, Tanaka C, Thakur A, Smith MT, Tkalčec Z, Valenzuela-Lopez N, Vanderkleij P, Verbeken A, Viana MG, Wang XW, Groenewald JZ (2017) Fungal Planet description sheets: 625-715. *Persoonia: Molecular Phylogeny and Evolution of Fungi* 39 (1): 270-467. <https://doi.org/10.3767/persoonia.2017.39.11>
 - Crous PW, Osieck ER, Jurjević Ž, Boers J, Iperen AL, Starink-Willemsse M, Dima B, Balashov S, Bulgakov TS, Johnston PR, Morozova OV, Pinruan U, Sommai S, Alvarado P, Decock CA, Lebel T, McMullan-Fisher S, Moreno G, Shivas RG, Zhao L, Abdollahzadeh J, Abrinbana M, Ageev DV, Akhmetova G, Alexandrova AV, Altés A,

- Amaral AG, Angelini C, Antonín V, Arenas F, Asselman P, Badali F, Baghela A, Bañares A, Barreto RW, Baseia IG, Bellanger JM, Berraf-Tebbal A, Biketova AY, Bukharova NV, Burgess TI, Cabero J, Câmara MP, Cano-Lira JF, Ceryngier P, Chávez R, Cowan DA, de Lima AF, Oliveira RL, Denman S, Dang QN, Dovana F, Duarte IG, Eichmeier A, Erhard A, Esteve-Raventós F, Fellin A, Ferisin G, Ferreira RJ, Ferrer A, Finy P, Gaya E, Geering AD, Gil-Durán C, Glässnerová K, Glushakova AM, Gramaje D, Guard FE, Guarnizo AL, Haelewaters D, Halling RE, Hill R, Hirooka Y, Hubka V, Iliushin VA, Ivanova DD, Ivanushkina NE, Jangsantear P, Justo A, Kachalkin AV, Kato S, Khamsuntorn P, Kirtsideli IY, Knapp DG, Kochkina GA, Koukol O, Kovács GM, Kruse J, Kumar TKA, Kušan I, Læssøe T, Larsson E, Lebeuf R, Levicán G, M L, P M, JJ L, EG L, Magaña-Dueñas V, Maggs-Kölling G, Malysheva EF, Malysheva VF, Martín B, Martín MP, Matočec N, McTaggart AR, Mehrabi-Koushki M, Mešić A, Miller AN, Mironova P, Moreau PA, Morte A, Müller K, Nagy LG, Nanu S, Navarro-Ródenas A, Nel WJ, Nguyen TH, Nóbrega TF, Noordeloos ME, Olariaga I, Overton BE, Ozerskaya AM, Palani P, Pancorbo F, Papp V, Pawłowska J, Pham TQ, Phosri C, Popov ES, Portugal A, Pošta A, Reschke K, Reul M, Ricci GM, Rodríguez A, Romanowski J, Ruchikachorn N, Saar I, Safi A, Sakolrak B, Salzmann F, Sandoval-Denis M, Sangwichein E, Sanhueza L, Sato T, Sastoque A, Senn-Irlert B, Shibata A, Siepe K, Somrithipol S, Spetik M, Sridhar P, Stchigel AM, Stuskova K, Suwannasai N, Tan YP, Thangavel R, Tiago I, Tiwari S, Tkalčec Z, Tomashevskaya MA, Tonegawa C, Tran HX, Tran NT, Trovão J, Trubitsyn VE, Van Wyk J, Vieira WA, Vila J, Visagie CM, Vizzini A, Volobuev SV, Vu DT, Wangsawat N, Yaguchi T, Ercole E, Ferreira BW, de Souza AP, Vieira BS, Groenewald JZ (2021) Fungal Planet description sheets: 1284-1382. *Persoonia: Molecular Phylogeny and Evolution of Fungi* 47 (1): 178-374. <https://doi.org/10.3767/persoonia.2021.47.06>
- de Hoog GS, Papendorf MC (1976) The genus *Phaeoisaria*. *Persoonia: Molecular Phylogeny and Evolution of Fungi* 8 (4): 407-414.
 - Deighton FC (1974) Four synnematosous hyphomycetes. *Transactions of British Mycological Society* 62: 243-252. [https://doi.org/10.1016/S0007-1536\(74\)80033-1](https://doi.org/10.1016/S0007-1536(74)80033-1)
 - Doyle JJ, Doyle JL (1987) A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochemical Bulletin* 19: 11-15.
 - Guarro J, Vieira LA, de Freitas D, Gené J, Zaror L, Hofling-Lima AL, Fischman O, Zorat-Yu C, Figueras MJ (2000) *Phaeoisaria clematidis* as a cause of keratomycosis. *Journal of Clinical Microbiology* 38 (6): 2434-2437. <https://doi.org/10.1128/JCM.38.6.2434-2437.2000>
 - Höhnelt FV (1909) Fragmente zur Mykologie (VI. Mitteilung, Nr. 182 bis 288). *Sitzungsberichten der Kaiserliche Akademie der Wissenschaften in Wien Mathematische-Naturwissenschaftliche Klasse, Abt I* 118 (182): 275-452.
 - Hopple JS, Vilgalys R (1999) Phylogenetic relationships in the mushroom genus *Coprinus* and dark-spored allies based on sequence data from the nuclear gene coding for the large ribosomal subunit RNA: Divergent domains, outgroups, and monophyly. *Molecular Phylogenetics and Evolution* 13 (1): 1-19. <https://doi.org/10.1006/mpev.1999.0634>
 - Hughes SJ (1958) Revisiones hyphomycetum aliquot cum appendice de nominibus rejiciendis. *Canadian Journal of Botany* 36 (6): 727-836. <https://doi.org/10.1139/b58-067>
 - Hyde KD, Fryar S, Tian Q, Bahkali AH, Xu JC (2016) Lignicolous freshwater fungi along a north-south latitudinal gradient in the Asian/Australian region; can we predict the

- impact of global warming on biodiversity and function? *Fungal Ecology* 19: 190-200.
<https://doi.org/10.1016/j.funeco.2015.07.002>
- Hyde KD, Norphanphoun C, Abreu VP, Bazzicalupo AL, Chethana KWT, Clericuzio M, Dayarathne MC, Dissanayake AJ, Ekanayaka AH, He MQ, Hongsanan S, Huang SK, Jayasiri SC, Jayawardena RS, Karunarathna A, Konta S, Kušan I, Lee HG, Li J, Lin C, Liu NG, Lu YZ, Luo ZL, Manawasinghe IS, Mapook A, Perera RH, Phookamsak R, Phukhamsakda C, Siedlecki I, Soares AM, Tennakoon DS, Tian Q, Tibpromma S, Wanasinghe DN, Xiao YP, Yang J, Zeng XY, Abdel-Aziz FA, Li WJ, Senanayake IC, Shang QJ, Daranagama DA, de Silva NI, Thambugala KM, Abdel-Wahab MA, Bahkali AH, Berbee ML, Boonmee S, Bhat DJ, Bulgakov TS, Buyck B, Camporesi E, Castañeda-Ruiz RF, Chomnunti P, Doilom M, Dovana F, Gibertoni TB, Jadan M, Jeewon R, Jones EB, Kang JC, Karunarathna SC, Lim YW, Liu JK, Liu ZY, Plautz HL, Lumyong S, Maharachchikumbura SS, Matočec N, McKenzie EH, Mešić A, Miller D, Pawłowska J, Pereira OL, Promputtha I, Romero AI, Ryvarden L, Su HY, Suetrong S, Tkalčec Z, Vizzini A, Wen TC, Wisitrassameewong K, Wrzosek M, Xu JC, Zhao Q, Zhao RL, Mortimer PE (2017) Fungal diversity notes 603-708: taxonomic and phylogenetic notes on genera and species. *Fungal Diversity* 87 (1): 1-235. <https://doi.org/10.1007/s13225-017-0391-3>
 - Hyde KD, Chaiwan N, Norphanphoun C, Boonmee S, Camporesi E, Chethana KW, Dayarathne MC, Silva NI, Dissanayake AJ, Ekanayaka AH, Hongsanan S, Huang SK, Jayasiri SC, Jayawardena RS, Jiang HB, Karunarathna A, Lin CG, Liu JK, Liu NG, Lu YZ, Luo ZL, Maharachchimbura SS, Manawasinghe IS, Pem D, Perera RH, Phukhamsakda C, Samarakoon MC, Senwana C, Shang QJ, Tennakoon DS, Thambugala KM, Tibpromma S, Wanasinghe DN, Xiao YP, Yang J, Zeng XY, Zhang JF, Zhang SN, Bulgakov TS, Bhat DJ, Cheewangkoon R, Goh TK, Jones EB, Kang JC, Jeewon R, Liu ZY, Lumyong S, Kuo CH, McKenzie EH, Wen TC, Yan J, Zhao Q (2018) Mycosphere notes 169-224. *Mycosphere* 9 (2): 271-430. <https://doi.org/10.5943/mycosphere/9/2/8>
 - Hyde KD, Tennakoon DS, Jeewon R, Bhat DJ, Maharachchikumbura SSN, Rossi W, Leonardi M, Lee HB, Mun HY, Houbraken J, Nguyen TTT, Jeon SJ, Frisvad JC, Wanasinghe DN, Luücking R, Aptroot A, Cáceres MES, Karunarathna SC, Hongsanan S, Phookamsak R, de Silva NI, Thambugala KM, Jayawardena RS, Senanayake IC, Boonmee S, Chen J, Luo ZL, Phukhamsakda C, Pereira OL, Abreu VP, Rosado AW, Bart B, Randrianjohany E, Hofstetter V, Gibertoni TB, da Silva Soares AM, Plautz Jr. HL, Sotão HMP, Xavier WKS, Bezerra JDP, de Oliveira TGL, de Souza Motta CM, Magalhães OMC, Bundhun D, Harishchandra D, Manawasinghe IS, Dong W, Zhang SN, Bao DF, Samarakoon MC, Pem D, Karunarathna A, Lin CG, Yang J, Perera RH, Kumar V, Huang SK, Dayarathne MC, Ekanayaka AH, Jayasiri SC, Xiao Y, Konta S, Niskanen T, Liimatainen K, Dai YC, Ji XH, Tian XM, Mešić A, Singh SK, Phutthacharoen K, Cai L, Sorvongxay T, Thiyagaraja V, Norphanphoun C, Chaiwan N, Lu YZ, Jiang HB, Zhang JF, Abeywickrama PD, Aluthmuhandiram JV, Brahmanage RS, Zeng M, Chethana T, Wei D, Réblová M, Fournier J, Nekvindová J, do Nascimento Barbosa R, dos Santos JE, de Oliveira NT, Li GJ, Ertz D, Shang QJ, Phillips AJL, Kuo CH, Camporesi E, Bulgakov TS, Lumyong S, Jones EBG, Chomnunti P, Gentekaki E, Bungartz F, Zeng XY, Fryar S, Tkalčec Z, Liang J, Li G, Wen TC, Singh PN, Gafforov Y, Promputtha I, Yasanthika E, Goonasekara ID, Zhao RL, Zhao Q, Kirk PM, Liu JK, Yan J, Mortimer PE, Xu JC (2019) Fungal diversity notes 1036-1150: taxonomic and

- phylogenetic contributions on genera and species of fungal taxa. *Fungal Diversity* 96 (1): 1-242. <https://doi.org/10.1007/s13225-019-00429-2>
- Kato H, Rozewicki J, Yamada KD (2019) MAFFT online service: multiple sequence alignment, interactive sequence choice and visualization. *Briefings in Bioinformatics* 20: 1160-1166. <https://doi.org/10.1093/bib/bbx108>
 - Liu JK, Hyde KD, Jones EBG, Ariyawansa HA, Bhat DJ, Boonmee S, Maharachchikumbura SS, McKenzie EHC, Phookamsak R, Phukhamsakda C, Shenoy BD, Abdel-Wahab MA, Buyck B, Chen J, Chethana KWT, Singtripop C, Dai DQ, Dai YC, Daranagama DA, Dissanayake AJ, Doilom M, D'souza MJ, Fan XL, Goonasekara ID, Hirayama K, Hongsanan S, Jayasiri SC, Jayawardena RS, Karunarathna SC, Li WJ, Mapook A, Norphanphoun C, Pang KL, Perera RH, Peršoh D, Pinruan U, Senanayake IC, Somrithipol S, Suetrong S, Tanaka K, Thambugala KM, Tian Q, Tibpromma S, Udayanga D, Wijayawardene NN, Wanasinghe D, Wisitrasameewong K, Zeng XY, Abdel-Aziz FA, Adamčík S, Bahkali AH, Boonyuen N, Bulgakov TS, Callac P, Chomnunti P, Greiner K, Hashimoto A, Hofstetter V, Kang JC, Lewis D, Li XH, Liu XZ, Liu ZY, Matsumura M, Mortimer PE, Rambold G, Randrianjohany E, Sato G, Sriindrasutdhi V, Tian CM, Verbeken A, Brackel WV, Wang Y, Wen TC, Xu JC, Yan JY, Zhao RL, Camporesi E (2015) Fungal diversity notes 1-110: taxonomic and phylogenetic contributions to fungal species. *Fungal Diversity* 72: 1-197. <https://doi.org/10.1007/s13225-015-0324-y>
 - Li XH, Liu YL, Song HY, Hu DM, Gao Y, Hu HJ, Zhou JP (2021) *Sporidesmiella lignicola* sp. nov., a new hyphomycetous fungus from freshwater habitats in China. *Biodiversity Data Journal* 9: 77414. <https://doi.org/10.3897/BDJ.9.e77414>
 - Luo ZL, Hyde KD, Bhat DJ, Maharachchikumbura SS, Bao DF, Li WL, Su XJ, Yang XY (2018) Morphological and molecular taxonomy of novel species Pleurotheciaceae from freshwater habitats in Yunnan, China. *Mycological Progress* 17 (5): 511-530. <https://doi.org/10.1007/s11557-018-1377-6>
 - Luo ZL, Hyde KD, Liu JK, Maharachchikumbura SS, Jeewon R, Bao DF, Bhat DJ, Lin CG, Li WL, Yang J, Liu NG, Lu YZ, Jaya-wardena RS, Li JF, Su HY (2019) Freshwater sordariomycetes. *Fungal Diversity* 99: 451-660. <https://doi.org/10.1007/s13225-019-00438-1>
 - Mason EW, Ellis MB (1953) British species of *Periconia*. *Mycological Paper* 56: 1-127.
 - Mel'nik VA (2012) *Phaeoisaria vietnamensis* sp. nov. and *P. clematidis* (hyphomycetes) from Vietnam. *Mycosphere* 3 (6): 957-960. <https://doi.org/10.5943/mycosphere/3/6/10>
 - Mercado-Sierra A (1984) Nuevas especies de *Deightoniella*, *Phaeoisaria*, *Sporidesmium*, Y *Taeniolella* (Hyphomycetes) de Cuba. *Acta Botánica Cubana* 21: 1-10.
 - Mercado-Sierra A, Figueras MJ, Gene J (1997) New or rare hyphomycetes from Cuba VIII. Species of *Lylea*, *Phaeisaria*, *Arxiella*, *Graphium*, *Periconia*, and *Ramichloridium*. *Mycotaxon* 63: 369-375.
 - Rambaut A (2018) FigTree v1.4.4: Tree figure drawing tool. URL: <https://github.com/rambaut/figtree/releases>
 - Réblová M, Seifert KA, Fournier J, Štěpánek V (2016) Newly recognized lineages of perithecial ascomycetes: the new orders Conioscyphales and Pleurotheciales. *Persoonia: Molecular Phylogeny and Evolution of Fungi* 37 (1): 57-81. <https://doi.org/10.3767/003158516X689819>
 - Siboe GM, Kipk PM, Cannon PF (1999) New dematiaceous hyphomycetes from Kenyan rare plants. *Mycotaxon* 73: 283-302.

- Stamatakis A, Alachiotis N (2010) Time and memory efficient likelihood-based tree searches on phylogenomic alignments with missing data. *Bioinformatics* 26 (12): 132-139. <https://doi.org/10.1093/bioinformatics/btq205>
- Sutton BC (1973) Hyphomycetes from Manitoba and Saskatchewan, Canada. *Mycological Papers* 132: 1-143.
- Sutton BC, Hodges CS (1976) Eucalyptus micofungi. *Microdochium* and *Phaeoisaria* species from Brazil. *Nova Hedwigia* 27: 215-222.
- Sutton BC (1993) Mitosporic fungi from Malawi. *Mycological Papers* 167: 1-93.
- Vijaykrishna D, Jeewon R, Hyde KD (2005) *Fusoidispora aquatica*: a new freshwater ascomycete from Hong Kong based on morphology and phylogeny inferred from rDNA gene sequences. *Sydowia* 57 (2): 267-280.
- White TJ, Bruns T, Lee SJ, Taylor J (1990) PCR protocols: a guide to methods and applications. In: Innis M, Gelfand D, Shinsky J, White T (Eds) *Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics*. Academic Press [In Academic Press, New York]. <https://doi.org/10.1016/B978-0-12-372180-8.50042-1>
- Zhai ZJ, Yan JQ, Li WW, Gao Y, Hu HJ, Zhou JP, Song HY, Hu DM (2022) Three novel species of *Distoseptispora* (Distoseptisporaceae) isolated from bamboo in Jiangxi Province, China. *MycKeys* 88: 35-54. <https://doi.org/10.3897/mycokeys.88.79346>
- Zhang D, Gao F, Jakovlić I, Zou H, Zhang J, Li WX, Wang GT (2020) PhyloSuite: An integrated and scalable desktop platform for streamlined molecular sequence data management and evolutionary phylogenetics studies. *Molecular Ecology Resources* 20 (1): 348-355. <https://doi.org/10.1111/1755-0998.13096>
- Zhaxybayeva O, Gogarten JP (2002) Bootstrap, Bayesian probability and maximum likelihood mapping: exploring new tools for comparative genome analyses. *BMC genomics* 3 (1): 1-15. <https://doi.org/10.1186/1471-2164-3-4>




Figure 1.

Phaeoisaria laianensis (HFJAU 10040, Holotype) **a, b** Colonies on wood **c, d** Conidiophores **e, f** Conidio-genous cells with conidia **g** Germinating conidium **h** Conidia **i, j** Colony on PDA for 26 days from above and reverse. Scale bars: a, b = 100 μm , c, d = 50 μm , e–h = 10 μm .




Figure 2.

Phylogenetic tree of Bayesian analysis based on a concatenated alignment of ITS, LSU and SSU sequences. Branch support is shown at the nodes, maximum likelihood bootstrap support (BS, black) $\geq 50\%$ and Bayesian posterior probability (PP, red) ≥ 0.90 . *Leotialubrica* (AFTOLID 1) and *Microglossum rufum* (OSC100641) are selected as the outgroup taxon. The new species is marked in red and ex-type strains are in bold.

Table 1.

Sequences used in this study. **Note** : Ex-type strains are in **bold**. The sequences of new species have been underlined. “ – ”, sequence is unavailable in GenBank.

Taxonomy	Strain	GenBank accession numbers		
		ITS	LSU	SSU
<i>Adelosphaeria catenata</i>	CBS 138679	KT278721	KT278707	KT278692
<i>Ascotaiwania fusiformis</i>	MFLUCC 15-0625	–	KX550894	KX550898
<i>Ascotaiwania fusiformis</i>	MFLU 15-1156	MG388215	NG-057114	–
<i>Ascotaiwania lignicola</i>	NIL 00005	HQ446341	HQ446364	HQ446284
<i>Ascotaiwania sawadae</i>	SS00051	HQ446340	HQ446363	HQ446283
<i>Bactrodesmiastrum obovatum</i>	FMR 6482	FR870264	FR870266	–
<i>Bactrodesmiastrum pyriforme</i>	FMR 10747	FR870263	FR870265	–
<i>Brachysporiella setosa</i>	HKUCC 3713	–	AF132334	–
<i>Canalisporium exiguum</i>	SS 00809	GQ390296	GQ390281	GQ390266
<i>Canalisporium grenadoideum</i>	BCC 20507	–	GQ390267	GQ390252
<i>Canalisporium pulchrum</i>	SS03982	GQ390292	GQ390277	GQ390262
<i>Conioscypha lignicola</i>	CBS 335.93	–	AY484513	JQ437439
<i>Conioscypha minutispora</i>	CBS 137253	–	MH878131	–
<i>Conioscypha peruviana</i>	CBS 137657	–	KF781539	–
<i>Conioscypha varia</i>	CBS 113653	–	AY484512	AY484511
<i>Fuscosporella pyriformis</i>	MFLUCC 16-0570	MG388217	KX550896	KX550900
<i>Helicoon farinosum</i>	DAOM 241947	JQ429145	JQ429230	–
<i>Leotia lubrica</i>	AFTOLID 1	DQ491484	AY544644	AY544746
<i>Melanotrigonum ovale</i>	CBS 138815	KT278722	KT278711	KT278698
<i>Melanotrigonum ovale</i>	CBS 138744	KT278725	KT278710	KT278697
<i>Melanotrigonum ovale</i>	CBS 138743	KT278724	KT278709	KT278696
<i>Melanotrigonum ovale</i>	CBS 138742	KT278723	KT278708	KT278695

<i>Microglossum rufum</i>	OSC100641	–	DQ470981	DQ471033
<i>Mucispora obscuriseptata</i>	MFLUCC 15-0618	MG388218	KX550892	KX550897
<i>Parafuscosporella moniliformis</i>	MFLUCC 15-0626	MG388219	KX550895	KX550899
<i>Phaeoisaria annesophieae</i>	CBS 143235	MG022180	MG022159	–
<i>Phaeoisaria annesophieae</i>	MFLU190531	MT559109	MT559084	–
<i>Phaeoisaria aquatica</i>	MFLUCC 16-1298	MF399237	MF399254	–
<i>Phaeoisaria clematidis</i>	MFLUCC 16-1273	MF399229	MF399246	–
<i>Phaeoisaria clematidis</i>	MFLUCC 17-1341	MF399230	MF399247	MF399216
<i>Phaeoisaria clematidis</i>	MFLUCC 17-1968	MG837022	MG837017	MG837027
<i>Phaeoisaria clematidis</i>	DAOM 226789	JQ429155	JQ429231	JQ429243
<i>Phaeoisaria dalbergiae</i>	CPC 39540	OK664703	–	–
<i>Phaeoisaria fasciculata</i>	CBS 127885	KT278719	KT278705	KT278693
<i>Phaeoisaria fasciculata</i>	DAOM 230055	KT278720	KT278706	KT278694
<i>Phaeoisaria filiformis</i>	MFLUCC 18-0214	MK878381	MK835852	MK834785
<i>Phaeoisaria guttulata</i>	MFLUCC 17-1965	MG837021	MG837016	MG837026
<u><i>Phaeoisaria laianensis</i></u>	<u>JAUCC4967</u>	<u>ON937559</u>	<u>ON937557</u>	<u>ON937562</u>
<i>Phaeoisaria laianensis</i>	<u>JAUCC4968</u>	<u>ON937560</u>	<u>ON937561</u>	<u>ON937558</u>
<i>Phaeoisaria lorantheaeum</i>	CBS 140009	KR611888	MH878676	–
<i>Phaeoisaria lorantheaeum</i>	BYCDW25	MG820097	–	–
<i>Phaeoisaria lorantheaeum</i>	BYCDW24	MG820098	–	–
<i>Phaeoisaria microspora</i>	MFLUCC 16-0033	MF671987	MF167351	–
<i>Phaeoisaria pseudoclematidis</i>	MFLUCC 11-0393	KP744457	KP744501	KP753962
<i>Phaeoisaria sedimenticola</i>	CGMCC3.14949	JQ074237	JQ031561	–
<i>Phaeoisaria sedimenticola</i>	S-908	MK878380	MK835851	–
<i>Phaeoisaria siamensis</i>	MFLUCC 16-0607	MK607610	MK607613	MK607612
<i>Phaeoisaria sparsa</i>	FMR 11939	–	HF677185	–

<i>Phaeoisaria synnematica</i>	NFCCI 4479	MK391494	MK391492	–
<i>Phragmocephala stemphylioides</i>	KAS 4277	KT278730	KT278717	–
<i>Plagiascoma frondosum</i>	CBS 139031	–	KT278713	KT278701
<i>Pleurotheciella centenaria</i>	DAOM 229631	JQ429151	JQ429234	JQ429246
<i>Pleurotheciella rivularia</i>	CBS 125237	JQ429161	JQ429233	JQ429245
<i>Pleurotheciella rivularia</i>	CBS 125238	JQ429160	JQ429232	JQ429244
<i>Pleurotheciella uniseptata</i>	KUMCC 15-0407	MF399231	MF399248	–
<i>Pleurothecium aquaticum</i>	MFLUCC 17-1331	MF399245	MF399263	–
<i>Pleurothecium aquaticum</i>	MFLUCC 21-0148	OM654775	OM654772	OM654807
<i>Pleurothecium floriforme</i>	MFLUCC 15-0628	KY697281	KY697277	KY697279
<i>Pleurothecium obovoideum</i>	CBS 209.95	EU041784	EU041841	–
<i>Pleurothecium pulneyense</i>	MFLUCC 16-1293	–	MF399262	MF399228
<i>Pleurothecium recurvatum</i>	CBS 138747	KT278728	KT278714	KT278703
<i>Pleurothecium recurvatum</i>	CBS 131272	JQ429149	JQ429237	JQ429251
<i>Pleurothecium recurvatum</i>	CBS 101581	JQ429148	AF261070	JQ429248
<i>Pleurothecium semifecundum</i>	CBS 131482	JQ429158	JQ429239	JQ429253
<i>Pleurothecium semifecundum</i>	CBS 131271	JQ429159	JQ429240	JQ429254
<i>Savoryella longispora</i>	SAT 00322	HQ446359	HQ446380	HQ446302
<i>Savoryella paucispora</i>	SAT 00866	–	HQ446381	HQ446303
<i>Savoryella verrucosa</i>	SS 00052	HQ446353	HQ446374	HQ446296
<i>Sterigmatobotrys macrocarpa</i>	DAOM 230059	JQ429154	GU017316	–
<i>Sterigmatobotrys macrocarpa</i>	PRM 915682	JQ429153	GU017317	JQ429255
<i>Sterigmatobotrys rudis</i>	DAOM 229838	JQ429152	JQ429241	JQ429256

Table 2.

Synopsis 1 of asexual morphological characteristics of *Phaeoisaria* species. **Note:** “-” is indeterminate or unavailable.

Species	synnemata (µm)	synnemata characteristics	Conidiophores (µm)	Conidiophores characteristics	Conidia (µm)
<i>Phaeoisaria laianensis</i>	290–848 × 9.3–30.7	erect, rigid, dark brown to black, velvety, smooth, composed of compactly and parallels adpressed conidiophores	116.2–491.1 × 2 - 3.2	macronematous, synnematos, septate, branched, brown to dark brown, smooth	5–7.2 × 1.7–2.9
<i>P. aguilerae</i>	-	-	-	-	18–29.5 × 4–5
<i>P. annesophieae</i>	-	-	Conidiophores indeterminate	sometimes grouping in strands of 2–4 hyphae, semi-macronematous reduced to a single conidiogenous cell, arising from aerial hyphae, cylindrical, hyaline to pale brown	4.5–9 × 2–3.5
<i>P. aquatica</i>	-	erect, rigid, dark brown to black, velvety, smooth	1028–1262 × 3.5–4.5	macronematous, synnematos, brown to dark brown, smooth	6.5–7.5 × 2.5–3.5
<i>P. bambusae</i>	-	erect, rigid, dark brown to black, velvety, smooth	-	macronematous, synnematos, brown to dark brown, smooth	-
<i>P. caffra</i>	-	Synnemata composed of at least 10 adpressed hyphae. Synnemata fertile over most of their surface	-	Conidiophores not tuberculate	7.5–12 × 2.5–3.5

<i>P. clavulata</i>	-	Stiff synnemata, composed of parallel hyphae, packed with slender, curved conidiogenous cells with very thin, fragile conidiogenous rachides	-	-	1–2 long
<i>P. clematidis</i>	1000–1500 × 20–80	Conidiomata scattered, indeterminate, erect, rigid, superficial, dark brown composed of compact appressed conidiophores.	312–568 × 2.5–3.5	macronematous, septate, branched, brown to dark brown, smooth	4–10 × 1.5–2.5
<i>P. curvata</i>	-	-	Conidiophores indeterminate	-	(4–)6–8(–11) × (1–)2–3
<i>P. dalbergiae</i>	-	-	10–50 × 1.5–2.5	Conidiophores indeterminate, erect, subcylindrical, hyaline, smooth, 0–2-septate, unbranched or branched at apex	0.5 µm diam, (5–)6–7 × (1.5–)2
<i>P. fasciculata</i>	-	Synnemata absent	25–65 × 3.0–3.5	macronematous, arising from brown, thick-walled cells, cylindrical, pale brown, subhyaline towards the apex, unbranched, smooth-walled	6.0–8.0(–9.0) × 2.0
<i>P. filiformis</i>	-	-	-	-	-
<i>P. glauca</i>	-	-	Conidiophores indeterminate	-	2.5–3.5 × 1.6–2.2

<i>P. guttulata</i>	-	erect, rigid, dark brown to black, velvety, smooth, composed of compact and parallel adpressed conidiophores	480–700 × 2–5	macronematous, synnematos, erect, septate, smooth, mid brown to dark brown.	3.5–5.5 × 2.5–4.8
<i>P. infrafertilis</i>	-	Synnemata narrow, composed of only 5-6 brown adpressed hyphae	-	-	19.5–22 × 2–3
<i>P. loranthacearum</i>	-	-	10–30 × 2–3	Conidiophores arising from superficial hyphae, erect, solitary, branched at base or not, subcylindrical, straight to geniculate-sinuuous, 1–3-septate, hyaline, becoming medium brown, smooth	(5)7–8(9) × (1.5)2(3)
<i>P. magnifica</i>	-	Synnemata brush-like, synnemata with flaring hyphae at the tip	-	the conidiophores growing well away from the column in the apical portion	5–6.5 × 4–4.5
<i>P. microspora</i>	35–238 µm long, 4–31 µm wide at the base, 5–35 µm wide at the ape	erect, straight or flexuous, dark brown at base, pale brown at apex	25–225 × 1–3	macronematous, synnematos, septate, branched at the apex, smooth, pale to dark brown	4.5–6.9 × 1.3–3.1
<i>P. muscariformis</i>	-	-	-	-	12–22 × 4

<i>P. pseudoclematidis</i>	200–500 µm long, 40–80 µm wide at the base, 40–60 µm wide in the middle, 20–30 µm wide at the apex	erect, rigid, dark brown, velvety, smooth, composed of compactly and parallels adpressed conidiophores	50–500 × 2–3	macronematous, synnematos, brown to dark brown, septate, branched, smooth, single conidiophores	5–8.5 × 3–4
<i>P. sedimenticola</i>	up to 4000 µm high or sometimes longer, 70–90 µm wide	erect, cylindrical to subulate, consisting of very regular, parallel, brown hyphae			aseptate (3.5–)4.5–5.5(–7.5) × (2.5–)3–4(–4.5) 1-septate (4.5–)5.5–6.5(–9) × (2–)2.5–3.5(–4.5)
<i>P. siamensis</i>	330–380 × 20–25(–30)	Conidiomata scattered, indeterminate, erect, rigid, superficial, dark brown composed of compact appressed conidiophores	2–2.5(–3) µm wide,	macronematous, in synnematos conidiomata, scattered, synnemata subulate or cylindrical, indeterminate, at the base 13–15 µm beneath the fertile portion with conidiogenous cells, composed of medium to dark brown, smooth, septate parallel hyphae, splaying out at the middle to apex	5–8 × 3–4
<i>P. sparsa</i>	-	Synnemata composed of at least 10 adpressed hyphae. Synnemata fertile only at the apex	-	Conidiophores not tuberculate	10–15.5 × 2.5–3.5
<i>P. sparsa var. cubensis</i>	-	-	-	-	(4–)7–11 (–17) × (1.5–)2–3(–4)

<i>P. synnemata</i>	399–960 × 12–30	synnematal, erect, rigid, dark brown to olivaceous brown, composed of compact parallel adpressed conidiophores, cylindrical to clavate	1.5–960 μm long, 1–3.5 μm wide.	macronematous to semi- macronematous, highly geniculate, dark brown to olivaceous brown, synnematus, simple to dichotomously branched, emerging out at the apex and along the sides of the upper half or two thirds of each synnema, dark brown at the base, brown to pale brown towards the apex	4–11 × 2–5
<i>P. tuberculata</i>	-	Synnemata composed of at least 10 adpressed hyphae	-	Conidiophores conspicuously tuberculate	8–13.5 × 1.5–2
<i>P. uniseptata</i>	-	-	-	-	(3.5–)5.5– 7.5(–10) × 1.5–3
<i>P. vietnamensis</i>	330–380 μm high, 20–25(–30) μm wide at the base	-	2–2.5(–3) μm wide	macronematous, in synnematus conidiomata, scattered, synnemata subulate or cylindrical, indeterminate, composed of medium to dark brown, smooth, septate parallel hyphae, splaying out at the apex	18.5–23.5 × 4.5–5

Table 3.

Synopsis 2 of asexual morphological characteristics of *Phaeoisaria* species. **Note:** “-” is indeterminate or unavailable.

Species	Conidia septation	Conidia characteristics	Host	District	References
<i>Phaeoisaria laianensis</i>	aseptate	ellipsoidal to obovoid, straight, rounded at the apex, obtuse and tapering towards base, hyaline, guttulate, smooth-walled	Decaying wood	China, Anhui Province	This study
<i>P. aguilerae</i>	1-septate, rarely 2–3-septate	clavate or cylindrical, curved, with obtuse, rounded apex, slightly uncinata, and truncate base, hyaline, smooth	Decaying twig submerged in river	Cuba	Castañeda Ruiz et al. (2002)
<i>P. annesophieae</i>	aseptate	ellipsoidal to obovoid, straight or slightly curved, rounded at the ends or sometimes tapering toward the base, hyaline, guttulate, smooth-walled	Isolated from soil	The Netherlands, Geldermalsen	Crous et al. (2017)
<i>P. aquatica</i>	aseptate	ellipsoidal to obovoidal, rounded at the apex, hyaline, with two guttules smooth-walled	Decaying wood submerged in Jinsha River	China, Yunnan Province	Luo et al. 2018
<i>P. bambusae</i>	aseptate or septate	ellipsoidal to obovoidal, fusiform-cylindrical to falcate, hyaline, straight, guttulate, smooth-walled	Unidentified submerged bamboo	Indonesia	Höhnel (1909), Hyde et al. (2019), Luo et al. (2019), Réblová et al. (2016)
<i>P. caffra</i>	aseptate, rarely 1-septate	conidia straight, ellipsoidal to clavate, obovoid, not attenuated at the apex, pale yellow brown, smooth	On decaying leaf of <i>Podocarpus</i>	Cape Province	Castañeda Ruiz et al. (2002), de Hoog and Papendorf (1976)

<i>P. clavulata</i>	aseptate	broadly ellipsoidal to ± spherical, subspherical, smooth, hyaline	On rotten decorticated wood	Great Britain	Castañeda Ruiz et al. (2002), de Hoog and Papendorf (1976), Mason and Ellis (1953)
<i>P. clematidis</i>	aseptate	obovoidal, rounded at the apex, obtuse and tapering towards base, hyaline, smooth-walled	Decaying wood submerged in Lancang River	China, Yunnan Province	Castañeda Ruiz et al. (2002), Hughes (1958), Luo et al. (2018)
<i>P. curvata</i>	aseptate	smooth, thin-walled, hyaline, clavate to obovoid and pointed at base, curved, occasionally sickle-shaped	rotten leaves of <i>Parinari capensis</i>	South West Africa	de Hoog and Papendorf (1976)
<i>P. dalbergiae</i>	aseptate	solitary, hyaline, smooth, thin-walled, guttulate, subcylindrical to obovoid, tapering towards both ends, apex subobtuse, base with truncate hilum	On bark of <i>Dalbergia armata</i>	South africa, Northern Province	Crous et al. (2021)
<i>P. fasciculata</i>	aseptate	ellipsoidal to obovoid, straight, rounded at the apex, obtuse and tapering towards base, hyaline, smooth-walled	Decorticated wood of <i>Sambucus nigra</i>	Canada, Ontario, Goulbourn Twp	Réblová et al. (2016)
<i>P. filiformis</i>	-	-	Decaying wood submerged in freshwater stream	Thailand, Sai khu Waterfall	Luo et al. (2019)
<i>P. glauca</i>	aseptate	smooth, thin-walled, hyaline, guttuliform to ellipsoidal, with pointed base, occasionally sickle-shaped	On rotten wood of <i>Quercus</i> sp.	America, Newfield	de Hoog and Papendorf (1976)
<i>P. guttulata</i>	aseptate	globose to obovoid, hyaline, smooth-walled, guttulate	Decaying wood submerged in Suoluo River	China, Guizhou Province	Hyde et al. (2018)

<i>P. infrafertilis</i>	aseptate, rarely 1-septate	conidia falcate, hyaline	On dead leaves of <i>Eucalyptus</i>	Brazil	de Hoog and Papendorf (1976), Sutton and Hodges (1976)
<i>P. loranthacearum</i>	-	solitary, hyaline, smooth, fusoidal-ellipsoidal with obtuse ends, straight to falcate, guttulate	On twigs of <i>Loranthus europaeus</i>	Germany	Crous et al. (2015)
<i>P. magnifica</i>	aseptate	straight, ellipsoidal to obovoid, clavate, very pale olivaceous, smooth	On <i>Bambusa</i>	New Caledonia	de Hoog and Papendorf (1976), Deighton (1974)
<i>P. microspora</i>	aseptate	solitary, fusiform, straight, smooth-walled, guttulate, hyaline	On decaying wood	Thailand, Krabi, Wat ThumSua	Hyde et al. (2017)
<i>P. muscariformis</i>	3-septate	cylindric –fusiform, subhyaline, smooth	On leaves of <i>Tiliacora kenyensis</i>	Kenya	Castañeda Ruiz et al. (2002), Siboe et al. (1999)
<i>P. pseudoclematidis</i>	aseptate	cylindric-ovate, straight, hyaline, smooth-walled, guttulate	On dead culm of bamboo (Bambusae)	Thailand, Chiang Rai	Liu et al. (2015)
<i>P. sedimenticola</i>	aseptate, 1-septate	smooth-walled, hyaline, with a pointed base, usually aseptate when attached to the conidiogenous cells, 0–1-septate after release; aseptate conidia, obovoid to ellipsoidal; 1-septate conidia, obovoid, slightly constricted at septum	Isolated from surface marine sediment in intertidal zone	China, Shandong Province	Cheng et al. (2014)
<i>P. siamensis</i>	aseptate	globose to subglobose, hyaline	Saprobic on decaying fruits	Thailand, Chiang Mai Province	Hyde et al. (2019)

<i>P. sparsa</i>	0-3-septate	fusiform to clavate, conidia straight, ellipsoidal to fusiform, hyaline, not attenuated at the apex	On bark of <i>Acer spicatum</i>	Saskatchewan	de Hoog and Papendorf (1976), Sutton (1973)
<i>P. sparsa</i> var. <i>cubensis</i>	0–1(–4)-septate	fusiform, cylindrical or clavate, hyaline, sometimes slightly curved	On dead branch	Cuba	Mercado-Sierra et al. (1997), Mel'nik (2012)
<i>P. synnematica</i>	0–1-septate	dimorphic, clavate to ellipsoidal, cylindrical to falcate, base narrowly truncate, tip obtuse, variable in size, sometimes constricted near septa, 1–2-guttulate, hyaline, smooth-walled	Dead bark of <i>Azadirachta indica</i> (Meliaceae)	India, Maharashtra	Boonmee et al. (2021)
<i>P. tuberculata</i>	aseptate, rarely 1-septate	conidia fusiform, straight, the apex attenuated, hyaline, smooth, guttulate	On Labiatae	Malawi	Castañeda Ruiz et al. (2002), de Hoog and Papendorf (1976), Sutton (1993)
<i>P. uniseptata</i>	mostly with a median septum	two-celled, fusiform, ellipsoid, hyaline, cylindrical or clavate	On dead branch	Cuba	de Hoog and Papendorf (1976), Mercado-Sierra (1984), Mel'nik (2012)
<i>P. vietnamensis</i>	a single median septum	fusiform-subcylindrical to short obovoid-subclavate, somewhat attenuated towards the base, apex obtuse, straight to slightly curved, not constricted, hyaline, smooth, often guttulate	On bark of a living unidentified liane	South Vietnam, Dong Nai Province	Mel'nik (2012)