

# Mediterranean predatory flies of the Dolichopodidae (Diptera, Empidoidea) family from Bulgaria and Turkey

Mihail Kechev<sup>1</sup>, Alper Tonguç<sup>2</sup>

<sup>1</sup>Forest Research Institute, Bulgarian Academy of Sciences, 132 St Kliment Ohridski Blvd, Sofia, Bulgaria, E-mail: mkechev@gmail.com

<sup>2</sup>Molecular Biology and Genetics Department, Faculty of Science, Muğla Sıtkı Kocman University, Muğla, Turkey, E-mail: alpertonguc@mu.edu.tr

Corresponding author: Mihail Kechev (mkechev@gmail.com)

---

Academic editor: Georgi Georgiev | Received 22 February 2024 | Accepted 2 March | Published 23 May 2024

Citation: Kechev M., Tonguç A. 2024. Mediterranean predatory flies of the Dolichopodidae (Diptera, Empidoidea) family from Bulgaria and Turkey. *Silva Balcanica* 25(1): 109-114. <https://doi.org/10.3897/silvabalcanica.25.e121464>

---

## Abstract

The paper presents information about some Mediterranean dolichopodid species and their habitat preferences, as well as distribution from the southern parts of Bulgaria and southeast Turkey. One new locality of *Sybistroma discipes* is given.

## Keywords

Diptera, Dolichopodidae, distribution, habitats, *Chrysotus hubenovii*, *Hercostomus thraciensis*, *Sybistroma discipes*.

## Introduction

The family Dolichopodidae (also called Long-legged flies) belongs to the superfamily Empidoidea of the suborder Brachycera, order Diptera. The Dolichopodidae family is widespread in all zoogeographic areas of the planet, without Antarctic. Today approximately 8300 species from 261 genera are described worldwide (Grichanov, 2023). About 700 species are known in Europe. The dolichopodids are a species with small to medium body size (1-10 mm). The color of the body is usually metallic green but some genera, for example, genus *Neurigona* are with yellow coloration. Most species of the family Dolichopodidae are found in moist habitats: on vegetation around rivers, streams, marshes, lakes, flooded forests, etc. A large number of representatives

of the genera Medetera and individual species of other genera can be found in dry places. As predators, the species of the family play an important role in the regulation of a number of pests in forests and agroecosystems.

The main purpose of this paper is to indicate the habitat preferences and distribution of some dolichopodid species in the southern, Mediterranean-influenced areas of Bulgaria, as well as in the same areas of Turkey.

## Material and methods

The species in this work were collected by means of sweeping net and Malaise traps in different years from 2007 to 2023. The species *Sybistroma discipes* was collected on 17.11.2023 with a sweeping net, leg. G. Zaemdzhikova. After collecting the specimen was put in 95% ethanol and separated from the other insects in the sample in laboratory conditions using a microscope.

## Results

The species *Chrysotus hubenovi* Kechev, Naglis, Negrobov, 2022, described from the West Rhodope Mountains from 1100 m altitude (Kechev et al., 2022), was found from Kaloferska Planina Mt. situated in the Balkan Mountains at an altitude of 530 m. The species inhabits humid grasses near river banks.

*Chrysotus pennatus* Lichtwardt, 1902 is not widespread in Bulgaria but is known with few specimens from several localities, mainly riverside and forest habitats.

*Hercostomus chetifer* (Walker, 1849) in Turkey was found in humid coniferous woodland (Tonguç, 2023). For Bulgaria the species was found near a small brook in deciduous forests (Kechev, 2021a).

*Hercostomus gracilis* (Stannius, 1831) is known from two localities in Bulgaria. The species could be found in humid habitats in deciduous forests (Kechev et al., 2014, Kechev, 2021b). In Turkey the species was collected from a wide range of habitats – streams, ponds, humid grassland, etc. (Tonguç, 2023).

*Hercostomus tanjusilus* Negrobov & Zurikov, 1988 known only from Georgia and Turkey was found in poplar and willow grove habitats (Küçükberber et al., 2017).

*Hercostomus thraciensis* Kechev & Negrobov, 2015 (Fig. 1) was first described in Bulgaria (Kechev, Negrobov, 2015), collected from the Gradina floodplain forest (situated to the west of Gradina Village) and a humid, wooded and marshy place called Basha (situated to the southwest of the town of Chirpan). In Turkey the species was found in 2017 (Küçükberber et al., 2017) in humid grassland. Grichanov & Freidberg (2018) found *H. thraciensis* in Israel.

*Poecilobothrus chrysozygos* (Wiedemann, 1817) for Turkey was found in fenlands and humid grasslands (Küçükberber et al., 2017). In Bulgaria the species was reported in the Gradina floodplain forest (Kechev, 2012a).



**Figure 1.** *Hercostomus thraciensis* Kechev & Negrobov, 2015, habitus.

*Poecilobothrus regalis* (Meigen, 1824) very abundant species in the investigated area in the lower parts, from 100 to 350 m a.s.l.

*Sybistroma discipes* (Germar, 1821) (Fig. 2) known in Bulgaria from two localities Markovo Village, situated in the West Rhodope Mountains and Shipkovo Village, in the Balkan Mountains. This study gives a new locality near the town of Gotse Delchev, along the banks of Mesta River.

*Sybistroma israelensis* (Grichanov, 2000) is known only in Israel and Turkey. In Turkey the species inhabits mixed woodland, humid coniferous woodland and humid grassland (Küçükberber et al., 2017).

*Sciapus bellus* Loew, 1873 was found in Kremen (473 m a.s.l.) and Dzherovo (275 m a.s.l.) villages with a large number of specimens near a river surrounded by a deciduous forest (Kechev, 2021b).

Known only in Italy, France, Morocco and Tunisia (Grichanov, Negrobov, 2014), *Sciapus costae* Mik, 1890 is an interesting species. It was found in two new localities in Bulgaria (Kechev, 2021a). The species was collected from the Balkan Mountains and Dzherovo village in the East Rhodopes along river banks in deciduous forests.

*Sciapus flavicinctus* (Loew, 1857) is a common species along the banks of rivers in lower areas of Bulgaria. In Turkey the species was known in the Antalya and Adana provinces (Naglis et al., 2021).



**Figure 2.** *Sybistroma discipes* (Germar, 1821), habitus.

*Teuchophorus chaetifemoratus* Pollet & Kechev, 2007 described in Bulgaria (Markovo village, West Rhodope Mts., 300 m a.s.l.) within a cool deciduous forest with clean shallow streams (Pollet, Kechev, 2007). During 2016 another locality was found near Naydenovo Village, situated in Sarnena Sredna Gora Mountain, and the habitat was the same as the first one (Kechev, 2016). In Turkey the species was collected from ponds and humid coniferous woodlands (Küçükberber et al., 2017).

*Teuchophorus cristulatus* Meuffels et Grootaert, 1990 is a typical Mediterranean species known in Sicily, Bulgaria and Turkey. Tonguc et al. (2010) announced one male specimen of this species from the Asian part of Turkey. In Bulgaria *T. cristulatus* was first recorded (Kechev, 2012b) with three males found on the banks of the Tekirska River in the town of Chirpan.

Other dolichopodids not mentioned above in the list of the investigated species in this study as *Dolichopus griseipennis* Stannius, 1831, *Campsicnemus curvipes* (Fallén, 1823), *Campsicnemus umbripennis* Loew, 1856, *Hydrophorus balticus* (Meigen, 1824), *Rhaphium caliginosum* Meigen, 1824, *Sympycnus pulicarius* (Fallén, 1823), *Sympycnus simplicipes* Becker, 1908, *Syntormon pallipes* (Fabricius, 1794), *Teuchophorus mona-*

*canthus* Loew, 1859 and *Teuchophorus spinigerellus* (Zetterstedt, 1843) are eurytopic species and could be found in a wide range of habitats and different altitude. For instance, *H. balticus* is collected from sea level to 2000 m a.s.l.

## Discussion

*Hercostomus thraciensis* is one of the most abundant species in spring, inhabiting flooded forests and humid woodlands situated along the banks of the Maritsa River in the Upper Thracian Plain. The adults are active in April and May. A large number of specimens were collected at the end of April and at the beginning of May. During the remaining months of the year, the species is not caught. As *H. thraciensis*, *P. regalis* is also a wide-spread species in this area. Found mostly around open sunny places like ponds and riverbanks in Bulgaria. In Turkey it was reported in streams, fenlands, reedbeds, humid mixed woodland, etc.

On the other hand, the species *H. chetifer* is rare for the investigated area. As mentioned above it was discovered in humid coniferous woodland in Turkey and in Bulgaria it is known with one male specimen collected near a small brook in a beech forest.

The Long-legged flies have been intensively studied in Bulgaria and Turkey for the last two decades. More than 100 species were reported for the fauna of Bulgaria and the number is the same for Turkey. Several new species were described for science. Three check-lists were published (Kechev, 2005, Tonguç, et al., 2016, Kechev, et al., 2020).

In conclusion, Dolichopodidae as a species-rich family is promising for future research in the eastern Mediterranean region. It is possible that more new species will emerge for the fauna, and as predators, it is good to study their habitats and the relationship with other organisms.

## References

- Grichanov I.Y. 2023. A new species of Thrypticus Gerstaecker, 1864 and new records of Dolichopodidae (Diptera) from Turkey. *Journal of insect biodiversity* 44 (2), 041-050. doi: 10.12976/JIB/2023.44.2.2
- Grichanov I.Y., Freidberg A. 2018. New species and new records of Dolichopodidae (Diptera) from Israel. *Russian Entomological Journal* 27 (3), 299-306. doi: 10.15298/rusentj.27.3.12
- Grichanov I., Negrobov O.P. 2014. Palaearctic species of the genus *Sciapus* Zeller (Diptera: Dolichopodidae). *VIZR RAAS (Plant Protection News Supplement)*, St Petersburg, Russia, 85 pp.
- Kechev M. 2005. Check-List of Dolichopodidae (Diptera, Brachycera) species in Bulgaria. *Animalia* 41, 47-62.
- Kechev M. 2012a. Diversity of Long-Legged Flies (Diptera: Dolichopodidae) in Gradina Floodplain Forest, Bulgaria. *Acta Zoologica Bulgarica. Suppl.*, 4, 43-46.
- Kechev M. 2012b. Long-legged flies (Diptera: Dolichopodidae) from the Upper Thracian Plain new to Bulgaria: Habitats and Distribution. *Acta zoologica bulgarica* 64 (2), 205-208.

- Kechev M. 2016. Forestry and riparian Long-legged flies (Diptera: Dolichopodidae) from Sarnena Sredan Gora Mountain with new records for Bulgaria. *Silva balcanica* 17 (2), 45-55.
- Kechev M. 2021a. Diversity of long-legged flies (Diptera, Dolichopodidae) of the Balkan Mountains (Bulgaria and Serbia). *Historia naturalis bulgarica* 42, 15-24. <https://doi.org/10.48027/hnb.42.041>
- Kechev M. 2021b. Data on the distribution of Dolichopodidae (Diptera: Empidoidea) in Bulgaria, with first records for the country. *Historia naturalis bulgarica* 42, 95-102. <https://doi.org/10.48027/hnb.42.131>
- Kechev M., Negrobov O. 2015. A new species of the genus *Hercostomus* Loew, 1857 (Diptera: Dolichopodidae) from Bulgaria. *Acta Zoologica Bulgarica* 67 (2), 199-202.
- Kechev M., Negrobov O., Grichanov I. 2014. Diversity of long-legged flies (Diptera: Dolichopodidae) along the banks of the Omurovska river (Bulgaria), with the description of a new species of the genus *Teuchophorus* Loew. *Acta Zoologica Bulgarica* 66 (3), 317-323.
- Kechev M., Naglis S., Negrobov O.P. 2022. A new species *Chrysotus hubenovii* and new data on the family Dolichopodidae (Diptera) of Bulgaria. *Zoosystematica Rossica* 31 (1), 27-41. <https://doi.org/10.31610/zsr/2022.31.1.27>
- Kechev M., Naglis S., Tonguc A., Pollet M. 2020. Checklist of the Dolichopodidae (Diptera, Empidoidea) of the Balkan Peninsula, with first records for Bulgaria, Montenegro, North Macedonia, and for the European part of Turkey. *Zootaxa* 4819 (3), 436-472.
- Küçükberber M., Tonguç A., Koç H. 2017. Dolichopodidae (Diptera) fauna of Spil Mountain with four new records. *Türkiye Entomoloji Bülteni* 7 (1), 23-29.
- Naglis S., Tonguç A., Barták M. 2021. New faunistic data on Dolichopodidae (Diptera) from Turkey. *Caucasian Entomological Bulletin* 7 (1), 121-128. DOI: 10.23885/181433262021171-121128
- Pollet M., Kechev M. 2007. Are view of Palaearctic *Teuchophorus*, with a new species from Bulgaria (Diptera, Dolichopodidae). *Zootaxa* 1592, 45-56.
- Tonguç A. 2023. Habitat preferences and abundances of the Dolichopodidae (Diptera) from inner west Anatolia. *Acta Entomologica Serbica* 28 (1), 37-43.
- Tonguç A., Grichanov I.Ya., Naglis S. 2016. Checklist of the Dolichopodidae (Diptera, Brachycera) of Turkey. *Turkish Journal of Zoology* 40, 14-26.
- Tonguç A., Grichanov I., Koç H., Özgül O., Barlas M. 2010. Contributions to the Dolichopodidae (Diptera) fauna of Turkey. *Journal of the Entomological Research Society* 12 (2), 103-107.