

# New data on the distribution of ground beetles (Coleoptera: Carabidae) in Kenya

Vladimir Sakalian<sup>1</sup>, Georgi Georgiev<sup>2</sup>

<sup>1</sup>*Institute of Biodiversity and Ecosystem Research, Bulgarian Academy of Sciences, 1 Tzar Osvoboditel Blvd., 1000 Sofia, Bulgaria*

<sup>2</sup>*Forest Research Institute, Bulgarian Academy of Sciences, 132, "St. Kliment Ohridski" Blvd.1756 Sofia, Bulgaria*

Corresponding author: Georgi Georgiev ([ggeorgiev.fri@gmail.com](mailto:ggeorgiev.fri@gmail.com))

---

Academic editor: Margarita Georgieva | Received 16 November 2022 | Accepted 23 November 2022 | Published 20 December 2022

**Citation:** Sakalian V., Georgiev G. 2023. New data on the distribution of ground beetles (Coleoptera: Carabidae) in Kenya. *Silva Balcanica* 23(2): 43-46. <https://doi.org/10.3897/silvabalcanica.22.e97628>

---

## Abstract

New data on the distribution of 12 species and subspecies of ground beetles (Coleoptera: Carabidae), which belong to 9 genera, 6 tribes and 5 subfamilies (Anthiinae, Carabinae, Cindolinae, Dryptinae and Panagaeinae) in Kenya are reported.

## Keywords

Carabidae, Anthiinae, Carabinae, Cindolinae, Dryptinae, Panagaeinae, distribution, Kenya

## Introduction

During the period 2003-2006, the first author collected many different coleopteran insect species in Kenya, mainly Buprestidae (Curletti, Sakalian, 2007, 2009; Sakalian, Georgiev, 2013; Bílý, Sakalian., 2014), but also Cerambycidae (Sakalian, Georgiev, 2011) and Tenebrionoidea (Ferrer et al., 2016). Among the other coleopteran material, some taxa belonging to Carabidae family were established.

The family Carabidae numbers more than 40,000 species worldwide. The fauna of the best studied area in the Ethiopian zoogeographical region – South Africa includes over 1,400 species (Schoeman et al., 2018), but its actual number in this country, as well as that of the Ethiopian region as a whole, is undoubtedly great-

er due to the still insufficient knowledge about ground beetles in the tropics and subtropics. A part of the carabid samples were identified by Dr. Augusto Vigna-Taglianti.

This note reports new data on the distribution of 12 carabid taxa belonging to 9 genera, 6 tribes and 5 subfamilies in Kenya.

## Materials and Methods

The ground beetles were collected by traditional entomological methods: hand collection on the ground and attracting to lamp light.

The current status, synonyms and combinations concerning Kenyan taxa are reported according to Lorenz (2005). The names of the subfamilies, tribes and species and subspecies are listed in alphabetical order.

The studied material is deposited in the Scientific Fund of Institute of Biodiversity and Ecosystem Research of Bulgarian Academy of Sciences (Sofia, Bulgaria).

## Results and Discussion

### Subfamily Anthiinae Bonelli, 1813

#### Tribe Anthiini Bonelli, 1813

##### *Anthia (Anthia) artemis* Gerstaecker, 1884 (Fig. 1A)

Elementaita Lake, 00°28'31"S, 36°15'46"E, 1820 m, 14-15.04.2006, 1 ex., V. Sakalian leg.

##### *Anthia (Termophilum) hexastictum hexastigmum* (Gerstaecker, 1866) (Fig. 1B)

Voi Wildlive Lodge, 600 m, 04-06.11.2005, 2 ex.; 27-28.04.2006, 1 ex., V. Sakalian leg.

##### *Chilanthia cavernosa* (Gerstaecker, 1866)

Voi Wildlive Lodge, 600 m, 04-06.11.2005, 1 ex.; Kipepeo Farm near Malindi, 24.04.2006, 1 ex., V. Sakalian leg.

##### *Cypholoba chanleri* (Linell, 1896)

Lower Tana River, Gamba Guest House, 25-27.10.2005, 2 ex.; 20-23.04.2006, 4 ex., V. Sakalian leg.

##### *Cypholoba cinereocincta alluaudi* (Sternberg, 1907)


Voi Wildlive Lodge, 600 m, 04-06.11.2005, 2 ex.; 27-28.04.2006, 4 ex., V. Sakalian leg.

##### *Cypholoba sparthulata* (Gerstaecker, 1866)

Lower Tana River, Gamba Guest House, 25-27.10.2005, 2 ex.; 20-23.04.2006, 1 ex.; Voi Wildlive Lodge, 600 m, 27-28.04.2006, 1 ex., V. Sakalian leg.

##### *Cypholoba tetrastigma tetrastigma* (Chaudoir, 1848)

Voi Wildlive Lodge, 600 m, 27-28.04.2006, 1 ex., V. Sakalian leg.



**Figure 1.** Ground beetles collected: A – *Anthia artemis*; B – *Anthia hexastictum hexastigmum*; C – *Triaenogenius gerstaeckeri*; D – *Calosoma planicolle*; E – *Megacephala regalis excelsa*; F – *Tefflus zebulianus transitionis*

**Tribe Helliunini Hope, 1838**

*Triaenogenius (Triaenogenius) gerstaeckeri* (Chaudoir, 1877) (Fig. 1C)

Voi Wildlive Lodge, 600 m, 04-06.11.2005, 2 ex., V. Sakalian leg.

**Subfamily Carabinae Latreille, 1802**

**Tribe Carabini Latreille, 1802**

*Calosoma (Ctenosta) planicolle* Chaudoir, 1869 (Fig. 1D)

Voi Wildlive Lodge, 600 m, 27-28.04.2006, 1 ex., V. Sakalian leg.

**Subfamily Cindelinae Latreille, 1802**

**Tribe Megacephalini Laporte, 1834**

*Megacephala regalis excelsa* Bates, 1874 (Fig. 1E)

Kipepeo Farm near Malindi, 24.04.2006, 2 ex., V. Sakalian leg.

**Subfamily Dryptinae Bonelli, 1810**

**Tribe Galeritini Leconte, 1853**

***Galerita (Galerita) angustipennis Gerstaecker, 1867***

Lower Tana River, Gamba Guest House, 20-23.04.2006, 1 ex., V. Sakalian leg.

**Subfamily Panagaeinae Bonelli, 1810**

**Tribe Panagaeini Bonelli, 1810**

***Tefflus (Tefflus) zebulianus transitionis Kolbe, 1903 (Fig. 1F)***

Voi Wildlive Lodge, 600 m, 27-28.04.2006, 1 ex., V. Sakalian leg.

We consider that this short list is only a small contribution to a better understanding of the species diversity of the Kenyan Carabidae. We expect a lot of new data to be established to improve our knowledge on the very interesting fauna of this country.

## Acknowledgements

We would like to express our gratitude to the late famous entomologist Dr. Augusto Vigna-Taglianti (Rome, Italy) for determining of our material.

## References

- Bilý S., Sakalian V.P. 2014. A revision of the *Anthaxia (Haplanthaxia) mashuna* species-group (Coleoptera: Buprestidae: Buprestinae). Acta entomologica musei nationalis Pragae 54(2), 605-621.
- Curletti G., Sakalian V. 2007. Two new species of the genus *Agrilus* Curtis, 1825 from Kenya (Coleoptera: Buprestidae). Deutsche Entomologische Zeitschrift, 54(1), 75-78.
- Curletti G., Sakalian V. 2009. Sixteen new species of the genus *Agrilus* Curtis, 1825 from East Africa (Coleoptera: Buprestidae). ZooKeys 24, 1-29.
- Ferrer J., Sakalian V., Georgiev G. 2016. Darkling and ironclad beetles (Coleoptera: Tenebrionoidea) of Kenya, with description of two new species. Acta zoologica bulgarica 68(2), 159-170.
- Lorenz W. 2005. Systematic List of Extant Ground Beetles of the World: (Coleoptera "Geadephaga": Trachypachidae and Carabidae, incl. Paussinae, Cicindelinae, Rhysodinae). 2nd ed., Tutzing, 530 pp.
- Sakalian V., Georgiev G. 2011. Contribution to the Knowledge of Longhorn Beetles (Coleoptera, Cerambycidae) of Kenya. Biodiversity Journal 2(2), 67-72.
- Sakalian V., G. Georgiev. 2013. New data about the diversity of jewel beetles (Coleoptera: Buprestidae) of Kenya. Acta zoologica bulgarica 65(4), 457-460.
- Schoeman C.S., Tshililo P., Foord S.H., Hamer M. 2018. Annotated checklist of Carabidae (Insecta: Coleoptera) of the Vhembe biosphere reserve, South Africa. Project: Patterns in ground-dwelling arthropod diversity in a biosphere reserve: implications for conservation. DOI: 10.13140/RG.2.2.33610.06089