

Molecular characterization and phylogenetic position of the giant deep-sea oyster *Neopycnodonte zibrowii* Gofas, Salas & Taviani, 2009

Matteo Garzia¹, Daniele Salvi¹

¹ Department of Health, Life & Environmental Sciences - University of L'Aquila, Via Vetoio snc, 67100 L'Aquila-Coppito, Italy

<https://zoobank.org/7DB92668-7DCA-4F0D-B6AB-8C74C3EC6B89>

Corresponding author: Daniele Salvi (danielesalvi.bio@gmail.com)

Academic editor: M. Glaubrecht ♦ Received 13 November 2023 ♦ Accepted 3 January 2024 ♦ Published 26 January 2024

Abstract

The giant deep-sea oyster *Neopycnodonte zibrowii* Gofas, C. Salas & Taviani, 2009 is a keystone deep-sea habitat builder species. Discovered about fifteen years ago in the Azores, it has been described and assigned to the genus *Neopycnodonte* Fischer von Waldheim, 1835 based on morphological features. In this study, we generated DNA sequence data for both mitochondrial (COI and 16S) and nuclear (ITS2 and 28S) markers based on the holotype specimen of *N. zibrowii* to establish a molecular phylogenetic framework for the systematic assessment of this species and to provide a reliable (i.e., holotype-based) reference sequence set for multilocus DNA barcoding approaches. Molecular data provide compelling evidence that the giant deep-sea oyster is a distinct species, rather than a deep-water ecophenotype of *Neopycnodonte cochlear* (Poli, 1795), with extremely high genetic divergence from any other gryphaeid. Multilocus phylogenetic analyses place the giant deep-sea oyster within the clade “*Neopycnodonte/Pycnodonte*” with closer affinity to *N. cochlear* rather than to *P. taniguchii* Hayami & Kase, 1992, thus supporting its assignment to the genus *Neopycnodonte*. Relationships within this clade are not well supported because mitochondrial variation is inflated by saturation that eroded phylogenetic signal, implying an old split between taxa within this clade. Finally, the set of reference barcode sequences of *N. zibrowii* generated in this study will be useful for a wide plethora of barcoding applications in deep-sea biodiversity surveys. Molecular validation of recent records of deep-sea oysters from the Atlantic Ocean and the Mediterranean Sea will be crucial to clarify the distribution of *N. zibrowii* and assess the phenotypic variation and ecology of this enigmatic species.

Key Words

Azores, DNA sequences, Gryphaeidae, holotype, molecular systematics, Mollusca, multilocus phylogeny, Natural History Museum

Introduction

Deep-sea is the Earth's largest biome but it is still one of the most underexplored regions (Ramirez-Llodra et al. 2010). Deep-sea biodiversity is mostly unknown due to the extreme environmental conditions that limits sampling capabilities (Rogers et al. 2015; Sinniger et al. 2016; Woodall et al. 2018). Along with advances in exploration technologies (Feng et al. 2022), new molecular technologies such as high-throughput sequencing and the molecular identification of multiple species in environmental DNA (eDNA metabarcoding; Taberlet et al. 2012) have boosted deep-sea biodiversity assessments (Guardi-

ola et al. 2016; Everett and Park 2018). However, the low number of reference sequences taxonomically validated in online repository databases (e.g. GenBank) limits the identifications of MOTUs (Molecular Operational Taxonomic Unit), thus reducing the taxonomic resolution of eDNA studies (Ruppert et al. 2019). Studies using an integrative taxonomic approach – combining molecular, morphological and environmental data – on new deep-sea taxa have been carried out in several groups of organisms, such as Anthozoa (López-González et al. 2022), Mollusca (Xu et al. 2019), and some others (Silva et al. 2016; Błażewicz et al. 2019). However, while molecular data are still not available for a great portion of known

deep-sea biodiversity, deep-sea exploration has continued, contributing to the discovery of new benthic ecosystems and associated communities. Therefore, there is a great need to constantly improve with reliable reference sequences the taxonomic coverage of deep-sea taxa in repository databases.

In this study, we focused on a keystone deep-sea habitat builder species discovered about fifteen years ago in the Azores Archipelago: the giant deep-sea oyster *Neopycnodonte zibrowii* Gofas, C. Salas & Taviani, 2009 (Gryphaeidae Vialov, 1936) (Wisshak et al. 2009b). This reef-forming oyster was first observed during a submersible dive along the Faial Channel (480–500 meter depth) (Wisshak et al. 2009b). Deep-sea reefs of *N. zibrowii* are built by both stacked living and dead specimens on vertical rocky substrate of seamounts, escarpments and in canyons (Beuck et al. 2016), and host peculiar deep-sea communities. Benthic associations between *N. zibrowii* and the cyrtocrinid *Cyathidium foresti* Chérbonnier & Guille, 1972 have been documented in the Atlantic Ocean (Wisshak et al. 2009a), and between *N. zibrowii* and cold-water corals in both the Atlantic Ocean (Van Rooij et al. 2010) and the Mediterranean Sea (Taviani et al. 2017, 2019). Recently, new records and observations on *N. zibrowii* in the Atlantic Ocean allowed updating its ecology and distribution (Beuck et al. 2016). The giant deep-sea oyster has been meticulously described in terms of external morphology, microstructures of shell and anatomy (Wisshak et al. 2009b). The systematic placement of this species in the genus *Neopycnodonte* was based on morphological characteristics such as the circular muscle scar, the enlarged vermiculate chomata (see ‘neopycnodontine chomata’ in Harry 1985) and the vesicular structures in the inner shell layer. *Neopycnodonte zibrowii* is morphologically different from the only extant congeneric species *Neopycnodonte cochlear* (Poli, 1795) in several characters such as the shell architecture and outline, the absence of the resilifer bulge in the latter species and the shape and thickness of the vesicular microstructures. On the other hand, 15 years on from its discovery, molecular data are still not available for this species, thus limiting the assessment of its phylogenetic position and systematic placement.

The taxonomic assessment of oysters based on morphology can be challenging due to a high shell variability and a low number of diagnostic characters (Lam and Morton 2006; Raith et al. 2015; Salvi et al. 2021). Molecular data have a key role in species delimitation and taxonomic identification of oyster species (Lam and Morton 2003; Bieler et al. 2004; Kirkendale et al. 2004; Al-Kandari et al. 2021; Salvi et al. 2022) and would provide compelling evidence that the giant deep-sea oyster *N. zibrowii* is a distinct species rather than a deep-water ecophenotype of *N. cochlear* (Wisshak et al. 2009b).

In this study, we generated DNA sequence data of the giant deep-sea oyster *N. zibrowii* for both mitochondrial and nuclear markers based on the holotype and performed

a multilocus phylogenetic analyses to establish its relationships with other gryphaeids. The main aims of this study are to provide: (i) a molecular phylogenetic framework for the systematic assessment of the giant deep-sea oyster, and (ii) a reliable (i.e., holotype-based) reference sequence set for multilocus DNA barcoding approaches.

Materials and methods

Specimens and sequence data gathering

We gathered tissue samples for molecular analyses from museum collections and by field collection. The holotype of *N. zibrowii* (MNHN-IM-2000-20888) and the specimen of *Hyotissa numisma* (Lamarck, 1819) (MNHN-IM-2013-13700) are deposited at the National Museum of Natural History (MNHN) of Paris, while the specimen of *Pycnodonte taniguchii* Hayami & Kase, 1992 (UF 280382) is preserved in the collection of Florida Museum of Natural History (FLMNH). *Neopycnodonte cochlear* (OS239) was collected during scuba diving off the coast of Civitavecchia (nearby Rome, Italy) and stored in pure ethanol. Total genomic DNA was extracted from adductor muscles following standard high-salt protocols (Sambrook et al. 1989). We amplified two mitochondrial – cytochrome oxidase subunit I (COI) and 16S rRNA (16S) – and two nuclear – 28S rRNA (28S) and ITS2 rRNA (ITS2) – gene fragments by polymerase chain reaction (PCR). Primers and conditions used for the amplification are reported in Table 2. Sequencing of PCR products was carried out by the company Genewiz® (<https://www.genewiz.com>), using the same primers employed for amplification. Sequences generated from these specimens were complemented with sequences obtained from GenBank for additional gryphaeid species. Localities and GenBank accession numbers of sequences used for molecular analyses are shown in Table 1. GenBank sequences were selected in order to minimise the use of chimeric sequences in concatenated alignments (i.e., sequences of different gene fragments obtained from different voucher specimens), therefore whenever possible for each species we selected mitochondrial (COI and 16S) and nuclear (28S and ITS2) sequences from the same voucher. Three specimens (*Hyotissa hyotis* #2, *Hyotissa imbricata* and *N. cochlear* #1) have GenBank sequences from different vouchers (chimeric concatenated sequences). We validated the taxonomic identification of each of these vouchers based on single-gene NJ trees. First, we built four single-gene datasets (COI, 16S, 28S and ITS2) including all the sequences of Gryphaeidae species in GenBank and our sequences. Then for each marker we selected GenBank sequences that clustered within the same clade of conspecific vouchers we sequenced (*H. hyotis* #1 and *N. cochlear* #1) or that have a congruent phylogenetic placement among the four single-gene datasets (*H. imbricata*) (results not shown).

Table 1. Details on the species and DNA sequence data used in this study. Asterisks indicate specimens sequenced in this study. GenBank data are as follows: ¹: Matsumoto 2003; ²: Matsumoto and Hashimoto unpublished; ³: Kirkendale et al. 2004; ⁴: Plazzi and Passamonti 2010; ⁵: Kim et al. 2009; ⁶: Plazzi et al. 2011; ⁷: Li et al. unpublished; ⁸: Ren et al. 2016; ⁹: Salvi et al. 2014; ¹⁰: Ip et al. 2022.

Specimen	Locality	Genbank accession number			
		COI	16S	28S	ITS2
<i>Hyotissa hyotis</i> #1	Madagascar	GQ166583 ⁶	GQ166564 ⁶	–	–
<i>Hyotissa hyotis</i> #2	Singapore (COI); Maldives (16S and ITS2)	OM946450 ¹⁰	LM993886 ⁸	–	LM993876 ⁹
<i>Hyotissa imbricata</i>	Japan: Okinawa (COI and ITS2); China: Beibu Bay (16S and 28S)	AB076917 ¹	KC847136 ⁷	KC847157 ⁷	AB102758 ²
<i>Hyotissa numisma</i> #1	Guam	–	AY376598 ⁴	AF137035 ³	–
<i>Hyotissa numisma</i> #2 *	Papua New Guinea: Rempia Area	–	PP070396	PP070400	–
<i>Neopycnodonte cochlear</i> #1	Italy: Mediterranean Sea (COI, 16S and ITS2)	JF496772 ⁶	JF496758 ⁶	–	LM993878 ⁹
<i>Neopycnodonte cochlear</i> #2 *	Italy: Civitavecchia	PP069758	PP070397	PP070401	PP074322
<i>Neopycnodonte zibrowii</i> *	Azores: Faial Channel	PP069759	PP070398	PP070402	PP074323
<i>Pycnodonte taniguchii</i> #1	Japan: Okinawa	AB076916 ¹	–	AB102759 ²	–
<i>Pycnodonte taniguchii</i> #2 *	Indonesia: Sulawesi Island	PP069760	PP070399	PP070403	PP082050
<i>Magallana gigas</i> (outgroup)	Japan (COI, 16S and 28S); South Korea (ITS2)	KJ855241 ⁸	KJ855241 ⁸	AB102757 ²	EU072458 ⁵

Table 2. Primers used in this study: forward primers are listed above and reverse primers below. For the COI and ITS2 gene fragments we designed new primers specific to *Ostreoidea Rafinesque, 1815*, and we used the following PCR cycling conditions: denaturation step: 94 °C / 3 min; 35 cycles of: 94 °C / 60 s, T° annealing (COI: 49 °C; ITS2: 50 °C) / 60 s, 72 °C / 60 s; final extension: 10 min at 72 °C.

Gene	Primer	Sequence	Reference	Notes
COI	MolIF	5' – ATAATYGGNGNTTTGGNAAAYTG – 3'	This study	Dr Zuccon D. (MNHN), pers. comm. Salvi et al., in prep
	osHCO998-R	5' – ACRGTIGCIGCICTRAARTAAGCICG – 3'		
16S	16Sar-L	5' – CGCCTGTTTATCAAAAACAT – 3'	Salvi et al. (2010)	
	16Sbr-H	5' – CCGGTCTGAACTCAGATCAC – 3'		
28S	D1F-OS	5' – GAGACTACGCCCTGAACTTAAGCAT – 3'	This study Salvi et al. (2022)	
	D6R-OS	5' – GCTATCCTGAGGAAACYTCAGAGG – 3'		
ITS2	its3d-OS	5' – GGGTCGATGAAGARCGCAGC – 3'	This study	Modified from Oliverio and Mariottini (2001)
	its4r-OS	5' – CCTAGTTAGTTTCTTTCTCTGC – 3'		

Phylogenetic analyses

Newly generated sequences for each marker were used as query in BLAST searches (blastn algorithm) using default settings to evaluate contaminants and to confirm the identification of the specimens from family to species level. Multiple sequence alignments of each marker were performed with MAFFT v.7 (Katoh et al. 2019) using the G-INS-I iterative refinement algorithm for the COI and the E-INS-i iterative refinement algorithm for the rRNA markers. GBlocks (Castresana 2000) was used to remove poorly aligned and ambiguous position of the hypervariable regions of the rRNA alignments using a relaxed selection of blocks (Talavera and Castresana 2007). Single-gene alignments were concatenated using the software SequenceMatrix (Vaidya et al. 2011).

Phylogenetic relationships were inferred using Maximum Likelihood (ML) and Bayesian Inference (BI) methods. We used the oyster *Magallana gigas* (Thunberg, 1793) as outgroup based on previous phylogenetic studies (Tëmkin 2010; Plazzi et al. 2011). ML analyses were performed in the W-IQ-TREE web server v.1.6.12 [<http://iqtree.cibiv.univie.ac.at/>]; (Trifinopoulos et al. 2016)] based on a partitioned substitution

model. For each gene partition, the best substitution model was calculated by the ModelFinder module (Kalyaanamoorthy et al. 2017) using an edge-linked model and the BIC criterion (COI: TPM2u+F+G4; 16S: HKY+F+G4; 28S: TN+F+G4; ITS2: K2P+G4). ML analysis was performed with 1,000 pseudo-replicates of ultrafast bootstrapping [uBS; (Minh et al. 2013)]. Bayesian analyses (BA) were carried out with MrBayes v.3.2.7 (Ronquist et al. 2012), using the substitution models selected by ModelFinder for each gene partition. We ran two Markov chains of two million generations each, with a sample frequency of 200 generations. Convergence of the runs (ESS values > 200) were checked with Tracer 1.7 (Rambaut et al. 2018) after a burn-in of 25%. Nodal support was estimated as Bayesian posterior probability (BPP). FigTree v.1.4.4 (<http://tree.bio.ed.ac.uk/software/figtree/>) was used to visualize both ML and BI trees.

Genetic divergence between species at each marker (COI, 16S, 28S and ITS2) were calculated using both uncorrected genetic distance (*p*-distance) and genetic distance corrected under the Kimura 2-parameter model (K2P-distance) using the software Mega11 and the option “Compute Between Groups Mean Distance” (Tamura et al. 2021).

Results

BLAST searches using mitochondrial sequences (COI and 16S) of the newly sequenced specimens of *H. numisma*, *N. cochlear* and *P. taniguchii* confirmed the taxonomic identifications of these species (sequence identity of 99–100%). BLAST searches using the mitochondrial sequences generated from the holotype of *N. zibrowii* recovered as best hits sequences belonging to Gryphaeidae species (COI: sequence identity of 73.2%/72.5%/73.1% with GenBank sequences of *Hyotissa* sp./*Neopycnodonte* sp./*Pycnodonte* sp. respectively; 16S: sequence identity of 87.3%/87.5% with GenBank sequences of *Hyotissa* sp./*Neopycnodonte* sp. respectively). This confirms the lack of contamination during the amplification and the affiliation of this species to Gryphaeidae.

The concatenated dataset included 2409 positions (COI: 455, 16S: 449, 28S: 1078, ITS2: 427 positions) and among the 828 variable positions 436 were phylogenetically informative (i.e., parsimony informative). Maximum likelihood and Bayesian trees show two main clades: one including *Hyotissa* species (uBS = 95; BPP = 1), and the other one including *Pycnodonte* and *Neopycnodonte* species (uBS = 83; BPP = 0.94) (Fig. 1). *Neopycnodonte zibrowii* is nested within the second clade with a sister relationship with *N. cochlear* (uBS = 56; BPP = 0.72), whereas *P. taniguchii* is sister to *Neopycnodonte* species.

The COI genetic distances (K2P/*p*-distance) between *N. zibrowii* and *N. cochlear* and between *N. zibrowii* and *P. taniguchii* are respectively 35.8%/28.2% and 35%/27.6% (Table 3). The 16S genetic distances (K2P/*p*-distance) between *N. zibrowii* and either *N. cochlear* or *P. taniguchii* are 13.5%/12.1% (Table 3). The mean interspecific genetic distances (K2P/*p*-distance) among the six gryphaeid species are $33.7\% \pm 4.6\%$ / $26.8\% \pm 3\%$ at the COI and $15.5\% \pm 4.6\%$ / $13.7\% \pm 3.7\%$ at the 16S. The 28S genetic distances (K2P/*p*-distance) between *N. zibrowii* and *N. cochlear* and between *N. zibrowii* and *P. taniguchii* are respectively 2.5%/2.4% and 9%/8.4% (Table 4). The ITS2 genetic distances (K2P/*p*-distance) between *N. zibrowii* and *N. cochlear* and between *N. zibrowii* and *P. taniguchii* are respectively 15.8%/14.9% and 38.2%/29.6% (Table 4). The mean interspecific genetic distances (K2P/*p*-distance) among the six gryphaeid species are $5.5\% \pm 2.5\%$ / $5.1\% \pm 2.2\%$ at the 28S and $27.4\% \pm 16.8\%$ / $22.0\% \pm 11.9\%$ at the ITS2.

Discussion

Benthic organisms such as oysters, with extensive phenotypic variation and few diagnostic characters, are prone to misidentification in morphological assessments. The utility of molecular characters for taxonomic identification




Figure 1. Bayesian phylogenetic tree of six Gryphaeidae species based on COI, 16S, 28S and ITS2 markers. Nodal supports indicate the values of uBS (upper) and the BPP (lower). The tree is rooted with *Magallana gigas* which belongs to the sister family Ostreidae Rafinesque, 1815. Specimens sequenced in this study are highlighted in bold.

Table 3. Mean genetic distance based on COI (lower triangular matrix) and 16S (upper triangular matrix) DNA sequences, calculated using the K2P model (first value) and uncorrected (*p*-distance: value inside brackets). The COI and 16S dataset are composed by 2 sequences for each species, except for *N. zibrowii* (one sequence for each marker), *H. imbricata* (one COI and one 16S sequence) and *P. taniguchii* (one 16S sequence), see Table 1; n. a.: not available.

	<i>Neopycnodonte zibrowii</i>	<i>Neopycnodonte cochlear</i>	<i>Pycnodonte taniguchii</i>	<i>Hyotissa hyotis</i>	<i>Hyotissa numisma</i>	<i>Hyotissa imbricata</i>
<i>Neopycnodonte zibrowii</i>	–	13.5% (12.1%)	13.5% (12.1%)	15.1% (13.4%)	23.5% (19.9%)	14.9% (13.3%)
<i>Neopycnodonte cochlear</i>	35.8% (28.2%)	–	11.2% (10.3%)	15.5% (13.8%)	22.9% (19.5%)	14.9% (13.4%)
<i>Pycnodonte taniguchii</i>	35.0% (27.6%)	35.4% (28.1%)	–	14.9% (13.3%)	22.3% (19.0%)	14.2% (12.7%)
<i>Hyotissa hyotis</i>	33.3% (26.7%)	39.6% (30.5%)	32.7% (26.5%)	–	15.6% (13.9%)	5.3% (5.1%)
<i>Hyotissa numisma</i>	n. a.	n. a.	n. a.	n. a.	–	15.1% (13.6%)
<i>Hyotissa imbricata</i>	34.0% (27.2%)	37.0% (28.8%)	31.8% (25.8%)	22.4% (19.1%)	n. a.	–

Table 4. Mean genetic distance based on 28S (lower triangular matrix) and ITS2 (upper triangular matrix) DNA sequences, calculated using the K2P model (first value) and uncorrected (*p*-distance: value inside brackets). The 28S and ITS2 dataset are composed by 2 sequences for each species, except for *N. zibrowii* (one sequence for each marker), *H. hyotis* (one ITS2 sequence and no 28S sequence), *N. cochlear* (one 28S sequence) and *P. taniguchii* (one ITS2 sequence), see Table 1. The ITS2 sequence of *H. imbricata* was not used for genetic distance calculation because it was too short; n. a.: not available.

	<i>Neopycnodonte zibrowii</i>	<i>Neopycnodonte cochlear</i>	<i>Pycnodonte taniguchii</i>	<i>Hyotissa hyotis</i>	<i>Hyotissa numisma</i>	<i>Hyotissa imbricata</i>
<i>Neopycnodonte zibrowii</i>	–	15.8% (14.9%)	38.2% (29.6%)	23.8% (20.4%)	n. a.	n. a.
<i>Neopycnodonte cochlear</i>	2.5% (2.4%)	–	43.5% (33.0%)	20.9% (19.4%)	n. a.	n. a.
<i>Pycnodonte taniguchii</i>	9.0% (8.4%)	4.4% (4.2%)	–	48.4% (35.4%)	n. a.	n. a.
<i>Hyotissa hyotis</i>	n. a.	n. a.	n. a.	–	n. a.	n. a.
<i>Hyotissa numisma</i>	6.8% (6.5%)	4.4% (4.2%)	7.3% (6.9%)	n. a.	–	n. a.
<i>Hyotissa imbricata</i>	6.8% (6.5%)	4.6% (4.4%)	7.7% (6.3%)	n. a.	1.2% (1.2%)	–

and systematic assessment of these organisms cannot be overstated and has been proven over and over by studies on true oysters (Lam and Morton 2006; Raith et al. 2015; Salvi et al. 2021), pearl oysters (Cunha et al. 2011), tree oysters (Garzia et al. 2022) as well as gryphaeid oysters (Li et al. 2023). Conchological convergence, phenotypic plasticity, and the occurrence of cryptic species make molecular taxonomic validation of new oyster species necessary to accurately estimate the diversity of these taxa.

Our molecular phylogenetic results clearly demonstrate that *N. zibrowii* is a distinct species with extremely high genetic divergence from any other gryphaeid at all the markers analysed (Tables 3, 4). *Neopycnodonte zibrowii* is nested within the clade “*Neopycnodonte/Pycnodonte*” with closer affinity to *N. cochlear* rather than *P. taniguchii* (Fig. 1), and thus supporting its assignment to the genus *Neopycnodonte* Fischer von Waldheim, 1835 based on morphological features (Wisshak et al. 2009b). Phylogenetic relationships within this clade are not well-supported, like in a previous phylogenetic

study including *N. cochlear* and *P. taniguchii* and based on COI and 28S markers (Li et al. 2021). However, the extended dataset of our study improved nodal support and allowed us to clarify the source of phylogenetic uncertainty. Indeed, at both mitochondrial markers, values of pairwise genetic distance between *N. cochlear* / *N. zibrowii* / *P. taniguchii* are similar and remarkably high (COI: 35.0–35.8%; 16S: 11.2–13.5%); whereas at nuclear markers the genetic distance between *P. taniguchii* and *N. zibrowii* is two-three times higher than between the latter and *N. cochlear* (Table 4). Such a pattern suggests that mitochondrial variation is inflated by saturation that eroded phylogenetic signal, implying an old split between taxa within this clade. Wisshak et al. (2009b) highlighted a low number of morphological and ecological differences between the genus *Neopycnodonte* and *Pycnodonte* Fischer von Waldheim, 1835 and pointed out the need for a systematic revision of the genera. Our results highlight that nuclear data will have a key role in further systematic assessment of these genera.

The availability of taxonomically validated reference sequence is a premise for DNA barcoding and metabarcoding approaches for large-scale, fast, and cost-effective molecular taxonomic identification (Hebert et al. 2003; Moritz and Cicero 2004; Schindel and Miller 2005; Salvi et al. 2020). The mitochondrial COI and 16S are the most common markers in DNA barcoding studies on Ostreidae and Gryphaeidae (Lam and Morton 2003, 2004, 2006; Kirkendale et al. 2004; Liu et al. 2011; Hsiao et al. 2016; Salvi et al. 2021). However, also nuclear rRNA markers such as 28S (Mazón-Suástegui et al. 2016) and ITS2 (Salvi et al. 2014; Salvi and Mariottini 2017) have proven useful for molecular taxonomic identification of oysters. Moreover, rRNA markers are frequently selected as target genes in eDNA metabarcoding projects (Ruppert et al. 2019). In this respect, the set of four reference (holotype-based) barcode sequences of *N. zibrowii* provided in this study will be useful for a wide plethora of barcoding applications in deep-sea biodiversity surveys. During the last decade, deep-sea oysters from a mounting number of regions across the Atlantic Ocean and the Mediterranean Sea have been morphologically identified as *N. zibrowii*: from Bay of Biscay (Van Rooij et al. 2010), Gulf of Cadiz (Gofas et al. 2010), Celtic Sea (Johnson et al. 2013), Angola and Mauritania (Beuck et al. 2016), southern Sardinia (Taviani et al. 2017), Sicilian Channel (Rueda et al. 2019) and Gulf of Naples (Taviani et al. 2019). Molecular validation of these records will be crucial to clarify the distribution of *N. zibrowii* and assess the phenotypic variation and ecology of this enigmatic species. Finally, given the high prevalence of cryptic species in oysters, it is not unlikely that future molecular assessments of deep-sea oysters will disclose new species.

Acknowledgements

The authors wish to thank Philippe Bouchet (Muséum National d'Histoire Naturelle, Paris) and Serge Gofas (Universidad de Málaga) for the access to holotype material, Paolo Mariottini for the specimen of *Neopycnodonte cochlear* and to Gustav Paulay (FLMNH) for the specimens of *Pycnodonte taniguchii*. MG is supported by the SYNTHESYS Project “Synthesys FR-TAF Call3_040” at The Muséum National d'Histoire Naturelle (MNHN, Paris) and the research grant of the Symposium in honour of Philippe Bouchet (World Congress of Malacology 2022 - LMU Munich). This work is part of the PhD thesis of MG under the supervision of DS (“Health and Environmental Sciences” PhD Program, University of L'Aquila).

References

Al-Kandari M, Oliver PG, Salvi D (2021) Molecular and morphological systematics of a new, reef forming, cupped oyster from the northern Arabian Gulf: *Talonostrea salpinx* new species. *ZooKeys* 1043: 1–20. <https://doi.org/10.3897/zookeys.1043.66992>

- Beuck L, Aguilar R, Fabri MC, Freiwald A, Gofas S, Hebbeln D, López Correa M, Ramos A, Ramil F, Sánchez F, Taviani M, Wienberg C, Wisshak M, Zibrowius H (2016) Biotope characterisation and compiled geographical distribution of the deepwater oyster *Neopycnodonte zibrowii* in the Atlantic Ocean and Mediterranean Sea. In: Centro Oceanográfico de Vigo. <http://www.repositorio.ieo.es/e-ieo/handle/10508/10677>
- Bieler R, Mikkelsen PM, Lee T, Foighil DÓ (2004) Discovery of the Indo-Pacific oyster *Hyotissa hyotis* (Linnaeus, 1758) in the Florida Keys (Bivalvia: Gryphaeidae). *Molluscan Research* 24(3): 149–159. <https://doi.org/10.1071/MR04013>
- Błażewicz M, Józwiak P, Jennings RM, Studzian M, Frutos I (2019) Integrative systematics and ecology of a new deep-sea family of tanaidacean crustaceans. *Scientific Reports* 9(1): 18720. <https://doi.org/10.1038/s41598-019-53446-1>
- Castresana J (2000) Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. *Molecular Biology and Evolution* 17(4): 540–552. <https://doi.org/10.1093/oxfordjournals.molbev.a026334>
- Cunha RL, Blanc F, Bonhomme F, Arnaud-Haond S (2011) Evolutionary patterns in pearl oysters of the genus *Pinctada* (Bivalvia: Pteriidae). *Marine Biotechnology* (New York, N.Y.) 13(2): 181–192. <https://doi.org/10.1007/s10126-010-9278-y>
- Everett MV, Park LK (2018) Exploring deep-water coral communities using environmental DNA. *Deep-sea Research, Part II, Topical Studies in Oceanography* 150: 229–241. <https://doi.org/10.1016/j.dsr2.2017.09.008>
- Feng J-C, Liang J, Cai Y, Zhang S, Xue J, Yang Z (2022) Deep-sea organisms research oriented by deep-sea technologies development. *Science Bulletin* 67(17): 1802–1816. <https://doi.org/10.1016/j.scib.2022.07.016>
- Garzia M, Furfaro G, Renda W, Rosati A-M, Mariottini P, Giacobbe S (2022) Mediterranean spreading of the bicolor purse oyster, *Isognomon bicolor*, and the chicken trigger, *Malleus* sp., vs. the Lessepsian prejudice. *Mediterranean Marine Science* 23(4): 777–788. <https://doi.org/10.12681/mms.29218>
- Gofas S, Rueda JL, Salas C, Díaz-Del-Río V (2010) A new record of the giant deep-sea oyster *Neopycnodonte zibrowii* in the Gulf of Cadiz (south-western Iberian Peninsula). *Marine Biodiversity Records* 3: e72. <https://doi.org/10.1017/S1755267210000618>
- Guardiola M, Wangensteen OS, Taberlet P, Coissac E, Uriz MJ, Turon X (2016) Spatio-temporal monitoring of deep-sea communities using metabarcoding of sediment DNA and RNA. *PeerJ* 4: e2807. <https://doi.org/10.7717/peerj.2807>
- Harry HW (1985) Synopsis of the supraspecific classification of the living oysters (Gryphaeidae and Ostreidae). *The Veliger* 28: 121–158.
- Hebert PDN, Cywinska A, Ball SL, deWaard JR (2003) Biological identifications through DNA barcodes. *Proceedings. Biological Sciences* 270(1512): 313–321. <https://doi.org/10.1098/rspb.2002.2218>
- Hsiao S-T, Chuang S-C, Chen K-S, Ho P-H, Wu C-L, Chen CA (2016) DNA barcoding reveals that the common cupped oyster in Taiwan is the Portuguese oyster *Crassostrea angulata* (Ostreoida; Ostreidae), not *C. gigas*. *Scientific Reports* 6(1): 34057. <https://doi.org/10.1038/srep34057>
- Ip YCA, Chang JJM, Oh RM, Quek ZBR, Chan YKS, Bauman AG, Huang D (2022) Seq' and ARMS shall find: DNA (meta)barcoding of Autonomous Reef Monitoring Structures across the tree of life uncovers hidden cryptobiome of tropical urban coral reefs. *Molecular Ecology* 32(23): 6223–6242. <https://doi.org/10.1111/mec.16568>

- Johnson MP, White M, Wilson A, Würzberg L, Schwabe E, Folch H, Allcock AL (2013) A vertical wall dominated by *Acesta excavata* and *Neopycnodonte zibrowii*, part of an undersampled group of deep-sea habitats. PLoS ONE 8(11): e79917. <https://doi.org/10.1371/journal.pone.0079917>
- Kalyanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS (2017) ModelFinder: Fast model selection for accurate phylogenetic estimates. Nature Methods 14(6): 587–589. <https://doi.org/10.1038/nmeth.4285>
- Katoh K, Rozewicki J, Yamada KD (2019) MAFFT online service: Multiple sequence alignment, interactive sequence choice and visualization. Briefings in Bioinformatics 20(4): 1160–1166. <https://doi.org/10.1093/bib/bbx108>
- Kim W-J, Lee J-H, Kim K-K, Kim Y-O, Nam B-H, Kong H-J, Jung H-T (2009) Genetic relationships of four Korean oysters based on RAPD and nuclear rDNA ITS sequence analyses. The Korean Journal of Malacology 25: 41–49.
- Kirkendale L, Lee T, Baker P, Ó Foighil D (2004) Oysters of the Conch Republic (Florida Keys): A molecular phylogenetic study of *Parahyotissa mcgintyi*, *Teskeyostrea weberi* and *Ostreola equestris*. Malacologia 46: 309–326.
- Lam K, Morton B (2003) Mitochondrial DNA and morphological identification of a new species of *Crassostrea* (Bivalvia: Ostreidae) cultured for centuries in the Pearl River Delta, Hong Kong, China. Aquaculture (Amsterdam, Netherlands) 228: 1–13. [https://doi.org/10.1016/S0044-8486\(03\)00215-1](https://doi.org/10.1016/S0044-8486(03)00215-1)
- Lam K, Morton B (2004) The oysters of Hong Kong (Bivalvia: Ostreidae and Gryphaeidae). The Raffles Bulletin Of Zoology 52: 11–28.
- Lam K, Morton B (2006) Morphological and mitochondrial-DNA analysis of the Indo-West Pacific rock oysters (Ostreidae: *Saccostrea* species). The Journal of Molluscan Studies 72(3): 235–245. <https://doi.org/10.1093/mollus/eyl002>
- Li C, Kou Q, Zhang Z, Hu L, Huang W, Cui Z, Liu Y, Ma P, Wang H (2021) Reconstruction of the evolutionary biogeography reveal the origins and diversification of oysters (Bivalvia: Ostreidae). Molecular Phylogenetics and Evolution 164: 107268. <https://doi.org/10.1016/j.ympev.2021.107268>
- Li F, Liu H, Heng X, Zhang Y, Fan M, Wang S, Liu C, Gu Z, Wang A, Yang Y (2023) The complete mitochondrial genome of *Hyotissa sinensis* (Bivalvia, Ostreidae) indicates the genetic diversity within Gryphaeidae. Biodiversity Data Journal 11: e101333. <https://doi.org/10.3897/BDJ.11.e101333>
- Liu J, Li Q, Kong L, Yu H, Zheng X (2011) Identifying the true oysters (Bivalvia: Ostreidae) with mitochondrial phylogeny and distance-based DNA barcoding. Molecular Ecology Resources 11(5): 820–830. <https://doi.org/10.1111/j.1755-0998.2011.03025.x>
- López-González PJ, Drewery J, López-González PJ, Drewery J (2022) When distant relatives look too alike: A new family, two new genera and a new species of deep-sea *Umbellula*-like sea pens (Anthozoa, Octocorallia, Pennatulacea). Invertebrate Systematics 36(3): 199–225. <https://doi.org/10.1071/IS21040>
- Matsumoto M (2003) Phylogenetic analysis of the subclass Pteriomorpha (Bivalvia) from mtDNA COI sequences. Molecular Phylogenetics and Evolution 27(3): 429–440. [https://doi.org/10.1016/S1055-7903\(03\)00013-7](https://doi.org/10.1016/S1055-7903(03)00013-7)
- Mazón-Suástegui JM, Fernández NT, Valencia IL, Cruz-Hernández P, Latisnere-Barragán H (2016) 28S rDNA as an alternative marker for commercially important oyster identification. Food Control 66: 205–214. <https://doi.org/10.1016/j.foodcont.2016.02.006>
- Minh BQ, Nguyen MAT, von Haeseler A (2013) Ultrafast Approximation for Phylogenetic Bootstrap. Molecular Biology and Evolution 30(5): 1188–1195. <https://doi.org/10.1093/molbev/mst024>
- Moritz C, Cicero C (2004) DNA Barcoding: Promise and Pitfalls. PLOS Biology 2(10): e354. <https://doi.org/10.1371/journal.pbio.0020354>
- Oliverio M, Mariottini P (2001) Contrasting morphological and molecular variation in *Coralliophila meyendorffii* (Muricidae, Coralliophilinae). The Journal of Molluscan Studies 67(2): 243–246. <https://doi.org/10.1093/mollus/67.2.243>
- Piazza F, Passamonti M (2010) Towards a molecular phylogeny of Mollusks: Bivalves' early evolution as revealed by mitochondrial genes. Molecular Phylogenetics and Evolution 57(2): 641–657. <https://doi.org/10.1016/j.ympev.2010.08.032>
- Piazza F, Ceregato A, Taviani M, Passamonti M (2011) A Molecular Phylogeny of Bivalve Mollusks: Ancient Radiations and Divergences as Revealed by Mitochondrial Genes. PLoS ONE 6(11): e27147. <https://doi.org/10.1371/journal.pone.0027147>
- Raith M, Zacherl DC, Pilgrim EM, Eernisse DJ (2015) Phylogeny and Species Diversity of Gulf of California Oysters (Ostreidae) Inferred from Mitochondrial DNA. American Malacological Bulletin 33(2): 263–283. <https://doi.org/10.4003/006.033.0206>
- Rambaut A, Drummond AJ, Xie D, Baele G, Suchard MA (2018) Posterior Summarization in Bayesian Phylogenetics Using Tracer 1.7. Systematic Biology 67(5): 901–904. <https://doi.org/10.1093/sysbio/syy032>
- Ramírez-Llodra E, Brandt A, Danovaro R, De Mol B, Escobar E, German CR, Levin LA, Martínez Arbizu P, Menot L, Buhl-Mortensen P, Narayanaswamy BE, Smith CR, Tittensor DP, Tyler PA, Vanreusel A, Vecchione M (2010) Deep, diverse and definitely different: Unique attributes of the world's largest ecosystem. Biogeosciences 7(9): 2851–2899. <https://doi.org/10.5194/bg-7-2851-2010>
- Ren J, Hou Z, Wang H, Sun M-A, Liu X, Liu B, Guo X (2016) Intraspecific variation in mitogenomes of five *Crassostrea* species provides insight into oyster diversification and speciation. Marine Biotechnology (New York, N.Y.) 18(2): 242–254. <https://doi.org/10.1007/s10126-016-9686-8>
- Rogers A, Brierley A, Croot P, Cunha M, Danovaro R, Devery C, Hoel AH, Ruhl H, Sarradin P-M, Trevisan S, van den Hove S, Vieira H, Visbeck M, McDonough N, Donaldson K, Larkin K (2015) Delving Deeper: Critical challenges for 21st century deep-sea research.
- Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP (2012) MrBayes 3.2: Efficient Bayesian Phylogenetic Inference and Model Choice Across a Large Model Space. Systematic Biology 61(3): 539–542. <https://doi.org/10.1093/sysbio/sys029>
- Rueda JL, Urra J, Aguilar R, Angeletti L, Bo M, García-Ruiz C, González-Duarte MM, López E, Madurell T, Maldonado M, Mateo-Ramírez Á, Megina C, Moreira J, Moya F, Ramalho LV, Rosso A, Sijà C, Taviani M (2019) 29 Cold-Water Coral Associated Fauna in the Mediterranean Sea and Adjacent Areas. In: Orejas C, Jiménez C (Eds) Mediterranean Cold-Water Corals: Past, Present and Future: Understanding the Deep-Sea Realms of Coral. Coral Reefs of the World. Springer International Publishing, Cham, 295–333. https://doi.org/10.1007/978-3-319-91608-8_29
- Ruppert KM, Kline RJ, Rahman MS (2019) Past, present, and future perspectives of environmental DNA (eDNA) metabarcoding: A

- systematic review in methods, monitoring, and applications of global eDNA. *Global Ecology and Conservation* 17: e00547. <https://doi.org/10.1016/j.gecco.2019.e00547>
- Salvi D, Mariottini P (2017) Molecular taxonomy in 2D: a novel ITS2 rRNA sequence-structure approach guides the description of the oysters' subfamily Saccostreinae and the genus *Magallana* (Bivalvia: Ostreidae). *Zoological Journal of the Linnean Society* 179: 263–276. <https://doi.org/10.1111/zoj.12455>
- Salvi D, Bellavia G, Cervelli M, Mariottini P (2010) The analysis of rRNA sequence-structure in phylogenetics: An application to the family Pectinidae (Mollusca: Bivalvia). *Molecular Phylogenetics and Evolution* 56(3): 1059–1067. <https://doi.org/10.1016/j.ympev.2010.04.025>
- Salvi D, Macali A, Mariottini P (2014) Molecular Phylogenetics and Systematics of the Bivalve Family Ostreidae Based on rRNA Sequence-Structure Models and Multilocus Species Tree. *PLoS ONE* 9: e108696. <https://doi.org/10.1371/journal.pone.0108696>
- Salvi D, Berrilli E, D'Alessandro P, Biondi M (2020) Sharpening the DNA barcoding tool through a posteriori taxonomic validation: The case of *Longitarsus* flea beetles (Coleoptera: Chrysomelidae). *PLoS ONE* 15(5): e0233573. <https://doi.org/10.1371/journal.pone.0233573>
- Salvi D, Berrilli E, Garzia M, Mariottini P (2021) Yet Another Mitochondrial Genome of the Pacific Cupped Oyster: The Published Mitogenome of *Alectryonella plicatula* (Ostreinae) Is Based on a Misidentified *Magallana gigas* (Crassostreinae). *Frontiers in Marine Science* 8: 741455. <https://doi.org/10.3389/fmars.2021.741455>
- Salvi D, Al-Kandari M, Oliver PG, Berrilli E, Garzia M (2022) Cryptic Marine Diversity in the Northern Arabian Gulf: An Integrative Approach Uncovers a New Species of Oyster (Bivalvia: Ostreidae), *Ostrea oleomargarita*. *Journal of Zoological Systematics and Evolutionary Research* 2022: e7058975. <https://doi.org/10.1155/2022/7058975>
- Sambrook J, Fritsch EF, Maniatis T (1989) *Molecular cloning: a laboratory manual*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York.
- Schindel DE, Miller SE (2005) DNA barcoding a useful tool for taxonomists. *Nature* 435(7038): 17. <https://doi.org/10.1038/435017b>
- Silva CF, Shimabukuro M, Alfaro-Lucas JM, Fujiwara Y, Sumida PYG, Amaral ACZ (2016) A new *Capitella* polychaete worm (Annelida: Capitellidae) living inside whale bones in the abyssal South Atlantic. *Deep-sea Research, Part I, Oceanographic Research Papers* 108: 23–31. <https://doi.org/10.1016/j.dsr.2015.12.004>
- Sinniger F, Pawlowski J, Harii S, Gooday AJ, Yamamoto H, Chevallon P, Cedhagen T, Carvalho G, Creer S (2016) Worldwide analysis of sedimentary DNA reveals major gaps in taxonomic knowledge of deep-sea benthos. *Frontiers in Marine Science* 3: 92. <https://doi.org/10.3389/fmars.2016.00092>
- Taberlet P, Coissac E, Hajibabaei M, Rieseberg LH (2012) Environmental DNA. *Molecular Ecology* 21(8): 1789–1793. <https://doi.org/10.1111/j.1365-294X.2012.05542.x>
- Talavera G, Castresana J (2007) Improvement of Phylogenies after Removing Divergent and Ambiguously Aligned Blocks from Protein Sequence Alignments. *Kjer K, Page R, Sullivan J (Eds) Systematic Biology* 56: 564–577. <https://doi.org/10.1080/10635150701472164>
- Tamura K, Stecher G, Kumar S (2021) MEGA11: Molecular Evolutionary Genetics Analysis Version 11. *Molecular Biology and Evolution* 38(7): 3022–3027. <https://doi.org/10.1093/molbev/msab120>
- Taviani M, Angeletti L, Canese S, Cannas R, Cardone F, Cau A, Cau AB, Follesa MC, Marchese F, Montagna P, Tassarolo C (2017) The “Sardinian cold-water coral province” in the context of the Mediterranean coral ecosystems. *Deep-sea Research, Part II, Topical Studies in Oceanography* 145: 61–78. <https://doi.org/10.1016/j.dsr2.2015.12.008>
- Taviani M, Angeletti L, Cardone F, Montagna P, Danovaro R (2019) A unique and threatened deep water coral-bivalve biotope new to the Mediterranean Sea offshore the Naples megalopolis. *Scientific Reports* 9(1): 3411. <https://doi.org/10.1038/s41598-019-39655-8>
- Tëmkin I (2010) Molecular phylogeny of pearl oysters and their relatives (Mollusca, Bivalvia, Pterioidea). *BMC Evolutionary Biology* 10(1): 342. <https://doi.org/10.1186/1471-2148-10-342>
- Trifinopoulos J, Nguyen L-T, von Haeseler A, Minh BQ (2016) W-IQ-TREE: A fast online phylogenetic tool for maximum likelihood analysis. *Nucleic Acids Research* 44(W1): W232–W235. <https://doi.org/10.1093/nar/gkw256>
- Vaidya G, Lohman DJ, Meier R (2011) SequenceMatrix: Concatenation software for the fast assembly of multi-gene datasets with character set and codon information. *Cladistics* 27(2): 171–180. <https://doi.org/10.1111/j.1096-0031.2010.00329.x>
- Van Rooij D, De Mol L, Le Guilloux E, Wisshak M, Huvenne VAI, Moeremans R, Henriët J-P (2010) Environmental setting of deep-water oysters in the Bay of Biscay. *Deep-sea Research, Part I, Oceanographic Research Papers* 57(12): 1561–1572. <https://doi.org/10.1016/j.dsr.2010.09.002>
- Wisshak M, Neumann C, Jakobsen J, Freiwald A (2009a) The ‘living-fossil community’ of the cyrtocrinid *Cyathidium foresti* and the deep-sea oyster *Neopycnodonte zibrowii* (Azores Archipelago). *Palaeogeography, Palaeoclimatology, Palaeoecology* 271(1–2): 77–83. <https://doi.org/10.1016/j.palaeo.2008.09.015>
- Wisshak M, López Correa M, Gofas S, Salas C, Taviani M, Jakobsen J, Freiwald A (2009b) Shell architecture, element composition, and stable isotope signature of the giant deep-sea oyster *Neopycnodonte zibrowii* sp. n. from the NE Atlantic. *Deep-sea Research, Part I, Oceanographic Research Papers* 56(3): 374–407. <https://doi.org/10.1016/j.dsr.2008.10.002>
- Woodall LC, Andradi-Brown DA, Brierley AS, Clark MR, Connelly D, Hall RA, Howell KL, Huvenne VAI, Linse K, Ross RE, Snelgrove P, Stefanoudis PV, Sutton TT, Taylor M, Thornton TF, Rogers AD (2018) A multidisciplinary approach for generating globally consistent data on mesophotic, deep-pelagic, and bathyal biological communities. *Oceanography (Washington, D.C.)* 31(3): 76–89. <https://doi.org/10.5670/oceanog.2018.301>
- Xu T, Feng D, Tao J, Qiu J-W (2019) A new species of deep-sea mussel (Bivalvia: Mytilidae: Gigantidas) from the South China Sea: Morphology, phylogenetic position, and gill-associated microbes. *Deep-sea Research, Part I, Oceanographic Research Papers* 146: 79–90. <https://doi.org/10.1016/j.dsr.2019.03.001>