

IMA Genome-F 7**Draft genome sequences for *Ceratocystis fagacearum*, *C. harringtonii*, *Grosmannia penicillata*, and *Huntiella bhutanensis***

Brenda D. Wingfield¹, Tuan A. Duong¹, Almuth Hammerbacher², Magriet A. van der Nest¹, Andi Wilson¹, Runlei Chang², Z. Wilhelm de Beer², Emma T. Steenkamp², P. Markus Wilken¹, Kershney Naidoo¹, and Michael J. Wingfield²

¹Department of Genetics, Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria, Pretoria, South Africa; corresponding author e-mail: Brenda.Wingfield@up.ac.za

²Department of Microbiology and Plant Pathology, Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria, Pretoria, South Africa

Abstract: Draft genomes for the fungi *Ceratocystis fagacearum*, *C. harringtonii*, *Grosmannia penicillata*, and *Huntiella bhutanensis* are presented. *Ceratocystis fagacearum* is a major causal agent of vascular wilt of oaks and other trees in the family Fagaceae. *Ceratocystis harringtonii*, previously known as *C. populincola*, causes disease in *Populus* species in the USA and Canada. *Grosmannia penicillata* is the causal agent of bluestain of sapwood on various conifers, including *Picea* spp. and *Pinus* spp. in Europe. *Huntiella bhutanensis* is a fungus in Ceratocystidaceae and known only in association with the bark beetle *Ips schmutzenhorferi* that infests *Picea spinulosa* in Bhutan. The availability of these genomes will facilitate further studies on these fungi.

Key words:

bluestain fungi
entomogenous fungi
Ips
Populus
tree diseases
vascular wilt

Article info: Submitted: 1 September 2016; Accepted: 6 November 2016; Published: 29 November 2016.

IMA Genome-F 7A**Draft genome sequence for the oak pathogen *Ceratocystis fagacearum***

Ceratocystis fagacearum (Microascales; Ceratocystidaceae) is a wilt pathogen of oak (*Quercus* spp.) and other Fagaceae in the eastern and north-central US (Henry *et al.* 1944, Billings 2000, Juzwik *et al.* 2004). Based on ecological and phylogenetic data, however, *C. fagacearum* falls outside of the genus *Ceratocystis* sensu stricto as defined by De Beer *et al.* (2014). Its taxonomic position as a discrete genus is currently being established (De Beer, unpublished).

Ceratocystis fagacearum causes a damaging and important vascular wilt disease known as oak wilt (Appel 1995). Infection occurs in spring through wounds commonly made during pruning operations. Trees die rapidly and the pathogen can pass from one tree to another via root grafts resulting in rows or patches of dying trees. As the trees die, pressure pads develop under the bark to expose spore-bearing mats where *C. fagacearum* produces a fruity odour attractive to insects (Lin & Phelan 1992, Cease & Juzwik 2001). These include sap-feeding nitidulid beetles that can transfer the pathogen to freshly made wounds on trees, thereby resulting in new infections (Cease & Juzwik 2001, Juzwik *et al.* 2004).

The aim of this study was to sequence the genome of *C. fagacearum*. The data are thus intended to compliment the

previously established genome resources for species in the Ceratocystidaceae (Wilken *et al.* 2013, van der Nest *et al.* 2014a, b, Wingfield *et al.* 2015a, b, Belbahri 2015, Wingfield *et al.* 2016), particularly in order to draw comparisons between them.

SEQUENCED STRAIN

USA: Iowa: West Des Moines, isol. *Quercus rubra*, Jan. 1991, S. Seegmueller (CMW 2656, CBS 138363, PREM 61535 - dried culture).

NUCLEOTIDE SEQUENCE ACCESSION NUMBER

The *Ceratocystis fagacearum* isolate CMW 2656 Whole Genome Shotgun project has been deposited in GenBank under the accession number MKGJ00000000.

MATERIALS AND METHODS

Ceratocystis fagacearum isolate CMW 2656 is obtainable from the culture collection (CMW) of the Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria, South Africa and the CBS-KNAW Fungal

Biodiversity Centre (CBS), Utrecht, The Netherlands. The fungus was grown at 25 °C on 2 % malt extract agar (MEA: 20 % w/v, Biolab, Midrand, South Africa) supplemented with 100 µg/L thiamine. Total genomic DNA was isolated using a phenol-chloroform method as described by Roux *et al.* (2004) and sequenced on the Genomics Analyzer IIx platform (Illumina) at the Genome Centre (University of California at Davis, CA). Two libraries (one with 350-bp and one with 600-bp paired-end inserts) were generated according to standard Illumina protocols and produced sequences with read lengths of approximately 100 bases. Reads with a limit of 0.05 and below were quality controlled and trimmed using CLC Genomics Workbench v. 7.5.1 (CLCBio, Aarhus). This software was also used to produce a draft genome assembly using the *de novo* assembly function under default settings. The assemblies were subsequently scaffolded using SSPACE v. 2.0 (Boetzer *et al.* 2011) and gaps filled using GapFiller v. 2.2.1 (Boetzer & Pirovano 2012). The online version of the *de novo* prediction software AUGUSTUS was employed to predict the putative open reading frames (ORFs) using *Fusarium graminearum* gene models (Stanke *et al.* 2004). Genome completeness was assessed with the Benchmarking Universal Single-Copy Orthologs (BUSCO v. 1.1b1) tool (Simão *et al.* 2015).

RESULTS AND DISCUSSION

The draft nuclear genome of *C. fagacearum* had an estimated size of 26 736 264 bases with 123x coverage. The assembly included 1257 contigs larger than 500 bases with an N50 value of 42 305 bases and an average GC content of 46.9 %. AUGUSTUS predicted 6703 ORFs, which correlated with an average gene density of 251 ORFs/Mb. This assembly had a high degree of completeness with a BUSCO score of 96 %, of which 1321 were Complete Single-Copy BUSCOs, 61 were Complete Duplicated BUSCOs, 47 were Fragmented BUSCOs, and only nine were missing BUSCO orthologs out of the 1438 BUSCO groups searched (Simao *et al.* 2015).

Comparison of these genome statistics revealed no striking differences between the genome of *C. fagacearum* and those determined in previous studies (Wilken *et al.* 2013, van der Nest *et al.* 2014a, b, Wingfield *et al.* 2015a, b, 2016, Belbahri 2015). This is despite its unique ecology and taxonomic position (de Beer *et al.* 2014). Future genome-based analyses will undoubtedly improve our understanding of the molecular and evolutionary processes underlying the biology of this pathogen.

Authors: M.A. van der Nest, P.M. Wilken, E.T. Steenkamp, A. Wilson, K. Naidoo, Al. Hammerbacher, M.J. Wingfield, and B.D. Wingfield

*Contact: Magriet.vanderNest@fabi.up.ac.za

IMA Genome-F 7B

Draft genome sequence of the poplar pathogen *Ceratocystis harringtonii*

The genome of black cottonwood (*Populus trichocarpa*) has been sequenced and is available for study, which has resulted in it becoming an important model tree for woody plant research (Tuskan *et al.* 2006). This tree species is well known for its resistance to fungal infection, and to date only a few pathogens are known to infect it (Royle & Ostry 1995). Fungal species currently causing severe disease on plantation-grown poplar include rust diseases caused by species of *Melampsora* (e. g. Newcombe *et al.* 2000) and leaf spots caused by *Marssonia* spp. (e.g. Erickson *et al.* 2004). In order to gain increased understanding of the adaptations allowing pathogens to infect poplar, the genome of the European poplar rust fungus, *Melampsora larici-populina*, was sequenced. This provided a powerful tool for studies in molecular plant-pathogen interactions in the phyllosphere of a perennial host (Duplessis *et al.* 2011). In contrast, little is known regarding stem-infecting pathogens in poplar, and particularly which resistance mechanisms exist to provide such extensive protection against fungal infection of its woody tissues. Furthermore, knowledge pertaining to strategies that pathogens employ to overcome this resistance has not been well documented.

Ceratocystis harringtonii resides in Ceratostidaceae (de Beer *et al.* 2013a, b), and was previously treated as *C. populicola* (Johnson *et al.* 2005). It causes target-shaped cankers on infected trunks and rooted cuttings (Wood & French 1963, Johnson *et al.* 2005). Furthermore, it is one of the few pathogens known to breach the effective defence barriers in poplar stems. This fungus occurs throughout the natural range of *Populus* species in the USA and Canada, as well as in hybrid poplar plantations in Poland (Gremmen & de Kam 1977). The pathogen is most aggressive on the North American poplar species *P. trichocarpa*, *P. balsamifera*, and *P. tremuloides*, while the European *P. nigra* is known to be almost entirely resistant to infection (Johnson *et al.* 2005). One mechanism by which *C. harringtonii* elicits host defences is by the production of cerato-populin, a pathogen-associated molecular pattern protein. This toxin is similar to the well-characterized ceratoplatatin and cerato-ulmin, which are known in *C. platani* and *Ophiostoma novo-ulmi*, respectively (Comparini *et al.* 2009, Lombardi *et al.* 2013, Martellini *et al.* 2013).

In order to extend our knowledge on plant-pathogen interactions in woody hosts and to understand the basis of resistance against fungal stem infection in poplar, the genome of *C. harringtonii* was sequenced. This genome sequence will provide a powerful tool to interrogate the mechanisms by which plant pathogens infect woody stems and cause canker disease in a highly resistant tree species.

SEQUENCED STRAIN

Poland: isol. ex hybrid poplar *Populus maximowiczii* x *P. laurifolia* x *P. nigra* 'Italica' (*P. xberolinensis*), 1977, J.

Gremmen (culture CBS 110.78 = CMW 14789 (ex-epitype); PREM 61533- dried culture).

NUCLEOTIDE SEQUENCE ASSESSMENT NUMBER

This Whole Genome Shotgun project of the *Ceratocystis harringtonii* genome has been deposited at DDBJ/EMBL/GenBank under accession number MKGM00000000; this is the first version described here.

MATERIALS AND METHODS

Total genomic DNA was isolated from the mycelium of a single-spore culture of isolate *C. harringtonii* CMW 14789 grown on 2 % malt extract agar (MEA: 2 % w/v, Biolab, Midrand, South Africa) supplemented with 100 µg/L thiamine for 10 d using the method of Barnes *et al.* (2001). The Genomics analyzer IIx platform (Illumina) at the Genome Centre (University of California at Davis, CA) was used for sequencing the genome. Two 350-bp and three 600-bp paired-end libraries were made using standard Illumina protocols. Sequence data was assessed for quality, trimmed and assembled with the software package CLC Genomics (CLCBio, Aarhus, Denmark) using default settings. Poor quality reads (limit of 0.05) and/or terminal nucleotides were discarded. The contigs were assembled into scaffolds using SSPACE v.2.0 (Boetzer *et al.* 2011) and gaps were filled using GapFiller v. 2.2.1 (Boetzer & Pirovano 2012). The genome assembly was verified and completeness assessed using the Benchmarking Universal Single-Copy Orthologs (BUSCO) software (Simão *et al.* 2015). Contigs smaller than 500 bp were removed from the final dataset.

RESULTS AND DISCUSSION

Sequencing of five *C. harringtonii* DNA libraries yielded a total of 31 057 034 paired-end reads with an average length of 101 bases. After trimming, 30 972 782 reads were recovered with an average length of 96 bases. The estimated size of the assembled draft genome was 26 Mb with a coverage of 110x. The genome had a mean GC content of 48.8 % and a N50 contig size of 66 kb. A total of 920 contigs were assembled of which 813 were larger than 500 bases (excluding the mitochondrial genome sequence data). BUSCO analysis revealed that out of 1438 possible BUSCO groups searched 1327 were single-copy, while 67 were duplicated and six were fragmented or missing. Analysis using AUGUSTUS (Stanke *et al.* 2006) revealed 6627 putative open reading frames.

Genome sizes differ widely among *Sordariomycetes*, for example the *Fusarium graminearum* and *Trichoderma reesii* genomes have approximate sizes of 36 Mb and 34 Mb, respectively (Martinez *et al.* 2008, King *et al.* 2015), while the genome of *Ceratocystiopsis minuta* is only 21 Mb in size (Wingfield *et al.* 2016). The estimated genome size of *C. harringtonii* (26 Mb) is marginally smaller than the genomes reported for other closely related members of

Ceratocystidaceae (28 – 32 Mb; Wingfield *et al.* 2016, Wilken *et al.* 2013, van der Nest *et al.* 2014). Despite the smaller genome size, the *C. harringtonii* genome contains similar numbers of predicted ORF's as other sequenced species of *Ceratocystis* (Wilken *et al.* 2013, van der Nest *et al.* 2014, Wingfield *et al.* 2016).

The *C. harringtonii* draft genome will be an important resource in studies of plant/pathogen interactions in woody tissues. This is especially true because this fungus is one of only a few reported pathogens that can overcome the defence mechanisms in the stem tissues of the model tree, *P. trichocarpa*. Furthermore, with growing numbers of genomes available for species in *Ceratocysticaceae*, the *C. harringtonii* genome will also be used for comparative genomic studies and those considering the mechanisms of host specialization in this fascinating group of plant pathogens.

Authors: A. Hammerbacher*, P.M. Wilken, M.A. van der Nest, A. Wilson, K. Naidoo, M.J. Wingfield, and B.D. Wingfield

*Contact: almuth.hammerbacher@fabi.up.ac.za

IMA Genome-F 7C

Draft genome sequence of *Grosmannia penicillata*

The asexual morph of *Grosmannia penicillata* was first described as *Leptographium penicillatum* (Grosmann 1931), the causal agent of blue stain of sapwood surrounding the galleries of the spruce bark beetle, *Ips typographus*. The sexual morph was discovered soon afterwards and described as *Ceratostomella penicillata* (Grosmann 1932), but Goidánich (1936) considered the species to be distinct from other *Ceratostomella* spp. and introduced a new generic name, *Grosmannia*, with *G. penicillata* as type species. Subsequently, *Grosmannia* was treated as a synonym of *Ophiostoma* (Siemaszko 1939, Arx 1952, Jacobs & Wingfield 2001) and *Ceratocystis* (Bakshi 1951, Hunt 1956, Upadhyay 1981) until Zipfel *et al.* (2006) re-instated the name to accommodate the sexual morphs of *Leptographium* spp. After the implementation of the one fungus one name principles (Hawksworth 2011), *Leptographium*, as the older generic name, currently has preference over *Grosmannia* (de Beer & Wingfield 2013). However, the type species of the two genera, *L. lundbergii* and *G. penicillata*, group in distinct lineages of which the generic status needs reconsideration (de Beer & Wingfield 2013). For the interim, the lineage that includes *G. penicillata* and 17 other *Leptographium* and *Grosmannia* species, are referred to as the *G. penicillata* species complex in *Leptographium* s. lat. (Six *et al.* 2011, Linnakoski *et al.* 2012, de Beer & Wingfield 2013).

Grosmannia penicillata occurs on various conifers including *Picea* and *Pinus* spp. in Europe (Jacobs & Wingfield 2001). It is vectored by various scolytine bark beetle species (Coleoptera; Curculionidae; Scolytinae) but most importantly, by the aggressive tree-killing bark beetle *Ips typographus* (Jacobs & Wingfield 2001, Linnakoski *et al.* 2012). Inoculation

studies have indicated that *G. penicillata* is not pathogenic to its hosts (Jankowiak *et al.* 2009, Repe *et al.* 2015). This is unlike *Endoconidiophora polonica*, as defined by de Beer *et al.* (2014), that is a common associate of *I. typographus* and is able to kill trees in inoculation tests (Horntvedt *et al.* 1983, Repe *et al.* 2015).

In this study, we determined the genome sequence of *G. penicillata*. The primary intention was to provide basal genomic data to enable further studies on the taxonomy and evolutionary relationships of this species and other genera and species in the *Ophiostomatales*.

SEQUENCED STRAIN

Norway: Akershus: Ås, isol. *Picea abies*, Jan. 1980, H. Solheim (culture CMW 2644 = CBS 116008; PREM 61536 – dried culture)

NUCLEOTIDE SEQUENCE ACCESSION NUMBER

The genomic sequence of *Grosmannia penicillata* (CMW 2644, CBS 116008) has been deposited at DDBJ/EMBL/GenBank under the accession number MLJV00000000. The version described in this paper is version MLJV01000000.

MATERIALS AND METHODS

Grosmannia penicillata isolate CMW 2644 was obtained from the culture collection (CMW) of the Forestry and Agricultural Biotechnology Institute (FABI), the University of Pretoria, South Africa. Genomic DNA was extracted from the freeze-dried mycelium using the method described by Duong *et al.* (2013). Two pair-end libraries (350 bp and 530 bp average insert size) were prepared and sequenced using the Illumina HiSeq 2000 platform with 100 bp read length. Obtained reads were subjected to quality and adapters trimming using Trimmomatic v. 0.36 (Bolger *et al.* 2014). The genome was assembled using the program SPAdes v. 3.9 (Bankevich *et al.* 2012). The scaffolds obtained from SPAdes were subjected to further scaffolding with SSPACE-standard v. 3.0 (Boetzer *et al.* 2011). Assembly gaps were filled with GapFiller V1-10 (Boetzer & Pirovano 2012). The completeness of the assembly was assessed with the Benchmarking Universal Single-Copy Orthologs (BUSCO v. 1.1b1) program using the fungal dataset (Simão *et al.* 2015). The number of protein coding genes was determined using Augustus v 3.2.2 (Stanke *et al.* 2006).

RESULTS AND DISCUSSION

Over 11 million read pairs and 1.4 million single reads were obtained after the quality trimming. *De-novo* assembly using SPAdes resulted in 293 scaffolds which were larger than 500 bp. The number of final scaffolds was reduced to 199 after scaffolding with SSPACE and filling gaps with GapFiller. The current assembly has an N50 of 316 Kb and size of 26.33

Mb, with an overall GC content of 58.73 %. When compared with other species of *Leptographium s. lat.* for which whole genome data are available, *G. penicillata* has a similar genome size to that of *Leptographium lundbergii* (26.6 Mb) (Wingfield *et al.* 2015), and slightly smaller than those of *G. clavigera* (29.8 Mb) (DiGuistini *et al.* 2011) and *L. procerum* (28.6 Mb) (van der Nest *et al.* 2014).

The assembly included 94.6 % complete, 4.2 % fragmented, and 1.2 % missing, BUSCOs. Augustus predicted a total of 8713 protein coding genes, of which 6718 are multi-exon and 1995 are single-exon genes. As the type species of *Grosmannia*, the genome sequence of *G. penicillata* will be a valuable resource to study the taxonomic relationships between species in the genus. The data will also be useful in comparisons between genera in *Ophiostomatales* where questions relating to their ecology and evolutionary biology are of particular interest.

Authors: T.A. Duong*, M.J. Wingfield, Z.W. de Beer, R. Chang, and B.D. Wingfield

*Contact: Tuan.Duong@fabu.up.ac.za

IMA Genome-F 7D

Draft genome sequence for the bark beetle-associated fungus *Huntiella bhutanensis*

Huntiella bhutanensis (Ascomycota; Microascales) is a filamentous fungus in Ceratocystidaceae. The species was previously accommodated in *Ceratocystis* as *C. bhutanensis* (van Wyk *et al.* 2004). However, a recent taxonomic review of this and related genera led to the re-circumscription of these fungi into distinct genera based on morphological, ecological and molecular characteristics (de Beer *et al.* 2014). Thus, *H. bhutanensis*, along with other cosmopolitan saprobes such as *Huntiella omanensis* and *H. moniliformis*, were assigned to the genus *Huntiella* (de Beer *et al.* 2014).

Huntiella bhutanensis was first isolated from adults of the bark beetle *Ips schmutzenhoferi* (Coleoptera; Curculionidae; Scolytinae) or their galleries found on *Picea spinulosa* in Bhutan (van Wyk *et al.* 2004). This was only the second report of any ophiostomatoid fungi from this country and represented the first species of Ceratocystidaceae to be reported in this locality (van Wyk *et al.* 2004). Despite its close relationship with the bark beetle, and unlike many other bark beetle-associates in the Ceratocystidaceae, *H. bhutanensis* is not considered a primary pathogen and in inoculation studies, it gave rise to only small necrotic lesions on *P. spinulosa* trees (Kiritsis *et al.* 2012).

The non-pathogenic nature of *H. bhutanensis* is consistent with other *Huntiella* species that are weak pathogens or saprobes (de Beer *et al.* 2014). It also shares many morphological characteristics with other *Huntiella* species, such as globose ascomatal bases, extended ascromatal necks, and hat-shaped ascospores (van Wyk *et al.* 2004). It is unusual amongst species in the genus in that it is able to grow at 4°C (van Wyk *et al.* 2004) and it gives off a

putrid odour unlike the sweet smelling odours typical of other *Huntiella* species (van Wyk *et al.* 2004).

The aim of this study was to produce a good quality draft genome assembly for *H. bhutanensis* for use in future comparative genomics studies, both between species and genera in *Ceratostidaceae*. Genome sequences are already publically available for *H. moniliformis* (van der Nest *et al.* 2014a) and *H. omanensis* (van der Nest *et al.* 2014b) and thus the availability of this genome assembly will aid in the elucidation and comparison of ecological strategies, sexual cycles, and other key life-style aspects of and between these species.

SEQUENCED STRAINS

Bhutan: isol. *Picea spinulosa*, July 2001, M. J. Wingfield (CMW 8217, CBS 114289, PREM 57807- dried culture).

NUCLEOTIDE SEQUENCE ACCESSION NUMBER

The *Huntiella bhutanensis* isolate CMW 8217 Whole Genome Shotgun project has been deposited at DDBJ/ENA/GenBank under the MJMS00000000. The version described in this paper is version MJMS01000000.

MATERIALS AND METHODS

Huntiella bhutanensis isolate CMW 8217 is preserved in the culture collection (CMW) of the Forestry and Agricultural Biotechnology Institute (FABI) at the University of Pretoria, South Africa. It can also be requested from the CBS-KNAW Fungal Biodiversity Centre (CBS), Utrecht, The Netherlands. Genomic DNA was isolated using the method of Barnes *et al.* (2001) and sequenced using the Genomics Analyzer IIx, Illumina platform at the UC Davis Genome Centre, University of California, Davis (CA). Paired-end libraries of both 350-bp and 600-bp insert sizes were prepared and sequenced following Illumina protocol. Quality trimming of the reads obtained was conducted using CLC Genomics Workbench v. 7.5.1 (CLCBio, Aarhus). This same program was used to assemble a draft genome sequence using the *de novo* assembly function, with default settings. Thereafter, SSPACE v. 2.0 (Boetzer *et al.* 2011) was used to scaffold the assemblies, and gaps created during this process were filled using GapFiller v. 2.2.1 (Boetzer & Pirovano 2012). The web-based *de novo* gene prediction software AUGUSTUS was used to predict the number of putative open reading frames (ORFs) in this assembly using the gene models of *Fusarium graminearum* (Stanke *et al.* 2004). Genome completeness was assessed with the Benchmarking Universal Single-Copy Orthologs (BUSCO v. 1.22) tool (Simão *et al.* 2015) using the fungal dataset.

RESULTS AND DISCUSSION

Sequencing of the *Huntiella bhutanensis* libraries yielded 35 886 298 raw reads with an average length of 101 bases. Trimming and quality control left 35 822 293 reads with an average length of 96 bases. The assembled nuclear genome of *H. bhutanensis* had a size of 26.77 Mb with an average coverage of 126X. This assembly had 448 scaffolds larger than 500 bases, an N50 value of 201 808 bases and an approximate GC content of 47.9 %. Web AUGUSTUS predicted 7 261 ORFs. This corresponded to an average gene density of 279 ORFs/Mb. The BUSCO analysis for this assembly also indicated a high level of completeness. Out of the 1438 fungal BUSCO groups searched, the genome contained 1315 (91.4 %) complete single copy BUSCOs, 62 (4.3 %) complete duplicated BUSCOs, 52 (3.6 %) fragmented BUSCOs and only nine (< 1 %) missing BUSCO orthologs.

The genome of *H. bhutanensis* is about 5 Mb smaller than that of its phylogenetically-close relative *H. omanensis* and possesses 1 134 fewer genes (van der Nest *et al.* 2014b). However, the genome assemblies of *H. bhutanensis* and *H. moniliformis* are more similar, at 25.4 Mb and 26.7 Mb with 6832 and 7261 ORFs, respectively (van der Nest *et al.* 2014a). The genome sequence of *H. bhutanensis* will serve as an essential resource in future genome comparisons between species of this genus and will aid in the characterization of their lifestyles, sexual cycles, and other key aspects of their biology.

Authors: A. Wilson*, M.A. van der Nest, P.M. Wilken,

K. Naidoo, M.J. Wingfield, and B.D. Wingfield

*Contact: Andi.Wilson@fabi.up.ac.za

ACKNOWLEDGEMENTS

This work was co-funded by the Genomics Research Institute (University of Pretoria), the University of Pretoria Research Development Programme, the DST/NRF Center of Excellence in Tree Health Biotechnology (FABI, University of Pretoria), and the National Research Foundation (NRF) (Grant number 87332). Sequencing the genomes of *Ceratostysis fagacearum*, *C. harringtonii*, *Huntiella bhutanensis* was made possible through the Improving Academics Qualifications (IAQ) grant from the NRF.

REFERENCES

- Appel DN (1995) The oak wilt enigma: perspectives from the Texas epidemic. *Annual Review of Phytopathology* **33**: 103–118.
- Appel DN, Kurdyla T, Lewis R (1990) Nitidulids as vectors of the oak wilt fungus and other *Ceratostysis* spp. in Texas. *European Journal of Forest Pathology* **20**: 412–417.
- Arx JA von (1952) Ueber die Ascomycetengattungen *Ceratostomella* Sacc., *Ophiostoma* Syd. und *Rostrella* Zimmermann. *Antonie van Leeuwenhoek* **18**: 201–213.
- Bakshi, BK (1951) Studies on four species of *Ceratostysis*, with a discussion on fungi causing sap-stain in Britain. *Mycological Papers* **35**: 1–16.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, *et al.* (2012) SPAdes: a new genome assembly algorithm and its

- applications to single-cell sequencing. *Journal of Computational Biology* **19**: 455–477.
- Barnes I, Gaur A, Burgess T, Roux J, Wingfield BD, Wingfield MJ (2001) Microsatellite markers reflect intra-specific relationships between isolates of the vascular wilt pathogen *Ceratocystis fimbriata*. *Molecular Plant Pathology* **2**: 319–25.
- Belbahri L (2015) Genome sequence of *Ceratocystis platani*, a major pathogen of plane trees. <http://www.ncbi.nlm.nih.gov/nuccore/814603118>.
- Billings RF (2000) State forest health problems: a survey of state foresters. *Journal of Forestry* **98**: 20–25.
- Boetzer M, Henkel CV, Jansen HJ, Butler D, Pirovano W (2011) Scaffolding pre-assembled contigs using SSPACE. *Bioinformatics* **27**: 578–579.
- Boetzer M, Pirovano W (2012) Toward almost closed genomes with GapFiller. *Genome Biology* **13**: R56.
- Bolger AM, Lohse M, Usadel B (2014) Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**: 2114–2120.
- Cease KR, Juzwik J (2001) Predominant nitidulid species (Coleoptera: Nitidulidae) associated with spring oak wilt mats in Minnesota. *Canadian Journal of Forest Research* **31**: 635–643.
- Comparini C, Carresi L, Pagni E, Sbrana F, Sebastiani F, et al. (2009) New proteins orthologous to cerato-platinin in various *Ceratocystis* species and the purification and characterization of cerato-populin from *Ceratocystis populicola*. *Applied Microbiology and Biotechnology* **84**: 309–322.
- de Beer ZW, Duong TA, Barnes I, Wingfield BD, Wingfield MJ (2014) Redefining *Ceratocystis* and allied genera. *Studies in Mycology* **79**: 187–219.
- de Beer ZW, Wingfield MJ (2013) Emerging lineages in the Ophiostomatales, In: *The Ophiostomatoid Fungi: expanding frontiers* (Seifert KA, de Beer ZW, Wingfield MJ eds): 21–46. [CBS Biodiversity Series no. 12.] Utrecht: CBS-KNAW Fungal Biodiversity Centre.
- DiGuistini S, Wang Y, Liao NY, Taylor G, Tanguay P, et al. (2011) Genome and transcriptome analyses of the mountain pine beetle-fungal symbiont *Grosmannia clavigera*, a lodgepole pine pathogen. *Proceedings of the National Academy of Sciences, USA* **108**: 2504–2509.
- Duong TA, de Beer ZW, Wingfield BD, Wingfield MJ (2013) Characterization of the mating-type genes in *Leptographium procerum* and *Leptographium profanum*. *Fungal Biology* **117**: 411–421.
- Duplessis S, Cuomo CA, Lin YC, Aerts A, Tisserant E, et al. (2011) Obligate biotrophy features unraveled by the genomic analysis of rust fungi. *Proceedings of the National Academy of Sciences, USA* **108**: 9166–9171.
- Erickson JE, Stanosz GR, Kruger EL (2004) Photosynthetic consequences of *Marssonina* leaf spot differ between two poplar hybrids. *New Phytologist* **161**: 577–583.
- Goidàñich G (1936) Il genere di Ascomiceti 'Grosmannia' G. Goid. *Bollettino della Stazione di Patologia Vegetale di Roma* **16**: 26–40.
- Gremmen J, Kam, MD (1977) *Ceratocystis fimbriata*, a fungus associated with poplar canker in Poland. *European Journal of Forest Pathology* **7**: 44–47.
- Grosmann H (1931) Beiträge zur Kenntnis der Lebensgemeinschaft zwischen Borkenkäfern und Pilzen. *Zeitschrift für Parasitenkunde* **3**: 56–102.
- Grosmann H (1932) Über die systematischen Beziehungen der Gattung *Leptographium* Lagerberg et Melin zur Gattung *Ceratostomella* Sacc. *Nova Hedwigia* **72**: 183–194.
- Hawksworth DL (2011) A new dawn for the naming of fungi: impacts of decisions made in Melbourne in July 2011 on the future publication and regulation of fungal names. *MycoKeys* **1**: 7–20; *IMA Fungus* **2**: 155–162.
- Henry BW, Moses CS, Richards CA, Riker AJ (1944) Oak wilt: Its significance, symptoms and cause. *Phytopathology* **34**: 636–647.
- Horntvedt, R, Christiansen, E, Solheim, H, Wang, S (1983) Artificial inoculation with *Ips typographus*-associated blue-stain fungi can kill healthy Norway spruce trees. *Meddelelser fra Norsk Institutt for Skogforskning* **38**: 1–20.
- Hunt J (1956) Taxonomy of the genus *Ceratocystis*. *Lloydia* **19**: 1–58.
- Jacobs K, Wingfield MJ (2001) *Leptographium* species: tree pathogens, insect associates and agents of blue-stain. St Paul, MN: American Phytopathological Society Press.
- Jankowiak R, Rossa R, Bilański P (2009) A preliminary study on the pathogenicity of three blue-stain fungi associated with *Tetropium* spp. to Norway spruce in Poland. *Forest Research Papers* **70**: 69–75.
- Johnson JA, Harrington TC, Engelbrecht CJB (2005) Phylogeny and taxonomy of the North American clade of the *Ceratocystis fimbriata* complex. *Mycologia* **97**: 1067–1092.
- Juzwik J, Skalbeck TC, Neuman MF (2004) Sap beetle species (Coleoptera: Nitidulidae) visiting fresh wounds on healthy oaks during spring in Minnesota. *Forest Science* **50**: 757–764.
- King R, Urban M, Hammond-Kosack MC, Hassani-Pak K, Hammond-Kosack KE (2015) The completed genome sequence of the pathogenic ascomycete fungus *Fusarium graminearum*. *BMC Genomics* **16**: 544.
- Kirisits T, Konrad H, Wingfield MJ (2012) Ophiostomatoid fungi associated with the Eastern Himalayan spruce bark beetle (*Ips schmutzenhoferi*) in Bhutan: species assemblage and phytopathogenicity. *Journal of Agricultural Extension and Rural Development* **4**: 266–268.
- Lin H, Phelan PL (1992) Comparison of volatiles from beetle-transmitted *Ceratocystis fagacearum* and four non-insect-dependent fungi. *Journal of Chemical Ecology* **18**: 1623–1632.
- Linnakoski R, de Beer ZW, Duong TA, Niemelä P, Pappinen A, Wingfield MJ (2012) *Grosmannia* and *Leptographium* spp. associated with conifer-infesting bark beetles in Finland and Russia, including *Leptographium taigense* sp. nov. *Antonie van Leeuwenhoek* **102**: 375–399.
- Lombardi L, Faoro F, Luti S, Bacelli I, Martellini F, et al. (2013) Differential timing of defense-related responses induced by cerato-platinin and cerato-populin, two non-catalytic fungal elicitors. *Physiologia Plantarum* **149**: 408–421.
- Martellini F, Faoro F, Carresi L, Pantera B, Bacelli I, et al. (2013) Cerato-populin and cerato-platinin, two non-catalytic proteins from phytopathogenic fungi, interact with hydrophobic inanimate surfaces and leaves. *Molecular Biotechnology* **55**: 27–42.
- Martinez D, Berka RM, Henrissat B, Saloheimo M, Arvas M, et al. (2008) Genome sequencing and analysis of the biomass-degrading fungus *Trichoderma reesei* (syn. *Hypocreah jecorina*). *Nature Biotechnology* **26**: 553–560.
- Mayers CG, McNew DL, Harrington TC, Roeper RA, Fraedrich SW, Biedermann PH, Castrillo LA, Reed SE (2015) Three genera in the *Ceratostomellaceae* are the respective symbionts of three independent lineages of ambrosia beetles with large, complex

- mycangia. *Fungal Biology* **119**: 1075–1092.
- Newcombe G, Stirling B, McDonald S, Bradshaw HD (2000) *Melampsora x columbiana*, a natural hybrid of *M. medusae* and *M. occidentalis*. *Mycological Research* **104**: 261–274.
- Repe A, Bojović S, Jurc M (2015) Pathogenicity of ophiostomatoid fungi on *Picea abies* in Slovenia. *Forest Pathology* **45**: 290–297.
- Roux J, van Wyk M, Hatting H, Wingfield MJ (2004) Ceratocystis species infecting stem wounds on *Eucalyptus grandis* in South Africa. *Plant Pathology* **53**: 414–421.
- Royle DJ, Ostry M (1995) Disease and pest control in the bioenergy crops poplar and willow. *Biomass and Bioenergy* **9**: 69–79.
- Siemaszko, W (1939) Zespol grzybów towarzyszących kornikom polskim. *Planta Polonica* **7**: 1–54.
- Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM (2015) BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics* **31**: 3210–3212.
- Six D, de Beer Z, Duong T, Carroll A, Wingfield M (2011) Fungal associates of the lodgepole pine beetle, *Dendroctonus murrayanae*. *Antonie van Leeuwenhoek* **100**: 231–244.
- Stanke M, Steinkamp R, Waack S, Morgenstern B (2004) AUGUSTUS: A web server for gene finding in Eukaryotes. *Nucleic Acids Research* **32**: W309–W312.
- Tuskan GA, Difazio S, Jansson S, Bohlmann J, Grigoriev I, et al. (2006) The genome of black cottonwood, *Populus trichocarpa* (Torr. & Gray). *Science* **313**: 1596–1604.
- Upadhyay HP (1981) *A monograph of Ceratocystis and Ceratocystiopsis*. Athens, GA: University of Georgia Press.
- van der Nest MA, Beirn LA, Crouch JA, Demers JE, de Beer ZW, et al. (2014b) IMA Genome-F 3: Draft genomes of *Amanita jacksonii*, *Ceratocystis albifundus*, *Fusarium circinatum*, *Huntiella omanensis*, *Leptographium procerum*, *Rutstroemia sydowiana*, and *Sclerotinia echinophila*. *IMA Fungus* **5**: 473–486.
- van der Nest MA, Bihon W, De Vos L, Naidoo K, Roodt D, et al. (2014a) IMA Genome-F 2: *Ceratocystis manginecans*, *Ceratocystis moniliformis*, *Diplodia sapinea*: Draft genome sequences of *Diplodia sapinea*, *Ceratocystis manginecans*, and *Ceratocystis moniliformis*. *IMA Fungus* **5**: 135–140.
- van Wyk M, Roux J, Barnes I, Wingfield BD, Chhetri DB, et al. (2004) *Ceratocystis bhutanensis* sp. nov., associated with the bark beetle *Ips schmutzenhoferi* on *Picea spinulosa* in Bhutan. *Studies in Mycology* **50**: 365–379.
- Wilken PM, Steenkamp ET, Wingfield MJ, de Beer ZW, Wingfield BD (2013) IMA Genome-F 1: *Ceratocystis fimbriata*: draft nuclear genome sequence for the plant pathogen, *Ceratocystis fimbriata*. *IMA Fungus* **4**: 357–358.
- Wingfield BD, Ades PK, Al-Naemi FA, Beirn LA, Bihon W, et al. (2015a) IMA Genome-F 4: Draft genome sequences of *Chrysoporthe austroafricana*, *Diplodia scrobiculata*, *Fusarium nygamai*, *Leptographium lundbergii*, *Limonomyces culmigenus*, *Stagonosporopsis tanaceti*, and *Thielaviopsis punctulata*. *IMA Fungus* **6**: 233–248.
- Wingfield BD, Barnes I, de Beer ZW, de Vos L, Duong TA, et al. (2015b) IMA Genome-F 5: Draft genome sequences of *Ceratocystis eucalypticola*, *Chrysoporthe cubensis*, *C. deuterocubensis*, *Davidsoniella virescens*, *Fusarium temperatum*, *Graphilbum fragrans*, *Penicillium nordicum*, and *Thielaviopsis musarum*. *IMA Fungus* **6**: 493–506.
- Wingfield BD, Ambler JM, Coetzee M, de Beer ZW, Duong TA, et al. (2016) Draft genome sequences of *Armillaria fuscipes*, *Ceratocystiopsis minuta*, *Ceratocystis adiposa*, *Endoconidiophora laricicola*, *E. polonica* and *Penicillium freii* DAOMC 242723. *IMA Fungus* **30**: 217–227.
- Wood FA, French DW (1963) *Ceratocystis fimbriata*, the cause of a stem canker of quaking aspen. *Forest Science* **9**: 232–235.
- Zipfel RD, de Beer ZW, Jacobs K, Wingfield BD, Wingfield MJ (2006) Multi-gene phylogenies define *Ceratocystiopsis* and *Grosmannia* distinct from *Ophiostoma*. *Studies in Mycology* **55**: 75–97.