

# The subfamily Dermestinae (Coleoptera, Dermestidae) from Saudi Arabia

Jiří Háva<sup>1</sup>, Mahmoud S. Abdel-Dayem<sup>2,3</sup>, Hathal M. Aldhafer<sup>2</sup>

**1** Private Entomological Laboratory and Collection, Rýznerova 37/37, CZ-252 62 Únětice u Prahy, Prague-West, Czech Republic **2** King Saud University Museum of Arthropods (KSMA), Plant Protection Department, College of Food and Agricultural Sciences, King Saud University, P.O. Box 2460 Riyadh 11451, Saudi Arabia **3** Entomology Department, Faculty of Science, Cairo University, Giza, 12613, Egypt

Corresponding author: Mahmoud S. Abdel-Dayem ([mseleem@ksu.edu.sa](mailto:mseleem@ksu.edu.sa))

Academic editor: Thomas Philips | Received 26 July 2022 | Accepted 18 December 2022 | Published 5 January 2023

<https://zoobank.org/E4C18E91-8B1E-4656-BC09-8D147E87082F>

**Citation:** Háva J, Abdel-Dayem MS, Aldhafer HM (2023) The subfamily Dermestinae (Coleoptera, Dermestidae) from Saudi Arabia. ZooKeys 1138: 161–173. <https://doi.org/10.3897/zookeys.1138.90338>

## Abstract

In this study, the fauna of Saudi Arabian Dermestinae (Coleoptera, Dermestidae) is summarised. Six *Dermestes* species and single species from two Marioutini genera, *Mariouta* and *Rhopalosilpha*, are reported. *Dermestes* (*Dermestinus*) *undulatus* Brahm, 1790 and *Dermestes* (*Dermestes*) *haemorrhoidalis* Küster, 1852 are newly recorded from Saudi Arabia. A list of Dermestinae species from the Arabian Peninsula is provided with their distributions.

## Keywords

Beetles, Dermestini, distribution, fauna, Marioutini, new records

## Introduction

Dermestinae is a subfamily of Dermestidae with a worldwide distribution, but concentrated in the Holarctic and Afrotropical areas. According to Háva (2015, 2022), there are approximately 95 species assigned to only five genera under two tribes: Dermestini with three genera, *Derbyana* Lawrence & Šlipiński, *Dermalius* Háva, and *Dermestes* Linnaeus, 1758, and Marioutini with two genera, *Mariouta* Pic and *Rhopalosilpha*

Arrow; additionally there is the fossil tribe Paradermestini with one genus, *Paradermestes* Deng, Ślipiński, Ren & Pang (Háva 2015, 2022). The genus *Dermestes* is the largest genus in Dermestinae and recently included 89 species and subspecies worldwide (Háva 2015, 2022). Members of the subfamily are generally recognised by their elongate body structure, lack of ocelli, and males with small tufts of erect setae on the abdominal ventrites (females are without tufts). Larvae are zoonecrophagous.

The first data concerning Dermestinae of Saudi Arabia date back to the second half of the 20<sup>th</sup> century. In the early 1960s, the Egyptian entomologist F. Shalaby (1961) was perhaps the first who catalogued data on *Dermestes maculatus* DeGeer, 1774. The work of the Polish entomologist M. Mroczkowski (1979) was the first important faunistic study on the Saudi Arabian Dermestidae fauna. His work was based on the collection made by W. Büttiker who intensively explored many areas of Saudi Arabia, and he recorded three *Dermestes* species. Mroczkowski and Ślipiński (1997) published their review and keys to world genera and species of the tribe Marioutini and reported *Mariouta stangei* Reitter, 1910 and *Rhopalosilpha wasmanni* Arrow, 1929 from Saudi Arabia.

From the beginning of the 21<sup>st</sup> century and during the last two decades, the forensic importance of dermestid beetles attracted the attention of many workers from Saudi Arabia (e.g., Abouzied 2014; Alajmi et al. 2016; Al-Shareef and Al-Mazyad 2017; Al-Shareef and Zaki 2017; Mashaly 2017; Shaalan et al. 2017; Mashaly et al. 2018, 2019; Al-Dakhil and Alharbi 2020; Al-Qahtni et al. 2020). However, the faunistic data on Dermestinae were published as part of general surveys of insects or beetles (Abdel-Dayem et al. 2017, 2020; Elgharbawy 2018). The systematic, faunistic, and distribution of Dermestinae in Saudi Arabia are still not well known, and few works have been published. This paper aims to summarise the known Saudi Arabian Dermestinae and update distribution data.

## Materials and methods

The data on the distribution of the species in the subfamily Dermestinae (Coleoptera, Dermestidae) in Saudi Arabia is based on three main sources. The first are the historical works of Shalaby (1961), Mroczkowski (1979), Mroczkowski and Ślipiński (1997), and additionally the recent publication of Abouzied (2014), Abdel-Dayem et al. (2017, 2020), Al-Shareef and Zaki (2017), Mashaly (2017), Elgharbawy (2018), Mashaly et al. (2019), Al-Dakhil and Alharbi (2020), and Al-Qahtni et al. (2020). The second source are specimens preserved in the insect collections of the King Saud University Museum of Arthropods (**KSMA**) in Riyadh, Saudi Arabia, the Florida State Collection and Arthropods (**FSCA**), and the collection of the first author. The third source is an extended field survey conducted by the second and third authors, which is still ongoing. The collected specimens were deposited in the collections of KSMA, unless otherwise indicated (JHAC: Jiří Háva). The nomenclature follows Motyka et al. (2022). A note entry summarises published and current data on the species distribution within

Saudi Arabia. The general range and the world distribution data were derived from the catalogues of Háva (2015, 2022).

For each material lot examined, the following label data are provided as follows: Country name (in capital letters) at the beginning. Then each record starts with a bullet point (•) followed by the number of examined specimens followed by sex (if determined) or “ex” (if the specimen sex could not be recognised because the abdomen was lost, damaged, or other reasons); Saudi Province followed by a comma (,), governorate, locality; geographical coordinates; elevation (m), collection date; collector(s) name followed by “leg.”; method of collection (bait trap (**BT**), handpicking (**HP**), light trap (**LT**), malaise trap (**MT**), pitfall trap (**PT**), sweeping net (**SW**)), the identifier name followed by “det.”, and the depository collection acronym. The material examined was arranged in alphabetical order with respect to the Saudi province, governorate, and locality name. Data were then arranged in chronological order according to the month of collection. Records with the same locality data, except for slight differences (such as date of collection, altitude, collector(s)), were reported together with the second label, given “same collection data as for preceding” and followed by a semicolon (;) and the different data.

The following acronyms of type depositories are used in the text:

- JHAC** Jiří Háva, Private Entomological Laboratory & Collection, Únětice u Prahy, Prague-West, Czech Republic;  
**FSCA** Florida State Collection and Arthropods, Gainesville, USA;  
**KSMA** King Saud University Museum of Arthropods, Plant Protection Department, College of Food and Agriculture Sciences, King Saud University, Riyadh, Saudi Arabia.

## Results

### Family Dermestidae Latreille, 1804

### Subfamily Dermestinae Latreille, 1804

### Tribe Dermestini Latreille, 1804

### Genus *Dermestes* Linnaeus, 1758

### Subgenus *Dermestes* s. str.

### *Dermestes (Dermestes) ater* DeGeer, 1774

Fig. 1A, B

**Material examined.** SAUDI ARABIA • 1 ♂; Eastern Province, An Nuayriah, Al Sarar; 27°25'45.5"N, 48°27'0.0"E; 60 m a.s.l.; 2 Mar. 2011; H. Al Dhafer; H. Setyaningrum & A. Al Ansi leg.; collected from carcasses on the road; J. Háva det.; KSMA • 1 ♀; Makkah Province, Jeddah, “Ras Halibah” [Ras Hatibah]; 7 May. 1982; W. Büttiker leg.; JHAC • 4 ex; Riyadh Province, Dirab, Al-Dhab Farm; 5 Oct. 1986; collected from

chicken farm waste, J. Háva det.; KSMA; • 1 ex; Riyadh, Al-Wahah Farm; 12 Oct. 1989; J. Háva det.; KSMA.

**Note.** This species was previously recorded in Eastern Province at Al Hofuf (Mroczkowski 1979); Dammam (Mroczkowski 1979), Dhahran (Mroczkowski 1979), and Riyadh Province at Riyadh (Mroczkowski 1979). The listed specimens were collected from low elevation areas (<600 m) in central, eastern, and southwestern Saudi Arabia (Fig. 4A).


**Distribution.** Cosmopolitan (Háva 2007, 2015, 2022).

***Dermestes (Dermestes) haemorrhoidalis* Küster, 1852**

Fig. 1C, D

**Material examined.** SAUDI ARABIA • Riyadh Province, 1 ♀; Al Zulfi, Rawdhat Al Sablh; 26°22.429'N, 44°58.241'E; 670 m a.s.l.; 26 Aug. 2015; H. Al Dhafer, M. Abdel-Deyem, A. El Torkey, A. El Gharbawy, & A. Solimanleg leg.; LT; J. Háva det.; KSMA.

**Note.** The female specimen was collected at a low elevation (670 m) in a sandy area in central Saudi Arabia (Fig. 4A). This represents a new record for Saudi Arabia.



**Figure 1.** Dorsal habitus and abdominal ventrites (photos by A. Herrmann) of *Dermestes* species **A, B** *D. ater* DeGeer, 1774 **C, D** *D. haemorrhoidalis* Küster, 1852 **E, F** *D. lardarius* Linnaeus, 1758.

**Distribution.** Nearly cosmopolitan (Háva 2015, 2022), where it is widely distributed in Europe; North Africa; Africa: Burundi, Congo, Madagascar, South Africa, Tanzania, Zambia; Asia: China (Liaoning), Iran, Japan, Mongolia, Oman, Russia, South Korea, Vietnam; Australia: New Zealand (introduced); North America: USA; South America: Argentina, Bolivia, Brazil, Peru, Uruguay.

***Dermestes (Dermestes) lardarius* Linnaeus, 1758**

Fig. 1E, F

**Material examined.** SAUDI ARABIA • 1 ♀; Makkah Province, Jeddah, “Ras Halibah” [Ras Hatibah]; 7 May. 1982; W. Büttiker leg.; JHAC.

**Note.** *Dermestes lardarius* was previously reported from Saudi Arabia without a specific locality (Hagstrum and Subramanyam 2009). The only known female representing this species in Saudi Arabia was collected from the coastal area in Jeddah (Makkah Province) (Fig. 4D).

**Distribution.** Cosmopolitan (Háva 2011, 2015, 2022).

**Subgenus *Dermestinus* Zhantiev, 1967**

***Dermestes (Dermestinus) frischii* Kugellan, 1792**

Fig. 2A–C

**Material examined.** SAUDI ARABIA • 2 ♂; Baha Province, Al Mandaq, Amadan; 20°12'11"N, 41°13'43"E; 14 Oct. 2010; H. Aldhafer & H. Fadl leg.; M.S. Abdel-Dayem det.; KSMA • 1 ex; Asir Province, Bareq, Thloth Al Mandhar, Wadi Baqrah; 18°47.476'N, 41°56.310'E; 331 m a.s.l.; 20 Apr. 2011; H. Fadl & H. Setyaningrum leg.; LT; M.S. Abdel-Dayem det.; KSMA • 1 ♀; Eastern Province, Al Jubail, Ras al Ghar; 26°15'34"N, 49°52'01"E; 16 Apr. 2010; H. Al Dhafer leg.; HP; J. Háva det.; KSMA • 3 ♂, 4 ♀; Eastern Province, An Nuayriyah, Al Sarar; 27°25'45.5"N, 48°27'00"E; 60 m a.s.l.; 2 Mar. 2011; H. Al Dhafer, H. Setyaningrum & A. Al Ansi leg.; collected from Carcasses; M.S. Abdel-Dayem det.; KSMA • 1 ex; same collection data as for preceding; J. Háva det.; KSMA • 5 ex; Eastern Province, Dammam, near shore; 26°21'3.744"N, 50°13'41.462"E; 3 m a.s.l.; 15 Oct. 2018; A. Alqurashi leg.; PT beside rabbit carcass, M.S. Abdel-Dayem det.; KSMA • 1 ♂; Makkah Province, Jeddah, Shoiba; 20°51'N, 39°24'E; 1 m a.s.l.; 19 Oct. 1982; W. Büttiker leg.; JHAC • 2 ♂, 1 ♀; Makkah Province, Taif, Al Wesam District; 21°12'17"N, 40°20'43"E; 11 Oct. 2010; H. Al Dhafer, B. Kondratieff, H. Fadl & A. El Gharbawy leg.; M.S. Abdel-Dayem det.; KSMA • 1 ♂, 1 ♀; Riyadh Province, Ad Diriah, Ad Diriah Desert; 6 May. 2010; H. Al Dhafer, A. El Gharbawy & A. El Torkey leg.; MT; M.S. Abdel-Dayem det.; KSMA • 1 ♂, 4 ♀; same collection data as for preceding; Al Amariyah, Animal Production Dept. Farm KSU; 31 Mar. 2008; LT; M.S. Abdel-Dayem det.; KSMA • 2 ♂, 3 ♀; same collection data as for preceding; Aljabilah, Prince Bander Farm; 26 Apr. 2008; M. Otybi leg.; LT; M.S. Abdel-Dayem det.; KSMA • 1 ♂; same collection data as for preceding; 3 May. 2008; M. Otybi leg.; LT; M.S. Abdel-Dayem det.; KSMA • 1 ♂; same collection data

as for preceding; 31 May. 2008; M. Otybi leg.; LT; M.S. Abdel-Dayem det.; KSMA • 1 ♀; same collection data as for preceding; Al Obaiteh, 50 km W. Riyadh, Obikan Farm; 7 May. 2007; M. Otybi leg.; LT; J. Háva det.; KSMA • 1 ♀; same collection data as for preceding; Thonyan Al Thonyan Farm; 28 Jul. 2007; H. Al Ayedh & H. Al Dhafer leg.; LT; M.S. Abdel-Dayem det.; KSMA • 3 ♂, 4 ♀; same collection data as for preceding; Education Farm KSU; 1 Apr. 2008; J. Háva det.; KSMA • 7 ♀; same collection data as for preceding; 2 Apr. 2008; M.S. Abdel-Dayem det.; KSMA • 4 ♂, 6 ♀; same collection data as for preceding; 3 Apr. 2008; M.S. Abdel-Dayem det.; KSMA • 14 ex; same collection data as for preceding; 5 Apr. 2008, M.S. Abdel-Dayem det.; KSMA • 7 ♂, 8 ♀; same collection data as for preceding; 7 Apr. 2008; M.S. Abdel-Dayem det.; KSMA • 2 ♂, 6 ♀; same collection data as for preceding; 9 Apr. 2008; M.S. Abdel-Dayem det.; KSMA • 1 ♂, 11 ♀; same collection data as for preceding 11 Apr. 2008; M.S. Abdel-Dayem det.; KSMA • 1 ♀; same collection data as for preceding; 20 Apr. 2011; H. Setyaningrum leg.; J. Háva det.; KSMA • 1 ♂; Riyadh Province, Alkharj, Al-Shahwan Farm; 24 Mar. 2010; A. Al-Hasbel leg.; SW; J. Háva det.; KSMA • 1 ♀; Riyadh Province, Huraymala, Wadi Huraymala; 770 m a.s.l.; 24 Nov. 1988; C.W. Mills leg.; J. Háva det.; FSCA • 1 ♀; Riyadh Province, Mozahmiya, Al Khararah; 24°24'21"N, 46°14'40"E; 17 Apr. 2012; H. Al Dhafer, H. Fadl, A. El Torkey, M. Abdel-Dayem & A. Al Ansi leg.; LT; M.S. Abdel-Dayem det.; KSMA • 1 ♂, 1 ♀; Riyadh Province, Rumah, Rawdhat khorim; 29 Apr. 2011; Y. Aldryhim leg.; LT; M.S. Abdel-Dayem det.; KSMA • 1 ex; same collection data as for preceding; 25°25.943'N, 47°13.863'E; 572 m a.s.l.; 6 Mar. 2012; PT; M.S. Abdel-Dayem det.; KSMA • 1 ex; same collection data as for preceding; 27 May. 2012; LT; M.S. Abdel-Dayem det.; KSMA.

**Note.** Mroczkowski (1979) documented this species in Jeddah. Recently it was collected at Jeddah from rabbit carcasses (Al-Shareef and Al-Mazyad 2017) and human remains (Al-Shareef and Zaki 2017), and at Riyadh from camel, dog, and goat carcasses (Mashaly et al. 2019) and human corpses (Alajmi et al. 2016). This species has also been collected from sheep carcasses in Riyadh, Jazan, and Arar (Mashaly et al. 2018). The listed specimens were collected at different elevations (7–1920 m) in the central, east, and lowlands and mountainous areas of southwest Saudi Arabia (Fig. 4B).

**Distribution.** Cosmopolitan (Háva 2007, 2015, 2022).


### *Dermestes (Dermestinus) maculatus* DeGeer, 1774

Fig. 2D–F

**Material examined.** SAUDI ARABIA • 2 ♀; Baha Province, Al Mandaq, Amadan; 20°12'11"N, 41°13'43"E; 14 Oct. 2010; H. Al Dhafer, B. Kondratieff, H. Fadl & A. El Gharbawy leg.; J. Háva det.; KSMA • 4 ♀; same collection data as for preceding; 14 Oct. 2010; H. Aldhafer & H. Fadl leg.; J. Háva det.; KSMA • 12 ex; Baha Province, Al Baha, Al-Baher Mountain; 15 Mar. 2010; J. Háva det.; KSMA • 1 ♀; Asir Province, Khamis Mushayt; 2050 m a.s.l.; 9 Jan. 1998; J. Háva det.; JHAC • 1 ex; Makkah Province; Taif; 21°12'17"N, 40°20'43"E; 11 Oct. 2010; H. Al Dhafer, B. Kondratieff, H. Fadl & A. El Gharbawy leg.; M.S. Abdel-Dayem det.; KSMA • 6 ex; Eastern Province; Dammam, near shore; 26°21'3.744"N, 50°13'41.462"E; 3 m a.s.l.; 15 Oct.



2018, A. Alqurashi leg.; PT beside rabbit carcass; M.S. Abdel-Dayem det.; KSMA • 1 ♀; Riyadh Province, Ad Diriyah, Al Amariyah, Animal Production Dept. Farm KSU; 23 Mar. 2011; H. Setyaningrum leg.; BT; M.S. Abdel-Dayem det.; KSMA • 1 ex; Riyadh Province, Riyadh; Oct.1989; M.S. Abdel-Dayem det.; KSMA • 4 ex; Riyadh Province, Ad Diriyah, Al Amariyah; 28 Jan. 2008; D. Boy Valenza leg., M.S. Abdel-Dayem det.; KSMA • 2 ex; same collection data as for preceding; Albeer Farm; 29 Oct. 2008; A. Al-Ahmari leg.; SW; M.S. Abdel-Dayem det.; KSMA • 1 ex, same collection data as for preceding; 8 Dec. 2010; SW; M.S. Abdel-Dayem det.; KSMA • 2 ♀; Riyadh Province, Ad Diriyah, Education Farm KSU; 1 Apr. 2008; J. Háva det.; KSMA • 1 ex; same collection data as for preceding; 31 Mar. 2008; M.S. Abdel-Dayem det.; KSMA • 1 ex; same collection data as for preceding; 2 Apr. 2008; M.S. Abdel-Dayem det.; KSMA • 1 ex; same collection data as for preceding; 3 Apr. 2008; M.S. Abdel-Dayem det.; KSMA • 3 ex; same collection data as for preceding; 5 Apr. 2008; M.S. Abdel-Dayem det.; KSMA • 1 ex; same collection data as for preceding; 7 Apr. 2008; M.S. Abdel-Dayem det.; KSMA • 3 ex; same collection data as for preceding; 9 Apr. 2008; M.S. Abdel-Dayem det.; KSMA • 2 ex; same collection data as for preceding; 11 Apr. 2008; M.S. Abdel-Dayem det.; KSMA • 1 ex; same collection data as for preceding; 3 Nov. 2009; H. Setyaningrum leg.; M.S. Abdel-Dayem det.; KSMA • 6 ex; Riyadh Province, Riyadh, Al-Wahah Farm; 15 May. 2021; M.S. Abdel-Dayem det.; KSMA • 1 ex; Riyadh Province, Shaqra; 21 May. 1978; HP from mill waste; J. Háva det.; KSMA.



**Figure 2.** Dorsal habitus, apical part of elytron and abdominal ventrites (photographs by A. Herrmann) of *Dermestes* species. **A–C** *D. frischii* Kugellan, 1792 **D–F** *Dermestes maculatus* DeGeer, 1774.

**Note.** *Dermestes maculatus* is the most common species within the subfamily Dermestinae in Saudi Arabia. It was previously collected from rabbit carcasses at Baha (Abouzied 2014), Al-Ahsa (Shaalan et al. 2017), τ Madinah (Al-Dakhil and Alharbi 2020), Jeddah (Al-Shareef and Al-Mazyad 2017), and Riyadh (Mashaly 2017). It has been collected from sheep carcasses in Jizan and Northern Border (Mashaly et al. 2018). Al-Qahtni et al. (2020) have reported the species from dead human bodies in Riyadh. Also, it was collected by other methods from Makkah Province, Jeddah (Shalaby 1961; Mroczkowski 1979) and Riyadh Province, Dierab (Elgharbawy 2018). The species was collected in both low- and highlands (10–2330 m) in the following provinces: Asir, Baha, Eastern Province, Madinah, Makkah, and Riyadh (Fig. 4C).

**Distribution.** Cosmopolitan (Háva 2011, 2015, 2022).

### *Dermestes (Dermestinus) undulatus* Brahm, 1790

Fig. 3A, B

**Material examined.** SAUDI ARABIA • 1 ♀; Asir Province, Khamis Mushayt; 2050 m a.s.l.; 9 Jan.1998; J. Háva det.; JHAC.

**Note.** The only specimen known (a female) was collected from the highlands in southwestern Saudi Arabia (Fig. 4A). This is a new record for Saudi Arabia.

**Distribution.** Holarctic species (Háva 2015, 2022).

### Tribe Marioutini Jacobson, 1913

#### Genus *Mariouta* Pic, 1898

### *Mariouta stangei* Reitter, 1910

Fig. 3C

**Record.** SAUDI ARABIA • Eastern Province, Al-Ahsa, Salwah, 248 km S (Rub al Khali) (Mroczkowski and Ślipiński 1997).

**Note.** This species is only known from a single specimen preserved in the NHMB collection. This specimen was collected by W. Büttiker in May 1985 at a location in the Empty Quarter (Rub al Khali), 248 km south of the town of Salwa in southeastern Saudi Arabia, located near the border with Qatar (Fig. 4D).

**Distribution.** This taxon is known from the Sultanate of Oman, Pakistan, Saudi Arabia, Sudan, and Turkmenistan (Háva 2015, 2022).


### Genus *Rhopalosilpha* Arrow, 1929

### *Rhopalosilpha wasmanni* Arrow, 1929


Fig. 3D

**Record.** SAUDI ARABIA • Eastern Province, Hofuf (Mroczkowski and Ślipiński 1997).





**Figure 3.** **A, B** *Dermestes undulatus* Brahm, 1790 (photographs by A. Herrmann) **A** dorsal habitus **B** Abdominal ventrites **C** *Mariouta stangei* Reitter, 1910 (photographs by K. Matsumoto) **D** *Rhopalosilpha wasmanni* Arrow, 1929 (photographs by J. Háva).



**Figure 4.** Distribution map of the Dermestinae species in Saudi Arabia **A** *Dermestes ater*, *D. haemorrhoidalis*, and *D. undulatus* **B** *D. frischii* **C** *D. maculatus* **D** *D. lardarius*, *Mariouta stangei*, and *Rhopalosilpha wasmanni*.

**Note.** *Rhopalosilpha wasmanni* is only known from a single specimen in the NHMB collection. It was collected from Hofuf in eastern Saudi Arabia by W. Büttiker (Mroczkowski and Ślipiński 1997) (Fig. 4D).

**Distribution.** This very rare species is known only from Iran, Jordan, and Saudi Arabia (Háva 2015, 2022).

## Discussion

The first forensic case being solved using insects was during the 13<sup>th</sup> century in China, while the first systematic studies of forensic entomology took place in Germany during the 19<sup>th</sup> century (Benecke 2001). Despite the earlier published data documenting the Saudi fauna of Dermestinae (Shalaby 1961; Mroczkowski 1979), forensic entomology in this country began only during the last two decades. Accordingly, several studies were conducted during this period in four areas: at the centre of the country (Alajmi et al. 2016; Mashaly 2017; Mashaly et al. 2018, 2019; Al-Qahtni et al. 2020), in the east (Shaalán et al. 2017), in the west (Al-Shareef and Al-Mazyad 2017; Al-Shareef and Zaki 2017; Al-Dakhil and Alharbi 2020), and in the southwest (Abouzied 2014). Taxonomic and faunal studies are needed to support this growing interest in forensic entomology in Saudi Arabia.

During the late stage of decay of animal remains, *Dermestes* species are one of the predominant taxa among forensic insects (Magni et al. 2015). *Dermestes frischii* and *D. maculatus* have been the most frequently documented dermestid beetles in forensic studies in Saudi Arabia (Alajmi et al. 2016; Mashaly et al. 2018). The current study listed eight species in three genera (*Dermestes*, *Mariouta*, and *Rhopalosilpha*) in two tribes (Dermestini and Marioutini) belonging to the subfamily Dermestinae. *Dermestes haemorrhoidalis* and *D. undulatus* are recorded for the first time from Saudi Arabia. Based on the world distribution range, the Saudi Dermestinae fauna is dominated by the widespread cosmopolitan or nearly cosmopolitan species, which includes all members of the tribe Dermestini (e.g., *Dermestes ater*, *D. haemorrhoidalis*, *D. lardarius*, *D. frischii*, *D. maculatus*, *D. undulatus*; Háva 2015, 2022) (Table 1), while the members of tribe Marioutini, *Mariouta stangei* and *Rhopalosilpha wasmanni* have a narrower distribution range and appear to have Saharo-Sindian elements (Háva 2015, 2022).

The analysis of data based on the examination of museum specimens and literature records revealed that *D. frischii* and *D. maculatus* are the most abundant and distributed over a fairly wide range in Saudi Arabia. These findings are consistent with what has been documented in several other studies (Shalaby 1961; Mroczkowski 1979; Abouzied 2014; Alajmi et al. 2016; Shaalan et al. 2017; Elgharbawy 2018; Al-Dakhil and Al-Harbi 2020). However, the remaining Dermestinae are rare or very rare species, either documented from a few specimens (e.g., *D. ater*) or a single specimen (e.g., *D. haemorrhoidalis*, *D. lardarius*, *D. undulatus*, *M. stangei*, and *R. wasmanni*). This may be due to different feeding behaviours or a rarity of these species in the Saudi fauna. Although *D. ater*, *D. haemorrhoidalis*, *D. lardarius*, and *D. undulatus* have been reported from human cadavers (Charabidze et al. 2014; Kadej et al. 2020), none of the forensic

**Table I.** List of *Dermestes* species from the Arabian Peninsula. Notes: recorded (\*) or not recorded (-).

	Kuwait	Saudi Arabia		Yemen		Oman	United Arab Emirates	Qatar
		Farasan Archipelago	Saudi Arabia mainland	Yemen mainland	Socotra Island			
<i>Dermestes (Dermestinus) maculatus</i> DeGeer, 1774	*	-	*	*	*	*	*	*
<i>Dermestes (Dermestinus) frischii</i> Kugellan, 1792	*	-	*	*	*	*	*	*
<i>Dermestes (Dermestinus) undulatus</i> Brahm, 1790	-	-	*	-	-	-	-	-
<i>Dermestes (Dermestes) ater</i> DeGeer, 1774	-	-	*	*	-	*	*	*
<i>Dermestes (Dermestes) haemorrhoidalis</i> Küster, 1852	-	-	*	-	-	-	-	-
<i>Dermestes (Dermestes) lardarius</i> Linnaeus, 1758	-	-	*	*	-	-	*	-

entomological studies in Saudi Arabia reported any of them. As for *Mariouta stangei* and *Rhopalosilpha wasmanni*, no information is available that documents their feeding habits.

Despite more than 60 years since the first faunistic study (Shalaby 1961), we may still have an imprecise idea about the actual species number and faunal composition of Dermestinae in Saudi Arabia. In conclusion, the few numbers of faunistic studies on the Dermestinae in Saudi Arabia and the registration of two new records in the current study indicate that there are more species that have yet to be discovered.

## Acknowledgements

We are grateful to Dr Iftekhar Rasool, Plant Protection Department, College of Food and Agriculture Sciences, King Saud University, Saudi Arabia, for processing and arranging specimens for dispatch. The authors extend their appreciation to the Deputyship for Research & Innovation, Ministry of Education in Saudi Arabia for funding this research work through the project no. IFKSURG-2-000.

## References

- Abdel-Dayem MS, Fad HH, El-Torkey AM, Elgharbawy AA, Aldryhim YN, Kondratieff BC, Al Ansi AN, Aldhafer HM (2017) The beetle fauna (Insecta, Coleoptera) of the Rawdhat Khorim National Park, Central Saudi Arabia. *ZooKeys* 653: 1–78. <https://doi.org/10.3897/zookeys.653.10252>
- Abdel-Dayem MS, Abu El-Ghiet UM, Elsheikh TM, Elgharbawy AA, Al-Fifi ZIA, Aldhafer HM (2020) The first survey of the Beetles (Coleoptera) of the Farasan Archipelago of the southern Red Sea, Kingdom of Saudi Arabia. *ZooKeys* 959: 17–86. <https://doi.org/10.3897/zookeys.959.51224>

- Abouzieid EM (2014) Insect colonization and succession on rabbit carcasses in Southwestern Mountains of the Kingdom of Saudi Arabia. *Journal of Medical Entomology* 51(6): 1168–1174. <https://doi.org/10.1603/ME13181>
- Al-Dakhil AA, Alharbi SA (2020) A preliminary investigation of the entomofauna composition of forensically important necrophagous insects in Al-Madinah Al Munawwarah region, Kingdom of Saudi Arabia. *Journal of Taibah University for Science* 14(1): 1127–1133. <https://doi.org/10.1080/16583655.2020.1805176>
- Alajmi RA, AlGhufaili H, Farrukh A, Aljohani H, Mashaly AM (2016) First report of necrophagous insects on human corpses in Riyadh, Saudi Arabia. *Journal of Medical Entomology* 53(6): 1276–1282. <https://doi.org/10.1093/jme/tjw113>
- Al-Qahni AH, Al-Khalifa MS, Mashaly AM (2020) Two human cases associated with forensic insects in Riyadh, Saudi Arabia. *Saudi Journal of Biological Sciences* 27(3): 881–886. <https://doi.org/10.1016/j.sjbs.2019.12.027>
- Al-Shareef LAH, Al-Mazyad MMF (2017) Beetles (Insecta, Coleoptera) associated with rabbit carcasses in three habitats in Jeddah, Kingdom of Saudi Arabia. *Australian Journal of Basic and Applied Sciences* 11(2): 139–145.
- Al-Shareef LAH, Zaki MK (2017) Arthropods associated with human remains and determination of postmortem interval in Jeddah, Kingdom of Saudi Arabia. *The Journal of American Science* 13(3): 106–114. <https://doi.org/10.7537/marsjas130317.11>
- Benecke M (2001) A brief history of forensic entomology. *Forensic Science International* 120(1–2): 2–14. [https://doi.org/10.1016/S0379-0738\(01\)00409-1](https://doi.org/10.1016/S0379-0738(01)00409-1)
- Charabidze D, Colard T, Vincent B, Pasquerault T, Hedouin V (2014) Involvement of larder beetles (Coleoptera: Dermestidae) on human cadavers: a review of 81 forensic cases. *International Journal of Legal Medicine* 128(6): 1021–1030. <https://doi.org/10.1007/s00414-013-0945-1>
- Elgharbawy AA (2018) Survey and population studies of Insects at Dierab Area, South Riyadh, Saudi Arabia. *Al Azhar Bulletin of Science* 29(1): 47–54. <https://doi.org/10.21608/absb.2018.33763>
- Hagstrum DW, Subramanyam B (2009) *Stored-product Insect Resource*. American Association of Cereal Chemists International, St. Paul, 509 pp.
- Háva J (2007) New species and new records of Dermestidae (Insecta: Coleoptera) from the Arabian Peninsula including Socotra Island. *Fauna of Arabia* 23: 309–317.
- Háva J (2011) Contribution to the Dermestidae (Coleoptera) from the Arabian Peninsula—1. *Latvijas Entomologs* 50: 5–8.
- Háva J (2015) *World Catalogue of Insects*. (Vol. 13). Dermestidae (Coleoptera). Brill, Leiden/Boston, [xxvi +] 419 pp. <https://doi.org/10.1163/9789004286610>
- Háva J (2022) *Dermestidae World (Coleoptera)* [Version 2018, updated January 2022]. <https://doi.org/10.24394/NatSom.2022.38.19> [Accessed on: 2022-12-19]
- Kadej M, Szleszkowski Ł, Thannhäuser A, Jurek T (2020) A mummified human corpse and associated insects of forensic importance in indoor conditions. *International Journal of Legal Medicine* 34(5): 1963–1971. <https://doi.org/10.1007/s00414-020-02373-2>
- Magni PA, Voss SC, Testi R, Borrini M, Dadour IR (2015) A biological and procedural review of forensically significant *Dermestes* species (Coleoptera: Dermestidae). *Journal of Medical Entomology* 52(5): 755–769. <https://doi.org/10.1093/jme/tjv106>

- Mashaly AM (2017) Carrion beetles succession in three different habitats in Riyadh, Saudi Arabia. *Saudi Journal of Biological Sciences* 24(2): 430–435. <https://doi.org/10.1016/j.sjbs.2016.02.015>
- Mashaly AM, Al-Ajmi RA, Al-Johani HA (2018) Molecular identification of the carrion beetles (Coleoptera) in selected regions of Saudi Arabia. *Journal of Medical Entomology* 55(6):1423–1430. <https://doi.org/10.1093/jme/tjy116>
- Mashaly AM, Al-Ajmi RA, Rady A, Al-Musawi Z, Farrukh A (2019) Species richness of scavenger insects on different carcass types. *Tropical Biomedicine* 36(3): 630–639.
- Motyka M, Kusy D, Háva J, Jahodářová E, Bílková R, Vogler AP, Bocak L (2022) Mitogenomic data elucidate the phylogeny and evolution of life strategies of Dermestidae (Coleoptera). *Systematic Entomology* 47(1): 82–93. <https://doi.org/10.1111/syen.12520>
- Mroczkowski M (1979) Insects of Saudi Arabia. (Coleoptera, Dermestidae). Part 1, faunistic data. *Fauna of Saudi Arabia* 1: 212–214.
- Mroczkowski M, Ślipiński SA (1997) Notes on the Marioutinae (Coleoptera: Dermestidae) with a review of the described species. *Annales Zoologici* 47: 11–16.
- Shalan EA, El-Moaty ZA, Abdelsalam S, Anderson GS (2017) A preliminary study of insect succession in Al-Ahsaa Oasis, in the Eastern region of the Kingdom of Saudi Arabia. *Journal of Forensic Sciences* 62(1): 239–243. <https://doi.org/10.1111/1556-4029.13252>
- Shalaby F (1961) A preliminary survey of the insect fauna of Saudi Arabia. *Bulletin de la Société Entomologique d’Egypte* 45: 211–228.