

A new solitary free-living species of the genus *Sphenopus* (Cnidaria, Anthozoa, Zoantharia, Sphenopidae) from Okinawa-jima Island, Japan

Takuma Fujii^{1,2}, James Davis Reimer^{3,4}

1 Research Center for the Pacific Islands Amami Station, Kagoshima University, Naze-Yanagimachi 2-1, Amami, Kagoshima 894-0032, Japan **2** Graduate School of Engineering and Science, University of the Ryukyus, 1 Senbaru, Nishihara-cho, Okinawa 903-0213, Japan **3** Molecular Invertebrate Systematics and Ecology Laboratory, Department of Biology, Chemistry & Marine Sciences, Faculty of Science, University of the Ryukyus, 1 Senbaru, Nishihara, Okinawa 903-0213, Japan **4** Tropical Biosphere Research Center, University of the Ryukyus, 1 Senbaru, Nishihara, Okinawa 903-0213, Japan

Corresponding author: Takuma Fujii (tfujii@cpi.kagoshima-u.ac.jp)

Academic editor: B.W. Hoeksema | Received 23 May 2015 | Accepted 29 June 2016 | Published 21 July 2016

<http://zoobank.org/EB98FE3B-665B-4CF2-8E8B-740D167BA2BB>

Citation: Fujii T, Reimer JD (2016) A new solitary free-living species of the genus *Sphenopus* (Cnidaria, Anthozoa, Zoantharia, Sphenopidae) from Okinawa-jima Island, Japan. ZooKeys 606: 11–24. doi: 10.3897/zookeys.606.9310

Abstract

A new species of free-living solitary zoantharian is described from Okinawa, Japan. *Sphenopus exilis* **sp. n.** occurs on silty seafloors in Kin Bay and Oura Bay on the east coast of Okinawa-jima Island. *Sphenopus exilis* **sp. n.** is easily distinguished from other *Sphenopus* species by its small polyp size and slender shape, although there were relatively few differences between *Sphenopus exilis* **sp. n.** and *S. marsupialis* in the molecular phylogenetic analyses. Currently, very little is known about the ecology and diversity of *Sphenopus* species. Thus, reviewing each species carefully via combined morphological and molecular analyses by using newly obtained specimens from type localities is required to clearly understand and distinguish the species within the genus *Sphenopus*.

Keywords

Zoantharia, *Sphenopus*, new species, free-living, enclosed bay, identification key

Introduction

The suborder Brachycnemina (Cnidaria: Anthozoa: Hexacorallia: Zoantharia) consists of zoantharians commonly found in shallow warm waters, as almost all species within this group contain endosymbiotic photosynthetic *Symbiodinium* spp. (e.g. Swain 2010). The genus *Sphenopus* belongs to the family Sphenopidae within Brachycnemina based on its brachycnemic mesenterial arrangement, mesogleal sphincter muscles, and heavy encrustation of granules into the body column. This genus is unique as *Sphenopus* individuals consist of free-living solitary polyps, in contrast to not only other Sphenopidae and Brachycnemina species but also to all other known zoantharians. This unique feature is considered to be an adaptation to *Sphenopus*' muddy/sandy sea floor habitats that are difficult for most sessile benthos to inhabit (e.g. Soong et al. 1999, Reimer et al. 2012). Here they risk burial, which would require them to shed sediments in order to survive as seen in free-living scleractinians (e.g. Schuhmacher 1977, Fisk 1982, Bongaerts et al. 2013, Sentoku et al. 2016).

Three species are currently considered valid within the genus *Sphenopus*; *S. marsupialis* (Gmelin, 1791), *S. arenaceus* Hertwig, 1882, and *S. pedunculatus* Hertwig, 1888. In contrast to *S. marsupialis* with a wide distribution in the Indo-Pacific (Soong et al. 1999, Reimer et al. 2012, 2014), there have been no further records of *S. arenaceus* and only one additional record of *S. pedunculatus* (in Reimer et al. 2014) after their original descriptions. Although *S. marsupialis* has been reported in some field guides, formal taxonomic studies based on specimens are limited (Soong et al. 1999, Reimer et al. 2012). Thus, comparatively very little is known about the species diversity of the genus *Sphenopus*.

Recently, we discovered comparatively small *Sphenopus* specimens (polyp lengths <2.5 cm) from the shallow silty seafloors of enclosed bays on the east coast of Okinawa-jima Island, Japan. Combined morphological and molecular phylogenetic analyses lead us to conclude that the specimens belong to a previously unknown species. Thus, in this paper, a new *Sphenopus* species is formally described and a dichotomous key to identify all known *Sphenopus* species is provided. This report represents only the second formal record of this genus in Japan after Reimer et al. (2016).

Material and methods

Sample collection. Specimens from Okinawa were collected by SCUBA. Prior to collecting, *in situ* images of expanded polyps were taken to assist in morphological analyses (colour, tentacle counts and size, polyp form). Half of the specimens collected were preserved in 99% EtOH for DNA analyses, and the other specimens were fixed for morphological analyses in 5 to 10% formalin sea water after anesthesia using MgCl₂, and subsequently transferred to 70% EtOH some days later.

Morphological analyses. The lengths, maximum widths (largest diameter of column) and minimum widths (width at the top of physa where the aboral ampullaceous ends) of the column of preserved polyps were measured using calipers to the nearest




Figure 1. Diagram of external morphology of a contracted polyp of *Sphenopus exilis* sp. n.. ca = capitulum; cr = capitular ridge; fo = foot; le = length; max = maximum width; min = minimum width; ph = physa; sc = scapus, column.

0.1 mm (Figure 1). Gross shape of polyps, color of polyps, and numbers and lengths of the tentacles were recorded utilizing in situ images. Internal morphology was observed from horizontally and longitudinally hand-cutting polyps through the actinopharynx using a dissecting microscope.

Cnidae. Undischarged cnidae were measured from small pieces of tissue from the tentacles, column (external portion), actinopharynx, and mesenterial filaments of specimen NSMT-Co1576 (MISE-TF-107; fixed in 5–10% formalin seawater). Images of cnidae were obtained by differential interference contrast microscopy, and measured using the software ImageJ (National Institute of Health, Bethesda, Maryland, USA). Cnidae nomenclature generally followed England (1991) and Ryland and Lancaster (2003). However, both Schmidt (1974) and Hidaka et al. (1987, 1992) have suggested basitrichs and microbasic b-mastigophores are the same type of nematocyst, and in this study, as in recent zoantharians studies (e.g. Kise and Reimer 2016, Ryland and Ward 2016), these two types were treated as the same.

DNA processing and amplification. DNA was extracted from ethanol preserved specimens by following a guanidine extraction protocol (Sinniger et al. 2010). PCR amplifications were performed for mitochondrial cytochrome oxidase subunit I (COI), mitochondrial 16S ribosomal DNA (mt 16S rDNA), and the internal transcribed spacer region of ribosomal DNA (ITS-rDNA) region using the primer pairs

HCO and LCO (Folmer et al. 1994), 16SarmL (modified primer for mt 16S rDNA used in Sinniger et al. 2008, see Fujii and Reimer 2011) and 16SBmoH (Sinniger et al. 2005), and ITSf and ITSr (Swain 2009), respectively. Amplified PCR products were sequenced in both directions by Fasmac (Atsugi, Kanagawa, Japan).

Phylogenetic analyses. New sequences obtained in this study were deposited in GenBank (accession numbers: COI, KX400760–KX400768; mt 16S rDNA, KX400756–KX400759; ITS-rDNA, KX400769–KX400772). Obtained DNA sequences were manually aligned using Bioedit ver. 7.1.3.0 (Hall 1999). The nucleotide sequences of mt 16S rDNA, COI, and the ITS-rDNA region from specimens were separately aligned with previously obtained Sphenopidae (*Palythoa* and *Sphenopus*) sequences deposited in GenBank. Some sequences that were too short in length were removed from the analyses. For outgroups, sequences of *Zoanthus sansibaricus* (suborder Brachycnemina, family Zoanthidae) were used for all three DNA alignments' trees. Indels were kept unedited in the alignments of mt 16S rDNA. All phylogenetic alignments are available from the corresponding author.

For phylogenetic analyses of mt 16S rDNA, COI, and ITS-rDNA the same methods were independently applied. The maximum-likelihood (ML) method was performed using MEGA5 (Tamura et al. 2011), with 500 replicates performed using an input tree generated by BIONJ with the general time-reversible model (Rodriguez et al. 1990) of nucleotide substitution incorporating invariable sites and a discrete gamma distribution (eight categories) (GTR+I+C). The proportion of invariable sites, a discrete gamma distribution, and base frequencies of the model were estimated from the dataset. Bayesian trees for nuclear ITS-rDNA region were made by Mr. Bayes 3.2.5 (Ronquist and Huelsenbeck 2003) under GTR+I+C. One cold and three heated Markov chains Monte Carlo (MCMC) with default-chain temperatures were run for 10 million generations, sampling log-likelihoods (InLs), and trees at 100-generation intervals (100,000 InLs and trees were saved during MCMC). The likelihood plots for ITS-rDNA datasets suggested that MCMC reached the stationary phase after the first 10,000 generations (standard deviation of split frequencies = 0.004361). Thus, the remaining 90,000 trees of ITS-rDNA were used to obtain clade probabilities and branch-length estimates.

Results

Suborder Brachycnemina Haddon and Shackleton, 1891

Family Sphenopidae Hertwig, 1882

Genus *Sphenopus* Steenstrup, 1856

Sphenopus exilis sp. n.

<http://zoobank.org/30C107C6-8104-4EC9-9DC9-CF2DEE4A9638>

Figures 2, 3

Holotype. Specimen number NSMT-Co1576 (MISE-TF-107): Kin Bay, Uruma, Okinawa-jima Island, Japan (26°22'25"N, 127°53'30"E), 15 m depth, collected by

Takuma Fujii, 29 October 2011, fixed in 5–10% SW formalin, deposited in National Museum of Nature and Science, Tokyo, Japan (NSMT). Polyp length 2.4 cm, maximum width 0.8 cm, minimum width 0.3 cm. Figure 2B.

Paratypes. Specimen number NSMT-Co1577 (MISE-TF-107), a lot of total 11 polyps collected on the same dive, collection data same as holotype, five polyps fixed in 5–10% formalin, six polyps fixed in 99% EtOH, polyp length 1.3 to 2.2 cm (average 1.7 ± 0.3 cm), maximum width 0.4 to 1.0 cm (average 0.5 ± 0.2 cm), minimum width 0.2 cm, deposited in NSMT. GenBank accession numbers: COI, KX400760–KX400768; mt 16S rDNA, KX400756–KX400759; ITS-rDNA, KX400769–KX400772. Figure 2A and B; Specimen number RMNH Coel. 42121 (MISE-TF-144): a lot of total 16 polyps collected on the same dive, Kin Bay, Uruma, Okinawa-jima Island, Japan ($26^{\circ}22'25''\text{N}$, $127^{\circ}53'30''\text{E}$), 15 m depth, collected by Takuma Fujii, 24 May 2012, 11 polyps fixed in 5–10% formalin, five polyps fixed in 99% EtOH, polyp length 1.1 to 2.2 cm (average 1.7 ± 0.4 cm), maximum width 0.4 to 0.5 cm (average 0.5 ± 0.1 cm), minimum width 0.1 to 0.3 cm (average 0.2 ± 0.1), deposited in Naturalis Biodiversity Center, Leiden, Netherlands (RMNH); Specimen number NSMT-Co1578 (MISE-TF-151), a lot of total six polyps collected on the same dive, Oura Bay, Nago, Okinawa-jima Island, Japan ($26^{\circ}32'29''\text{N}$, $128^{\circ}3'16''\text{E}$), 17 m depth, collected by Takuma Fujii, 13 November 2012, five polyps fixed in 5–10% formalin, 1 polyp fixed in 99% EtOH, polyp length 1.0 to 1.9 cm (average 1.5 ± 0.4 cm), maximum width 0.3 to 1.1 cm (average 0.7 ± 0.3 cm), minimum width 0.1 to 0.3 cm (average 0.2 ± 0.1 cm), deposited in NSMT. Figure 2C and D.

Diagnosis: external morphology. Solitary, cylindrical polyp. Length of polyps 1.0 to 2.4 cm (average 1.7 ± 0.3 cm), maximum width 0.3 to 1.1 cm (average 0.6 ± 0.2 cm), minimum width 0.1 to 0.3 cm (average 0.2 ± 0.1 cm) ($n=34$). Tentacles longer than half diameter of the expanded oral disc (Figure 2A). Oral disc gently hollowing into mouth, with stellate grooves as many as tentacles (Figure 2A, C). Capitular ridges present but not strongly pronounced when polyps closed (Figure 1). The upper part of the polyp between capitulum and the column slightly constricted (the width of the most constricted region approximately 0.1 cm to 0.4 cm thinner than the width of contracted capitulum) when polyp contracted (Figures 1, 2B, D). Upper part of the column generally thick and oval (Figures 1, 2B, D). Aboral narrow bottom portion of column extended (=foot), thinner than upper portion of column, like a cone (Figures 1, 2B, D), with the distal portion round and thicker than the extended foot (=physa) (Figures 1, 2B, D). Column smooth, with encrusted fine dense sand particles. Occasionally broken piece(s) of bivalve shells attached to the aboral end (Figure 2B).

Diagnosis: internal morphology. Fine sand particles heavily encrusted into ectoderm and mesoglea. Mesenteries in brachycnemic arrangement. Mesentery number 36, complete 18, incomplete 18 (Figure 3A; $n=6$ polyps). Single siphonoglyph apparent. Mesogleal sphincter muscle well developed, visible under dissecting microscope (Figure 3B). Endosymbiotic *Symbiodinium* spp. (zooxanthellae) absent (=azooxanthellate).

Diagnosis: cnidae. Basitrichs and spirocysts in tentacles and actinopharynx. Basitrichs, holotrichs, microbasic p-mastigophores and basitrichs in mesenterial filaments. Holotrichs in column (Table 1).










Figure 2. Polyps of *Sphenopus exilis* sp. n. **A** In situ image of *S. exilis* sp. n., polyp with no black patterns, from the type locality in Kin Bay, Okinawa, Japan on 29 October 2011 **B** Polyps of NSMT-Co1576 & NSMT-Co1577 from Kin Bay, Okinawa-jima Island, Japan. The white circle points to the holotype **C** In situ image of NSMT-Co1578 from Oura Bay, Okinawa-jima Island, Japan, on 13 November 2012. Faint black patterns and bands appear on the oral disc and the tentacles **D** Polyps of lot number NSMT-Co1578 showing phenotypic variation with black stripes on the upper part of the polyps. Scale bars: 1 cm.

Habitat. Specimens were found at approximately 10 to 20 m depths on the slopes of silty seafloors in enclosed bays. Most polyps semi-burrowed in silt, with only the open oral disc visible and protruding out from the seafloor.

Colour. Tentacles and oral disc whitish and translucent in life. Faint black narrow horizontal bands appear on tentacles, and similar faint patterns on the oral disc of a few polyps (Figure 2C). Column colour of encrusted sand particles, a few polyps with 2 to 6 faint black vertical stripes approximately 15 mm wide on the upper part of the column, reaching from oral end to aboral end (Figure 2C, D).

Table 1. Cnidae types and sizes in different tissue sections of the holotype of *Sphenopus exilis* sp. n.

| | | Image (Scale bars: 50 μ m) | Length* | Width* | Frequency** |
|---------------|------------------------|---|---------------------|---------------------|---------------------|
| Tentacle | Basitrich |  | 10.4 (21.4–19.1) | 3.0 (2.6–3.3) | Occasional (n=8) |
| | Spirocyst |  | 13.1 (12.0–14.5) | 2.6 (2.4–2.8) | Numerous (n=20) |
| Column | Holotrich |  | 26.7 (16.3–34.5) | 14.7 (12.5–16.2) | Rare (n=3) |
| Actinopharynx | Holotrich |  | 28.0 (21.0–34.9) | 10.5 (5.5–15.5) | Rare (n=2) |
| | Basitrich |  | 26.9 (26.4–27.4) | 4.4 (3.9–5.0) | Numerous (n=20) |
| | Spirocyst |  | 13.4 (10.8–15.2) | 2.9 (2.8–2.9) | Common (n=13) |
| Filaments | Holotrich |  | 21.4 (17.0–25.6) | 10.4 (5.9–15.25) | Occasional (n=4) |
| | Basitrich |  | 31.1 (30.6–31.6) | 3.2 (2.8–3.5) | Numerous (n=20) |
| | <i>p</i> -mastigophore |  | 15.4 (14.6–16.2) | 4.4 (3.7–4.9) | Occasional (n=5) |

*Length and width: average, minimum–maximum, all sizes in μ m.

**Frequency: n=number of examined cnidae in these analyses. Frequency in decreasing order; numerous, common, occasional, rare.

Etymology. Named from latin ‘exilis’ meaning ‘slender’ or ‘small’, as polyps have an elongate and narrow foot more slender than other known species in this genus to the exception of *S. pedunculatus*. Polyps of this species are also much smaller than those of all three other species in the genus.

Common name. Hime-daruma-sunaginchaku (new Japanese name)

Molecular phylogeny. The results of the phylogenetic analyses of both mitochondrial cytochrome oxidase subunit I (COI) and 16S rDNA showed very few differences between sequences of our specimens and those of *S. marsupialis*, as well as compared with those of various *Palythoa* species. These results are not incongruous with previous studies on the molecular phylogeny of family Sphenopidae, where intra-family variation levels of mitochondrial DNA sequences were relatively low (Reimer et al. 2006, 2012).




Figure 3. Morphological features of *Sphenopus exilis* sp. n. **A** Cross section of holotype NSMT-Co1576 through the actinopharynx showing the mesenterial arrangement and dense sand encrustations **B** Well-developed mesogleal sphincter muscles visible on a hand-cut longitudinal section of the holotype NSMT-Co1576 **C** Comparison of polyp shape between *S. exilis* sp. n. NSMT-Co1577 and *S. marsupialis* (from Brunei, refer to Reimer et al. 2012).

The results of the phylogenetic analyses of nuclear internal transcribed spacer rDNA region showed *S. exilis* sp. n. forming a well-supported clade in the maximum likelihood and Bayesian analyses (Figure 4; ML=94%, Bayes=0.91). As well, together with sequences of *S. marsupialis*, *S. exilis* sp. n. formed a strongly supported *Sphenopus* clade (Figure 4; ML=99%, Bayes=1.00). In comparing the ITS-rDNA sequences between *S. exilis* sp. n. and *S. marsupialis*, there were 12 to 27 b.p. differences over a total 470 b.p. (=2.5–5.7% difference).




Figure 4. Maximum likelihood tree of nuclear internal transcribed spacer of ribosomal DNA (ITS-rDNA) region for newly obtained sequences from *Sphenopus exilis* sp. n. in this study along with previously published GenBank sequences of family Sphenopidae. Bootstrap values of ML >60% are shown at respective nodes. Nodes supported by Bayesian posterior probabilities >0.90 are marked with asterisks. Species names of sequences obtained from GenBank follow with accession numbers. The subtree shown in **b**) shows only the clade formed by genus *Sphenopus*, *Palythoa mizigama* and *P. umbrosa*, delineated by the gray square in **a**).

Remarks. Until now three species have been considered valid within *Sphenopus*; *S. marsupialis* (Gmelin, 1791), *S. arenaceus* Hertwig, 1882, and *S. pedunculatus* Hertwig, 1888. *S. exilis* sp. n. is easily distinguished from these other species by its small polyp size (length of *S. exilis* sp. n. <2.5 cm and width <1 cm), and by the shape of its elongated foot and physa. Polyps of both *S. marsupialis* and *S. arenaceus* are round on the aboral end, and not elongated as in *S. exilis* sp. n. (Figure 3C). Soong et al. (1999) examined various sized *S. marsupialis* collected from around Taiwan including small polyps without any narrow elongated foot (length < 2 cm). Additionally, Reimer et al. (2016) recently reported on a *S. marsupialis* specimen of the typical rounded shape and large size (~9 cm in height) from Okinawa-jima Island. No polyps with intermediate morphology between *S. marsupialis* and *S. exilis* sp. n. have ever been found. Thus, the specimens collected in this study cannot be considered to be immature polyps of *S. marsupialis*. The morphologically most similar species to *S. exilis* sp. n. is *S. pedunculatus* as it also has a narrow foot, but *S. pedunculatus* is much larger than *S. exilis* sp. n., with polyp lengths of 2.4 to 3.2 cm and widths of 2 to 2.4 cm, and with approximately 60 mesenteries. As well, the aboral end of *S. pedunculatus* is shaped like a clasping disc, different from that of *S. exilis* sp. n. with a narrow rounded shape (Hertwig 1888, Reimer et al. 2014).

In contrast to the morphological differentiation from other *Sphenopus* species, only a few differences were found in molecular analyses. The COI sequences of *S. exilis* sp. n. were identical to those of *S. marsupialis*, *Palythoa tuberculosa* (Esper, 1805), and

P. umbrosa Irei, Singer & Reimer, 2015. However, it is known that the evolutionary rate of mitochondrial DNA markers is quite slow in most Anthozoa (Shearer et al. 2004; Huang et al. 2008; Stampar et al. 2014), and the nuclear ITS-rDNA region is currently the fastest evolving DNA marker that has been utilized for species-level analyses of suborder Brachycnemina (Reimer et al. 2007). Although there are only relatively few differences between the ITS-rDNA sequences of *S. exilis* sp. n. and *S. marsupialis* (2.5–5.7% sequence divergence), the formation of a supported monophyletic clade confirms the results of our morphological analyses that the specimens collected in this study belong to a species different from *S. marsupialis* (Figure 4). Moreover, these results suggest the possibility of the presence of multiple, cryptic species within *S. marsupialis* as previously mentioned by Soong et al. (1999).

Currently, very little is known about the ecology and species diversity of the genus *Sphenopus*, demonstrated by the fact that there have been no or few records of both *S. arenaceus* and *S. pedunculatus* within the last 100 years. Thus, morphological and molecular analyses of newly obtained specimens from type localities followed by reviewing each species' description carefully are required to clearly understand the species distinction of *Sphenopus* species. As mentioned in previous studies, the phylogenetic results of this study indicate a need to re-examine the validity of the genus *Sphenopus* as it is positioned within the genus *Palythoa*, and by extension the definition of genera within the family Sphenopidae should be reconsidered (Reimer et al. 2012, Irei et al. 2015).

In the ITS-rDNA molecular phylogeny, it is notable that two recently described azooxanthellate *Palythoa* species from caves, *P. umbrosa* and *P. mizigama*, form a well-supported subclade with *S. exilis* sp. n. and *S. marsupialis*. As the phylogenetic relationship between *Sphenopus* and *Palythoa* is not yet clear, and likely does not reflect the traditional taxonomy (Reimer et al. 2012), construction of a large ITS-rDNA phylogeny with additional sequences from other *Palythoa* and *Sphenopus* species is needed. At the same time, investigation with additional DNA markers besides from the mt DNA and ITS-rDNA currently utilized in zoantharian phylogeny may be helpful.

Key to species of genus *Sphenopus*

- 1 Aboral end rounded, column shape oval, never having a very narrow stalk ...2
- Aboral end elongated and narrow, forming a foot3
- 2 Polyp colored earthy gray ***S. marsupialis***
- Polyp colored rusty red ***S. arenaceus***
- 3 Aboral end forms clasping disc, or the narrow stalk part of younger polyps very short compared to the oval part of the column. Comparatively large polyps (=polyp lengths > 2.4 cm, width > 2 cm) with approximately 60 mesenteries ***S. pedunculatus***
- Aboral end forms a rounded anchor. Comparatively small polyps (length < 2.4 cm, width < 2 cm), and approximately 36 mesenteries ***S. exilis* sp. n.**

Discussion

Unlike some other recently described zoantharian taxa such as Nanozoanthidae and Microzoanthidae, *S. exilis* sp. n. is not very small in size and does not inhabit a cryptic habitat. However, silty, sandy and rubble habitats are often overlooked in biodiversity surveys in favor of coral reef habitats, and this sampling bias has resulted in a relative lack of understanding of the diversity of these habitats and the evolutionary position of their inhabitants (Sheppard 1981, Obuchi et al. 2010, Fujii and Reimer 2011, Hoeksema 2012). In addition, not only are silty and muddy habitats less understood, but their ecosystem service value is often underestimated as well. For example, silty and muddy ecosystems on Okinawa-jima Island have been degraded by landfill dredging, and other development (Coral Reefs of Japan 2004), and over 35% of muddy tidal flats and shallow waters in Kin Bay have been lost (Uruma City Cultural Sea Museum 2007). Currently, *S. exilis* sp. n. is only known from two bays on the east coast of Okinawa-jima Island. Kin Bay, the type locality of *S. exilis* sp. n., has undergone ecological degradation over the past approximately 40 years (Reimer et al. 2015). The other locality, Oura Bay, is currently a center of controversy over proposed landfill and military base construction, while it is also known to house a unique benthic community (Fujii et al. 2015).

Thus, not only *S. exilis* sp. n., but also the diverse and various organisms that exclusively inhabit soft substrates in coral reef regions in the world face issues of decreasing habitat despite our lack of knowledge of their biodiversity (e.g. Cnidaria: Ceriantharia: Spier et al. 2012; Annelida: Polychaeta: Godet et al. 2008). *Sphenopus exilis* sp. n. serves as a clear reminder of how little we know of these ecosystems. It also provides a clear reason for better conservation and more exploration of the remaining silty and muddy areas.

Acknowledgements

We thank Tadashi Yamada (Yamato Co., Ltd.), Shin Nishihira (Diving Team Snack Snuffkin), and Dr. Masami Obuchi (University of the Ryukyus) for supporting collection of specimens, and we also thank Fumihito Iwase (Shikoku Marine Life Laboratory) for providing valuable information and comments. We are also grateful to the reviewer and the editor for improving this manuscript. The first author was supported by a research fellowship from the Japan Society for the Promotion of Science for young scientists, JSPS KAKENHI (24-3048), and the Special Budget of MEXT in 2015 (Establishment of Research and Education Network on Biodiversity and its Conservation in the Satsunan Islands). The second author was funded by a Japan Society for the Promotion of Science (JSPS) 'Zuno-Junkan' grant entitled 'Studies on origin and maintenance of marine biodiversity and systematic conservation planning', and a JSPS 'Kiban B' grant entitled 'Global evolution of Brachycnemina and their *Symbiodinium*'.

References

- Bongaerts P, Hoeksema BW, Hay KB, Hoegh-Guldberg O (2012) Mushroom corals overcome live burial through pulsed inflation. *Coral Reefs* 31(2): 399. doi: 10.1007/s00338-011-0862-z
- Coral Reefs of Japan (2004) Japan Ministry of Environment & Japan Coral Reef Society, Tokyo.
- England KW (1991) Nematocysts of sea anemones (Actiniaria, Ceriantharia and Corallimorpharia: Cnidaria): nomenclature. *Hydrobiologia* 216(1): 691–697. doi: 10.1007/BF00026532
- Fisk DA (1982) Sediment shedding and particulate feeding in two free-living, sediment-dwelling corals (*Heteropsarnmia cochlea* and *Heterocyathus aequicostatus*) at Wistari Reef, Great Barrier Reef. *Proceedings of the 4th International Coral Reef Symposium 2*: 21–26.
- Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R (1994) DNA primers for amplification of mitochondrial cytochrome oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology* 3(5): 294–299.
- Fujii T, Reimer JD (2011) Phylogeny of the highly divergent zoanthid family Microzoanthidae Anthozoa, Hexacorallia) from the Pacific. *Zoologica Scripta* 40(4): 418–431. doi: 10.1111/j.1463-6409.2011.00479.x
- Fujii T, Watanabe K, Nishihara C, Obuchi M, Ohba H (2015) The giant brown alga *Sargassum carpophyllum* on a nearshore coral reef in Okinawa Island, Japan. *Marine Biodiversity* 45(4): 603–604. doi: 10.1007/s12526-014-0280-x
- Godet L, Toupoint N, Oliver F, Fournier J, Retière C (2008) Considering the functional value of common marine species as a conservation stake: the case of sandmason worm *Lanice conchilega* (Pallas 1766) (Annelida, Polychaeta) beds. *A Journal of the Human Environment* 37(5): 347–355. doi: 10.1579/07-A-317.1
- Haddon AC, Shackleton AM (1891) Revision of the British Actinae, Part II. The Zoantheae. *Scientific Transactions of the Royal Dublin Society* 4: 609–672.
- Hall TA (1999) BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98 / NT. *Nucleic Acids Symposium Series* 41: 95–98.
- Hertwig R (1882) Report on the Actiniaria dredged by H.M.S. Challenger during the years 1873–1876. *Reports on the Scientific Results of the Exploring Voyage of H.M.S. Challenger during the years 1873–1876, Zoology* 6: 1–134.
- Hertwig R (1888) Report on the Actiniaria dredged by H.M.S. Challenger during the years 1873–1876, Supplement. *Reports on the Scientific Results of the Exploring Voyage of H.M.S. Challenger during the years 1873–1876, Zoology* 26: 1–56.
- Hidaka M (1992) Use of nematocyst morphology for taxonomy of some related species of scleractinian corals. *Galaxea* 11(1): 21–28.
- Hidaka M, Miyazaki I, Yamazato K (1987) Nematocysts characteristic of the sweeper tentacles of the coral *Galaxea fascicularis* (Linnaeus). *Galaxea* 6(1): 195–207.
- Hoeksema BW (2012) Evolutionary trends in onshore-offshore distribution patterns of mushroom coral species (Scleractinia: Fungiidae). *Contributions to Zoology* 81(4): 199–221.
- Huang D, Meier R, Todd PA, Chou LM (2008) Slow mitochondrial COI sequence evolution at the base of the Metazoan tree and its implications for DNA barcoding. *Journal of Molecular Evolution* 66(2): 164–174. doi: 10.1007/s00239-008-9069-5

- Irei Y, Sinniger F, Reimer JD (2015) Descriptions of two azooxanthellate *Palythoa* species (Subclass Hexacorallia, Order Zoantharia) from the Ryukyu Archipelago, southern Japan. *Zookeys* 478: 1–26. doi: 10.3897/zookeys.478.8512
- Kise H, Reimer JD (2016) Unexpected diversity and a new species of *Epizoanthus* (Anthozoa, Hexacorallia) attached to eunicid worm tubes from the Pacific Ocean. *ZooKeys* 562: 49–71. doi: 10.3897/zookeys.562.6181
- Obuchi M, Fujita Y, Nakano Y, Uehara T, Motokawa T (2010) Reproductive biology and early life history of the hermaphroditic feather star *Dorometra sesokonis* (Echinodermata: Crinoidea). *Marine Biology* 157(6): 1191–1201. doi: 10.1007/s00227-010-1399-3
- Reimer JD, Ono S, Takishita K, Tsukahara J, Maruyama T (2006) Molecular evidence suggesting species in the zoanthid genera *Palythoa* and *Protospalythoa* (Anthozoa: Hexacorallia) are congeneric. *Zoological Science* 23(1): 87–94. doi: 10.2108/zsj.23.87.
- Reimer JD, Takishita K, Ono S, Maruyama T (2007) Diversity and evolution in the zoanthid genus *Palythoa* (Cnidaria: Hexacorallia) utilizing nuclear ITS-rDNA. *Coral Reefs* 26(2): 399–410. doi: 10.1007/s00338-007-0210-5
- Reimer JD, Lin M, Fujii T, Lane DJW, Hoeksema BW (2012) The phylogenetic position of the solitary zoanthid genus *Sphenopus* (Cnidaria: Hexacorallia). *Contributions to Zoology* 81(1): 43–54.
- Reimer JD, Poliseno A, Hoeksema BW (2014) Shallow-water zoantharians (Cnidaria, Hexacorallia) from the Central Indo-Pacific. *ZooKeys* 444: 1–57. doi: 10.3897/zookeys.444.7537
- Reimer JD, Yang S, White KN, Asami R, Fujita K, Hongo C, Ito S, Kawamura I, Maeda B, Mizuyama M, Obuchi M, Sakamaki T, Tachihara K, Tamura M, Tanahara A, Yamaguchi A, Jenke-Kodama H (2015) Effects of causeway construction on environment and biota of subtropical tidal flats in Okinawa, Japan. *Marine Pollution Bulletin* 94(1–2): 153–167. doi:10.1016/j.marpolbul.2015.02.037
- Reimer JD, Higashiji T, Nonaka M (2016) First record of the genus *Sphenopus* (Anthozoa: Hexacorallia: Zoantharia) from Japan. *Fauna Ryukyana* 29: 37–40.
- Ronquist F, Huelsenbeck JP (2003) MrBayes version 3.0: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19(12): 1572–1574. doi: 10.1093/bioinformatics/btg180
- Ryland JS, Lancaster JE (2003) Revision of methods for separating species of *Protospalythoa* (Hexacorallia: Zoanthidea) in the tropical West Pacific. *Invertebrate Systematics* 17(3): 407–428. doi: 10.1071/IS02008
- Ryland JS, Ward H (2016) Carcinocidium-forming *Epizoanthus* (Hexacorallia: Zoantharia) and the biology of *E. papillosus* in the eastern Atlantic, with special reference to the cnidom. *Zootaxa* 4088(4): 489–514. doi: 10.11646/zootaxa.4088.4.2
- Sentoku A, Tokuda Y, Ezaki Y (2016) Burrowing hard corals occurring on the sea floor since 80 million years ago. *Scientific Reports* 6: 24355. doi: 10.1038/srep24355
- Schmidt H (1974) On evolution in the Anthozoa. *Proceedings of the 2nd International Coral Reef Symposium* 1: 533–560.
- Schuhmacher H (1977) Ability in fungiid corals to overcome sedimentation. *Proceedings of the 3rd International Coral Reef Symposium* 1: 503–509.

- Shearer TL, Van Oppen MJ, Romano SL, Wörheide G (2002) Slow mitochondrial DNA sequence evolution in the Anthozoa (Cnidaria). *Molecular Ecology* 11(12): 2475–2487. doi: 10.1046/j.1365-294X.2002.01652.x
- Sheppard CRC (1981) The reef and soft-substrate coral fauna of Chagos, Indian Ocean. *Journal of Natural History* 15(4): 607–621. doi: 10.1080/00222938100770431
- Sinniger F, Montoya-Burgos JI, Chevaldonné P, Pawlowski J (2005) Phylogeny of the order Zoantharia (Anthozoa, Hexacorallia) based on the mitochondrial ribosomal genes. *Marine Biology* 147(5): 1121–1128. doi: 10.1007/s00227-005-0016-3
- Sinniger F, Reimer JD, Pawlowski J (2008) Potential of DNA sequences to identify zoanthids (Cnidaria: Zoantharia). *Zoological Science* 25(12): 1253–1260. doi: 10.2108/zsj.25.1253
- Sinniger F, Reimer JD, Pawlowski J (2010) The Parazoanthidae (Hexacorallia: Zoantharia) DNA taxonomy: description of two new genera. *Marine Biodiversity* 40(1): 57–70. doi: 10.1007/s12526-009-0034-3
- Soong K, Shiau Y, Chen C (1999) Morphological and life history divergence of the zoanthid, *Sphenopus marsupialis* off the Taiwanese coast. *Zoological Studies* 38(3): 333–343.
- Spier D, Stampar SN, Prantoni AL (2012) New record of the endangered cerianthid *Cerianthomorpha brasiliensis* (Cnidaria: Hexacorallia) in Paranaguá Bay, southern Brazil. *Marine Biodiversity Records* 5: e119. doi: 10.1017/S1755267212001078
- Stampar SN, Maronna MM, Kitahara MV, Reimer JD, Morandini AC (2014) Fast-evolving mitochondrial DNA in Ceriantharia: A reflection of Hexacorallia paraphyly? *PLoS ONE* 9(1): e86612. doi:10.1371/journal.pone.0086612
- Steenstrup JJS (1856) *Sphenopus marsupialis* (Gmelin) – Saerskilt Aftryk. af det Kongelige danske Videnskabernes Selkabs Forhandling 1856 (Mit Autograph des Verfassers). Lunos, Kjøbenhavn.
- Swain TD (2009) Phylogeny-based species delimitations and the evolution of host associations in symbiotic zoanthids (Anthozoa, Zoanthidea) of the wider Caribbean region. *Zoological Journal of the Linnean Society* 156(2): 223–238. doi: 10.1111/j.1096-3642.2008.00513.x
- Swain TD (2010) Evolutionary transitions in symbioses: dramatic reductions in bathymetric and geographic ranges of Zoanthidea coincide with loss of symbioses with invertebrates. *Molecular Ecology* 19(12): 2587–2598. doi: 10.1111/j.1365-294X.2010.04672.x
- Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S (2011) MEGA5: Molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular Biology and Evolution* 28(10): 2731–2739. doi: 10.1093/molbev/msr121
- Uruma City Cultural Sea Museum (2007) The secrets of ancient Uruma City – the blessing of sea and interchange. Uruma City Cultural Sea Museum, Uruma. [In Japanese]