

# Description of three new species previously identified as *Stolephorus bengalensis* (Dutt & Babu Rao, 1959) or *Stolephorus insularis* Hardenberg, 1933 and a re-description of *S. bengalensis* (Chordata, Osteichthyes, Clupeiformes, Engraulidae)

Harutaka Hata<sup>1</sup>, Sébastien Lavoué<sup>2</sup>, Hiroyuki Motomura<sup>3</sup>

**1** National Museum of Natural History, Smithsonian Institution, 10<sup>th</sup> and Constitution Ave, NW Washington, DC 20560, USA **2** School of Biological Sciences, Universiti Sains Malaysia, 11800, Penang, Malaysia **3** The Kagoshima University Museum, 1-21-30 Korimoto, Kagoshima 890-0065, Japan

Corresponding author: Hata Harutaka ([k2795502@kadai.jp](mailto:k2795502@kadai.jp))

Academic editor: Tihomir Stefanov | Received 22 March 2022 | Accepted 22 August 2022 | Published 15 September 2022

<https://zoobank.org/1D6F2518-F6B7-4BEE-A41F-86F417DBD423>

**Citation:** Hata H, Lavoué S, Motomura H (2022) Description of three new species previously identified as *Stolephorus bengalensis* (Dutt & Babu Rao, 1959) or *Stolephorus insularis* Hardenberg, 1933 and a re-description of *S. bengalensis* (Chordata, Osteichthyes, Clupeiformes, Engraulidae). ZooKeys 1121: 145–173. <https://doi.org/10.3897/zookeys.1121.84171>

## Abstract

Examination of numerous specimens characterised by predorsal scute, long maxilla, indented preopercle and pelvic scute lacking a spine and previously identified as *Stolephorus bengalensis* (Dutt & Babu Rao, 1959) or *Stolephorus insularis* Hardenberg, 1933, revealed four distinct species, true *S. bengalensis* (distributed from the Bay of Bengal to Pakistan) and three new species, viz., *Stolephorus eldorado* **sp. nov.** (Taiwan to Java, Indonesia), *Stolephorus diabolus* **sp. nov.** (Strait of Malacca, from Penang, Malaysia, to Singapore) and *Stolephorus eclipsis* **sp. nov.** (Bintan Island, Riau Archipelago, Indonesia). Characters separating the four species include numbers of gill rakers on each gill arch and vertebrae and pelvic fin and dorsal-fin ray lengths. Two molecular markers (mitochondrial cytochrome *b* and cytochrome oxidase I genes) demonstrated the distinction of three of the species examined morphologically and enabled a reconstruction of their phylogenetic relationships. Each species was genetically divergent from the others by 3.5%–7.7% mean uncorrected distance in the mitochondrial cytochrome oxidase I gene.

## Keywords

Actinopterygii, Clupeomorpha, phylogenetics, *Stolephorus tri*, taxonomy


## Introduction

The anchovy genus *Stolephorus* Lacepède, 1803 (Teleostei: Clupeiformes: Engraulidae), diagnosed by the presence of prepelvic scutes and an embedded urohyal and lack of postpelvic scutes, currently includes 37 valid species that preferentially inhabit marine and/or estuarine waters in the Indo-Pacific region (Wongratana 1983, 1987a, b; Whitehead et al. 1988; Wongratana et al. 1999; Kimura et al. 2009; Hata and Motomura 2018a, b, c, d, e, 2021a, b, c, 2022; Hata et al. 2019, 2020a, b, 2021; Gangan et al. 2020). Amongst them, species with a predorsal scute, paired dark lines on the dorsum behind the dorsal fin, a long maxilla (posterior tip well beyond the preopercle posterior margin), the preopercle posterior margin concave and pelvic scute without a posteriorly projecting spine (Fig. 1) are regarded as *Stolephorus insularis* Hardenberg, 1933 by Whitehead et al. (1988), who reviewed the genus. Hata et al. (2019) revised the taxonomy of seven nominal species of *Stolephorus*, treating Whitehead et al.'s (1988) *S. insularis* as *Stolephorus bengalensis* (Dutt & Babu Rao, 1959) and regarding the nominal species *S. insularis* as a junior synonym of *Stolephorus tri* (Bleeker, 1852). However, subsequent re-examination of specimens, identified as *S. bengalensis*, in fact revealed the presence of four species.

The aim of this study is to re-describe *S. bengalensis* and describe three new species of *Stolephorus* from specimens previously regarded as *S. insularis* or *S. bengalensis*. In addition to the morphological comparisons, complete mitochondrial cytochrome *b* gene and partial mitochondrial cytochrome oxidase I (COI) gene sequences from 31 specimens were used to estimate the genetic distinction of three of the latter (the fourth species unavailable) plus one unidentified, but related species from Segara Anakan Lagoon, Central Java, Indonesia (Nuryanto et al. 2017).

## Materials and methods

Counts and proportional measurements followed Hata and Motomura (2017). Counts of fin rays and vertebrae followed Hubbs and Lagler (1947), the last two rays of dorsal and anal fins being counted separately, unless they originated from the same base, in which case they were counted as one ray. Vertebrae counts includes urostyle. All measurements were made with digital calipers to the nearest 0.01 mm. “Pelvic scute” refers to a scute joined to the pelvic girdle and “prepelvic scute”, “postpelvic scute” and “predorsal scute” to hard spine-like scutes anterior to the pelvic fin, posterior to the pelvic fin and just anterior to the dorsal-fin origin, respectively. Osteological characters, including vertebral counts, were determined from radiographs of 32, 2, 14 and 45 specimens of *S. bengalensis*, *S. diabolus* sp. nov., *S. eclipsis* sp. nov. and *S. eldorado* sp. nov., respectively. Abbreviations are as follows – SL: standard length; HL: head length; and UGR, LGR and TGR: rakers on upper limb, lower limb and total gill rakers, respectively, with associated numbers indicating the specific gill arch. Institutional codes generally follow Sabaj (2020). USMFC stands for Universiti Sains Malaysia Fish Collection, School of Biological Sciences, Penang, Malaysia.



**Figure 1.** Diagnostic characters of species previously identified as *Stolephorus bengalensis* **A** lateral view of whole body **B** dorsal-fin origin (triangle indicates predorsal scute, located just anterior to dorsal-fin origin) **C** dorsal view of dorsum behind dorsal fin (triangle indicates paired dark lines) **D** lateral surface of head (triangle indicates posterior tip of maxilla, posteriorly well beyond posterior margin of pre-opercle) **E** preopercle with concave posterior margin (supramaxilla removed) and **F** ventral view of pelvic fin (triangle indicates pelvic scute, lacking spine) (**A** KAUM-I. 94521, paratype of *S. eldorado* sp. nov. in fresh condition, 43.4 mm SL, Ha Long Bay, northern Vietnam **B**, **E**, **F** KAUM-I. 113148, paratype of *S. eldorado* sp. nov., 55.3 mm SL, Ke-tzu-liao, south-western Taiwan **C** ZUMT 62056, paratype of *S. diabolus* sp. nov., 38.4 mm SL, Singapore **D** KAUM-I. 94509, paratype of *S. eldorado* sp. nov., 41.4 mm SL, Ha Long Bay, northern Vietnam) (**B**, **D**, **E** and **F** alizarin stain).

The mitochondrial genotypes of 31 specimens comprising three (out of four) species of *Stolephorus* examined in this study, plus one closely related, but unidentified species, were compared using the complete (1140 base pairs [bp]) cytochrome *b* gene and partial (648 bp) COI gene. The cytochrome *b* gene sequences were published in Hata et al (2019; 2020b) and are available in GenBank (Table 1). The COI gene was newly sequenced for 19 specimens of *S. eldorado*, including the holotype and several paratypes (Table 1) and the resulting data combined with COI sequences (available in GenBank) of *S. diabolus* (two specimens from West Peninsular Malaysia, including the holotype), *S. bengalensis* (eight specimens from India), *S. eldorado* (one specimen from China; Pang et al. 2019) and a single specimen of an unidentified *Stolephorus* species (from Segara Anakan Lagoon, Central

**Table 1.** Taxonomic treatment and molecular marker sampling of 32 specimens of *Stolephorus* examined in the molecular section of the present study. “-” indicates corresponding sequence not determined. Bold accession numbers indicate sequences determined during the study. (Abbreviations: Cytb, cytochrome b; COI, cytochrome oxidase I; Gb, GenBank; “\*\*\*”, holotype; “\*”, paratype).

Species	Voucher	Origin	Cytb	COI
<i>S. eldorado</i>	KAUM-I. 94509*	Ha Long Bay, northern Vietnam	MH380318	<b>OM672421</b>
sp. nov.	KAUM-I. 94517**	Ha Long Bay, northern Vietnam	MH380319	<b>OM672422</b>
	KAUM-I. 94519*	Ha Long Bay, northern Vietnam	MH380320	<b>OM672423</b>
	KAUM-I. 94520*	Ha Long Bay, northern Vietnam	MH380321	<b>OM672424</b>
	KAUM-I. 94521*	Ha Long Bay, northern Vietnam	MH380322	<b>OM672425</b>
	KAUM-I. 113142*	off Dong-gang, Pingtung, Taiwan	MH380323	<b>OM672426</b>
	KAUM-I. 113143*	off Dong-gang, Pingtung, Taiwan	MH380324	<b>OM672427</b>
	KAUM-I. 113144*	off Dong-gang, Pingtung, Taiwan	MH380325	<b>OM672428</b>
	KAUM-I. 113145*	off Dong-gang, Pingtung, Taiwan	MH380326	<b>OM672417</b>
	KAUM-I. 113146*	off Dong-gang, Pingtung, Taiwan	MH380327	<b>OM672418</b>
	KAUM-I. 113147*	off Dong-gang, Pingtung, Taiwan	MH380328	<b>OM672419</b>
	KAUM-I. 113148*	off Dong-gang, Pingtung, Taiwan	MH380329	<b>OM672429</b>
	KAUM-I. 113149*	off Dong-gang, Pingtung, Taiwan	MH380330	<b>OM672420</b>
	KAUM-I. 113150*	off Dong-gang, Pingtung, Taiwan	MH380331	<b>OM672430</b>
	KAUM-I. 113151*	off Dong-gang, Pingtung, Taiwan	MH380332	<b>OM672431</b>
	NTUM12426 (Bgk15)	Bangkok, Thailand	MH380652	<b>OM672414</b>
	NTUM12426 (Bgk17)	Bangkok, Thailand	MH380653	<b>OM672415</b>
	- (Bgk39)	Bangkok, Thailand	MH380333	<b>OM672416</b>
	- (HK01)	Hong Kong	MH380334	<b>OM672413</b>
	20180725PZ25	Zhangzhou city, China (24.26N, 118.11°E) (Gb)	MH732976	<b>MH732976</b>
<i>S. diabolus</i>	IPMB-I 13.00001**	Telok Bahang, Penang Island, Malaysia	MT080882	MT080410
sp. nov.	- (larvae not preserved)	Klang Strait, West Peninsular Malaysia (Gb)	-	MH673948
<i>S. bengalensis</i>	-	off Kochi, Kerala, India (9.97°N, 76.28°E) (Gb)	-	KU871055
(Dutt &	-	off Kochi, Kerala, India (9.97°N, 76.28°E) (Gb)	-	KU871061
Babu Rao,	-	off Kochi, Kerala, India (9.97°N, 76.28°E) (Gb)	-	KU894592
1959)	-	off Kochi, Kerala, India (9.97°N, 76.28°E) (Gb)	-	KU894597
	-	off Kochi, Kerala, India (9.97°N, 76.28°E) (Gb)	-	KU894598
	-	off Kochi, Kerala, India (9.97°N, 76.28°E) (Gb)	-	KU894599
	-	off Kochi, Kerala, India (9.97°N, 76.28°E) (Gb)	-	KU894600
	-	off Kochi, Kerala, India (9.97°N, 76.28°E) (Gb)	-	KU894601
<i>Stolephorus</i>	- (larvae not preserved)	Segara Anakan lagoon, Central Java (Gb)	-	KY944580
sp.				
<b>Outgroup:</b>				
<i>S. andhraensis</i>	NTUM12328 (Bg14)	Bangkok, Thailand	MH380656	MH380744
Babu Rao,				
1966				

Java) (Table 1). One specimen of *Stolephorus acinaces* Hata, Lavoué & Motomura, 2020 was selected as the outgroup.

DNA was extracted from ethanol-preserved tissue samples. Polymerase chain reaction (PCR) amplification and sequencing of the COI gene followed standard protocols (Ward et al. 2005), with annealing at 55 °C. Amplification of the partial COI gene used the following primers: forward COI\_FishF1 (5'-TCA ACC AAC CAC AAA GAC ATT GGC AC-3') and reverse COI\_FishR2 (5'-ACT TCA GGG TGA

CCG AAG AAT CAG AA-3') (Ward et al. 2005). PCR products were purified and sequenced in both directions by Sanger Sequencing technology using the same PCR primers. Sequences generated in this study have been deposited in the GenBank database (accession numbers given in Table 1).

Alignments of the cytochrome *b* and COI sequences were determined separately by eye, requiring neither insertions nor deletions. The final alignment combining the two genes (for 31 specimens plus one outgroup) comprised 1788 nucleotide positions. Uncorrected pairwise genetic distances (i.e. p-distances) amongst species were calculated with MEGA X (Stecher et al. 2020). The relationships between specimens were inferred by the Maximum Likelihood (ML) method of phylogenetic reconstruction using the general time-reversible model of nucleotide substitution with rate heterogeneity following a discrete gamma distribution (GTR +  $\Gamma$ ), using the software RAxML-NG (Kozlov et al. 2019) as implemented in the graphical interface raxmlGUI 2.0 (Edler et al. 2020). The tree was rooted using a specimen of *S. acinaces* and the robustness of each node determined by bootstrap support (500 replicates).

## Results and discussions

### *Stolephorus bengalensis* (Dutt & Babu Rao, 1959)

[English name: Hardenberg's Anchovy]

Figs 2, 3; Tables 2–4

*Anchoviella baganensis bengalensis* Dutt & Babu Rao, 1959: 160 [original locality: Waltair and Kakinada, Andhra Pradesh, east coast of India; type locality: Kilakarai, Gulf of Mannar, India, based on the neotype designated by Hata et al. (2019)].

*Stolephorus baganensis macrops* (lapsus memoriae for *Stolephorus megalops*) (not of Delsman): Whitehead 1967 (in part): 18 (Bay of Bengal).

*Stolephorus insularis* (not of Delsman): Whitehead et al. 1988 (in part): 413 (northern part of Indian Ocean); Young et al. 1994: 222, fig. 7 (Wangkun and Fangliao, Taiwan); Wongratana et al. 1999 (in part): 1736 (northern part of Indian Ocean); Gangan et al. 2020: 562, fig. 5 (Kochi, Kerala State, India).

*Stolephorus bengalensis*: Hata et al. 2019 (in part): 24, fig. 12 (Pakistan and India; neotype designation).

**Neotype.** USNM 276476, 45.8 mm SL, Kilakarai, Gulf of Mannar, India, 20 Feb 1964, J. W. Reintjes and P. S. B. R. James.

**Non-type specimens.** 46 specimens, 30.8–58.7 mm SL. **INDIA:** BMNH 1969.5.30.34–45, 12 specimens, Chennai, Tamil Nadu State; OCF-P 10435, 4 specimens, 30.8–38.1 mm SL, estuary of Hooghly River, West Bengal State (purchased in fish market in Kolkata, West Bengal State); USNM 204227, 21 specimens, 42.7–51.8 mm SL, Sassan Docks, Mumbai, Maharashtra State. **PAKISTAN:** KAUM–I. 69286, 50.0 mm SL, KAUM–I. 69287, 58.7 mm SL, KAUM–I. 69288, 50.5 mm SL,



**Figure 2.** **A** lateral **B** dorsal and **C** ventral views of neotype of *Stolephorus bengalensis* (USNM 276476, 45.6 mm SL, Gulf of Mannar, India). Scale bars indicate 2 mm.

KAUM-I. 69289, 54.4 mm SL, KAUM-I. 69290, 49.0 mm SL, KAUM-I. 69291, 53.1 mm SL, KAUM-I. 69292, 47.3 mm SL, KAUM-I. 69294, 58.6 mm, KAUM-I. 69295, 58.6 mm, West Wharf, Karachi.

**Diagnosis.** A species of *Stolephorus* with the following combination of characters: 1UGR 16–19 (modally 18), 1LGR 23–27 (25), 1TGR 40–45 (44); 2UGR 11–15 (13), 2LGR 21–25 (23), 2TGR 33–39 (36); 3UGR 10–12 (11), 3LGR 13–15 (13), 3TGR 23–27 (24); 4UGR 7–9 (8), 4LGR 10–12 (11), 4TGR 17–21 (19); prepelvic scutes 5–7 (6); total vertebrae 40 or 41 (40); long maxilla, posterior tip just reaching or slightly short of posterior margin of opercle; predorsal scutes present; pelvic scute without spine; body scales deciduous; posterior border of pre-opercle concave, indented; paired dark patch on parietal area with little following pigmentation; distinct double pigment lines along dorsum posterior to dorsal fin; black spots below eye and on lower-jaw tip absent; anal-fin base long, 19.0–21.3% (20.2%) of SL; maximum orbit diameter 7.3–8.6% (8.1%) of SL; third dorsal-fin ray long, 18.5–19.9% (19.0%) of SL; pelvic fin rather long, 9.4–11.0% (10.2%) of SL, its posterior tip not reaching to vertical through dorsal-fin origin when depressed in specimens > 50 mm SL; distance between posterior ends of supramaxilla and maxilla 5.3–6.6% (5.8%) of SL.

**Description.** Data for neotype presented first, followed by data for non-type specimens in parentheses (if different). Counts and measurements, expressed as percentages of SL or HL, given in Tables 2 and 3. Body laterally compressed, elongate, deepest at

**Table 2.** Meristics of specimens of *Stolephorus bengalensis* and *Stolephorus eldorado* sp. nov.

	<i>Stolephorus bengalensis</i>			<i>Stolephorus eldorado</i> sp. nov.		
	Neotype of <i>Anchoviella</i> <i>baganensis</i> <i>bengalensis</i>	Non- types		Holotype	Paratypes	
	USNM 276476	<i>n</i> = 46		KAUM-I. 94517	<i>n</i> = 57	
Standard length (mm)	45.8	30.8–58.7	Modes ± SD	44.4	37.5–58.8	Modes ± SD
Dorsal-fin rays (unbranched)	3	3	3 ± 0	3	3	3 ± 0
Dorsal-fin rays (branched)	12	11–14	13 ± 0.7	13	11–14	13 ± 0.6
Anal-fin rays (unbranched)	3	3	3 ± 0	3	3*	3 ± 0.1
Anal-fin rays (branched)	18	16–20	18 ± 0.9	18	16–19	18 ± 0.6
Pectoral-fin rays (unbranched)	1	1	1 ± 0	1	1	1 ± 0
Pectoral-fin rays (branched)	11	10–12	11 ± 0.7	12	9–13	11 ± 0.8
Pelvic-fin rays (unbranched)	1	1	1 ± 0	1	1	1 ± 0
Pelvic-fin rays (branched)	6	6	6 ± 0	6	6	6 ± 0
Gill rakers on 1 <sup>st</sup> gill arch (upper)	16	17–19	18 ± 0.8	18	16–21	18 ± 1.1
Gill rakers on 1 <sup>st</sup> gill arch (lower)	24	23–27	25 ± 1.1	26	23–28	25 ± 1.0
Gill rakers on 1 <sup>st</sup> gill arch (total)	40	40–45	44 ± 1.5	44	40–47	42 ± 1.7
Gill rakers on 2 <sup>nd</sup> gill arch (upper)	13	11–15	13 ± 0.8	12	10–14	13 ± 0.8
Gill rakers on 2 <sup>nd</sup> gill arch (lower)	22	21–25	23 ± 0.9	24	20–24	23 ± 1.0
Gill rakers on 2 <sup>nd</sup> gill arch (total)	35	33–39	36 ± 1.5	36	30–38	36 ± 1.5
Gill rakers on 3 <sup>rd</sup> gill arch (upper)	10	10–12	11 ± 0.6	10	8–12	10 ± 0.8
Gill rakers on 3 <sup>rd</sup> gill arch (lower)	13	13–15	13 ± 0.6	13	12–14	13 ± 0.6
Gill rakers on 3 <sup>rd</sup> gill arch (total)	23	23–27	24 ± 1.1	23	20–26	23 ± 1.1
Gill rakers on 4 <sup>th</sup> gill arch (upper)	9	7–9	8 ± 0.6	8	7–10	8 ± 0.7
Gill rakers on 4 <sup>th</sup> gill arch (lower)	10	10–12	11 ± 0.5	11	9–12	11 ± 0.8
Gill rakers on 4 <sup>th</sup> gill arch (total)	19	17–21	19 ± 1.0	19	16–22	18 ± 1.3
Gill rakers on posterior face of 3 <sup>rd</sup> gill arch	6	4–7	5 ± 0.7	5	4–7	5 ± 0.7
Prepelvic scutes	7	5–7	6 ± 0.5	6	5–7	6 ± 0.6
Scale rows in longitudinal series	35	34–36	35 ± 0.7	34	34–36	35 ± 0.7
Transverse scales	8	8	8 ± 0	8	8–9	8 ± 0.3
Pseudobranchial filaments	broken	13–18	16 ± 1.3	17	14–18	16 ± 1.2
Total vertebrae	40	40–41	40 ± 0.4	39	38–40	39 ± 0.7

\*one specimen with 4 unbranched anal-fin rays.

dorsal-fin origin. Dorsal profile of head and body slightly convex from snout tip to dorsal-fin origin, gently lowering to uppermost point of caudal-fin base. Ventral profile of head and body slightly convex from lower jaw tip to pelvic-fin insertion, thereafter, slowly rising to lowermost point of caudal-fin base. Single spine-like scute just anterior to dorsal-fin origin. Abdomen somewhat rounded, covered with seven (four to seven) spine-like prepelvic scutes. Pelvic scute without spine. Postpelvic scutes absent. Anus just anterior to anal-fin origin. Snout tip rounded; snout length less than eye diameter. Mouth large, inferior, ventral to body axis, extending backwards beyond posterior margin of eye. Maxilla long, its posterior tip pointed, just reaching (or slightly short of) opercle posterior margin. Lower jaw slender. Single row of conical teeth on both jaws

**Table 3.** Morphometrics of specimens of *Stolephorus bengalensis* and *Stolephorus eldorado* sp. nov.

	<i>Stolephorus bengalensis</i>			<i>Stolephorus eldorado</i> sp. nov.		
	Neotype of <i>Anchoviella</i> <i>baganensis</i> <i>bengalensis</i>	Non- types		Holotype	Paratypes	
	USNM 276476	n = 46	Means ± SD	KAUM-I. 94517	n = 57	
Standard length (mm)	45.8	30.8–58.7	Means ± SD	44.4	37.5–58.8	Means ± SD
As % SL						
Head length	25.8	23.0–26.1	24.7 ± 0.8	26.1	22.8–27.8	25.7 ± 1.3
Body depth	20.7	19.8–22.9	21.5 ± 0.7	17.9	17.3–22.0	20.3 ± 1.3
Pre-dorsal-fin length	56.8	52.3–57.1	54.5 ± 1.3	53.7	51.6–56.5	54.0 ± 1.2
Snout tip to pectoral-fin insertion	29.0	25.1–27.9	26.4 ± 0.8	28.0	22.5–29.2	26.9 ± 1.6
Snout tip to pelvic-fin insertion	47.6	42.4–49.4	45.1 ± 1.3	45.4	43.9–48.6	46.1 ± 1.2
Snout tip to anal-fin origin	66.3	61.3–66.5	64.2 ± 1.2	61.4	61.3–66.5	63.6 ± 1.2
Dorsal-fin base length	13.4	13.3–15.6	14.5 ± 0.6	14.3	13.2–15.7	14.6 ± 0.5
Anal-fin base length	20.8	19.0–21.3	20.2 ± 0.6	20.4	19.0–22.3	20.4 ± 0.8
Caudal-peduncle length	17.4	16.0–20.0	18.0 ± 1.1	18.1	16.4–19.8	18.2 ± 1.0
Caudal-peduncle depth	10.0	9.3–11.2	10.3 ± 0.4	9.3	8.7–10.9	9.7 ± 0.6
D–P1	37.2	33.9–38.1	35.9 ± 1.1	34.7	34.2–39.6	36.3 ± 1.2
D–P2	23.0	21.3–25.9	23.8 ± 1.0	20.4	19.1–26.1	23.2 ± 1.7
D–A	22.0	21.3–24.2	22.7 ± 0.8	20.2	19.2–23.2	21.6 ± 1.1
P1–P2	22.1	17.3–22.2	19.3 ± 1.4	19.6	16.9–23.8	20.3 ± 1.8
P2–A	19.7	15.9–20.3	18.6 ± 1.1	17.3	16.1–20.3	18.2 ± 1.0
Pectoral-fin length	broken	15.9–16.9	16.4 ± 0.4	18.0	14.9–18.5	16.5 ± 0.8
Pelvic-fin length	broken	9.4–11.0	10.2 ± 0.4	10.6	9.1–11.0	10.0 ± 0.5
Maxilla length	broken	19.7–22.3	21.0 ± 0.7	21.5	19.6–22.9	21.4 ± 0.9
Lower-jaw length	17.1	15.3–17.6	16.3 ± 0.5	17.3	14.6–17.9	16.7 ± 0.7
Supramaxilla end to maxilla end	broken	5.3–6.6	5.8 ± 0.3	5.1	5.0–6.3	5.6 ± 0.4
Maximum orbit diameter	8.5	7.3–8.6	8.1 ± 0.3	9.3	8.2–9.9	8.9 ± 0.4
Eye diameter	6.9	6.1–7.6	6.9 ± 0.3	8.0	6.4–8.6	7.5 ± 0.5
Snout length	4.4	3.4–4.0	3.7 ± 0.2	3.7	3.1–4.3	3.7 ± 0.3
Interorbital width	6.74	5.2–6.3	5.9 ± 0.3	5.8	4.9–6.2	5.8 ± 0.3
Postorbital length	12.8	11.8–14.1	13.0 ± 0.5	12.3	11.6–14.9	12.9 ± 0.7
1 <sup>st</sup> dorsal-fin ray length	0.9	0.9–2.1	1.5 ± 0.3	1.9	0.9–2.2	1.5 ± 0.3
2 <sup>nd</sup> dorsal-fin ray length	7.3	6.6–8.9	7.7 ± 0.6	broken	5.9–8.1	7.3 ± 0.6
3 <sup>rd</sup> dorsal-fin ray length	18.8	18.5–19.9	19.0 ± 0.4	broken	15.9–18.6	17.4 ± 0.8
1 <sup>st</sup> anal-fin ray length	1.9	1.0–2.0	1.6 ± 0.3	1.5	0.8–2.2	1.6 ± 0.3
2 <sup>nd</sup> anal-fin ray length	5.2	4.6–6.3	5.4 ± 0.5	6.5	4.1–6.5	5.2 ± 0.8
3 <sup>rd</sup> anal-fin ray length	13.0	14.0–16.5	15.0 ± 0.8	14.4	13.3–15.5	14.1 ± 1.8

Abbreviations: D–P1 (distance from dorsal-fin origin to pectoral-fin insertion); D–P2 (distance from dorsal-fin origin to pelvic-fin insertion); D–A (distance between origins of dorsal and anal fins); P1–P2 (distance between insertions of pectoral and pelvic fins); P2–A (distance between pelvic-fin insertion and anal-fin origin).

and palatine. Patch of fine conical teeth on pterygoid. Several distinct conical teeth on vomer. Several rows of conical teeth on upper edges of basihyal and basibranchial. Eye large, round, covered with adipose eyelid, positioned laterally on head dorsal to horizontal through pectoral-fin insertion, visible in dorsal view. Pupil round. Orbit elliptical. Nostrils close to each other, anterior to orbit. Posterior margin of pre-opercle

**Table 4.** Frequency distribution of total vertebral numbers of *Stolephorus bengalensis*, *Stolephorus diabolus* sp. nov., *Stolephorus eclipis* sp. nov. and *Stolephorus eldorado* sp. nov.

		Total vertebrae			
		38	39	40	41
<i>Stolephorus bengalensis</i>	<i>n</i> = 32			27	5
<i>Stolephorus diabolus</i> sp. nov.	<i>n</i> = 2		2		
<i>Stolephorus eclipis</i> sp. nov.	<i>n</i> = 14	6	8		
<i>Stolephorus eldorado</i> sp. nov.	<i>n</i> = 45	11	24	10	

concave, indented. Subopercle and opercle with smoothly rounded posterior margins. Gill membrane without serrations. Interorbital space flat, width less than eye diameter. Pseudobranchial filaments present, length of longest filament less than eye diameter. Gill rakers long, slender, rough, visible from side of head when mouth opened. Single row of asperities on anterior surface of gill rakers. Isthmus muscle long, reaching anteriorly to posterior margin of gill membranes. Urohyal hidden by isthmus muscle, not visible without dissection. Gill membrane on each side joined distally, most of isthmus muscle exposed, not covered by gill membrane. Scales on lateral surface of body thin, cycloid, deciduous, those on lateral body surface with several centrally continuous vertical grooves and several longitudinal striae anteriorly (Fig. 3). Head scales absent. Fins scaleless, except for broad triangular sheath of scales on caudal fin. Dorsal-fin origin posterior to vertical through base of last pelvic-fin ray, slightly posterior to middle of body. Dorsal and anal fins with three anteriormost rays unbranched. First dorsal and anal-fin rays minute. Anteriormost three rays of both dorsal and anal fins closely spaced. Anal-fin origin just below base of tenth (ninth to eleventh) dorsal-fin ray. Posterior tip of depressed anal fin not reaching caudal-fin base. Uppermost pectoral-fin ray unbranched, inserted below body axis. Posterior tip of pectoral fin not reaching to pelvic fin insertion. Dorsal, ventral and posterior margins of pectoral fin nearly linear. Pelvic fin shorter than pectoral fin, insertion anterior to vertical through dorsal-fin origin. Posterior tip of depressed pelvic fin not reaching to vertical through dorsal-fin origin. Caudal fin forked, posterior tips pointed.

**Colour of preserved specimens.** Body uniformly pale ivory. A pair of distinct dark patches on parietal region, with little pigmentation on occipital area. Double pigmented lines dorsally posterior to dorsal fin. A few melanophores scattered anteriorly on snout. No black spots below eye and on lower-jaw tip. Melanophores scattered along bases of dorsal and anal fins. All fins transparent, melanophores scattered along fin rays of caudal fin and anterior parts of dorsal and anal fins.


**Distribution.** *Stolephorus bengalensis* is distributed in the northern Indian Ocean from Pakistan to the Bay of Bengal (Fig. 4). It is abundantly landed and marketed along the coast of the Bay of Bengal.

**Morphological comparisons.** *Stolephorus bengalensis* has been considered conspecific with the three new species described herein, the four species being easily separable from all other congeners, except for *Stolephorus acinaces*, *Stolephorus andhraensis* Babu Rao, 1966, *Stolephorus carpentariae* (De Vis, 1882), *Stolephorus hindustanensis* Hata & Motomura, 2022, *Stolephorus holodon* (Boulenger, 1900),



**Figure 3.** Stained scale removed from right side of mid-body (above anal fin) of *Stolephorus bengalensis*, BMNH 1969.5.30.34–45, 47.8 mm SL, Chennai, India (left-right inverted; scale bar indicates 0.5 mm).

*Stolephorus ronquilloi* Wongratana, 1983 and *Stolephorus tamilensis* Gangan, Pavan-Kumar, Jahageerdar & Jaiswar, 2020, the former having a concavely indented pre-opercular margin and lacking a spine on the pelvic scute (Whitehead et al. 1988; Wongratana et al. 1999; Kimura et al. 2009; Hata and Motomura 2018a, b, c, d, e, 2021a, b, c, 2022; Hata et al. 2019, 2020a, b, 2021; Gangan et al. 2020). However, the former four species are distinguished from the other seven by having a predorsal scute (vs. absent in the latter) and double dark lines on the dorsum posterior to the dorsal fin (vs. no lines on the dorsum, except in *S. hindustanensis* and *S. ronquilloi*). Moreover, *S. carpentariae* also differs from *S. bengalensis* and the three new species in having 19 or 20 branched anal-fin rays [16–18 (rarely 19 or 20) in the remaining five species] and the anal-fin origin located below the origin of the second to sixth dorsal-



**Figure 4.** Map of the northern part of the Indo-West Pacific region showing distributional records of *Stolephorus bengalensis* (red circles), *S. diabolus* (yellow squares) sp. nov., *S. eclipsis* (green diamond) sp. nov. and *S. eldorado* (blue triangles) sp. nov., based on specimens examined in this study. Land masses outlined with black lines.

fin ray (vs. eighth to eleventh) (Whitehead et al. 1988; Wongratana et al. 1999; Gangan et al. 2020; Hata et al. 2020b). *Stolephorus bengalensis*, *S. diabolus* sp. nov., *S. eclipsis* sp. nov. and *S. eldorado* sp. nov. resemble *Stolephorus baganensis* Delsman, 1931, *Stolephorus dubiosus* Wongratana, 1983 and *Stolephorus tri* (Bleeker, 1852) in having a predorsal scute and double pigment lines on the dorsum behind the dorsal fin, but differ in having deciduous body scales (vs. body scales not deciduous) and lacking a spine on the pelvic scute (pelvic scute with a hard posteriorly projecting spine) (Whitehead et al. 1988; Wongratana et al. 1999; Hata et al. 2019). Comparisons of *S. bengalensis* with *S. diabolus* sp. nov., *S. eclipsis* sp. nov. and *S. eldorado* sp. nov. are given in “Comparisons” under each new species.

**Molecular comparisons.** *Stolephorus bengalensis*, *S. diabolus* sp. nov. and *S. eldorado* sp. nov. were divergent from each other by at least 3.5% COI-based mean uncorrected genetic distance (min-max = 3.5–7.7%) (Fig. 5). In contrast, each species was genetically uniform, with intraspecific differentiation not exceeding 1% (note: *Stolephorus* sp. represented by a single specimen – see below), forming clear intraspecific versus interspecific genetic gaps. The ML phylogenetic tree using COI and cytochrome *b* markers (Fig. 5) was fully resolved, with interspecific relationships supported by bootstrap values above 80%. Each species formed a well-supported monophyletic group, in agreement with their genetic distinction, thereby confirming their taxonomic status, which was further supported by the morphological observations. The COI se-

quence of an unidentified larva collected from the Segara Anakan Lagoon, Central Java (Nuryanto et al. 2017), indicated either a range extension of *S. eclipsis* sp. nov. or the presence of an unidentified species of *Stolephorus*, related to *S. bengalensis*.

***Stolephorus diabolus* sp. nov.**

<https://zoobank.org/09515D76-5020-4C13-9391-7F325335C74C>

[New English name: Demon Anchovy]

Figs 1C, 6; Tables 4–6

*Stolephorus bengalensis* (not of Dutt and Babu Rao): Hata et al. 2022: (in part) 34 (Singapore).

**Holotype.** IPMB-I 13.00001, 49.7 mm SL, Teluk Bahang, Penang, Malaysia.

**Paratypes.** 14 specimens, 28.5–43.7 mm SL. USMFC 82-0017, 43.7 mm SL, collected with the holotype; USMFC 82-0057, 4 specimens, 40.1–41.1 mm SL, estuary of Merbok River, Jeti Semeling, Malaysia; ZUMT 62056, 5 specimens, 28.5–38.4 mm SL, KAUM-I. 163702, 36.3 mm SL, KAUM-I. 163703, 36.4 mm SL, NSMT-P 143554, 36.4 mm SL, NSMT-P 143555, 36.6 mm SL, Singapore.

**Diagnosis.** A species of *Stolephorus* with the following combination of characters: 1UGR 14–16 (modally 16), 1LGR 20–23 (22), 1TGR 35–38 (38); 2UGR 10 or 11 (11), 2LGR 19 or 20 (20), 2TGR 30 or 31 (31); 3UGR 8 or 9 (9), 3LGR 11 or 12 (12), 3TGR 20 or 21 (21); 4UGR 6 or 7 (7), 4LGR 9 or 10 (9), 4TGR 15–17 (17); prepelvic scutes 5–7 (6); total vertebrae 39; long maxilla, posterior tip just reaching or slightly short of posterior margin of opercle; predorsal scute present; pelvic scute without spine; body scales deciduous; posterior border of pre-opercle concave, indented; paired dark patch on parietal area with little following pigmentation; distinct double pigment lines along dorsum posterior to dorsal fin; black spots below eye and on lower-jaw tip absent; anal-fin base long, 19.8–22.3% (mean 20.7%) of SL; maximum orbit diameter 8.1–8.7% (8.3%) of SL; third dorsal-fin ray short, 17.0–18.5% (18.0%) of SL; pelvic fin rather long, 9.6–11.3% (10.0%) of SL, its posterior tip not reaching to vertical through dorsal-fin origin when depressed in specimens > 40 mm SL; distance between posterior ends of supramaxilla and maxilla 5.7–6.4% (6.1%) of SL.

**Description.** Data for holotype presented first, followed by data for paratypes in parentheses (if different). Counts and measurements, expressed as percentages of SL or HL, given in Tables 5 and 6. Body laterally compressed, elongate, deepest at dorsal-fin origin. Dorsal profile of head and body slightly convex from snout tip to dorsal-fin origin, gently lowering to uppermost point of caudal-fin base. Ventral profile of head and body slightly convex from lower jaw tip to pelvic-fin insertion, thereafter, slowly rising to lowermost point of caudal-fin base. Single spine-like scute just anterior to dorsal-fin origin. Abdomen somewhat rounded. Scutes on ventrum broken in holotype (five to seven spine-like prepelvic scutes on ventrum in paratypes). Pelvic scute without spine. Postpelvic scutes absent. Anus just anterior to anal-fin origin. Snout

**Table 5.** Meristics of specimens of *Stolephorus diabolus* sp. nov. and *Stolephorus eclipsis* sp. nov.

	<i>Stolephorus diabolus</i> sp. nov.			<i>Stolephorus eclipsis</i> sp. nov.		
	Holotype	Paratypes		Holotype	Paratypes	
	IPMB-I 13.00001	n = 14		MZB 26452	n = 28	
Standard length (mm)	49.7	28.5–43.7	Modes ± SD	40.3	32.0–43.7	Modes ± SD
Dorsal-fin rays (unbranched)	3	3	3 ± 0	3	3	3 ± 0
Dorsal-fin rays (branched)	12	12–13	13 ± 0.5	12	11–13	12 ± 0.4
Anal-fin rays (unbranched)	3	3	3 ± 0	3	3	3 ± 0
Anal-fin rays (branched)	16	16–18	17 ± 0.7	17	16–18	17 ± 0.7
Pectoral-fin rays (unbranched)	1	1	1 ± 0	1	1	1 ± 0
Pectoral-fin rays (branched)	11	10–13	11 ± 0.8	10	10–12	11 ± 0.6
Pelvic-fin rays (unbranched)	1	1	1 ± 0	1	1	1 ± 0
Pelvic-fin rays (branched)	6	6	6 ± 0	6	6	6 ± 0
Gill rakers on 1 <sup>st</sup> gill arch (upper)	16	14–16	16 ± 0.6	20	19–21	20 ± 0.7
Gill rakers on 1 <sup>st</sup> gill arch (lower)	22	20–23	22 ± 0.7	28	26–30	28 ± 0.8
Gill rakers on 1 <sup>st</sup> gill arch (total)	38	35–38	38 ± 1.0	48	47–51	47 ± 1.1
Gill rakers on 2 <sup>nd</sup> gill arch (upper)	10	11	11 ± 0.2	13	13–16	14 ± 0.7
Gill rakers on 2 <sup>nd</sup> gill arch (lower)	20	19–20	20 ± 0.4	25	24–27	25 ± 0.8
Gill rakers on 2 <sup>nd</sup> gill arch (total)	30	30–31	31 ± 0.4	38	37–42	39 ± 1.4
Gill rakers on 3 <sup>rd</sup> gill arch (upper)	9	8–9	9 ± 0.2	11	10–13	12 ± 0.7
Gill rakers on 3 <sup>rd</sup> gill arch (lower)	12	11–12	12 ± 0.5	14	14–16	15 ± 0.6
Gill rakers on 3 <sup>rd</sup> gill arch (total)	21	20–21	21 ± 0.5	25	25–28	27 ± 1.1
Gill rakers on 4 <sup>th</sup> gill arch (upper)	7	6–7	7 ± 0.4	8	8–11	9 ± 0.9
Gill rakers on 4 <sup>th</sup> gill arch (lower)	9	9–10	9 ± 0.5	11	11–13	12 ± 0.5
Gill rakers on 4 <sup>th</sup> gill arch (total)	16	15–17	17 ± 0.8	19	19–24	21 ± 1.2
Gill rakers on posterior face of 3 <sup>rd</sup> gill arch	3	3–5	4 ± 0.5	4	4–7	5 ± 0.7
Prepelvic scutes	broken	5–7	6 ± 0.5	6	5–7	6 ± 0.5
Scale rows in longitudinal series	34	34–35	35 ± 0.5	35	35–36	35 ± 0.4
Transverse scales	8	8–9	8 ± 0.2	8	8	8 ± 0
Pseudobranchial filaments	broken	14–16	15 ± 0.7	14	14–18	15 ± 1.2
Total vertebrae	39	39	39 ± 0	38	38–39	39 ± 0.5


tip rounded; snout length less than eye diameter. Mouth large, inferior, ventral to body axis, extending backwards beyond posterior margin of eye. Maxilla long, its posterior tip broken in holotype (posterior pointed, just reaching or slightly short of opercle posterior margin in paratypes). Lower jaw slender. Single row of conical teeth on both jaws and palatine. Patch of fine conical teeth on pterygoid. Several distinct conical teeth on vomer. Several rows of conical teeth on upper edges of basihyal and basibranchial. Eye large, round, covered with adipose eyelid, positioned laterally on head dorsal to horizontal through pectoral-fin insertion, visible in dorsal view. Pupil round. Orbit elliptical. Nostrils close to each other, anterior to orbit. Posterior margin of pre-opercle concave, indented. Subopercle and opercle with smoothly rounded posterior margins. Gill membrane without serrations. Interorbital space flat, width less than eye diameter. Pseudobranchial filaments present, length of longest filament less than eye diameter. Gill rakers long, slender, rough, visible from side of head when mouth opened. Single row of asperities on anterior surface of gill rakers. Isthmus

**Table 6.** Morphometrics of specimens of *Stolephorus diabolus* sp. nov. and *Stolephorus eclipsis* sp. nov.

	<i>Stolephorus diabolus</i> sp. nov.			<i>Stolephorus eclipsis</i> sp. nov.		
	Holotype	Paratypes		Holotype	Paratypes	
	IPMB-I 13.00001	n = 14		MZB 26452	n = 28	
Standard length (mm)	49.7	28.5–43.7	Means ± SD	40.3	32.0–43.7	Means ± SD
As % SL						
Head length	24.8	25.0–25.9	25.4 ± 0.3	25.4	23.6–26.7	24.8 ± 0.8
Body depth	21.7	19.8–21.9	20.9 ± 0.7	20.6	18.4–20.8	19.6 ± 0.6
Pre-dorsal-fin length	51.8	51.3–52.9	52.1 ± 0.5	52.8	51.3–54.9	53.4 ± 1.0
Snout tip to pectoral-fin insertion	25.7	26.2–28.4	27.2 ± 0.6	26.8	24.8–28.5	26.5 ± 0.9
Snout tip to pelvic-fin insertion	48.2	45.8–49.0	47.2 ± 0.8	47.0	44.8–47.3	46.2 ± 0.7
Snout tip to anal-fin origin	65.4	63.0–66.0	64.4 ± 0.9	63.1	62.8–65.8	64.1 ± 0.9
Dorsal-fin base length	15.0	13.9–16.6	15.0 ± 0.7	13.6	13.1–14.5	13.8 ± 0.4
Anal-fin base length	19.8	19.9–22.3	20.7 ± 0.8	19.3	17.6–19.3	18.6 ± 0.5
Caudal-peduncle length	18.8	16.4–19.4	17.8 ± 0.9	17.4	14.7–18.5	17.1 ± 0.9
Caudal-peduncle depth	9.8	9.4–10.3	9.8 ± 0.2	9.9	9.2–10.6	9.8 ± 0.4
D–P1	36.5	33.8–36.4	35.3 ± 0.8	38.0	34.0–38.8	36.5 ± 1.3
D–P2	23.5	21.9–24.5	23.4 ± 0.6	24.1	21.0–23.9	22.5 ± 0.8
D–A	22.6	20.7–23.1	22.2 ± 0.8	21.3	20.0–21.9	20.9 ± 0.5
P1–P2	24.1	19.4–20.0	21.0 ± 1.1	21.3	19.3–22.8	21.0 ± 0.9
P2–A	18.0	16.3–19.5	18.0 ± 0.9	18.5	17.5–20.6	19.0 ± 0.8
Pectoral-fin length	16.2	15.4–17.1	16.2 ± 0.5	16.4	15.5–17.7	16.7 ± 0.6
Pelvic-fin length	9.8	9.6–11.3	10.0 ± 0.4	8.7	8.8–9.9	9.4 ± 0.3
Maxilla length	broken	20.9–21.9	21.4 ± 0.3	20.7	19.9–22.5	21.1 ± 0.7
Lower-jaw length	16.2	16.2–17.5	16.6 ± 0.3	16.4	15.8–17.7	16.6 ± 0.5
Supramaxilla end to maxilla end	broken	5.7–6.4	6.1 ± 0.2	5.2	4.7–5.4	5.1 ± 0.2
Maximum orbit diameter	8.2	8.1–8.7	8.3 ± 0.2	8.8	7.9–9.6	8.7 ± 0.4
Eye diameter	6.5	6.1–7.7	6.9 ± 0.5	6.8	6.7–8.3	7.3 ± 0.4
Snout length	3.7	3.6–4.2	3.8 ± 0.2	4.0	3.4–4.2	3.8 ± 0.2
Interorbital width	5.6	5.5–5.9	5.7 ± 0.2	5.7	5.5–6.2	5.8 ± 0.2
Postorbital length	12.8	12.9–14.2	13.4 ± 0.4	12.1	11.5–12.9	12.1 ± 0.4
1 <sup>st</sup> dorsal-fin ray length	1.5	0.8–2.2	1.6 ± 0.4	1.5	1.0–2.2	1.5 ± 0.3
2 <sup>nd</sup> dorsal-fin ray length	broken	7.0–9.8	8.1 ± 0.7	broken	5.1–5.7	7.6 ± 0.6
3 <sup>rd</sup> dorsal-fin ray length	broken	17.0–18.5	18.0 ± 0.5	17.1	16.5–18.8	17.6 ± 0.7
1 <sup>st</sup> anal-fin ray length	1.6	0.9–2.2	1.5 ± 0.4	1.9	1.2–2.2	1.7 ± 0.3
2 <sup>nd</sup> anal-fin ray length	5.3	4.8–7.2	5.5 ± 0.6	5.2	5.1–5.7	5.3 ± 0.2
3 <sup>rd</sup> anal-fin ray length	broken	14.6–16.0	15.4 ± 0.4	14.4	13.4–15.0	14.2 ± 0.5

Abbreviations: D–P1 (distance from dorsal-fin origin to pectoral-fin insertion); D–P2 (distance from dorsal-fin origin to pelvic-fin insertion); D–A (distance between origins of dorsal and anal fins); P1–P2 (distance between insertions of pectoral and pelvic fins); P2–A (distance between pelvic-fin insertion and anal-fin origin).

muscle long, reaching anteriorly to posterior margin of gill membranes. Urohyal hidden by isthmus muscle, not visible without dissection. Gill membrane on each side joined distally, most of isthmus muscle exposed, not covered by gill membrane. Body scales deciduous, completely lacking on specimens, except for prepelvic scutes. Head



**Figure 5.** Maximum-likelihood phylogenetic tree of four species of *Stolephorus* related to *S. bengalensis*, based on the cytochrome *b* and cytochrome oxidase I genes (total: 1,788 base pairs) of 31 specimens, each species forming a monophyletic group. Each specimen identified by Museum Registration Number or GenBank number (see text and Table 1 for details). Type status (either holotype or paratype), specimen code, gene used (mt or COI) and sequence origin (GB) are indicated in brackets where necessary. Tree rooted by a specimen of *Stolephorus andhraensis*. Branch lengths proportional to number of substitutions. Bootstrap proportions indicated at nodes.

scales absent. Fins scaleless, except for broad triangular sheath of scales on caudal fin. Dorsal-fin origin posterior to vertical through base of last pelvic-fin ray, slightly posterior to middle of body. Dorsal and anal fins with three anteriormost rays unbranched. First dorsal- and anal-fin rays minute. Anteriormost three rays of both dorsal and anal fins closely spaced. Anal-fin origin just below base of eighth (eighth to eleventh) dorsal-fin ray. Posterior tip of depressed anal fin not reaching caudal-fin base. Uppermost pectoral-fin ray unbranched, inserted below body axis. Posterior tip of pectoral fin not reaching to pelvic fin insertion. Dorsal, ventral and posterior margins of pectoral fin nearly linear. Pelvic fin shorter than pectoral fin, insertion anterior to vertical through



**Figure 6.** **A** lateral **B** dorsal and **C** ventral views of preserved holotype of *S. diabolus* sp. nov., IPMB-I 13.00001, 49.7 mm SL, Teluk Bahang, Penang, Malaysia **D** lateral view of preserved paratype of *S. diabolus* sp. nov., ZUMT 62056, 37.3 mm SL, Singapore. Scale bars: 2 mm.


dorsal-fin origin. Posterior tip of depressed pelvic fin not reaching to vertical through dorsal-fin origin (reaching to vertical through first to sixth dorsal-fin ray origin in some paratypes smaller than 40 mm SL). Caudal fin forked, posterior tips pointed.

**Colour of preserved specimens.** Body uniformly pale white. A pair of distinct dark patches on parietal region, with little pigmentation on occipital area. No black spots below eye and on lower-jaw tip. Melanophores scattered on posterior margins of scale pockets on dorsum. Double pigmented lines dorsally posterior to dorsal fin. Melanophores scattered along bases of dorsal and anal fins. All fins transparent, melanophores scattered along fin rays of caudal fin and anterior parts of dorsal and anal fins.

**Distribution.** *Stolephorus diabolus* sp. nov. is currently known only from the western coast of the Peninsular Malaysia (Merbok River Estuary and Penang) and Singapore (Fig. 4).

**Etymology.** The specific name “*diabolus*” is derived from Latin meaning “demon”, in reference to the hard spine on the dorsum of the species.

**Comparisons.** The new species is distinguished from *S. bengalensis*, *S. eclipsis* and *S. eldorado* by lower gill raker counts: 1TGR, 35–38 in *S. diabolus* (vs. 40 or more in the other three species); 2TGR, 30 or 31 in *S. diabolus* [vs. 33 or more (rarely 30 or 31 in *S. eldorado*)]; 3TGR, 20 or 21 in *S. diabolus* [vs. 22 or more in the other three



**Figure 7.** Relationships of total gill raker numbers (TGR) on **A** first gill arch (1GA) **B** second gill arch (2GA) **C** third gill arch (3GA) and **D** fourth gill arch (4GA) to standard length in *Stolephorus bengalensis* (red circles), *S. diabolus* sp. nov. (yellow squares), *S. eclipsis* sp. nov. (green diamonds) and *S. eldorado* sp. nov. (blue triangles).

species (rarely 21 in *S. eldorado*]; and 4TGR, 15–17 in *S. diabolus* (vs. 17 or more) (Fig. 7). Moreover, *S. diabolus* has a shorter orbit diameter than *S. eldorado* [maximum orbit diameter 8.1–8.7% (mean 8.3%) of SL in *S. diabolus* vs. 8.2–9.9% (8.9%) in *S. eldorado*; Fig. 8A]. Furthermore, *S. diabolus* is distinguished from *S. bengalensis* by having a shorter third dorsal-fin ray [17.0–18.5% (mean 18.0%) of SL in *S. diabolus* vs. 18.5–19.9% (19.0%) in *S. bengalensis* (Fig. 8B)] and lower total vertebral numbers [39 vs. 40 or 41 (modally 40) (Table 4)]. Detailed comparisons of *S. diabolus* with *S. eclipsis* and *S. eldorado* are given in “Comparisons” under each species.

### *Stolephorus eclipsis* sp. nov.


<https://zoobank.org/1556E6AA-0531-4361-874E-3FE6DB1FEA10>

[New English name: Eclipse Anchovy]

Fig. 9; Tables 4–6

**Holotype.** MZB 26452, 40.3 mm SL, Bintan Island, Riau Archipelago, Indonesia.

**Paratypes.** 28 specimens, 32.0–43.7 mm SL. LBRC-F 5039, 35.4 mm SL, LBRC-F 5040, 35.3 mm SL, LBRC-F 5041, 36.1 mm SL, Tanjungpinang, Bintan Island, Riau Archipelago, Indonesia; MZB 26440, 32.0 mm SL, MZB 26441, 36.1 mm SL, MZB 26442, 35.1 mm SL, MZB 26443, 34.7 mm SL, MZB 26444, 34.5 mm SL,



**Figure 8.** Relationships of **A** orbit diameter (as % of standard length; SL) **B** third dorsal-fin ray length (as % of SL) **C** anal-fin base length (as % of SL) **D** pelvic-fin length (as % of SL) and **E** distance between posterior ends of supramaxilla and maxilla (SMX–MX) in *Stolephorus bengalensis* (red circles), *S. diabolus* sp. nov. (yellow squares), *S. eclipsis* sp. nov. (green diamonds) and *S. eldorado* sp. nov. (blue triangles).

MZB 26445, 36.3 mm SL, MZB 26446, 36.2 mm SL, MZB 26447, 38.8 mm SL, MZB 26448, 39.2 mm SL, MZB 26449, 37.7 mm SL, MZB 26450, 40.0 mm SL, MZB 26451, 36.8 mm SL, 26453, 36.4 mm SL, MZB 26454, 39.0 mm SL, MZB 26455, 41.3 mm SL, MZB 26456, 43.7 mm SL, MZB 26457, 39.8 mm SL, MZB 26458, 40.7 mm SL, MZB 26459, 43.2 mm SL, MZB 26460, 43.7 mm SL, MZB 26461, 5 specimens, 38.3–42.4 mm SL, collected with the holotype.

**Diagnosis.** A species of *Stolephorus* with the following combination of characters: 1UGR 19–21 (modally 20), 1LGR 26–30 (28), 1TGR 47–51 (47); 2UGR 13–16 (14), 2LGR 24–27 (25), 2TGR 37–42 (39); 3UGR 10–13 (12), 3LGR 14–16 (15), 3TGR 25–28 (27); 4UGR 8–11 (9), 4LGR 11–13 (12), 4TGR 19–24 (21); prepelvic scutes 5–7 (6); total vertebrae 38–39 (39); long maxilla, posterior tip just reaching or slightly



**Figure 9.** **A** lateral **B** dorsal and **C** ventral views of preserved holotype of *S. eclipsis* sp. nov., MZB 26452, 40.3 mm SL, Bintan Island, Riau Archipelago, Indonesia. Scale bars indicate 2 mm.

short of posterior margin of opercle; predorsal scute present; pelvic scute without spine; body scales deciduous; posterior border of pre-opercle concave, indented; paired dark patch on parietal area with little following pigmentation; distinct double pigment lines along dorsum posterior to dorsal fin; black spots below eye and on lower-jaw tip absent; anal-fin base short, 17.6–19.3% (mean 18.6%) of SL; third dorsal-fin ray 16.5–18.8% (17.6%) of SL; pelvic fin short, 8.7–9.9% (9.4%) of SL, its posterior tip usually not reaching to vertical through dorsal-fin origin when depressed; distance between posterior ends of supramaxilla and maxilla 4.7–5.4% (5.1%) of SL; pre-dorsal-fin length 51.3–54.9% (53.4%) of SL; dorsal-fin base short, 13.1–14.5% (13.8%) of SL.

**Description.** Data for holotype presented first, followed by data for paratypes in parentheses (if different). Counts and measurements, expressed as percentages of SL or HL, given in Tables 5 and 6. Body laterally compressed, elongate, deepest at dorsal-fin origin. Dorsal profile of head and body slightly convex from snout tip to dorsal-fin origin, gently lowering to uppermost point of caudal-fin base. Ventral profile of head and body slightly convex from lower jaw tip to pelvic-fin insertion, thereafter, slowly rising to lowermost point of caudal-fin base. Single spine-like scute just anterior to dorsal-fin origin. Abdomen somewhat rounded, covered with six (five to seven) spine-like prepelvic scutes. Pelvic scute without spine. Postpelvic scutes absent. Anus just anterior to anal-fin origin. Snout tip rounded; snout length less than eye diameter. Mouth large, inferior, ventral to body axis, extending backwards beyond posterior margin of eye. Maxilla long, its posterior tip pointed, just reaching (or slightly short of) opercle posterior margin. Lower jaw slender. Single row of conical teeth on both jaws and palatine. Patch of fine conical teeth on pterygoid. Several distinct conical teeth on vomer. Several rows of conical teeth on upper edges of basihyal and basibranchial. Eye large, round, covered with adipose eyelid, positioned laterally on head dorsal to

horizontal through pectoral-fin insertion, visible in dorsal view. Pupil round. Orbit elliptical. Nostrils close to each other, anterior to orbit. Posterior margin of pre-opercle concave, indented. Subopercle and opercle with smoothly rounded posterior margins. Gill membrane without serrations. Interorbital space flat, width less than eye diameter. Pseudobranchial filaments present, length of longest filament less than eye diameter. Gill rakers long, slender, rough, visible from side of head when mouth opened. Single row of asperities on anterior surface of gill rakers. Isthmus muscle long, reaching anteriorly to posterior margin of gill membranes. Urohyal hidden by isthmus muscle, not visible without dissection. Gill membrane on each side joined distally, most of isthmus muscle exposed, not covered by gill membrane. Body scales deciduous, completely lacking on specimens, except for prepelvic scutes. Head scales absent. Fins scaleless, except for broad triangular sheath of scales on caudal fin. Dorsal-fin origin posterior to vertical through base of last pelvic-fin ray, slightly posterior to middle of body. Dorsal and anal fins with three anteriormost rays unbranched. First dorsal- and anal-fin rays minute. Anteriormost three rays of both dorsal and anal fins closely-spaced. Anal-fin origin just below base of eighth (eighth to eleventh) dorsal-fin ray. Posterior tip of depressed anal fin not reaching caudal-fin base. Uppermost pectoral-fin ray unbranched, inserted below body axis. Posterior tip of pectoral fin not reaching to pelvic fin insertion. Dorsal, ventral and posterior margins of pectoral fin nearly linear. Pelvic fin shorter than pectoral fin, insertion anterior to vertical through dorsal-fin origin. Posterior tip of depressed pelvic fin not reaching to vertical through dorsal-fin origin (reaching to vertical through first to third dorsal-fin ray origin in some paratypes). Caudal fin forked, posterior tips pointed.

**Colour of preserved specimens.** Body uniformly pale ivory. A pair of distinct dark patches on parietal region, with little pigmentation on occipital area. Double pigmented lines dorsally posterior to dorsal fin. A few melanophores scattered anteriorly on snout. No black spots below eye and on lower-jaw tip. Melanophores scattered along bases of dorsal and anal fins. All fins transparent, melanophores scattered along fin rays of caudal fin and anterior parts of dorsal and anal fins.

**Distribution.** *Stolephorus eclipsis* sp. nov. is currently known only from Bintan Island, Riau Archipelago, Indonesia (Fig. 4).

**Etymology.** The specific name “*eclipsis*” refers to eclipse, reminiscent of the concave pre-opercle of the new species.

**Comparisons.** The new species differs from *S. bengalensis*, *S. diabolus* and *S. eldorado* in having higher gill raker counts [1TGR, 47–51 or more in *S. eclipsis* (vs. 47 or fewer in the other three species); 2TGR, 37–42 in *S. eclipsis* (vs. 39 or fewer); 3TGR, 25–28 in *S. eclipsis* (vs. 27 or fewer); and 4TGR, 19–24 in *S. eclipsis* (vs. 22 or fewer) (Fig. 7)], a shorter anal-fin base (17.6–19.3% of SL in *S. eclipsis* vs. 19.0–21.3% in *S. bengalensis*, 19.8–22.3% in *S. diabolus* and 19.0–22.3% in *S. eldorado*; Fig. 8C) and pelvic fin [8.7–9.9% (mean 9.4%) of SL in *S. eclipsis* vs. 9.4–11.0% (10.2%) in *S. bengalensis*, 9.6–11.3% (10.0%) in *S. diabolus* and 9.1–11.0% (10.0%) in *S. eldorado*; Fig. 8D) and shorter distance between the posterior ends of the supramaxilla and maxilla [4.7–5.4% (5.1%) of SL in *S. eclipsis* vs. 5.3–6.6% (5.8%) in *S. bengalensis*, 5.7–6.4% (6.1%)

in *S. diabolus* and 5.0–6.3% (5.6%) in *S. eldorado*; Fig. 8E]. *Stolephorus eclipsis* also differs from *S. bengalensis* in having a shorter third dorsal-fin ray (16.5–18.8% of SL in *S. eclipsis* vs. 18.5–19.9% in *S. bengalensis*; Fig. 8B) and lower total vertebral number [38–39 (modally 39) vs. 40 or 41 (40) (Table 4)]. Moreover, *S. eclipsis* is distinguished from *S. diabolus* by a greater pre-dorsal-fin distance [51.3–54.9% (mean 53.4%) of SL in *S. eclipsis* vs. 51.3–52.9% (52.1%) in *S. diabolus*; Fig. 10A] and shorter dorsal-fin base (13.1–14.5% of SL vs. 13.9–16.6%; Fig. 10B) and postorbital head length (11.5–12.9% of SL vs. 12.8–14.2%; Fig. 10C).

***Stolephorus eldorado* sp. nov.**

<https://zoobank.org/0A916EDA-E70A-4EAF-85C2-7AC51C588BC4>

[New English name: El Dorado Anchovy]

Figs 1A, B, D, E, F, 11; Tables 2–4


*Stolephorus insularis* (not of Delsman): Whitehead et al. 1988 (in part): 413 unnumbered fig. (Taiwan to Java Sea); Young et al. 1999: 222, fig. 7 (western coast of Taiwan); Wongratana et al. 1999 (in part): 1736, unnumbered fig. (Taiwan to Java Sea); Hata 2018: 41, unnumbered figs (Ha Long Bay, northern Vietnam).

*Stolephorus tri* (not of Bleeker): Zhang 2001: 129, fig. II-59 (Beihai City, Guangxi Province, China).


*Stolephorus bengalensis* (not of Dutt and Babu Rao): Hata et al. 2019 (in part): 24, fig. 12a, b (Taiwan; Hainan Island, China; Ha Long Bay, Vietnam; Gulf of Thailand; Songkhla, Thailand; Kuala Terengganu, Terengganu, Malaysia); Hata 2019: 206, unnumbered figs (Ke-tzu-liao, Ziguan District, Kaohsiung, Taiwan); Hata et al. 2022: (in part) 34 (Wenzhou City, Zhejiang Province, China).

**Holotype.** KAUM–I. 94517, 44.4 mm SL, Ha Long Bay, Ha Long City, Quang Ninh District, Vietnam (purchased at fish market in Ha Long City), 24 Oct 2016; coll. by H. Hata and M. Matsunuma.

**Paratypes.** 57 specimens, 37.5–58.8 mm SL. **TAIWAN:** ASIZP 73957, 51.8 mm SL, Fangyan, Changhua (23°57'42.8"N, 120°17'39.8"E); KAUM–I. 110282, 49.5 mm SL, KAUM–I. 113142, 54.0 mm SL, KAUM–I. 113143, 45.5 mm SL, KAUM–I. 113144, 44.3 mm SL, KAUM–I. 113145, 46.3 mm SL, KAUM–I. 113146, 37.5 mm SL, KAUM–I. 113147, 47.3 mm SL, KAUM–I. 113148, 55.3 mm SL, KAUM–I. 113149, 49.4 mm SL, KAUM–I. 113150, 45.9 mm SL, KAUM–I. 113151, 47.3 mm SL, off Ke-tzu-liao, Ziguan District, Kaohsiung. **CHINA:** BMNH 1965.4.1.981–983, 3 specimens, 58.1–58.8 mm SL, Stanley, Hong Kong. **VIETNAM:** FRLM 49725, 46.9 mm SL, KAUM–I. 67322, 46.7 mm SL, KAUM–I. 67405, 45.6 mm SL, KAUM–I. 94509, 41.4 mm SL, KAUM–I. 94518, 43.7 mm SL, KAUM–I. 94519, 38.8 mm SL, KAUM–I. 94520, 41.7 mm SL, KAUM–I. 94521, 43.4 mm SL, Ha Long Bay, Ha Long, Quang Ninh Province. **THAILAND:** CAS 46931, 8 specimens, 44.4–46.7 mm SL, between Bangsaen and Chol Buri, Chol Buri, Gulf of Thailand; CAS 230414,



**Figure 10.** Relationships of **A** pre-dorsal-fin length (as % of standard length; SL) **B** dorsal-fin base length (as % of SL), and **C** postorbital length (as % of SL) in *Stolephorus diabolus* sp. nov. (yellow squares) and *S. eclipsis* sp. nov. (green diamonds).



**Figure 11.** Holotype of *S. eldorado* sp. nov. (KAUM-I. 94517, 44.4 mm SL, Ha Long Bay, northern Vietnam) **A** lateral view (fresh) **B** lateral **C** dorsal, and **D** ventral views (preserved); paratypes of *S. eldorado* sp. nov. (KAUM-I. 67322, 46.7 mm SL, Ha Long Bay, northern Vietnam) **E** lateral view (fresh), (KAUM-I. 110282, 49.5 mm SL, Ke-tzu-liao, southwestern Taiwan) **F** lateral **G** dorsal, and **H** ventral views (fresh). Scale bars indicate 2 mm.

4 specimens, 39.9–45.8 mm SL, Lem Nam Point, south tip of Lem Nam Peninsula, Gulf of Thailand (12°02'55"N, 102°35'35"E), approx. 0.6 m depth; KAUM-I. 23190, 48.2 mm SL, Gulf of Thailand (obtained at fish market in Mahachai, Samut Prakan Province), trawl; NSMT-P 142790, 47.9 mm SL, Ko Maeo Island, off Songkhla; URM-P 12398, 3 specimens, 43.0–45.5 mm SL, Song Khula; URM-P 13635, 11 specimens, 46.2–49.5 mm SL, Ang Sila. **INDONESIA:** BMNH 1965.10.20.42–47, 6 specimens, 40.6–43.5 mm SL, 20 miles (approx. 32 km) east of Tegal, Java.

**Diagnosis.** A species of *Stolephorus* with the following combination of characters: 1UGR 16–21 (modally 18), 1LGR 23–28 (25), 1TGR 40–47 (42); 2UGR 10–14 (13), 2LGR 20–24 (23), 2TGR 33–38 (rarely 30) (modally 36); 3UGR 8–12 (modally 10), 3LGR 12–14 (13), 3TGR 20–26 (23); 4UGR 7–10 (8), 4LGR 9–12 (11), 4TGR 16–22 (18); prepelvic scutes 5–7 (6); total vertebrae 38–40 (39); long maxilla, posterior tip just reaching or slightly short of posterior margin of opercle; predorsal scutes present; pelvic scute without spine; body scales deciduous; posterior border of pre-opercle concave, indented; paired dark patch on parietal area with little following pigmentation; distinct double pigment lines along dorsum posterior to dorsal fin; black spots below eye and on lower-jaw tip absent; anal-fin base long, 19.0–22.3% (20.4%) of SL; orbit rather long, 8.2–9.9% (8.9%) of SL; third dorsal-fin ray short, 15.9–18.6% (17.4%) of SL; pelvic fin rather long, 9.1–11.0% (10.0%) of SL, its posterior tip usually not reaching to vertical through dorsal-fin origin when depressed in individuals > 50 mm SL; distance between posterior ends of supramaxilla and maxilla 5.0–6.3% (5.6%) of SL.

**Description.** Data for holotype presented first, followed by data for paratypes in parentheses (if different). Counts and measurements, expressed as percentages of SL or HL, given in Tables 2 and 3. Body laterally compressed, elongate, deepest at dorsal-fin origin. Dorsal profile of head and body slightly convex from snout tip to dorsal-fin origin, gently lowering to uppermost point of caudal-fin base. Ventral profile of head and body slightly convex from lower jaw tip to pelvic-fin insertion, thereafter, slowly rising to lowermost point of caudal-fin base. Single spine-like scute just anterior to dorsal-fin origin. Abdomen somewhat rounded, covered with six (five to seven) spine-like prepelvic scutes. Pelvic scute without spine. Postpelvic scutes absent. Anus just anterior to anal-fin origin. Snout tip rounded; snout length less than eye diameter. Mouth large, inferior, ventral to body axis, extending backwards beyond posterior margin of eye. Maxilla long, its posterior tip pointed, just reaching (or slightly short of) opercle posterior margin. Lower jaw slender. Single row of conical teeth on both jaws and palatine. Patch of fine conical teeth on pterygoid. Several distinct conical teeth on vomer. Several rows of conical teeth on upper edges of basihyal and basibranchial. Eye large, round, covered with adipose eyelid, positioned laterally on head dorsal to horizontal through pectoral-fin insertion, visible in dorsal view. Pupil round. Orbit elliptical. Nostrils close to each other, anterior to orbit. Posterior margin of pre-opercle concave, indented. Subopercle and opercle with smoothly rounded posterior margins. Gill membrane without serrations. Interorbital space flat, width less than eye diameter. Pseudobranchial filaments present, length of longest filament less than eye diameter. Gill rakers long, slender, rough, visible from side of head when mouth opened. Single row of asperities on anterior surface of gill rakers. Isthmus muscle long, reaching anteriorly to posterior margin of gill membranes. Urohyal hidden by isthmus muscle, not visible without dissection. Gill membrane on each side joined distally, most of isthmus muscle exposed, not covered by gill membrane. Body scales deciduous, completely lacking on all specimens, except for prepelvic scutes. Head scales absent. Fins scaleless, except for broad triangular sheath of scales on caudal fin. Dorsal-fin origin posterior to vertical through base of last pelvic-fin ray, slightly posterior to middle of body. Dorsal and anal fins with three anteriormost rays unbranched. First dorsal- and anal-fin rays minute. Anteriormost three rays of both dorsal and anal fins closely spaced. Anal-fin origin just below base of ninth (eighth to tenth) dorsal-fin ray. Posterior tip of depressed anal fin not reaching caudal-fin base. Uppermost pectoral-fin ray unbranched, inserted below body axis. Posterior tip of pectoral fin not reaching to pelvic fin insertion. Dorsal, ventral and posterior margins of pectoral fin nearly linear. Pelvic fin shorter than pectoral fin, insertion anterior to vertical through dorsal-fin origin. Posterior tip of depressed pelvic fin not reaching to vertical through dorsal-fin origin (reaching to vertical through first to fourth dorsal-fin ray origin in some paratypes smaller than 50 mm SL). Caudal fin forked, posterior tips pointed.

**Colour of fresh specimens.** (based on colour photographs of KAUM–I. 67322, 46.7 mm SL, KAUM–I. 67405, 45.6 mm SL, KAUM–I. 94517, 44.4 mm SL, KAUM–I. 94521, 43.4 mm SL and KAUM–I. 110282, 49.5 mm SL). Body yellowish milky-white, a silver longitudinal band, of width slightly less than pupil diameter, extending from just above posterior tip of pectoral fin to caudal-fin base. Caudal fin yellow with black posterior margin. Melanophores scattered along caudal-fin rays, ventral surface of caudal pe-

duncle and bases of dorsal and anal fins. Fin rays of dorsal and anal fins yellow. A few melanophores scattered on snout and fin rays of anterior part of dorsal fin. Fin rays and fin membrane of pectoral and pelvic fins transparent whitish, lacking melanophores. A pair of dark patches on parietal region, with little pigmentation on occipital area. Distinct double pigment lines on dorsum from end of dorsal-fin base to caudal-fin base. Body wholly yellowish when freshly caught (Fig. 11E), quickly becoming white after death (Figs 11F–H).

**Colour of preserved specimens.** Body uniformly pale white. A pair of distinct dark patches on parietal region, with little pigmentation on occipital area. Melanophores scattered on posterior margins of scale pockets on dorsum. Double pigmented lines dorsally posterior to dorsal fin. A few melanophores scattered anteriorly on snout. No black spots below eye and on lower-jaw tip. Melanophores scattered along bases of dorsal and anal fins. All fins transparent, with melanophores scattered along fin rays of caudal fin and anterior parts of dorsal and anal fins.

**Distribution.** *Stolephorus eldorado* sp. nov. is distributed in the western Pacific from Taiwan to Java, Indonesia (Fig. 4). The species is abundantly caught by trawl and marketed fresh in northern Vietnam. It is a set net bycatch in south-western Taiwan.

**Etymology.** The specific name “*eldorado*”, referring to the mythical city of gold, reflects the bright yellow colouration of the new species.

**Morphological comparisons.** *Stolephorus eldorado* sp. nov. has been previously identified as *S. insularis* or *S. bengalensis* (together with *S. bengalensis*, *S. diabolus* and *S. eclipsis* as recognised here) (e.g. Whitehead et al. 1988; Wongratana et al. 1999; Hata et al. 2019). However, *S. eldorado* is distinguished from *S. diabolus* and *S. eclipsis* by having an intermediate number of gill rakers on each gill arch (Table 2; Fig. 7). More detailed comparisons of *S. eldorado* with the latter two species are given in “Comparisons” under each species.

Although *S. eldorado* sp. nov. closely resembles *S. bengalensis* in having very similar numbers of gill rakers on each gill arch, the former differs from the latter in having a greater orbit diameter [maximum orbit diameter 8.2–9.9% (mean 8.9%) of SL vs. 7.3–8.6% (8.1%) in *S. bengalensis* (Fig. 8A)], shorter third dorsal-fin ray [15.9–18.6% (mean 17.4%) of SL vs. 18.5–19.9% (19.0%)] (Fig. 8B) and fewer total vertebrae [38–40 (modally 39) vs. 40 or 41 (40)] (Table 4).

### Key to species previously identified as *Stolephorus insularis* by Whitehead et al. (1988) or *Stolephorus bengalensis* by Hata et al. (2019)

- 1 1TGR  $\leq$  38.....  
.....*S. diabolus* (western coast of Malay Peninsula to Singapore)
- 1TGR  $\geq$  41..... 2
- 2 1TGR  $\geq$  47; anal-fin base short, less than 19.3% of SL; pelvic fin short, 8.7–9.9% of SL; distance between posterior ends of supramaxilla and maxilla less than 5.4% of SL.....*S. eclipsis* (Bintan Island, Indonesia)
- 1TGR  $\leq$  47; anal-fin base rather long, more than 19.0% of SL; pelvic fin rather long, 9.1–10.1% of SL; distance between posterior ends of supramaxilla and maxilla more than 5.0% of SL..... 3

- 3 Third dorsal-fin ray short, 15.9–18.6% (mean 17.5%) of SL; maximum orbit diameter 8.2–9.9% (8.9%) of SL..... ***S. eldorado* (Taiwan to Java)**
- Third dorsal-fin ray long, 18.5–19.9% (mean 19.0%) of SL; maximum orbit diameter 7.3–8.6% (8.1%) of SL.....  
 ..... ***S. bengalensis* (Pakistan to Bay of Bengal)**

## Acknowledgements

We thank K.-T. Shao and S.-P. Huang (ASIZP), O. Crimmen, J. Maclaine and N. Martin (BMNH), D. Catania and M. Hoang (CAS), S. Kimura (FRLM), T. Peristywady (LBRC), K. Wibowo and G. Wahyudewantoro (MZB), K. Matsuura, G. Shinohara and M. Nakae (NSMT), K. Miyamoto (OCF), M. Manjaji-Matsumoto, A. A. Flora and S. R. M. Shaleh (Universiti Malaysia Sabah), K. Koeda (URM) J. Williams, K. Murphy, S. Raredon and D. Pitassy (USNM) and M. Aizawa, K. Sakamoto and R. Ueshima (ZUMT) for opportunities to examine specimens of *Stolephorus*. We also thank Y. Haraguchi and other volunteers and students of KAUM and NSMT for their curatorial assistance and G. Hardy (Ngunguru, New Zealand), for reading the manuscript and providing help with English. Vietnamese specimens were collected with the support of the Institute of Marine Environment and Resources (Haiphong) and the Ha Long Bay Management Department (Ha Long), with permission for the use of specimens granted by the Biodiversity Conservation Agency, Ministry of Natural Resources and Environment (Hanoi). Pakistan specimens were collected by the Marine Fisheries Department (MFD), Government of Pakistan, Karachi and donated from MFD via P. N. Psomadakis (Fisheries and Aquaculture Department, FAO, Rome) (permission no: MFD/UTF/Field guide/2014/J-1). Finally, we thank two anonymous reviewers and the editor for their insightful comments which helped to improve the manuscript. This study was supported in part by JSPS Overseas Research Fellowships (202160519), JSPS KAKENHI Grant Number 19K23691, JSPS Fellows (DC2: 29-6652) and the Sasakawa Scientific Research Grant from the Japan Science Society (28–745) to HH; JSPS KAKENHI Grant Numbers 20H03311 and 21H03651; the JSPS Core-to-core CREPSUM JPJSCCB2020009; and “Establishment of Glocal Research and Education Network in the Amami Islands” project of Kagoshima University adopted by the Ministry of Education, Culture, Sports, Science and Technology, Japan to HM; and short term research grant 304/PBIOLOGI/6315400 from Universiti Sains Malaysia to SL.

## References

- Babu Rao M (1966) A new species of *Stolephorus* Lacépède from the east coast of India (Pisces: Engraulidae). *Annals & Magazine of Natural History* 13(9): 101–110. <https://doi.org/10.1080/00222936608651641>
- Bleeker P (1852) Bijdrage tot de kennis der Haringachtige visschen van den Soenda-Molukschen Archipel. *Verhandelingen van het Bataviaasch Genootschap van Kunsten en Wetenschappen* 24(8): 1–52.

- Boulenger GA (1900) Descriptions of new fishes from the Cape of Good Hope. Marine Investigations in South Africa 8: 10–12[+ pls. 1–3].
- De Vis CW (1882) Description of three new fishes of Queensland. Proceedings of the Linnean Society of New South Wales 7(3): 318–320. <https://doi.org/10.5962/bhl.part.22754>
- Dutt S, Babu Rao M (1959) Occurrence of *Anchoviella baganensis* Hardenberg off east coast of India. Current Science 28: 160–161.
- Edler D, Klein J, Antonelli A, Silvestro D (2020) raxmlGUI 2.0: A graphical interface and toolkit for phylogenetic analyses using RAxML. Methods in Ecology and Evolution 12(2): 373–377. <https://doi.org/10.1111/2041-210X.13512>
- Gangan SS, Pavan-Kumar A, Jahageerdar S, Jaiswar AK (2020) A new species of *Stolephorus* (Clupeiformes: Engraulidae) from the Bay of Bengal. Zootaxa 4743(3): 561–574. <https://doi.org/10.11646/zootaxa.4743.4.6>
- Hardenberg JDF (1933) New *Stolephorus* species of the Indo-Australian seas. Natuurkundig Tijdschrift voor Nederlandsch Indië 93: 258–263.
- Hata H (2018) *Stolephorus insularis* Hardenberg, 1933. In: Kimura S, Imamura H, Quan NV, Duong PT (Eds) Fishes of Ha Long Bay, the World Natural Heritage Site in Northern Vietnam. Fisheries Research Laboratory, Mie University, Shima, 41.
- Hata H (2019) Family Engraulidae. In: Koeda K, Ho H-C (Eds) Fishes of southern Taiwan. National Museum of Marine Biology & Aquarium, Pingtung, 199–210.
- Hata H, Motomura H (2017) A new species of anchovy, *Encrasicholina auster* (Clupeiformes: Engraulidae), from Fiji, southwestern Pacific Ocean. New Zealand Journal of Zoology 44(2): 122–128. <https://doi.org/10.1080/03014223.2016.1268177>
- Hata H, Motomura H (2018a) *Stolephorus continentalis*, a new anchovy from the north-western South China Sea and redescription of *Stolephorus chinensis* (Günther 1880) (Clupeiformes: Engraulidae). Ichthyological Research 65(3): 374–382. <https://doi.org/10.1007/s10228-018-0621-z>
- Hata H, Motomura H (2018b) First record of the anchovy, *Stolephorus teguhi* (Engraulidae) from the Philippines. Philippine Journal of Systematic Biology 11: 20–24. <https://doi.org/10.26757/pjsb.2017b11017>
- Hata H, Motomura H (2018c) Additional specimens of the poorly known anchovy *Stolephorus multibranchus* (Clupeiformes: Engraulidae) from Kosrae, Caroline Islands. Biogeography 20: 78–84.
- Hata H, Motomura H (2018d) Redescription and distributional range extension of the poorly known anchovy *Stolephorus nelsoni* (Clupeiformes: Engraulidae). Acta Ichthyologica et Piscatoria 48(4): 381–386. <https://doi.org/10.3750/AIEP/02501>
- Hata H, Motomura H (2018e) *Stolephorus insignis*, a new anchovy from the western Pacific, and redescription of *Stolephorus apiensis* (Jordan and Seale 1906) (Clupeiformes: Engraulidae). Ichthyological Research 66(2): 280–288. <https://doi.org/10.1007/s10228-018-00675-5>
- Hata H, Motomura H (2021a) Two new species of *Stolephorus* (Teleostei: Clupeiformes: Engraulidae) from the western Pacific. The Raffles Bulletin of Zoology 69: 109–117. <https://doi.org/10.26107/RBZ-2021-0009>
- Hata H, Motomura H (2021b) *Stolephorus grandis*, a new anchovy (Teleostei: Clupeiformes: Engraulidae) from New Guinea and Australia. Zootaxa 5004(3): 481–489. <https://doi.org/10.11646/zootaxa.5004.3.5>

- Hata H, Motomura H (2021c) A new species of the anchovy genus *Stolephorus* Lacepède 1803 from North Sumatra, Indonesia, and redescrptions of *Stolephorus pacificus* Baldwin 1984 and *Stolephorus teguhi* Kimura, Hori and Shibukawa 2009 (Teleostei: Clupeiformes: Engraulidae). *Zoological Studies* 60: 65. <https://doi.org/10.6620/ZS.2021.60-65>
- Hata H, Motomura H (2022) Redescription of *Stolephorus ronquilloi* Wongratana, 1983 and description of *Stolephorus hindustanensis*, new anchovy from the western coast of India (Teleostei: Clupeiformes: Engraulidae). *Taxonomy* 2(1): 124–135. <https://doi.org/10.3390/taxonomy2010010>
- Hata H, Lavoué S, Motomura H (2019) Taxonomic status of seven nominal species of the anchovy genus *Stolephorus* described by Delsman (1931), Hardenberg (1933), and Dutt and Babu Rao (1959), with redescrptions of *Stolephorus tri* (Bleeker 1852) and *Stolephorus waitei* Jordan and Seale 1926 (Clupeiformes: Engraulidae). *Ichthyological Research* 67(1): 7–38. <https://doi.org/10.1007/s10228-019-00697-7>
- Hata H, Lavoué S, Motomura H (2020a) *Stolephorus babarani*, a new species of anchovy (Teleostei: Clupeiformes: Engraulidae) from Panay Island, central Philippines. *Zootaxa* 4178(4): 509–520. <https://doi.org/10.11646/zootaxa.4718.4.5>
- Hata H, Lavoué S, Motomura H (2020b) *Stolephorus acinaces*, a new anchovy from the northern Borneo, and redescription of *Stolephorus andhraensis* (Babu Rao 1966) (Clupeiformes: Engraulidae). *Marine Biodiversity* 50(6): 102. <https://doi.org/10.1007/s12526-020-01115-2>
- Hata H, Lavoué S, Motomura H (2021) Taxonomic status of nominal species of the anchovy genus *Stolephorus* previously regarded as synonyms of *Stolephorus commersonii* Lacepède 1803 and *Stolephorus indicus* (van Hasselt 1823), and descriptions of three new species (Clupeiformes: Engraulidae). *Ichthyological Research* 68(3): 327–372. <https://doi.org/10.1007/s10228-020-00792-0>
- Hata H, Koeda K, Aizawa M, Sakamoto K, Ueshima R (2022) A list of Clupeiformes (Actinopterygii: Teleostei) specimens deposited in the Department of Zoology, The University Museum, The University of Tokyo. *The University Museum, The University of Tokyo. Bulletin* 128: 17–58.
- Hubbs CL, Lagler JF (1947) *Fishes of the Great Lakes region*. Cranbrook Institute of Science Bulletin 26: [i–xi +]1–186.
- Kimura K, Hori K, Shibukawa K (2009) A new anchovy, *Stolephorus teguhi* (Clupeiformes: Engraulidae) from North Sulawesi, Indonesia. *Ichthyological Research* 56(3): 292–295. <https://doi.org/10.1007/s10228-009-0103-4>
- Kozlov AM, Darriba D, Flouri T, Morel B, Stamatakis A (2019) RAxML-NG: A fast, scalable and user-friendly tool for maximum likelihood phylogenetic inference. *Bioinformatics* 35(21): 4453–4455. <https://doi.org/10.1093/bioinformatics/btz305>
- Nuryanto A, Pramono H, Sastranegara MH (2017) Molecular Identification of Fish Larvae from East Plawangan of Segara Anakan, Cilacap, Central Java, Indonesia. *Biosaintifika* 9(1): 33–40. <https://doi.org/10.15294/biosaintifika.v9i1.9191>
- Pang Z, Liu L, Liu B, Gong L, Jiang L, Lü Z (2019) The complete mitochondrial genome of *Stolephorus insularis* (Clupeiformes: Engraulidae) and phylogenetic studies of Engraulidae. *Mitochondrial DNA Part B* 4(1): 921–922. <https://doi.org/10.1080/23802359.2018.1536470>

- Sabaj MH (2020) Codes for natural history collections in Ichthyology and Herpetology. *Copeia* 108(3): 593–669. <https://doi.org/10.1643/ASIHCODONS2020>
- Stecher G, Tamura K, Kumar S (2020) Molecular Evolutionary Genetics Analysis (MEGA) for macOS. *Molecular Biology and Evolution* 37(4): 1237–1239. <https://doi.org/10.1093/molbev/msz312>
- Ward RD, Zemlak TS, Innes BH, Last PR, Hebert PDN (2005) DNA barcoding Australia's fish species. *Philosophical Transactions of the Royal Society B Biological Sciences* 360: 1847–1857. <https://doi.org/10.1098/rstb.2005.1716>
- Whitehead PJP (1967) Indian Ocean anchovies collected by the Anton Bruun and Te Vega, 1963–1964. *Journal of the Marine Biological Association of India* 9: 13–37.
- Whitehead PJP, Nelson GJ, Wongratana T (1988) FAO species catalogue. Vol. 7. Clupeoid fishes of the world (suborder Clupeoidei). An annotated and illustrated catalogue of the herrings, sardines, pilchards, sprats, shads, anchovies and wolf-herrings. Pt 2 – Engraulidae. FAO Fisheries Synopsis No 125 7(2):[i–viii +] 305–579.
- Wongratana T (1983) Diagnoses of 24 new species and proposal of a new name for a species of Indo-Pacific clupeoid fishes. *Japanese Journal of Ichthyology* 29: 385–407.
- Wongratana T (1987a) Four new species of clupeoid fishes (Clupeidae and Engraulidae) from Australian waters. *Proceedings of the Biological Society of Washington* 100: 104–111.
- Wongratana T (1987b) Two new species of anchovies of the genus *Stolephorus* (Engraulidae), with a key to species of *Engraulis*, *Encrasicholina*, and *Stolephorus*. *American Museum Novitates* 2876: 1–8.
- Wongratana T, Munroe TA, Nizinski MS (1999) Order Clupeiformes. Engraulidae. Anchovies. In: Carpenter KE, Niem VH (Eds) FAO species identification guide for fishery purposes. The living marine resources of the western central Pacific. Vol. 3. Batoid fishes, chimaeras and bony fishes pt 1 (Elopidae to Linophrynidae). FAO, Rome, 1698–1753.
- Young S-S, Chiu T-S, Shen S-C (1994) A revision of the family Engraulidae (Pisces) from Taiwan. *Zoological Studies* 33(3): 217–227.
- Zhang S (2001) Fauna Sinica Osteichthyes. Acipenseriformes, Elopiformes, Clupeiformes, Gonorhynchiformes. Science Press, Beijing, 209 pp.