

A new species of *Patania* from the Hainan Island, China (Lepidoptera, Crambidae)

Dan Xu¹, Xi-Cui Du¹

¹ College of Plant Protection, Southwest University, Chongqing, China

Corresponding author: Xi-Cui Du (duxicui@hotmail.com)

Academic editor: Colin Plant | Received 11 April 2016 | Accepted 15 August 2016 | Published 1 September 2016

<http://zoobank.org/4BA9FEA0-2187-4133-A315-F9C34B938A19>

Citation: Xu D, Du X-C (2016) A new species of *Patania* from the Hainan Island, China (Lepidoptera, Crambidae). ZooKeys 614: 129–135. doi: 10.3897/zookeys.614.8810

Abstract

Patania clava sp. n. is described from the Diaoluo Mountain of Hainan Island, China. The new species is distinguished from its most similar congener, *P. iopasalis* (Walker, 1859), by the following features: wingspan 33.0–35.0 mm (vs. 21.0–30.0 mm), ventral cilia of the male antenna as long as the diameter of flagellomere (vs. 1/4), the thick finger-like gnathos (vs. the short broad sheet-like), a long thick needle-like cornutus stretching out from (vs. embedded in) a cluster of spicular cornuti near apex. Images of adult and genitalia of the new species are provided.

Keywords

Diaoluo Mountain, Pyraloidea, Spilomelinae, taxonomy

Introduction

The genus *Patania* was established by Moore (1888) based on the type species, *Botys concatenalis* Walker, 1866, from Darjeeling, India. Warren described another new *Patania* species in 1896, *Patania verecunda* (Warren, 1896). Meanwhile, Hampson (1896) regarded *Patania* and *Pleuroptya* Meyrick, 1890 as synonyms of *Sylepta* Hübner, 1823. Hampson's opinion was followed by most researchers (Hampson 1898; Shibuya 1928, 1929; Klima 1939; Bae et al. 2008) except for Rose and Singh (1989) who published a new species of *Patania* separately. Inoue (1980) revalidated *Pleuroptya*, and some new

species, sub-species, and combinations of *Pleuroptya* were recorded since (Yamanaka 1995, 1998; Munroe 1983; Shaffer and Munroe 1989; Leraut 2005). Kirti and Gill (2007) considered *Patania* and *Pleuroptya* congeneric by comparing the morphological characters and male genitalia of their respective type species. The generic name *Pleuroptya* Meyrick, 1890 was suppressed as subjective synonym of *Patania* Moore, 1888 by Kirti and Gill (2007).

Globally, there are approximately 50 described species of *Patania* (Kirti and Gill 2007; Li et al. 2009, Nuss et al. 2003–2015), 28 species being recorded in China (Li et al. 2009; Xu 2015). In this paper, *Patania clava* sp. n., collected from the Diaoluo Mountain, Hainan Island, China, is described as new to science.

Materials and methods

Specimens were collected by 250-W high-pressure mercury lamps. They were hand-collected alive and killed by ethyl acetate. The type specimens of *Patania* species deposited in the Natural History Museum of London (NHM) have been examined by corresponding author. All the type specimens of the new species are deposited in the College of Plant Protection, Southwest University, Chongqing, China (SWUCPP).

The terminology mainly follows Kristensen (2003) and Slamka (2013).

Genitalia preparation followed the procedure of Li and Zheng (1996), using boiling 10% NaOH solutions to digest internal tissues; after careful cleaning and removal of scales and content of coelom, genitalia were examined, compared, and described before being mounted on microscope slides by the first author. The images of the adults were taken with a digital camera (Nikon P7700). The illustrations of the genitalia were prepared with a digital camera Leica DFC 450 attached to a digital microscope Leica M205 A.

Taxonomic account

Patania clava sp. n.


<http://zoobank.org/3DEA6F10-06C7-4ABB-BE34-97C03A4E4729>

Figs 1–4, 9–10

Holotype. male, China, Hainan, Mt. Diaoluo, 109.87°E, 18.72°N, 900 m, 23.V.2014, leg. Li-Jun Xu & Dan Xu, pinned, slide number XD15056. Original label: “Hainan, Diaoluo Mountain, vocational village, 900 m, 23.V.2014, Li-jun Xu & Dan Xu”.

Paratype. 1 female, same data as holotype, slide number XD15050.


Diagnosis. This species can be distinguished by the wingspan of 33.0–35.0 mm (Fig. 1); the brown patch near base of the labial palpus (Fig. 2), the male antenna with ventral cilia nearly as long as the diameter of flagellomere; in the male genitalia by the thick finger-like gnathos bearing long setae on the apex (Fig. 3), the phallus with a dor-



Figures 1–4. *Patania clava* sp. n., Holotype. **1** Adult **2** Head **3** Gnathos **4** Cornuti.

sally protruding sclerotized structure with a slant nailhead-like apex, and a long thick needle-like cornutus stretching out from a cluster of spicular cornuti near apex (Fig. 4).

Description. Adult (Figs 1–2). Wingspan 33.0–35.0 mm. Body pale yellow. Antenna pale yellow dorsally, orange ventrally. Male antenna with ventral golden cilia nearly as long as the diameter of flagellomere. Labial palpus upturned, pale yellow with a brown patch near base. Maxillary palpus small, pale yellow. Patagium pale yellow, with large fuscous spots. Tegula pale yellow and fuscous. Forewing pale yellow, with fuscous lines, spots and patches; a spot at inner edge near basal line; a large elliptical patch between basal line and antemedial line; orbicular stigma distinct; discoidal stigma reniform, its posterior part overlapping postmedial line partly; a large patch outspread between discoidal stigma and inner margin; postmedial line oblique inward from costa, excurved vertically from M_1 to CuA_2 , with a distinct punctiform pattern between M_2 and CuA_2 , then sharply incurved along CuA_2 as an indistinct line before curving towards the dorsum; a large patch between anterior half of postmedial line and marginal line, marginal line consisting of a line of spots. Hindwing pale yellow; discoidal stigma brown; a brown slant stripe below end of discal cell and a pale brown fuzzy band beyond it; postmedial line oblique inward from costa, excurved vertically from M_1 to CuA_2 , with a distinct punctiform pattern between M_2 and CuA_2 , then sharply



Figures 5–8. *Patania iopasalis* (Walker). **5** Adult **6** Head **7** Gnathos **8** Cornuti. Slide number XLJ14083.

incurved along CuA_2 as an indistinct line before curving towards the dorsum; marginal line fuscous; brown blot near apical angle. Fringes of forewing and hindwing white at base, pale brown terminally. Fore coxa yellow, with two big black spots; femur yellow except for black distal end; tibia pale yellow with distal half black, tibial comb orange. Mid femur white, with a black spot near centre; tibia pale yellow, black distally, a black spot near the base. Hind femur white, with a brown spot near the distal end; tibia white, somewhat pale brown near the base. Abdomen orange dorsally; anterior edge of 2nd segment with two lateral black spots flanking the centre, 7th segment with a big black spot separated slightly at the center; pale yellow ventrally.

Male genitalia. (Figs 3–4, 9). Uncus triangular, blunt on posterior margin. Gnathos thick finger-like, apex circularly widened and dorsally studded with long thin simple setae. Valva ligulate. Fibula near base of valva, flat, triangular, with long dense setae. Transtilla triangular, connected medially, with sparse setae. Saccus oblong, with the anterior apex rounded. Juxta ovate. Phallus cylindrical, with a dorsally protruding sclerotized structure with a slant nailhead-like apex, a long thick needle-like cornutus stretching out from a cluster of spicular cornuti.

Female genitalia. (Fig. 10). Apophysis anterioris about 1.5 times length of apophysis posterioris, triangular extension near base unilaterally. Ductus bursae about



Figures 9–10. Genitalia of *Patania clava* sp. n. **9** Male, Holotype **a** Without phallus **b** Phallus, Holotype **10** Female, paratype.

4 times length of corpus bursae, middle ductus bursae widened, area posterior of widening slightly sclerotized; ductus seminalis originating at posterior end of ductus bursae. Corpus bursae elliptical, without signum. Both ductus bursae and corpus bursae densely studded with tiny granules.

Etymology. The specific name is derived from the Latin *clavus* (nail), in reference to the nail head-like apex of sclerotized structure of the phallus.

Distribution. This species is only known from the Diaoluo Mountain of Hainan Island, China.

Natural history. Unknown except that the moths fly late May and are attracted to light. The habitat in which this species has been collected is located at an altitude of 900 m. The vegetation of the habitat is a blend of shrubs, conifer trees and broad leaved trees.

Remarks. The most similar congener to *P. clava* sp. n. is *Patania iopasalis* (Walker, 1859). However, the wingspan of *P. iopasalis* is smaller with 21.0–30.0 mm (Fig. 5); the labial palpus has no brown patch near the base but a big brown patch near the apex (Fig. 6), the male antenna has ventral cilia about one-fourth the length of the diameter

of flagellomere; in the male genitalia the short broad sheet-like gnathos lacks setae on the apex (Fig. 7), the phallus has an ovate sclerotized structure protruding from apex, and a long thick needle-like cornutus is embedded in a cluster of spicular cornuti (Fig. 8). *Patania clava* sp. n. is also similar to *Patania obfuscalis* (Yamanaka, 1998) in appearance. The latter can, however, be distinguished by its serrated postmedial line of forewing; in the male genitalia by the middle costa slightly inflated and bearing a cluster of setae, the vestigial gnathos, the board-like sclerotized, apically tapering structure protruding from apex, and a short broad thorn-like cornutus near a cluster of spicular cornuti.

The new species is only known from Diaoluo Mountain of Hainan Island at present. *Patania iopasalis* is widely distributed in south Asia, southeast Asia, east Asia, including Hainan Island, China, the Caroline Islands, New Guinea, Australia and Guatemala (Hampson 1896, 1898; Klima 1939; Xu 2015). *Patania obfuscalis* is distributed in Nepal and some areas of China, excluding Hainan Island.

Acknowledgements

We are grateful to Prof. Houhun Li (Nankai University) for lending specimens and to BMNH (the Natural History Museum, London, UK) for providing access to the corresponding author in examining specimens. Cordial thanks are given to Prof. Shuxia Wang (Nankai University) and Dr Dandan Wei (Southwest University) for reviewing the manuscript and providing suggestions and to Richard Mally (University Museum of Bergen, Norway) for carefully revising the manuscript. This research was supported by the National Natural Science Foundation of China (31101663).

References

- Bae YS, Byun BK, Paek MK (2008) Pyralid Moths of Korea (Lepidoptera, Pyraloidea). Korea National Arboretum, Seoul, Korea, 78–181.
- Hampson GF (1896) Moths. The Fauna of British India, including Ceylon and Burma 4: 328.
- Hampson GF (1898) A revision of the moths of the subfamily Pyraustinae and family Pyralidae. Proceedings of the Zoological Society of London 1898(4): 712–725.
- Hübner J (1823) Beiträge zur Sammlung exotischer Schmettlinge [sic]: bestehend in Bekundigung einzelner Fliegmuster neuer oder rarer nichteuropäischer Gattungen. Augsburg, 1–40.
- Inoue H (1980) Some unrecorded and rarely known Pyralidae from Japan. Japan Heterocerists' Journal 106: 90–92.
- Kirti JS, Gill NS (2007) Revival of genus *Patania* Moore and reporting of a new species *menoni* (Pyraustinae: Pyralidae: Lepidoptera). Journal of Entomological Research 31(3): 265–275.
- Klima A (1939) Pyralidae: subfamily: Pyraustinae I. In: Bryk F (Ed.) Lepidopterorum Catalogous 89. Dr. W. Junk, Gravenhage, 186–224.
- Kristensen NP (2003) Skeleton and muscles: adults. Handbook of Zoology 4: 39–132. doi: 10.1515/9783110893724.39

- Leraut P (2005) Contribution to the study of *Pyralis* Linnaeus, *Pleuroptya* Meyrick and *Hariatalodes* Warren (Lepidoptera, Pyraloidea). *Revue Française d'Entomologie* 27(2): 77–94.
- Li HH, Zheng ZM (1996) Methods and techniques of specimens of microlepidoptera. *Journal of Shaanxi Normal University (natural science edition)* 24(3): 63–70.
- Li HH, Ren YD, Zhang DD, Du XC, Li WC, You P (2009) *Insect fauna of Henan: Lepidoptera: Pyraloidea*. Science Press, Beijing, China, 1271 pp.
- Meyrick E (1890) On the classification of the Pyralidina of the European fauna. *Transactions of the Entomological Society of London* 38: 443–444. doi: 10.1111/j.1365-2311.1890.tb02704.x
- Moore F (1888) Descriptions of new Indian lepidopterous insects from the collection of the late Mr. W.S. Atkinson. Heterocera (continued) (Pyralidae, Crambidae, Geometridae, Tortricidae, Tineidae). In: Hewitson WC, Moore F (Eds) *Descriptions of new Indian lepidopterous insects from the collection of the late Mr. W.S. Atkinson*. Asiatic Society of Bengal/ Taylor & Francis, Calcutta / London, 199–299.
- Munroe EG (1983) Pyralidae (except Crambinae). In: Hodges RW et al. (Eds) *Check List of the Lepidoptera of America north of Mexico including Greenland*. E. W. Classey Ltd. and the Wedge Entomological Research Foundation, London, 67–76, 78–85.
- Nuss M, Landry B, Vegliante F, Tränkner A, Mally R, Hayden J, Segerer A, Li H, Schouten R, Solis MA, Trofimova T, De prins J, Speidel W (2003–2015) *Global Information System on Pyraloidea*. <http://www.pyraloidea.org/> [accessed 16 January 2016]
- Rose HS, Singh AP (1989) A new species of genus *Patania* Moore (Lepidoptera, Pyralidae, Pyraustinae) from India. *Association for Advancement of Entomology* 14(1–2): 39–43.
- Shaffer JC, Munroe EG (1989) Type material of two African species of *Herpetogramma* and one of *Pleuroptya* (Lepidoptera: Crambidae: Pyraustinae). *Proceedings of the Entomological Society of Washington* 91(3): 414–420.
- Shibuya J (1928) The systematic study on the Formosan Pyralidae. *Journal of the Faculty of Agriculture, Hokkaido Imperial University* 22: 217–229.
- Shibuya J (1929) On the known and unrecorded species of the Japanese Pyraustinae (Lepid.). *Journal of the Faculty of Agriculture, Hokkaido Imperial University* 25: 182–183.
- Slamka F (2013) *Pyraloidea of Europe, Volume 3*. Bratislava, Slovakia, 7–8.
- Walker F (1859) Pyralides. *List of the Specimens of Lepidopterous Insects in the Collection of the British Museum*, London 18: 652–691.
- Walker F (1866) Supplement 4. *List of the Specimens of Lepidopterous Insects in the Collection of the British Museum*, London 34: 1121–1533.
- Warren W (1896) New species of Pyralidae from Khasia Hills. *Annals and Magazine of Natural History, including Zoology, Botany and Geology* (6) 18: 107–232.
- Xu LJ (2015) *Taxonomic Study on Patania* Moore, 1888 and *Syllepte* Hubner, 1823 from China (Lepidoptera: Pyraloidea: Spilomelinae). Master's thesis, Southwest University.
- Yamanaka H (1995) Pyralidae of Nepal (I). *Tinea* 14: 182–193.
- Yamanaka H (1998) Pyralidae of Nepal (II). *Tinea* 15(suppl. I): 99–115.