

# A new species and a newly recorded subgenus of *Lipotriches* Gerstaecker, 1858 (Hymenoptera, Apoidea, Halictidae, Nomiinae) from China

Dan Zhang<sup>1,2</sup>, Ze-Qing Niu<sup>1</sup>, Alain Pauly<sup>3</sup>, Wa Da<sup>4</sup>, Chao-Dong Zhu<sup>1,2,5</sup>

**1** Key Laboratory of Zoological Systematics and Evolution, Institute of Zoology, Chinese Academy of Sciences, 1 Beichen West Road, Chaoyang District, Beijing, 100101, China **2** College of Biological Sciences, University of Chinese Academy of Sciences, No.19A Yuquan Road, Shijingshan District, Beijing, 10049, China **3** Royal Institute of Natural Sciences of Belgium, Entomology, Rue Vautier-29, B-1000 Brussels, Belgium **4** Tibet Plateau Institute of Biology, Lhasa, 850001, Tibet, China **5** State Key Laboratory of Integrated Pest Management, Institute of Zoology, Chinese Academy of Sciences, 1 Beichen West Road, Chaoyang District, Beijing, 100101, China

Corresponding author: Chao-Dong Zhu ([zhucd@ioz.ac.cn](mailto:zhucd@ioz.ac.cn))

Academic editor: Andreas Köhler | Received 28 September 2021 | Accepted 2 March 2022 | Published 24 March 2022

<http://zoobank.org/A0C1571F-3D9F-4F47-8DC9-0083FBE7FAD1>

**Citation:** Zhang D, Niu Z-Q, Pauly A, Da W, Zhu C-D (2022) A new species and a newly recorded subgenus of *Lipotriches* Gerstaecker, 1858 (Hymenoptera, Apoidea, Halictidae, Nomiinae) from China. ZooKeys 1090: 103–111. <https://doi.org/10.3897/zookeys.1090.75872>

## Abstract

Two Chinese species of the genus *Lipotriches* Gerstaecker, 1858 are treated in this paper. *Lipotriches* (*Lipotriches*) *guihongyi* Zhang & Niu, **sp. nov.** is recognized as a new species and *Lipotriches* (*Maynenomia*) *nanensis* (Cockerell, 1929) is a new species and subgenus record for China. The number of Chinese species of the subfamily Nomiinae and genus *Lipotriches* are updated to 47 and 15, respectively.

## Keywords

Anthophila, Apiformes, bee pollinator, description, morphology, taxonomy

## Introduction

The genus *Lipotriches* (subfamily Nomiinae) was described by Gerstaecker (1858) with *Lipotriches abdominalis* as the type species, including more than 340 species (Ascher and Pickering 2021). *Lipotriches* has a high diversity of species occurring throughout

the Old World (Africa, Asia and Australia). Bee species of this genus are important pollinators for plants, especially for grasses (Pauly 2014a). Over the last few decades, various groups and species in this genus have been revised (Pauly 1984a, 2009, 2014a, b; Astafurova and Pesenko 2005; Michener 2007; Huang 2008; Niu et al. 2018). However, much more taxonomic work is needed for this group, especially in Asia.

In general, the body of *Lipotriches* species is relatively slender compared to most other nomiine species (Pauly 1990; Michener 2007). Morphological identification of the subgenus within *Lipotriches* strongly relies on adult characters: especially the pronotum with continuous, or medially, or laterally notched transverse carina in both sexes, and the basitibial plate of females with a carina only along the posterior margin (Michener 2007; Huang 2008). Pauly (1990) separated *Lipotriches* into several genera, such as *Austronomia* Michener, 1965, *Afronomia* Pauly, 1990, *Macronomia* Cockerell, 1917, and *Trinomia* Pauly, 1980. Later, Michener (2007) revised all the above groups as subgenera of the genus *Lipotriches*, dividing *Lipotriches* into nine subgenera in total.

Pauly (1984b) described *Maynenomia* as a genus with *Nomia maynei* Cockerell, 1937 as the type species. Subsequently, he described nine new species and transferred five *Nomia* species to this group, giving a total of 15 species of *Maynenomia* in Africa and Asia (Pauly 2009). Michener (2007) treated *Maynenomia* as a subgenus within the genus *Lipotriches*, and suspected this subgenus probably to be a synonym of the subgenus *Austronomia*. While Pauly treated *Maynenomia* at genus level, based on its “oval” head shape, we follow Ascher and Pickering (2021) and Michener (2007) in treating *Maynenomia* as a subgenus within *Lipotriches*.

Herein, we reported two Chinese species of the genus *Lipotriches*, including one newly described species and one newly recorded species. To date, the Chinese species of subfamily Nomiinae is increased to 47 in total (Niu et al. 2018; Zhang et al. 2020).

## Materials and methods

In this study, a total of 50 specimens were examined, all of them were deposited in the Collection of the Institute of Zoology, Chinese Academy of Sciences, Beijing, China (IZCAS). The specimens were examined with Nikon SMZ 1500 stereomicroscope. Photographs were taken with Nikon D7000 digital camera and were stacked with Helicon Focus and Zerene Stacker. Final images were edited for clarity and mounted into plates by Photoshop CS6.

The morphological terminology follows Pesenko (1983) and Michener (2007) in this study. Absolute measurements were taken in millimeters (mm) for body length. The following abbreviations are used: BL, body length which was measured from basal antennal socket to the metasomal apex; HL, head length which represented the widest point of the head in frontal view; T1–5, the first to fifth metasomal terga; S1–8, the first to eighth metasomal sterna; F1–11, the first to eleventh flagellar segments. We measured the punctation density, punctation diameter (d) and the space between them (i), such as  $i = 1d$  or  $i < d$ .

## Taxonomy

### Genus *Lipotriches* Gerstaecker, 1858

#### Subgenus *Lipotriches* Gerstaecker

*Lipotriches* Gerstaecker, 1858: 460. Type species: *Lipotriches abdominalis* Gerstaecker, 1857 = *Sphecodes cribrosa* Spinola, 1843, monobasic.

*Rhopalomelissa* Alfken, 1926: 267. Type species: *Rhopalomelissa xanthogaster* Alfken, 1926, by designation of Sandhouse (1943: 596).

*Nomia* (*Epinomia*) Alfken, 1939: 113, not Ashmead, 1899. Type species: *Nomia andreoides* Vachal, 1903 = *Nomia andrei* Vachal, 1897, by original designation.

*Alfkenomia* Hirashima, 1956: 33, replacement for *Epinomia* Alfken, 1939. Type species: *Nomia andreoides* Vachal, 1903 = *Nomia andrei* Vachal, 1897, autobasic.

*Rhopalomelissa* (*Lepidorhopalomelissa*) Wu, 1985: 58. Type species: *Nomia burmica* Cockerell, 1920, by original designation.

*Rhopalomelissa* (*Trichorhopalomelissa*) Wu, 1985: 58. Type species: *Rhopalomelissa hainanensis* Wu, 1985, by original designation.

*Rhopalomelissa* (*Tropirhopalomelissa*) Wu, 1985: 58. Type species: *Rhopalomelissa nigra* Wu, 1985, by original designation.

**Diagnosis.** Small-sized, body length about 5–12 mm; metasoma slender, with petiolate, T1 longer than broad in male (most species); pronotum with continuous or medially or laterally transverse carina or lamella anterior to the scutum; metasoma partly or wholly red for some species (i.e., *Lipotriches floralis*, *Lipotriches esakii* and *Lipotriches mediorufa*).

#### *Lipotriches* (*Lipotriches*) *guihong* Zhang & Niu, sp. nov.


<http://zoobank.org/F11C65CF-98CB-4016-8AE6-4D101D75D5E8>

Figs 1, 2

**Type material. Holotype:** CHINA: 1♂, Xizang, Jilong County, Jilong Town, Jipu Village, 28°37'N, 85°32'E, 2744 m, 9 Aug. 2019, Dan Zhang, Qing-Tao Wu leg.

**Paratypes:** 5♀21♂, Jilong County, Jilong Town, Jipu Village, 28°37'N, 85°32'E, 2744m, 7–9 Aug. 2019, Dan Zhang, Qing-Tao Wu leg.; 16♀1♂, Xizang, Jilong County, Jilong Town, Xinjiang Village, 28°22'N, 85°21'E, 2727m, 6 Aug. 2019, Dan Zhang, Qing-Tao Wu leg.

**Diagnosis.** Males of *L. guihong* sp. nov. differ from other species of subgenus *Lipotriches* by the following combination of features: mesoscutum, metapostnotum and center disc of metasomal terga with dense and large punctures (Fig. 1c, d), S4 with dense short white hairs (Fig. 1f); S5 with a pair of circular protruding thickness on the disc, a pair of triangular feathery bristles connected on the apical margin (Fig. 1g);



**Figure 1.** *Lipotriches (Lipotriches) guihongi* Zhang & Niu sp. nov., male **a** habitus in lateral view **b** head in frontal view **c** mesoscutum in dorsal view **d** propodeum in posterior view **e** metasoma in dorsal view **f** metasoma in lateral view **g** S5 in dorsal view **h** genitalia in dorsal view **i** genitalia in lateral view. Scale bars: 1 mm (**a–f**); 0.5mm (**g–i**).

*L. guihongi* is most similar to *Lipotriches yasumatsui* Hirashima, 1961 and *Lipotriches ceratina* (Smith, 1857), however, the male of the new species can be distinguished from the two latter by a pair of large, dense, erect and brownish longitudinal tufted hairs on S5. In addition, S5 structure of *L. guihongi* sp. nov. is similar to *L. acanthospermi* Pauly, 2014b. We have found that the both species have dense tomentum on S4 and a pair triangular feathery bristles on the apical margin of S5, while the latter lacks a pair of circular protruding thickness on the disc of S5, and has only been found in Africa.

**Description. Male** (measurements are only from the holotype). BL = 8 mm (Fig. 1a), body black. **Head.** HL: HW = 0.89, head broader than long (Fig. 1b); clypeus broader than long (Fig. 1b); paraocular area with dense large punctures ( $i = 0.1-0.2d$ ; Fig. 1b); vertex with sparser large punctures ( $i = 0.5-1d$ ; Fig. 1b); vertex behind shiny; posterior margin of vertex rounded (Fig. 1a, b); mandible blackish-brown; frons with dense and minute punctures, medial frontal line smooth (Fig. 1b); antenna reaching the posterior margin of T1; F1-11 almost equal in length, nearly 2.3 times as long as broad (Fig. 1a, b); ocelli normal (Fig. 1b). **Mesosoma.** Mesoscutum, scutellum, and metanotum dull, without reflections (Fig. 1c). Mesoscutum with large and mostly confluent punctures ( $i = 0.2-0.3d$ ; Fig. 1c); metanotum normal, without process, with large punctures which are sparser than punctures on the mesoscutum (Fig. 1c); posterior surface of propodeum with more larger and denser ( $i = 0.1-0.2d$ ) punctures than those on the mesosoma (Fig. 1d); metapostnotum shiny, with broad longitudinal wrinkles (Fig. 1d); fore wing with three submarginal cells, the 1<sup>st</sup> and 3<sup>rd</sup> submarginal cell nearly equal in length, almost two times as long as 2<sup>nd</sup> submarginal cells; tegula oval, yellow-brown, not enlarged (Fig. 1c); femur and tibia black (Fig. 1a); basitarsus, mediotarsus yellow-brown with tarsal claw (Fig. 1f); hind femur and tibia normal, not enlarged (Fig. 1c). **Metasoma.** Surface of metasomal terga shiny, center of disc with dense punctures ( $i = 0.2-0.3d$ ), apical of disc with sparser punctures than on the center ( $i = 0.5-1d$ ); apical margin of T1-5 transparent (Fig. 1e, f); S5 with a pair of circular protruding thickness on the disc, a pair triangular feathery bristles on the apical margin connected (Fig. 1g); gonostylus as shown in Fig. 1h (in dorsal view) and Fig. 1i (in lateral view). **Pubescence.** Clypeus, supraclypeal area and frons with white setae (Fig. 1a, b); scutellum with sparse long yellowish hairs (Fig. 1c); metanotum with dense pale tomentum (Fig. 1c); upper lateral surface of propodeum with sparse long white hairs (Fig. 1d); legs with sparse white hairs (Fig. 1a, d-f); apical area of metanotum with sparse long hairs (Fig. 1c); apical margin of T1-4 with white hair band, interrupted in middle (Fig. 1e); S4 covered with dense pale tomentum (Fig. 1f).

**Female.** BL = 10-12 mm. Similar to male, except the following: metapostnotum with punctures sparser ( $i = 0.4-0.5d$ ) and smaller than male's (Fig. 2d); head, mesosoma and metasoma with sparse yellowish hairs (Fig. 2b); T1-2 with densely and minutely punctures ( $i = 0.2-0.3d$ ; Fig. 2d).

**Etymology.** The name "*guihong*" is dedicated to Prof. Hong Gui. He is a famous entomologist in China, who advised and encouraged Chao-Dong Zhu to continue his study on insects.

**Floral association.** Unknown.

**Distribution.** China (Xizang).



**Figure 2.** *Lipotriches (Lipotriches) guihongi* Zhang & Niu sp. nov., female **a** habitus in lateral view **b** head in frontal view **c** mesoscutum in dorsal view **d** propodeum in posterior view. Scale bars: 1 mm.

### Subgenus *Maynenomia* Pauly, 1984

*Maynenomia* Pauly, 1984b: 698. Type species: *Nomia maynei* Cockerell, 1937 = *Nomia testacea* Friese, 1914, by original designation.

**Diagnosis.** Glossa slender; apical of basitibial plate opened, delimited on both sides in female; scape short, not reaching median ocellus in females.

**Distribution.** China (Yunnan); India (Uttarakhand); Indonesia; Laos; Malawi; Myanmar; South Africa; Tanzania; Thailand.

### *Lipotriches (Maynenomia) nanensis* (Cockerell, 1929)


Fig. 3

*Nomia nanensis* Cockerell, 1929:133, ♀. Holotype, ♀, Thailande, Nan, USNM.

*Maynenomia nanensis* (Cockerell, 1929): Pauly 2009.

**Material examined.** CHINA: 2♀, Yunnan, Xishuangbanna, Naban River, 22°04'N, 100°22'E, 1303 m, 16 Jun. 2014, Xiu-wei Liu leg..

**Diagnosis.** T1–2 mainly reddish, and T2 with large black spot at each side basally (Fig. 3e, f); legs reddish except fore femur black (Fig. 3c); mandibles subapically red,



**Figure 3.** *Lipotriches (Maynenomia) nanensis*, female **a** head in frontal view **b** head in dorsal view **c** habitus in lateral view, showing the glossa **d** mesoscutum in dorsal view **e** metasoma in dorsal view **f** habitus in dorsal view. Scale bars: 1mm.

and apically black (Fig. 3c, f); hind tibia with basitibial plate (Fig. 3f); mesoscutum, scutellum and propodeum dull (Fig. 3d); mesoscutum with extremely dense and minute punctures (Fig. 3d); posterior surface of propodeum almost without punctures (Fig. 3d, e); metanotum with dense pale ochreous tomentum (Fig. 3d, e); surface of metasomal terga highly polished, with extremely sparse and minute punctures (Fig. 3e, f).

**Distribution.** China (Yunnan); India (Maharashtra); Laos; Myanmar; Thailand.

**Remark.** This species was recorded from China for the first time in this study, increasing the number of Chinese species of *Lipotriches* and Nomiinae to 15 and 47, respectively.

## Acknowledgements

For the foremost, we sincerely express our thanks to Mr Qing-Tao Wu and Dr Xiu-Wei Liu who provided important assistance for collection. We thank Dr Michael Orr for English language editing. We also thank Mr Ye-Jie Lin, who helped to take pictures of S5. In addition, we would like to show our special thanks to Lasa Office of the Second Tibetan Plateau Scientific Expedition and Research, who provided important help with field work in Xizang.

This study was mainly supported by the Second Tibetan Plateau Scientific Expedition and Research (STEP) program (2019QZKK05010605). WD was supported by the Xizang Science and Technology Projects (XZ202101ZD0003G). CDZ was supported by the National Natural Science Foundation for Distinguished Young Scholars (31625024) and the grant (Y229YX5105) from the Key Laboratory of Zoological Systematics and Evolution, Chinese Academy of Sciences.

## References

- Alfken JD (1926) Fauna Buruana, Apidae. *Treubia* 7(3): 259–275.
- Alfken JD (1939) Hymenoptera, Apidae, in *Missione Biologica nel Paese dei Borana*, Vol. 3, Raccolte Zoologiche, Parte 2. Reale Accademia d'Italia, Roma.
- Ascher JS, Pickering J (2021) Discover Life bee species guide and world checklist. Available from (Hymenoptera: Apoidea: Anthophila). [http://www.discoverlife.org/mp/20q?guide=Apoidea\\_species](http://www.discoverlife.org/mp/20q?guide=Apoidea_species) [accessed September 2021]
- Astafurova YV, Pesenko YA (2005) Contributions to the halictid fauna of the Eastern Palaearctic Region: subfamily Nomiinae (Hymenoptera: Halictidae). *Far Eastern Entomologist [Dal'nevostochnyi Entomolog]* 154: 1–16. <https://doi.org/10.31610/zsr/2006.15.1.133>
- Cockerell TDA (1929) Descriptions and Records of Bees – CXVII. *Annals & Magazine of Natural History* 4(10): 132–141. <https://doi.org/10.1080/00222932908673035>
- Gerstaecker A (1858) [Bees and wasps collected in Mozambique] *Monatsberichte, Akademie der Wissenschaften*, Berlin 29 October 1857, 460–464.
- Hirashima Y (1956) Some bees of the genus *Nomia* Latreille from Formosa. *Insecta atsumurana* 20(1–2): 29–33.
- Hirashima Y (1961) Monographic study of the subfamily Nomiinae of Japan (Hymenoptera, Apoidea). *Acta hymenopterologica* 1(3): 241–303.
- Huang HR (2008) A taxonomic study of the subfamily Nomiinae (Hymenoptera, Apoidea) from China. Beijing Forestry University Master Dissertation, Beijing.
- Michener CD (2007) *The Bees of the World*. The John Hopkins University Press, Baltimore and London, [xvi +] 953 pp.
- Niu ZQ, Yuan F, Zhu CD (2018) Apoidea (Apidae, Melittidae, Halictidae). In: *Species Catalogue of China*. Vol. II. Insect (IV). Science press, Beijing, 190 pp.
- Pauly A (1984a) Mission entomologique en Afrique occidentale (1979–1980): Renseignements-séco-biologiques concernant les Hyménoptères. *Notes Fauniques de Gembloux* 11: 1–43.

- Pauly A (1984b) Contribution à l'étude des genres afrotropicaux de Nomiinae. Revue de Zoologie Africaine 98: 693–702.
- Pauly A (1990) Classification des Nomiinae africains. Musée Royal de l'Afrique Centrale Tervuren, Belgique. Annales Sciences Zoologiques 261: 1–206.
- Pauly A (2009) Classification des Nomiinae de la Région Orientale, de Nouvelle Guinée et des îles de l'Océan Pacifique (Hymenoptera: Apoidea: Halictidae). Bulletin de l'Institut Royal des Sciences Naturelles de Belgique 79: 151–229.
- Pauly A (2014a) Les Abeilles des Graminées ou *Lipotriches* Gerstaecker, 1858, sensu stricto (Hymenoptera: Apoidea: Halictidae: Nomiinae) de l'Afrique subsaharienne. Belgian Journal of Entomology 20: 1–393.
- Pauly A (2014b) Les Abeilles des Graminées ou *Lipotriches* Gerstaecker, 1858, sensu stricto (Hymenoptera Apoidea Halictidae Nomiinae) de la Région Orientale. Belgian Journal of Entomology 21: 1–94.
- Pesenko YuA (1983) Tribe Nomioidini (in the palearctic fauna). In: Fauna of the USSR. New Series No. 129. Hymenopterous Insects. Vol. XVII. No. 1. Halictid bees (Halictidae). Subfamily Halictinae. Tribe Nomioidini (in the Palaeartic Region). Nauka, Leningrad, 1–198. [In Russian]
- Sandhouse GA (1943) The type species of the genera and subgenera of bees. Proceedings of the United States National Museum 92: 519–619. <https://doi.org/10.5479/si.00963801.3156.519>
- Smith F (1857) Catalogue of the hymenopterous insects collected at Sarawak, Borneo; Mount Ophir, Malacca; and at Singapore, by A.R. Wallace. Journal of the Proceedings of the Linnean Society of London. Zoology (Jena, Germany) 2: 42–88. <https://doi.org/10.1111/j.1096-3642.1857.tb01759.x>
- Wu YR (1985) A study on the genus *Rhopalomelissa* of China with descriptions of new subgenus and new species (Apoidea, Halictidae). Zoological Research 6(1): 57–68. [In Chinese]
- Zhang D, Niu ZQ, Orr MC, Ascher JS, Zhu CD (2020) Chinese species of *Nomia* (*Gnathonomia*) Pauly, 2005 (Hymenoptera: Apoidea: Halictidae: Nomiinae). Zootaxa 4768(1): 76–94. <https://doi.org/10.11646/zootaxa.4768.1.5>