

A new species of the genus *Anteon* Jurine (Hymenoptera, Dryinidae) from Laos

Massimo Olmi¹, Zai-fu Xu², Adalgisa Guglielmino³, Stefano Speranza³

1 Tropical Entomology Research Center, Viterbo, Italy **2** Department of Entomology, South China Agricultural University, Guangzhou, Guangdong, P.R. China **3** Department of Agriculture and Forestry Sciences (DAFNE), University of Tuscia, Viterbo, Italy

Corresponding author: Massimo Olmi (olmi@unitus.it)

Academic editor: M. Engel | Received 5 December 2015 | Accepted 4 January 2016 | Published 8 February 2016

<http://zoobank.org/8446A28C-D48A-45CB-AA2E-C2746A341E3A>

Citation: Olmi M, Xu Z, Guglielmino A, Speranza S (2016) A new species of the genus *Anteon* Jurine (Hymenoptera, Dryinidae) from Laos. ZooKeys 561: 31–38. doi: 10.3897/zookeys.561.7417

Abstract

A new species of *Anteon* Jurine, 1807 is described from Laos, Houaphanh Province: *A. holzschuhi* sp. n. Morphologically the new species is similar to *A. semipolatum* Olmi, 2008, but it is distinguished by the sculpture of the face partly reticulate rugose and partly with deep punctures similar to areolae; in *A. semipolatum* the face is completely punctate and unsculptured among punctures. In addition, in the new species the distance from the outer edge of a lateral ocellus to the compound eye (OOL) is about 3.3 times as long as the distance between the inner edges of a lateral ocellus and the median ocellus (OL); in *A. semipolatum* OOL is less than twice as long as OL. Published identification keys to the Oriental species of *Anteon* are modified to include the new species.

Keywords

Taxonomy, *Anteon holzschuhi*, Oriental region, key, Houaphanh Province, Anteoninae

Introduction

Dryinidae (Hymenoptera, Chrysidoidea) are parasitoids of leafhoppers, planthoppers and treehoppers (Hemiptera, Auchenorrhyncha) (Carcupino et al. 1998; Guglielmino and Bückle 2003, 2010; Guglielmino et al. 2006, 2013, 2015; Guglielmino and Virla

1998). *Anteon* Jurine, 1807 is a genus that is present in all zoogeographical regions (Olmi 1984; Xu et al. 2013; Olmi and Virla 2014; Olmi and Xu 2015). In total 423 species have been described from all continents (Olmi and Xu 2015) and the genus was revised at the world level by Olmi (1984, 1991) and more recently in the Oriental, Neotropical and Eastern Palaearctic regions by Xu et al. (2013), Olmi and Virla (2014) and Olmi and Xu (2015) respectively.

The species of *Anteon* inhabiting the Oriental region were studied by Xu et al. (2013). More recently, Guglielmino and Olmi (2013) and Olmi et al. (2015) described further new species respectively from Indonesia (*Anteon seramense* Guglielmino & Olmi) and Thailand (*Anteon huettingeri* Olmi, Xu & Guglielmino). In total, 150 *Anteon* species have been described from the Oriental region (Xu et al. 2013; Guglielmino and Olmi 2013; Olmi et al. 2015).

Anteon species are parasitoids of leafhoppers belonging to the Cicadellidae (Guglielmino et al. 2013). As in almost all dryinids, females of *Anteon* have a chelate pronotus. Chelae are used to capture and restrain the host during oviposition and host-feeding (Olmi 1984, 1994).

In 2015 we examined additional specimens of *Anteon* from Laos and discovered a new species described in this paper.

Material and methods

The descriptions follow the terminology used by Olmi (1984) and Xu et al. (2013). The measurements reported are relative, except for the total length (head to abdominal tip, without antennae), which is expressed in millimetres. The following abbreviations are used in the descriptions: POL is the distance between the inner edges of the two lateral ocelli; OL is the distance between the inner edges of a lateral ocellus and the median ocellus; OOL is the distance from the outer edge of a lateral ocellus to the compound eye; OPL is the distance from the posterior edge of a lateral ocellus to the occipital carina; TL is the distance from the posterior edge of an eye to the occipital carina.

The types of all Oriental species of *Anteon* have been previously examined by the authors.

The type specimen described in this paper is deposited in the collection of the Oberösterreichisches Landesmuseum, Linz, Austria (OLL).

The description of the new species is based on the study of a single specimen. The authors are aware that descriptions of new taxa should normally be based on more individuals. However, Dryinidae are so rare that it is uncommon to collect more than one specimen of each species. In addition, on the basis of the experience and knowledge of the authors, the new species is sufficiently delimited by unique characters to justify its description.

Results

Genus *Anteon* Jurine, 1807

Anteon Jurine, 1807: 302. Type species: *Anteon jurineanum* Latreille, 1809, by subsequent monotypy.

Diagnosis. Female: Fully winged; rarely brachypterous; occipital carina complete; palpal formula 6/3; antenna without rhinaria; forewing with three cells enclosed by pigmented veins (costal, median and submedian); forewing with stigmal vein and pterostigma; distal part of stigmal vein much shorter than proximal part, occasionally slightly shorter, as long as, or longer than proximal part; propodeum usually with transverse keel between dorsal and posterior surface; protarsus chelate; inner side of enlarged claw with proximal prominence bearing one long bristle; tibial spurs 1/1/2. Male: Fully winged; rarely brachypterous; occipital carina complete; vertex of head usually without two oblique keels connecting posterior ocelli to occipital carina; palpal formula 6/3; forewing with three cells enclosed by pigmented veins (costal, median and submedian); forewing with stigmal vein and pterostigma; distal part of stigmal vein much shorter than proximal part, occasionally slightly shorter, as long as, or longer than proximal part; pterostigma less than four times as long as broad; propodeum usually with transverse keel between dorsal and posterior surface; paramere usually without inner branch wrapping penis; tibial spurs 1/1/2.

Anteon holzschuhi Olmi, Xu, Guglielmino & Speranza, sp. n.

<http://zoobank.org/80257611-458A-45B3-B633-292B862C17A3>

Diagnosis. Male with antenna filiform; face partly reticulate rugose and partly sculptured by deep punctures similar to areolae (Fig. 1B); OOL about 3.3 times as long as OL (Fig. 1A); notauli reaching about 0.8 length of scutum (Fig. 1A); posterior surface of propodeum with two complete longitudinal keels and median area unsculptured; distal part of stigmal vein much shorter than proximal part; paramere about as long as penis, without papillae on inner side, without distal inner process (Fig. 2); distivolsella not provided with two lateral processes (Fig. 2).

Description. Male. Fully winged (Fig. 1A). Length 4.5 mm. Head black, except mandible testaceous. Antenna brown-testaceous, except proximal half of segment 1 testaceous. Mesosoma black. Metasoma brown. Legs testaceous, except metacoxa basally brown. Antenna filiform. Antennal segments in following proportions: 17:10:16:15:14:14:14:14:13 (segment 10 missing in holotype). Head (Fig. 1A, B) shiny. Face partly rugose and partly strongly punctate, with deep punctures similar to areolae, unsculptured among punctures. Vertex and temple with deep punctures similar to areolae, unsculptured among punctures. Frontal line complete. Vertex with POL = 7; OL = 3; OOL = 10; OPL = 7; TL = 7; greatest breadth of posterior ocelli shorter




Figure 1. *Anteon holzschuhi* Olmi, Xu, Guglielmino & Speranza, sp. n., holotype: **A** habitus in dorsal view **B** head in frontal view. Scale bar: 2.77 mm (**A**), 2.20 mm (**B**).

than OPL (5:7). Occipital carina complete. Pronotum short and strongly punctate. Scutum and scutellum shiny, punctate, unsculptured among punctures. Notauli incomplete, reaching approximately 0.8 x length of scutum (Fig. 1A). Metanotum shiny, unsculptured. Propodeum with strong transverse keel between dorsal and posterior surface. Dorsal surface of propodeum reticulate rugose. Posterior surface of propodeum with two complete longitudinal keels, median area unsculptured, and lateral areas rugose. Forewing hyaline, without dark transverse bands. Distal part of stigmal vein much shorter than proximal part (7:16). Paramere (Fig. 2) about as long as penis, without distal inner pointed process and papillae, with long and broad dorsal proximal membranous process. Tibial spurs 1/1/2.

Female. Unknown.

Material examined. Holotype: male, Laos, Houaphanh Province, Phou Pan, Ort Ban Saleui environs, 20°13.30'N 103°59.26'E, 1350–1900 m, 6–11.iv.2014, C. Holzschuh and locals leg. (OLL).

Distribution. Laos.

Hosts. Unknown.

Etymology. The species is named after the collector, Mr Carolus Holzschuh (Villach, Austria).

Remarks. The new species is similar to *A. semipolatum* Olmi, 2008, by having the antenna filiform, notauli reaching about 0.8 x length of scutum (Fig. 1A), posterior surface of the propodeum with two complete longitudinal keels and unsculptured median area, distal part of stigmal vein much shorter than proximal part, paramere about as long as penis, without papillae on inner side, without distal inner process (Fig. 2) and distivolsella not provided with two lateral processes (Fig. 2). The main difference between the two species is in the facial sculpture (face partly reticulate rugose and partly sculptured by deep punctures similar to areolae in *A. holzschuhi* (Fig. 1B); face




Figure 2. *Anteon holzschuhi* Olmi, Xu, Guglielmino & Speranza, sp. n., holotype: male genitalia (right half removed). Scale bar = 0.26 mm.

punctate and unsculptured among punctures in *A. semipolatum*). In addition, OOL is about three times as long as OL in *A. holzschuhi*, less than twice in *A. semipolatum*. In the key to the males of Oriental *Anteon* published by Xu et al. (2013), the new species can be included by replacing couplet 43 as follows:

- 43 Head partly or totally reticulate rugose (Fig. 1A, B).....**43'**
- Head completely punctate and unsculptured among punctures **44**
- 43' Paramere with many papillae along inner side (Plate 38H in Xu et al. 2013)....
.....***A. papillum* Xu, He & Olmi**
- Paramere without papillae on inner side (Fig. 2; plate 42F in Xu et al. 2013)..... **43''**
- 43'' Face punctate and unsculptured among punctures; OOL less than twice as long as OL.....***A. semipolatum* Olmi**
- Face partly reticulate rugose and partly sculptured by deep punctures similar to areolae (Fig. 1B); OOL about 3.3 times as long as OL (Fig. 1A)
..... ***A. holzschuhi* Olmi, Xu, Guglielmino & Speranza, sp. n.**

Conclusion

Mita and Okajima (2011), Xu et al. (2013) and Olmi et al. (2015) recorded from Laos 41 species of Dryinidae belonging to the following subfamilies and genera: Aphelopininae: *Aphelopus* Dalman, 1823 (five species); Anteoninae: *Anteon* Jurine, 1807

(nine species), *Deinodryinus* Perkins, 1907 (one species); Bocchinae: *Bocchus* Ashmead, 1893 (four species); Dryininae: *Dryinus* Latreille, 1804 (16 species); Gonatopodinae: *Neodryinus* Perkins, 1905 (four species), *Echthrodolphax* Perkins, 1903 (two species). With the description of the above new species the number of species now known in Laos is 42. No hosts of Laotian Dryinidae are known.

In comparison with the 77 species listed in the Chinese province of Guangdong (total area: 177900 km²) (Xu et al. 2012c), the dryinid fauna of Laos (total area: 237800 km²) is poorly known. Some common genera such as *Gonatopus* Ljungh, 1810 (no species listed in Laos) are clearly understudied. Further evidence of this is the fact that on the small island of Hainan (total area: 33210 km²; 1/7 that of Laos) 56 species of Dryinidae are recorded (Xu et al. 2011a, 2011c). In contrast, in the more northern Chinese provinces of Shaanxi (total area: 205800 km²) and Hunan (total area: 210000 km²) only 36 (Xu et al. 2012a) and 17 (Xu et al. 2011b, 2012b, 2013) dryinid species have been recorded, respectively.

Acknowledgements

Many thanks to Dr. Martin Schwarz (Oberösterreichisches Landesmuseum, Linz, Austria) for sending the specimen of *Anteon* described in the present paper. We are also grateful to all curators of collections who have sent us type material on loan. Many thanks to Dr. Robert C. Copeland (International Centre of Insect Physiology and Ecology (ICIPE), Nairobi, Kenya) for checking the English language of this paper. This paper was supported by the National Basic Research Program of China (No. 2013CB127600).

References

- Ashmead WH (1893) Monograph of the North American Proctotrypidae. Bulletin of the United States National Museum 45: 1–472. doi: 10.5479/si.03629236.45.1
- Carcupino M, Guglielmino A, Mazzini M, Olmi M (1998) Morphology and ultrastructure of the cephalic vesicles in two species of the *Gonatopus* genus: *Gonatopus camelinus* Kieffer and *Gonatopus clavipes* (Thunberg) (Hymenoptera, Dryinidae, Gonatopodinae). Invertebrate Reproduction and Development 34: 177–186. doi: 10.1080/07924259.1998.9652651
- Dalman CR (1823) Analecta entomologica. Typis Lindhianis, Holmiae, Sweden, 104 pp. doi: 10.5962/bhl.title.66069
- Guglielmino A, Bückle C (2003) Description of larval instars of *Neodryinus typhlocybae* (Ashmead, 1893) (Hymenoptera Dryinidae), with remarks on its biology. Mitteilungen aus dem Museum fuer Naturkunde in Berlin - Deutsche Entomologische Zeitschrift 50(1): 143–150. doi: 10.1002/mmnd.20030500114
- Guglielmino A, Bückle C (2010) Description of larval instars of *Mystrophorus formicaeformis* Ruthe (Hymenoptera: Dryinidae). Zootaxa 2602: 57–66.

- Guglielmino A, Bückle C, Moya-Raygoza G (2006) Description of the larval instars of *Gonatopus bartletti* Olmi, 1984 (Hymenoptera: Dryinidae). *Zootaxa* 1226: 51–60.
- Guglielmino A, Olmi M (2013) Description of *Anteon seramense* (Hymenoptera: Dryinidae), a new species from Indonesia. *Florida Entomologist* 96(2): 598–601. doi: 10.1653/024.096.0226
- Guglielmino A, Olmi M, Bückle C (2013) An updated host-parasite catalogue of world Dryinidae (Hymenoptera: Chrysidoidea). *Zootaxa* 3740: 1–113. doi: 10.11646/zootaxa.3740.1.1
- Guglielmino A, Parise G, Bückle C (2015) Description of larval instars of *Dryinus tarraconensis* Marshall, 1868 and *Gonatopus baeticus* (Ceballos, 1927) (Hymenoptera: Chrysidoidea: Dryinidae), parasitoids of the genus *Dictyophara* Germar (Hemiptera: Auchenorrhyncha: Dictyopharidae). *Zootaxa* 4032(1): 42–54. doi: 10.11646/zootaxa.4032.1.2
- Guglielmino A, Virla EG (1998) Postembryonic development of *Gonatopus lunatus* Klug (Hymenoptera: Dryinidae: Gonatopodinae), with remarks on its biology. *Annales de la Société entomologique de France (NS)* 34(3): 321–333.
- Jurine L (1807) Nouvelle méthode de classer les Hyménoptères et les Diptères, 1. Hyménoptères. Paschoud, Genève, Switzerland, 319 pp.
- Latreille PA (1804) Nouvelle dictionnaire d'Histoire naturelle, 24. F. Dufart, Paris, 104 pp.
- Ljungh SJ (1810) *Gonatopus*, novum insectorum genus. *Beiträge zur Naturkunde* 2: 161–163.
- Mita T, Okajima S (2011) *Dryinus* species collected from Laos (Hymenoptera, Dryinidae, Dryininae). *Japanese Journal of Systematic Entomology* 17(2): 153–154.
- Olmi M (1984) A revision of the Dryinidae (Hymenoptera). *Memoirs of the American Entomological Institute* 37: 1–1913.
- Olmi M (1991) Supplement to the revision of the world Dryinidae (Hymenoptera Chrysidoidea). *Frustula Entomologica (NS)* 12(25): 109–395.
- Olmi M (1994) The Dryinidae and Embolemidae (Hymenoptera: Chrysidoidea) of Fennoscandia and Denmark (Fauna Entomologica Scandinavica 30). EJ Brill, Leiden, 100 pp.
- Olmi M, Virla EG (2014) Dryinidae of the Neotropical Region (Hymenoptera: Chrysidoidea). *Zootaxa* 3792(1): 1–534. doi: 10.11646/zootaxa.3792.2.1
- Olmi M, Xu Z (2015) Dryinidae of the Eastern Palaearctic region. *Zootaxa* 3996(1): 1–253. doi: 10.11646/zootaxa.3996.1.1
- Olmi M, Xu Z, Guglielmino A (2015) A new species of the genus *Anteon* Jurine (Hymenoptera, Dryinidae) from Thailand. *ZooKeys* 504: 141–147. doi: 10.3897/zookeys.504.9333
- Perkins RCL (1903) The leafhopper of the sugar cane. Territory of Hawaii, Board of Agriculture and Forest, Division of Entomology, Bulletin 1: 1–38. doi: 10.3897/zookeys.504.9333
- Perkins RCL (1905) Leafhoppers and their natural enemies (Pt. i. Dryinidae). Report of Work of the Experiment Station of the Hawaiian Sugar Planters' Association, Division of Entomology, Bulletin 1(1): 1–69.
- Perkins RCL (1907) Parasites of leaf-hoppers. Report of Work of the Experiment Station of the Hawaiian Sugar Planters' Association, Division of Entomology, Bulletin 4: 5–59.
- Xu Z, Olmi M, Guglielmino A, Chen H (2011a) A new species of Dryinidae (Hymenoptera: Chrysidoidea) from China. *Florida Entomologist* 94(4): 848–852.
- Xu Z, Olmi M, Guglielmino A, Chen H (2011b) Description of *Aphelopus mangshanensis*, a new species of Dryinidae from China. *Bulletin of Insectology* 64(2): 243–246.

- Xu Z, Olmi M, Guglielmino A, Chen H (2011c) Description of *Anteon diaoluoshanense* sp. nov. from China (Hymenoptera: Dryinidae). *Pan-Pacific Entomologist* 87(3): 172–176. doi: 10.3956/2011-31.1
- Xu Z, Olmi M, Guglielmino A, Chen H (2012a) Checklist of Dryinidae (Hymenoptera: Chrysidoidea) from Shaanxi Province, China, with descriptions of two new species. *Zootaxa* 3164: 1–16.
- Xu Z, Olmi M, Guglielmino A, Chen H (2012b) Descriptions of two new species of *Anteon* (Hymenoptera: Dryinidae) from China. *Entomological News* 121(5): 409–415.
- Xu Z, Olmi M, Guglielmino A, Chen H (2012c) Checklist of Dryinidae (Hymenoptera) from Guangdong Province, China, with descriptions of two new species. *Zootaxa* 3231: 1–28.
- Xu Z, Olmi M, He J (2013) Dryinidae of the Oriental region (Hymenoptera: Chrysidoidea). *Zootaxa* 3614: 1–460. doi: 10.11646/zootaxa.3900.1.1