

Taxonomic recovery of the ant cricket *Myrmecophilus albicinctus* from *M. americanus* (Orthoptera, Myrmecophilidae)

Takashi Komatsu¹, Munetoshi Maruyama¹

¹ *Kyushu University, Hakozaki 6-10-1, Higashi-ku, Fukuoka 812-8581 Fukuoka, Japan*

Corresponding author: *Takashi Komatsu* (corocoro1232000@yahoo.co.jp)

Academic editor: *F. Montealegre-Z* | Received 8 January 2016 | Accepted 12 April 2016 | Published 16 May 2016

<http://zoobank.org/9956EB10-A4CE-4933-A236-A34D809645E8>

Citation: Komatsu T, Maruyama M (2016) Taxonomic recovery of the ant cricket *Myrmecophilus albicinctus* from *M. americanus* (Orthoptera: Myrmecophilidae). ZooKeys 589: 97–106. doi: 10.3897/zookeys.589.7739

Abstract

Myrmecophilus americanus and *M. albicinctus* are typical myrmecophilous insects living inside ant nests. These species are ecologically important due to the obligate association with tramp ant species, including harmful invasive ant species. However, the taxonomy of these “white-banded ant crickets” is quite confused owing to a scarcity of useful external morphological characteristics. Recently, *M. albicinctus* was synonymized with *M. americanus* regardless of the apparent host use difference. To clarify taxonomical relationship between *M. albicinctus* and *M. americanus*, we reexamined morphological characteristics of both species mainly in the viewpoint of anatomy. Observation of genitalia parts, together with a few external body parts, revealed that *M. albicinctus* showed different tendency from them of *M. americanus*. Therefore, we recover *M. albicinctus* as a distinct species on the basis of the morphology.

Keywords

Formicidae, host specificity, myrmecophily, symbiont, synonymy

Introduction

Myrmecophilus (*Myrmophilina*) *americanus* Saussure, 1877 (Orthoptera: Myrmecophilidae) (Figs 1a, 2a) is a typical example of an ant guest that lives inside ant nests. This species, similar to its congeners, eats food found inside the ant nest, either by itself or via mouth-to-mouth feeding by the ants (Wetterer and Hugel 2008). Its body color is totally black except for a single white band on the mesonotum. *Myrmecophilus americanus* was first described on the basis of a single female specimen collected in Colombia (Saussure 1877). The species is currently known to be distributed across tropical Asia, including on small islands, northern Africa, and the Neotropics (Wetterer and Hugel 2008). Its host-ant-species specificity is quite high; it has been collected exclusively (Wetterer and Hugel 2008) from nests of the longhorn crazy ant, *Paratrechina longicornis* (Latreille 1802). Because of its broad distribution, however, specimens of *M. americanus* from several localities have been given different species names. For example, Wasmann (1905) described *M. prenolepidis* from India, and Gorochov (1994) described *M. (Eumyrmecophilus) microscopicus* from Seychelles, but both these species have since been synonymized with *M. americanus*, the former by Schimmer (1909) and the latter by Hugel (2006). On the other hands, Ebner (1956) described *M. robustus* from Egypt though it has been also synonymized with *M. americanus* by Chopard (1968).

Myrmecophilus albicinctus (Chopard 1924) (Figs 1b, 2b) was first described on the basis of four females (the holotype and three paratypes) collected from India, but Ingrisch (2010) considered this species to be indistinguishable from *M. americanus* due




Figure 1. Specimens of the two *Myrmecophilus* species in dorsal view: *M. americanus* (a) and *M. albicinctus* (b).




Figure 2. Specimens (same as those shown in Fig. 1) of the two *Myrmecophilus* species in lateral view: *M. americanus* (a) and *M. albicinctus* (b).

Table 1. Past literatures including host ant record of *Myrmecophilus americanus* and *M. albicinctus*. For records of *M. americanus*, Wetterer and Hugel (2008) have written up in detail.

Recorded species	Host ant species	Author
<i>Myrmecophilus prenolepidis</i>	<i>Prenolepis</i> (= <i>Paratrechina</i>) <i>longicornis</i>	Wasmann (1905)
<i>Myrmecophilus albicinctus</i> (= <i>albicinctus</i>)	<i>Camponotus mitis</i>	Chopard (1924)
<i>Myrmecophilus robustus</i>	<i>Camponotus</i> sp.	Ebner (1956)
<i>Myrmecophilus albicinctus</i>	<i>Anoplolepis gracilipes</i>	Ichikawa et al. 2000
<i>Myrmecophilus microscopicus</i>	<i>Paratrechina longicornis</i>	Hugel and Blard (2005)
<i>Myrmecophilus albicinctus</i>	<i>Anoplolepis gracilipes</i> , <i>Pheidole</i> spp. (the latter is quite rare case)	Maruyama (2006)
<i>Myrmecophilus americanus</i>	<i>Paratrechina longicornis</i> , <i>Camponotus</i> sp. (the latter is only single record)	Wetterer and Hugel (2008)
<i>Myrmecophilus albicinctus</i>	<i>Anoplolepis gracilipes</i>	Komatsu et al. (2009)
<i>Myrmecophilus albicinctus</i>	<i>Anoplolepis gracilipes</i>	Murai and Ito (2011)
<i>Myrmecophilus americanus</i>	<i>Paratrechina longicornis</i>	Wetterer and Hugel 2014

to the similarity of their morphological characteristics. In contrast to *M. americanus*, however, *M. albicinctus* is known from only tropical Asia, including small islands (Maruyama 2006, Murai and Ito 2011). Moreover, recent studies have indicated that it is exclusively found in nests of the yellow crazy ant, *Anoplolepis gracilipes* (Maruyama 2006, Komatsu et al. 2009, Murai and Ito 2011), although the holotype specimen was collected from a *Camponotus mitis* nest (Chopard 1924). Laboratory experiments have shown that, from the perspective of behavioral ecology, *M. albicinctus* is closely dependent on *A. gracilipes* (Komatsu et al. 2009). With regard to the taxonomy of this species, Ichikawa (2001), independently of Hugel and Blard (2005) and Hugel (2006), synonymized *M. microscopicus* with *M. albicinctus*. Thus, the taxonomic status of *M. americanus* and *M. albicinctus* is quite confused (Ingrisch 2010). The host ant species of both *M. americanus* and *M. albicinctus* are well-known tramp ants, and *A. gracilipes* in particular is known to be a highly destructive invasive species. From the viewpoint of pest control, therefore, the taxonomy of parasites and myrmecophilous insects associated with this invasive ant species is of fundamental interest.

Recently, Ingrisch (2010) synonymized *M. albicinctus* with *M. americanus* because no morphological characteristic except body size was found to clearly distinguish them. However, we have previously suggested, following Wetterer and Hugel (2008), that each of these two species depends strictly on a different host ant species, and, moreover, we showed by a preliminary molecular phylogenetic analysis that they can be genetically differentiated (Komatsu et al. 2009, Komatsu et al. unpublished). In addition, we have found clear morphological differences between these two *Myrmecophilus* species.

For recovery of "*Myrmecophilus albicinctus*", there is problem of validity to use the name toward the species. As above mentioned, *M. albicinctus* have once synonymized with *M. americanus*. In addition, there is an older synonym of *M. americanus*; that is, *M. prenolepidis*. Under normal circumstances, it should be used the name of *M. prenolepidis* toward the recovered species. However, *M. prenolepidis* was described on the basis of specimens collected from nest of *Prenolepis longicornis* (Roger, 1863) that is synonymized

as *Paratrechina longicornis* (Emery 1925). The host specificity of *M. albicinctus* toward *A. gracilipes* is strong in principle so it is unlikely that it is collected from nests of *P. longicornis*. Given this, the specimens that formerly regards as *M. prenolepidis* can be regarded as not *M. albicinctus* but *M. americanus* which we call in present paper. Therefore, we apply the name of *M. albicinctus* for the recovered species. A series of old and recent host ant species records for *M. americanus* and *M. albicinctus* are listed in Table 1.

Methods

Sampling

Field sampling of *Myrmecophilus americanus* and *M. albicinctus* in the Ryukyu Islands and in southeast Asia was conducted from 2005 to 2015. Ant crickets were collected from nests of *Anoplolepis gracilipes* and *Paratrechina longicornis* by locating nest entrances, turning over stones, or breaking up decayed logs and stumps. Whenever ant crickets were found, as many as possible were collected and preserved in absolute alcohol.

Examination of samples

One of us (TK) examined specimens that he collected or were collected by colleagues. In addition, he visited the Muséum national d'Histoire naturelle (MNHN) to examine both type specimens (*M. americanus* and *M. albicinctus*).

The collected ethanol-preserved specimens were used for morphological observation. Specimens were dissected to observe their genitalia (abdominal terminalia). Each specimen was softened before dissection by warming (60 °C for 30–60 min) it in a small ceramic bowl (2.5 cm in diameter) with a small amount of water. Then, the specimen was dissected in water at high magnification under a stereomicroscope (Olympus SZ-40, ×6.7–80). The abdominal apex was removed from each specimen and dissected. Body parts were soaked in a warm 5–8% solution of potassium chloride (60 °C, 20–60 min), cleaned in 30% ethanol (5 min), and dehydrated in 99% ethanol (5 min). The dehydrated materials were mounted in Euparal (Chroma-Gesellschaft) on glass slides for detailed observation.

Results

Taxonomy

Myrmecophilus (Myrmophilina) americanus Saussure, 1877

Material examined. 3♂ and 1♀, collected from 50 Ngamwongwan Rd. ChatuChak Bangkok, Thailand, 6-X-2007, Komatsu T.; 1♀, Plot 256, Tingkat Perusahaan 5, Ka-




Figure 3. Type specimens: *M. americanus* (a) and *M. albicinctus* (b).

wasan Perindustrian Perai 2, Perai, Penang, Malaysia, 28-I-2011, Sumino T.; 1♀, Andalas University, Jl. Limau Manis, Kecamatan Pauh, Padang, Sumatera Barat 25163, Indonesia, 27-XI-2013, Komatsu T.; 1♂, Lembaga Ilmu Pengetahuan Indonesia, JL. Raya Jakarta Bogor km 46, Cibinong 16911, Indonesia, 20 VI 2013, Komatsu T.

Type material. Syntype 1♀: Barkuda Id., Chilka Lake, Ganjam dist., Madras Pres. 4-19-1919. F. H. Gravery, Zool. Surv. Ind. (MNHN) (Fig. 3a)

Diagnosis. Hind tarsus is relatively short (less than 1 mm, Fig. 4a); male phallic complex with pseudepiphallic ancora short and roughly rounded with no dorsal branch. Ventral appendage of pseudepiphallic ancora somewhat predominant with both ends roughly square (Fig. 4b); male tenth abdominal tergite bituberculate, with scarce hair but without long strong spines (Fig. 4c); female ovipositor notably short and spoon-shaped in lateral view. Apical valves on both dorsal and ventral margins rounded, more than in other *Myrmecophilus* species (both *M. americanus* and *M. albicinctus* have rounded valves, with those of latter being more rounded) (Fig. 4d, e).

***Myrmecophilus albicinctus* Chopard, 1924, sp. rev.**

Material examined. 1♀, collected from Koshidake, Iheya-jima, Okinawa, Japan, 5-IV-1996, Inada S.; 2♂, Gusukube-sunagawa, Miyakojima-shi, Miyako-jima, Okinawa, Japan, 8-VI-1996, Inada S.; 1♂ and 1♀, collected from Urasoe, Okinawa-jima, Okinawa, Japan, 24-VII-2007, Komatsu T.; 3♂ and 1♀, Yona, Kunigami-son, Okinawa-jima, Okinawa, Japan, 6-VIII-2007, Komatsu T.; 2♀, Field Studies Centre of the University of




Figure 4. *Myrmecophilus americanus*. Hind tibia and tarsus, male inferior view (a); male pseudepiphallallic ancora (b); male abdominal apex showing tubercles of the last abdominal tergite (c); female ovipositor, lateral view, dorsal margin (d); female ovipositor, lateral view, ventral margin (e).

Malaya, Ulu Gombak, Selangor, Malaysia, 25-X-2012, Komatsu T.; 3♀, Bogor Botanical Gardens, Jalan Ir. Haji Juanda No.13, 16122, Indonesia, 22-XI-2013, Komatsu M.; 1♂, Andalas University, Jl. Limau Manis, Kecamatan Pauh, Padang, Sumatera Barat 25163, Indonesia, 1-XII-2013, Komatsu M.; 1♂, 16 km Point, Kaeng Krachan National Park, Phetchaburi, Thailand, 28-VI-2014, Komatsu T.; 4♂ and 2♀, Jalan Universiti, 50603 Kuala Lumpur, Wilayah Persekutuan Kuala Lumpur, Malaysia, 28-XI-2005, Komatsu T.; 2♂ and 1♀, Daruma-yama, Kume-jima, Okinawa, Japan, 9-XII-2014, Komatsu T.

Type material. Paratype 2♂2♀: Pattambi, Molabas Dist., F. H. Gravely V. 30 and *Anoplolepis longipes*. (Fig. 3b).




Figure 5. *M. albicinctus*. Hind tibia and tarsus, male inferior view (a); male pseudepiphallalic ancora (b); male abdominal apex showing tubercles of the last abdominal tergite (c); female ovipositor, side view, dorsal margin (d); female ovipositor, side view, ventral margin (e).

Diagnosis. Hind tarsus is relatively long (more than 1 mm, Fig. 5a); male phallic complex with pseudepiphallalic ancora straightly elongate with no dorsal branch. Ventral appendage of pseudepiphallalic ancora considerably reduced with both ends angular (Fig. 5b); male tenth abdominal tergite bituberculate, with rich hair and long strong spines (Fig. 5c); female ovipositor closely resembles that of *M. americanus*, except the apical valve on the dorsal margin is more rounded in lateral view (Fig. 5d, e).

Remark. This species can be clearly discriminated from *M. americanus* on the basis of the described diagnostic characteristics. Therefore, we recognize *M. albicinctus* as a distinct species.

Discussion

With regard to the taxonomy of *Myrmecophilus* ant crickets, Ingrisch (2010) has stated that better characteristics than host specificity are needed to differentiate species. In fact, some species of *Myrmecophilus* are host-generalists and do not show any apparent host specificity (Komatsu et al. 2009) whereas other *Myrmecophilus* species, including *M. americanus* and *M. albicinctus*, are characterized by strict host-species specificity (Wetterer and Hugel 2008; Komatsu et al. 2009). It has been suggested that host-species differentiation is one cause of speciation (Schönrogge et al. 2002, Ugelvig et al. 2011). Given the similarities of *M. americanus* and *M. albicinctus*, they may represent a transitional phase of speciation via host switching.

Acknowledgements

We thank Mr. Tomoki Sumino (Malaysia) and Satoshi Inada (Japan) for collecting material; Prof. Rosli Hasim (University of Malaya), Dr. Watana Sakchoowong (Department of National Parks, Wildlife and Plant Conservation), Dr. Wara Asfiya (Division of Zoology, Research Center for Biology, Indonesian Institute of Sciences), Dr. Sri Hartini (Zoology Division, Research Center for Biology, Indonesian Institute of Science–LIPI), and Dr. Henny Herwina (Universitas Andalas) for obtaining permission to conduct this research in their institutions; and Dr. Laure Desutter-Grandcolas (Muséum national d'Histoire naturelle) for permission to examine the type specimens. This work was supported by the Japan Society for the Promotion of Science KAKENHI Grant Number 26·931 to TK.

References

- Chopard L (1924) The fauna of an island in the Chilka Lake. The Dermaptera and Orthoptera of Barkuda Island. Records of the Indian Museum Calcutta 26: 165–191.
- Chopard L (1968) Fam. Gryllidae: Subfam. Mogoplistinae, Myrmecophilinae, Scleropterinae, Cachoplistinae, Pteroplistinae, Pentacentrinae, Phalangopsinae, Trigonidiinae, Eneopterinae; Fam. Oecanthidae, Gryllotalpidae. In: Beier M (Ed.) Orthopterorum Catalogus 12: 213–500.
- Ebner R (1956) Ueber einige fuer Aegypten neue oder seltene Orthopteren. Bulletin de la Societe Entomologique d’Egypte 40: 11–20.
- Emery C (1925) Hymenoptera, Formicidae, Formicinae. In: Wytzman MP (Ed.) Genera Insectum fasc 183. Nabu Press, Charleston, 302 pp.

- Gorochov AV (1994) Contribution to the knowledge of Grylloidea (Orthoptera) of the Seychelles. *Entomologicheskoe Obozrenie* 73: 95–102. [Russian; English translation in *Entomological Review* 73: 103–110]
- Hugel S (2006) Note synonymique sur *Myrmecophilus americanus* Saussure, 1877, et présence de *M. quadrispina* Perkins, 1899, à l'île Maurice (Orth., Myrmecophilinae). *Bulletin de la Société Entomologique de France* 111(4): 448.
- Hugel S, Blard F (2005) Présence de Grillons du genre *Myrmecophilus* à l'île de la Réunion (Orthoptera, Myrmecophilinae). *Bulletin de la Société Entomologique de France* 110: 387–389.
- Ichikawa A (2001) New species of Japanese crickets (Orthoptera, Grylloidea) with notes on certain taxa. *Tettigonia, Memoirs of the Orthopterological Society of Japan* 3: 45–58.
- Ichikawa A, Murai T, Honda E (2000) Monograph of Japanese crickets (Orthoptera; Grylloidea). *Bulletin of the Hoshizaki Green Foundation* 4: 257–332.
- Ingrisch S (2010) Review of the ant loving crickets from South-East Asia (Orthoptera: Gryllidae, Myrmecophilinae). *Entomologische Zeitschrift* 120(1): 3–12.
- Komatsu T, Maruyama M, Itino T (2009) Behavioral differences between two ant cricket species in Nansei Islands: host-specialist versus host-generalist. *Insectes Sociaux* 56(4): 389–396. doi: 10.1007/s00040-009-0036-y
- Latreille PA (1802) *Histoire Naturelle des Fourmis, et recueil de mémoires et d'observations sur les abeilles, les faucheurs, et autres insectes*. Pierre-Théophile Barrois, Paris, 445 pp.
- Maruyama M (2006) Family Myrmecophilidae Saussure, 1870. In: *Orthopterological Society of Japan (Ed.) Orthoptera of the Japanese Archipelago in Color*. Hokkaido University Press, Sapporo, 490–492.
- Murai T, Ito F (2011) Myrmecophilidae. In: *Orthopterological Society of Japan (Ed.) A Field Guide to the Orthoptera of Japan*. Hokkaido University Press, Sapporo, 307–310.
- Saussure HD (1877) *Mélanges Orthoptérologiques*, V^{ème} fascicule. Gryllides. *Mémoires de la Société de Physique et d'Histoire Naturelle de Genève* 25(1): 1–352.
- Schimmer F (1909) Beitrag zu einer Monographie der Gryllodeengattung *Myrmecophila* Latr. *Zeitschrift für Wissenschaftliche Zoologie* 93: 409–534. [pls 22–24]
- Ugelvig LV, Vila R, Pierce NE, Nash DR (2011) A phylogenetic revision of the *Glaucopsyche* section (Lepidoptera: Lycaenidae), with special focus on the *Phengaris–Maculinea* clade. *Molecular phylogenetics and evolution* 61(1): 237–243. doi: 10.1016/j.ympev.2011.05.016
- Wasmann E (1905) Zur Lebensweise einiger in- und ausländischer Ameisengäste (148. Beitrag zur Kenntnis der Myrmecophilen und Termitophilen). *Zeitschrift für Wissenschaftliche Insektenbiologie* 9(1): 329–336, 384–390, 418–428.
- Wetterer JK, Hugel S (2008) Worldwide spread of the ant cricket *Myrmecophilus americanus*, a symbiont of the longhorn crazy ant *Paratrechina longicornis*. *Sociobiology* 52: 157–165.
- Wetterer JK, Hugel S (2014) First North American records of the old world ant cricket *Myrmecophilus americanus* (Orthoptera, Myrmecophilidae). *Florida Entomologist* 97(1): 126–129.