

A new species of *Suwallia* Ricker, 1943 (Plecoptera, Chloroperlidae) from southwestern China, with an updated key to male *Suwallia* species

Abdur Rehman¹, Qing-Bo Huo¹, Yu-Zhou Du^{1,2}

1 School of Horticulture and Plant Protection & Institute of Applied Entomology, Yangzhou University, Yangzhou 225009, China **2** Joint International Research Laboratory of Agriculture and Agri-Product Safety, the Ministry of Education, Yangzhou University, Yangzhou 225009, China

Corresponding author: Yu-Zhou Du (yzdu@yzu.edu.cn)

Academic editor: Sven Bradler | Received 2 August 2021 | Accepted 12 February 2022 | Published 18 March 2022

<http://zoobank.org/95B980B1-7650-4823-8AEA-0E376B4A81D5>

Citation: Rehman A, Huo Q-B, Du Y-Z (2022) A new species of *Suwallia* Ricker, 1943 (Plecoptera, Chloroperlidae) from southwestern China, with an updated key to male *Suwallia* species. ZooKeys 1089: 169–180. <https://doi.org/10.3897/zookeys.1089.72485>

Abstract

A new species of the genus *Suwallia* Ricker, 1943 (Plecoptera, Chloroperlidae), *Suwallia dengba* sp. nov., is described from Tibet and Yunnan, southwestern China. A diagnosis and description of the adult habitus and aedeagal structure are illustrated with color images. Similarities in the terminalia with closely related species are discussed. In addition, an updated key to adult males of the *Suwallia* species of China is provided.

Keywords

Distribution, *Suwallia dengba* sp. nov., Tibet, Yunnan Province

Introduction

The family Chloroperlidae belongs to the superfamily Perloidea and is frequently referred to as “green stoneflies”. It consists of two subfamilies: Chloroperlinae Okamoto, 1912 and Paraperlinae Ricker, 1943. Presently, more than 29 species of the family Chloroperlidae are reported from China, belonging to six genera, namely: *Alloperla* Banks, 1906, *Alaskaperla* Stewart & DeWalt, 1991, *Haploperla* Navás, 1934, *Suwallia* Ricker, 1943, *Sweltsa* Ricker, 1943 and *Utaperla* Ricker, 1952 (Wu 1938; Nelson and Hanson 1968; Du 1999;

Li and Wang 2011; Li et al. 2013, 2014, 2015a, b; Chen and Du 2015, 2016a, b, 2017; Dong et al. 2018; Yang and Li 2018; Chen 2019; Mo et al. 2020; Shi et al. 2022).

The genus *Suwallia* Ricker, 1943 belongs to tribe Suwalliini Surdick, 1985 of the subfamily Chloroperlinae. It is distributed in the East Palearctic and Nearctic regions (DeWalt et al. 2021). Most species of the genus *Suwallia* were revised and recorded by Alexander and Stewart (1999). *Suwallia* is mainly distributed in Russia, Mongolia, Japan, and North America (Alexander and Stewart 1999; Teslenko and Zhiltzova 2009; Judson and Nelson 2012). In China, the first species of *Suwallia* was reported by Li et al. (2015a), and until now seven species of this genus had been reported for the country: *Suwallia errata* Li & Li, 2021, *Suwallia decolorata* Zhiltzova & Levanidova, 1978, and *Suwallia talalajensis* Zhiltzova, 1976 were reported by Li et al. (2015a, b) and Li et al. (2021) from the Inner Mongolia Autonomous Region, northern China (Fig. 7), whereas *Suwallia wolongshana* Du & Chen, 2015 and *Suwallia jibuae* Chen, 2019 were reported by Chen and Du (2015) and Chen (2019) from the Sichuan Province of southwestern China. Recently, *Suwallia kuandian* Shi, Wang & Li, 2022 and *Suwallia asiatica* Zhiltzova & Levanidova, 1978 were reported by Shi et al. (2022) from Liaoning Province, northeastern China. In the current paper, a new species of *Suwallia* is described from Tibet and the Yunnan Province of southwestern China. This is the first record of the *Suwallia* genus from both regions. Tibet is also known as Xizang in Chinese and is positioned on the Tibetan plateau, known as the world's highest and largest plateau. The Yunnan Province lies adjacent to the Tibet, Sichuan, Guizhou, and Guangxi provinces of China and borders with Myanmar, Laos, and Vietnam. The taxonomy of the new species is discussed, a distributional map, and a key to the known species of *Suwallia* from China are provided.

Materials and methods

All specimens were collected by aerial net or hands and preserved in 75% ethanol. Terminalia were examined and illustrated by KEYENCE VHX-5000 and the final images were prepared using Adobe Photoshop CS6. The type specimens of the new species were placed in the insect collection of Yangzhou University (ICYZU), Jiangsu Province, China. Data for the key and distribution map were extracted from the published literature (Chen and Du 2015; Li et al. 2015a, b; Chen 2019; Shi et al. 2022).

Results

Suwallia dengba sp. nov.

<http://zoobank.org/51F6012D-7AB2-4F16-9095-2B1B9E7CE5BE>

Figs 1–8

Type material. *Holotype*, 1♂, China, Tibet Autonomous Region, Dengba village, Mangkam County, Qamdo city, 3437 m, 29°32.406'N, 98°13.425'E, 18.IX.2019, Leg. Huo Qing-Bo (ICYZU). *Paratypes*, 6♂♂, 6♀♀, data same as holotype (Figs 7, 8);

5♂♂, 17♀♀, Yunnan Province, Diqing Tibetan Autonomous Prefecture, Shangri-la city, on the way from Diqing to Gezan Township, 3445 m, 27°45.656'N, 99°56.374'E, 7.IX.2019. Leg. Huo Qing-Bo (ICYZU); 2♂♂, 4♀♀, China, Yunnan Province, Diqing Tibetan Autonomous Prefecture, on national highway (G214) near Tongduishui and Deiyong Benglao, 3432 m, 28°18.282'N, 99°8.472'E, 9.IX.2019, Leg. Huo Qing-Bo (ICYZU); 1♂, 2♀♀, China, Yunnan Province, Diqing Tibetan Autonomous Prefecture, on national highway (G214) near Zhubagong, Deqin County (Fig. 7), 4027 m, 28°23.885'N, 98°59.143'E, 10.IX.2019, Leg. Huo Qing-Bo (ICYZU).

Diagnosis. The new species is characterized by the sclerotized median sclerite of tergum X and its aedeagus armature. The shape of the median sclerite of tergum X resembles a turtle or a hexagonal star. The aedeagus, with a large distinct sclerite divided into an eagle-shaped trifurcate structure, the large median sclerite, and one pair of wing-shaped lateral sclerites on both sides, is diagnostic (Figs 2–4).

Description. Adult habitus (Fig. 1A). Adult body length 8.5–9.5 mm (N = 10), forewing length 6.5–7.5 mm, hindwing length 5.5–6.5 mm. General color of body pale yellow in alcohol. Triocellate, head yellowish-white to yellowish-brown. Ocellar



Figure 1. *Suwallia dengba* sp. nov. **A** male habitus **B** female habitus.

triangle and frontoclypeal area pale yellowish-brown, antenna pale brown, covered with small brown to dark brown setae. Pronotum disc margins covered with dark brown bands and with a thin dark medial stripe (Fig. 2A). Legs pale brown, mesonotum and metanotum with a distinct dark brown U-shaped marking, wings hyaline with yellow venation. Abdominal terga I–VIII with a wide medial trapezoidal dark brown stripe, slightly constricted medially on terga VII and VIII (Figs 1A, 2C–D).




Figure 2. *Suwallia dengba* sp. nov. Holotype male **A** head and prothorax, dorsal view **B** head and prothorax, ventral view **C** terminalia, dorsal view **D** terminalia, ventral view.




Figure 3. *Suwallia dengba* sp. nov. Male paratype. **A** terminalia with aedeagus, dorsal view **B** aedeagus everted, dorsal view **C** aedeagus, caudal ventral view **D** aedeagus, ventral view.

Male (Figs 2–4). Tergum IX concave medially with semicircular stripe anteriorly, posteriorly covered with dark brown, thick hairs. Tergum X divided, median portion with a distinct dark brown sclerite resembling a turtle or hexagonal star in dorsal view (Figs 2C, 6A). Hemitergal processes sclerotized, with tiny hairs, finger-shaped and curved forward. Epiproct membranous, circular, knob-like, covered with minute hairs. Sternum IX ventrally extended anteriorly (Fig. 2D). Aedeagus membranous with a distinct sclerotized sclerite after eversion. Aedeagal sclerite resembling an eagle, divided into a trifurcate structure, a large median sclerite, and one pair of lateral sclerites




Figure 4. *Suwallia dengba* sp. nov. **A** aedeagus **B** terminalia, lateral view **C** aedeagal sclerite, dorsal view **D** aedeagal sclerite, ventral view.

(Figs 3A, 4A–D, 6B). Lateral sclerites armed with minute scales. Membranous part of aedeagus with fine cuticular asperities (Fig. 3A–D).

Female. Adult habitus (Fig. 1B). Body length 9.0–10 mm (N = 10), forewing length 7.5–8.5 mm, hindwing length 6.5–7.5 mm. General body color, shape and appearance similar to those of male. Head and pronotum similar. Dorsal segment of abdomen with trapezoidal dark brown stripe extended to sternum VIII, subgenital plate large, extending to posterior portion of sternum IX, constricted from base, expanded medially, then slightly tapering toward posterior margins. Subgenital plate covered with minute, fine hairs. Tergum X not produced posteriorly. Paraproct in the shape of a small triangle, bearing small hairs (Fig. 5A–C).

Egg and nymph. Unknown.

Distribution. Southwestern China (Tibet and Yunnan Province).

Etymology. The species is named after the type locality, Dengba village.

Remarks. The new species is closely related to *Suwallia talalajensis*, but can be distinguished by the sclerotized portion between the hemitergal processes, the




Figure 5. *Suwallia dengba* sp. nov. Female paratype. **A** terminalia, ventral view **B** terminalia, ventral view **C** terminalia, dorsal view.




Figure 6. *Suwallia dengba* sp. nov. **A** male terminalia, dorsal view **B** aedeagal sclerite.



Figure 7. Revised map showing distribution of *Suwallia* species in China (modified from www.tianditu.gov.cn).

pigmentation of tergum IX, the armature of the aedeagus and the well-developed, membranous, knob-like epiproct. *Suwallia talalajensis* does not have a distinct aedeagal sclerite (Li et al. 2015b: fig. 5), whereas the new species has a distinct sclerite (Figs 4A–D, 6B). Tergum IX of the new species is covered with abundant, thick hairs, and its body pigmentation is different from that of *Suwallia talalajensis*. The new species also shows similar characteristics to *Suwallia errata* (Li et al. 2021), but it can be easily differentiated by the sclerotized portion between the hemitergal process and the shape of the aedeagus. *Suwallia errata* has a V-shaped aedeagal sclerite (Li et al. 2015a: figs 1–6), but the new species has the aedeagal sclerite of a different shape. The new species lives in fast-flowing rivers (width = 5 m), where a large gravel substrate is present. The adults occur on leaves of trees or shrubs near the river (Fig. 8).



Figure 8. Habitat at the type locality of *Suwallia dengba* sp. nov. Specimens were collected from the small trees and grasses near the stream (photograph Huo Qing-Bo).

Key to adult males of *Suwallia* species from China (modified from Chen 2019)

- 1 Epiproct reduced, tergum X with two median sclerites 2
- Epiproct well developed, tergum X with undivided median sclerite 3
- 2 Tergum X with two longitudinal median sclerites (see Chen and Du 2015: figs 1–8)..... *Suwallia wolongshana*
- Tergum X with H-shaped median sclerite (see Chen 2019: fig. 3).....
..... *Suwallia jibuae*
- 3 Tergum X with V-shaped median sclerite, aedeagus membranous, without spines or structures (see Shi et al. 2022: fig. 2) *Suwallia asiatica*
- Tergum X median sclerite triangular or subrectangular in shape, aedeagus with spines or structures 4
- 4 Tergum X median sclerite triangular in shape, epiproct small, aedeagus with triangular spines forming T-shaped structure (see Li et al. 2015b: fig. 2).....
..... *Suwallia decolorata*
- Tergum X median sclerite not as above, epiproct well developed and knob-like..... 5
- 5 Tergum X medial sclerite subrectangular, anterior margins with two separate sclerites 6
- Tergum X median sclerite of turtle or hexagonal shape 7

- 6 Tergum X anterior margins divided into two sclerites, epiproct with long hairs and without posterolateral bifurcation, aedeagus with V-shaped sclerite (see Li et al. 2015a: figs 1–6)..... *Suwallia errata*
- Tergum X anterior margins with two separate paramedial sclerites, arch-shaped in lateral view, epiproct with stout posterolateral bifurcation, aedeagus with triangular sclerite, lateral margins darker (see Shi et al. 2022: fig. 1).....
.....*Suwallia kuandian*
- 7 Tergum X median sclerite turtle-like, aedeagus membranous, without distinct armature or sclerite (see Li et al. 2015b: fig. 2) ... *Suwallia talalajensis*
- Tergum X median sclerite hexagonal star-shaped, pointed posteriorly, aedeagus with distinct trifurcate sclerite (Figs 2–4)*Suwallia dengba* sp. nov.

Acknowledgements

This study was supported by the National Natural Science Foundation of China (No. 31872266, 31572295) and the Biodiversity Survey and Assessment Project of the Ministry of Ecology and Environment, China (No. 2019HJ2096001006). We express our deep gratitude to Xing-Min Wang, Wei-Dong Huang, Qing-Heng Tian (South China Agricultural University) and Yan-Dong Chen (Institute of Zoology, Chinese Academy of Sciences) for help collecting the specimens. Moreover, we express our heartfelt thanks to the reviewers for providing comments that significantly improved the manuscript.

References

- Alexander K-D, Stewart K-W (1999) Revision of the genus *Suwallia* Ricker (Plecoptera: Chloroperlidae). Transactions of the American Entomological Society 125(3): 185–250. <https://www.jstor.org/stable/25078681>
- Banks N (1906) On the perlid genus *Chloroperla*. Entomological News 17(5): 174–175.
- Chen Z-T, Du Y-Z (2015) A new species of *Suwallia* (Plecoptera: Chloroperlidae) from China. Zootaxa 4018(2): 297–300. <https://doi.org/10.11646/zootaxa.4018.2.9>
- Chen Z-T, Du Y-Z (2016a) A new species of *Haploperla* from China (Plecoptera, Chloroperlidae). ZooKeys 572: 1–6. <https://doi.org/10.3897/zookeys.572.6270>
- Chen Z-T, Du Y-Z (2016b) Two new species of *Haploperla* (Plecoptera: Chloroperlidae) from China. Zootaxa 4196(3): 415–422. <https://doi.org/10.11646/zootaxa.4196.3.5>
- Chen Z-T, Du Y-Z (2017) A new species of *Sweltsa* (Plecoptera: Chloroperlidae) from China, with a key to the *Sweltsa* males of China. Zootaxa 4337(2): 291–293. <https://doi.org/10.11646/zootaxa.4337.2.8>
- Chen Z-T (2019) Review of the genus *Suwallia* (Plecoptera: Chloroperlidae) from China with description of *Suwallia jibhuae* sp. nov. from Sichuan Province. Zootaxa 4603(3): 583–588. <https://doi.org/10.11646/zootaxa.4603.3.11>

- DeWalt R-E, Maehr M-D, Hopkins H, Neu-Becker U, Stueber G (2021) Plecoptera Species File online. Version 5.0/5.0. <http://plecoptera.speciesfile.org> [Accessed on: 2021-7-28]
- Dong W-B, Cui J-X, Li W-H (2018) A new species of *Sweltsa* (Plecoptera: Chloroperlidae) from Sichuan Province of southwestern China. *Zootaxa* 4418(4): 388–392. <https://doi.org/10.11646/zootaxa.4418.4.5>
- Du Y-Z (1999) A taxonomic study on Plecoptera from China. Zhejiang University, Hangzhou, 324 pp.
- Judson S-W, Nelson C-R (2012) A guide to Mongolian stoneflies (Insecta: Plecoptera). *Zootaxa* 3541(1): 1–118. <https://doi.org/10.11646/zootaxa.3541.1.1>
- Li W-H, Wang R (2011) A new species of *Alloperla* (Plecoptera: Chloroperlidae) from China. *Zootaxa* 3040(1): 29–33. <https://doi.org/10.11646/zootaxa.3040.1.4>
- Li W-H, Yao G, Qin X (2013) *Haploperla choui* sp. n. (Plecoptera: Chloroperlidae), a remarkable new stonefly from Qinling Mountains of China. *Zootaxa* 3640(4): 550–556. <https://doi.org/10.11646/zootaxa.3640.4.3>
- Li W-H, Yang J, Yao G (2014) Review of the genus *Sweltsa* (Plecoptera: Chloroperlidae) in China. *Journal of Insect Science* 14(1): e286. <https://doi.org/10.1093/jisesa/ieu148>
- Li W-H, Murányi D, Shi L (2015a) The first record of genus *Suwallia* Ricker, 1943 (Plecoptera: Chloroperlidae) from China. *Illiesia* 11(03): 23–28. <http://illiesia.speciesfile.org/papers/Illiesia11-03.pdf>
- Li W-H, Murányi D, Shi L (2015b) New species records of *Suwallia* Ricker, 1943 (Plecoptera: Chloroperlidae) from China, with description of the nymph of *S. decolorata* Zhiltzova & Levanidova, 1978. *Zootaxa* 3994(4): 556–564. <https://doi.org/10.11646/zootaxa.3994.4.4>
- Li W-L, Wang Y-Y, Wang Y, Li W (2021) A new species of *Suwallia* Ricker, 1943 from Japan, and the identity of *Alloperla telekojensis* Šámal, 1939 (Plecoptera: Chloroperlidae). *Zootaxa* 5040(4): 575–581. <https://doi.org/10.11646/zootaxa.5040.4.7>
- Mo R-R, Ye J-P, Wang G-Q, Li W-H (2020) The first record of the family Chloroperlidae (Plecoptera) from the Guangxi Zhuang Autonomous Region of southern China, with description of a new species of *Sweltsa* Ricker, 1943. *Zootaxa* 4853(2): 275–282. <https://doi.org/10.11646/zootaxa.4853.2.8>
- Navás R-P-L (1934) Nevropteres et insectes voisins (Chine et Pays environnements). Notes d'Entomologie Chinoise. Musée Heude. Shanghai 2: 1–16.
- Nelson C-H, Hanson J-F (1968) Two new species of *Alloperla* (Plecoptera: Chloroperlidae) from China. *Journal of the Kansas Entomological Society* 41: 425–428.
- Okamoto H (1912) Erster Beitrag zur Kenntnis der Japanischen Plecopteren. *Transactions of the Sapporo Natural History Society* 4: 105–170.
- Ricker W-E (1943) Stoneflies of southwest British Columbia. Indiana University Publications, Science Series 12: 1–145.
- Ricker W-E (1952) Systematic studies on Plecoptera. Bloomington, Indiana University, 1–200.
- Shi W-J, Wang H-L, Li W-H (2022) A new species and three new records of Chloroperlidae (Plecoptera) from northeastern China. *Zootaxa* 5093(5): 584–592. <https://doi.org/10.11646/zootaxa.5000.5.7>

- Surdick R-F (1985) Nearctic genera of Chloroperlinae (Plecoptera: Chloroperlidae). Illinois Biological Monographs, 54. University of Illinois Press, Urbana and Chicago, Illinois, 146 pp.
- Teslenko V-A, Zhiltzova L-A (2009) Key to the stoneflies (Insecta, Plecoptera) of Russia and adjacent countries. Imagines and nymphs. Dalnauka, Vladivostok, 382 pp.
- Wu C-F (1938) Plecopterorum sinensium: A monograph of the stoneflies of China (Order Plecoptera). Yenching University, Beijing, 225 pp.
- Yang D, Li W-H (2018) Species catalogue of China. Volume 2 Animals, Insecta (III), Plecoptera. Science Press, Beijing, 71 pp.
- Zhiltzova L-A, Levanidova I-M (1978) Novie vidi vesnianok (Plecoptera) s Dalnego Vostoka. Novie Vidi Zivotnih, Aka-demia Nauk SSSR Trudy Zoologiceskogo Instituta 61: 3–29.