

A new species of *Ampithoe* (Amphipoda, Ampithoidae) from Korea, with a redescription of *A. tarasovi*

Myung-Hwa Shin¹, Charles Oliver Coleman²

1 National Marine Biodiversity Institution of Korea, Seochun, 325–902, Republic of Korea **2** Museum für Naturkunde Berlin, Invalidenstraße 43, 10115 Berlin, Germany

Corresponding author: Charles Oliver Coleman (oliver.coleman@mfn-berlin.de)

Academic editor: Alan Myers | Received 24 August 2021 | Accepted 5 November 2021 | Published 22 December 2021

<http://zoobank.org/FDE9C8D9-5A99-4BBE-BAD1-F432729D9B9A>

Citation: Shin M-H, Coleman CO (2021) A new species of *Ampithoe* (Amphipoda, Ampithoidae) from Korea, with a redescription of *A. tarasovi*. ZooKeys 1079: 129–143. <https://doi.org/10.3897/zookeys.1079.73443>

Abstract

An ampithoid species, previously known as *A. tarasovi* in Korea, is assigned as a new species, *A. changbaensis* sp. nov. based on the reexamination of the holotype of *A. tarasovi*. The new species shows different morphological characteristics compared to *A. tarasovi*, especially in male gnathopods 1 and 2. The type material of *A. tarasovi* is redescribed and illustrated here and a key to the Korean species of *Ampithoe* is provided.

Keywords

Ampithoe changbaensis sp. nov., Crustacea, new species, Russia, taxonomy

Introduction

The genus *Ampithoe* Leach, 1814 is a herbivorous amphipod group associated with algae and seagrasses in coastal regions worldwide (Myers and Lowry 2003; Shin et al. 2015; Peart and Ahjong 2016). *Ampithoe* is the most speciose genus of the family Ampithoidae and includes more than 70 species worldwide (Horton et al. 2021).

To date, eight species of the genus *Ampithoe* have been reported from Korea: *A. akuolaka* J.L. Barnard, 1970; *A. brevipalma* Kim & Kim, 1988; *A. lacertosa* Bate, 1858; *A. ramondi* Audouin, 1826; *A. shimizuensis* Stephensen, 1944; *A. tarasovi* Bullycheva, 1952; *A. valida* Smith, 1873; and *A. youngsanensis* Kim & Kim, 1988 (Kim and Kim 1988; Shin et al. 2010; Kim 2011; Jung and Yoon 2014; Peart and Ahyong 2016). Among them, *A. lacertosa*, collected in Korea, was described by Kim and Kim (1987, 1988). After the study, the material of the Korean specimens of *A. lacertosa* was stored in the collections of the Seoul National University. Later, the deposited Korean material identified as *A. lacertosa* was reexamined in a taxonomic study of Shin et al. in 2010, and in this material a second species identified as *A. tarasovi* was found. These two species, however, were identified based on the original descriptions and other published records only. However, the type specimens of these two species were not examined by Shin et al. (2010).

For a precise identification of species, type specimens and detailed original descriptions are essential for taxonomy and a flawless identification. If type material is lost and the original texts and illustrations are short and poor in quality, it may lead to misidentifications of species.

In this study, the holotype of *A. tarasovi* collected in Russia was reexamined. Through the examination, the Korean material of *A. tarasovi* was identified as a distinct species having morphological characteristics differing from the type material of *A. tarasovi*. The examined specimens (previously known as *A. tarasovi* in Korea) are assigned as a new species, *A. changbaensis* sp. nov., which is described based on the specimens previously misidentified as *A. tarasovi* by Shin et al. (2010). Moreover, the type material of *A. tarasovi* is redescribed and illustrated.

Materials and methods

To designate the type material of the new species, the specimens which have been deposited at the Laboratory of Systematics and Molecular Evolution in the Seoul National University were used. Other material was collected in Korea among algae in tide pools, in the intertidal zone, and in shallow water at low tide. The holotype of *Ampithoe tarasovi* was loaned from the Moscow Museum, Russia, and examined at the crustacean department of Museum für Naturkunde Berlin, Germany.

The specimens were analyzed and pencil drawn under a dissection microscope (Leica M250C), and appendages and mouthparts were drawn under a Leica DMLB; both microscopes were equipped with a camera lucida. The line drawings were made using the technique described by Coleman (2003, 2009). Body length was measured along the mid-body line from the tip of the rostrum to the posterior end of urosomite 3. All examined material is currently being deposited at the Marine Arthropod Depository Bank of Korea (MADBK). The descriptions were produced from a DELTA (Dallwitz 2005) database to the ampithoid genera and species (initially compiled by our colleague Dr Jim Lowry).

Results

Systematics

Ampithoidae Boeck, 1871

Ampithoe Leach, 1814

Ampithoe changbaensis sp. nov.

<http://zoobank.org/14B275AB-4C6A-4390-BA22-E844D6DC044E>

Figures 1–3

Ampithoe lacertosa: Kim and Kim 1987: 3, fig. 2. Kim and Kim 1988: 109, fig. 2A [not *Ampithoe lacertosa* Bate, 1858].

Ampithoe tarasovi: Shin et al. 2010: 300, figs 4–6 [not *Ampithoe tarasovi* Bulycheva, 1952].

Type locality. Hamo beach, Jeju-do, South Korea.

Type material. Holotype. Male, 17.6 mm (MABIK CR00248547), Hamo beach, Daejeong-eup, Seogwipo-si, Jeju-do, Korea (33°12'37.01"N, 126°15'44.34"E), 30 May 2007, coll. Shin and Hong.

Paratype. Female, 22.3 mm (MABIK CR00248548); male and female, 16–21 mm (MABIK CR00248549); 2 males and 1 female, 17–20 mm (MABIK CR00248550), same data as the holotype.

Additional material examined. 3 males, Hamo beach (MABIK CR00248551), Daejeong-eup, Seogwipo-si, Jeju-do, Korea (33°12'37.01"N, 126°15'44.34"E), 30 May 2007; 3 males and female (MABIK CR00248552), Gujwa-eup, Jeju-si, Jeju-do, Korea (33°32'2.58"N, 126°50'27.25"E), 15 Mar. 2017; male and 3 females (MABIK CR00248553), Deajin port, Hyeonnae-myeon, Goseong-gun, Gangwon-do, Korea (38°29'55.42"N, 128°25'35.53"E), 21 Jun. 2019.

Etymology. The new species is named in honor to Prof. Chang Bae Kim, an early amphipodologist of Korea, who collected and described the species firstly from Jeju, Korea in 1987.

Description. Based on holotype male, 17.6 mm. Body (Figs 1, 2) heavily covered with dark pigmentation spots creating bands on head, coxae, pereon, and pleon.

Head. Antenna 1 longer than antenna 2; peduncular article 1 subequal in length to article 2; article 2 longer than article 3 (2.3 times article 3); article 3 shorter than article 1 (0.5 times article 1).

Antenna 2 slender, similar to antenna 1; peduncular article 4 subequal in length to article 5; flagellum longer than peduncular article 5.

Upper lip with midlateral notch on margins.

Mandible molar well developed, triturating; palp apically setose, 3-articulate; mandibular palp article 1 shorter than article 2 (0.5 times article 2); article 2 shorter than article 3 (0.7 times article 3); article 3 longer than article 1 (3 times article 1).



Figure 1. Photograph of the *Ampithoe changbaensis* sp. nov., habitus of the male paratype. Scale bar: 5 mm. Photograph by Jin-Ho Park.

Lower lip outer plates forming a medial excavation, lateral lobe slightly longer than medial lobe; mandibular lobe curved laterally, subacute apically.

Maxilla 1 inner plate with 1 slender seta; palp well developed, with apical robust setae.

Maxilla 2 inner plate narrower than outer plate, with oblique setal row.

Maxilliped outer plate with developed row of large robust setae along medial margin.

Pereon. Gnathopod 1 (Fig. 3A) sexually dimorphic, smaller than gnathopod 2, carpus and propodus with numerous plumose setae on both anterior and posterior margins; coxa broader than deep, anterior margin slightly convex, anteroventral corner produced, rounded; basis longer than coxa, expanded anterodistally, anterodistal lobe large and subrounded; ischium anterior margin with small subrounded lobe; merus posterodistal corner subacute, produced; carpus about 2 times as long as broad, longer than propodus (1.3 times propodus), with posterodistal lobe slightly overlapping propodus, posterior margin slightly convex; propodus broad, 1.4 times as wide as long, subovoid; palm acute, convex, defining corner rounded with 1 robust seta; dactylus subequal in length to palm.

Gnathopod 2 (Fig. 3B) sexually dimorphic; basis longer than coxa, anterodistal lobe large and rounded, not reaching beyond ischium; ischium with anterior rounded lobe; carpus much shorter than propodus (0.4 times propodus), subtriangular; propodus narrow, 1.8 times as long as wide, subrectangular; palm transverse, with a sloped quadrate midmedial hump and an apically rounded tooth on posterodistal corner; dactylus slightly overreaching palm, curved, robust, apically blunt.




Figure 2. *Ampithoe changbaensis* sp. nov., male holotype. Habitus. Scale bar: 5 mm.

Pereopod 3 basis narrow; merus narrow; carpus about twice as long as broad.

Pereopod 4 basis similar to pereopod 3.

Pereopod 5 (Fig. 3C) basis subovoid, without posterodistal lobe; merus subrectangular.

Pereopod 6 basis posterior margin rounded proximally, straight distally, with marginal robust setae; merus subrectangular.

Pereopod 7 (Fig. 3D) similar to pereopod 6; basis with marginal robust setae.

Pleon. Epimera 1–3 with lateral ridges; epimera 2 and 3 subrounded posterodistally, with rounded tooth on each posteroventral angle. Epimeron 1 rounded posterodistally, with tooth on posteroventral angle; epimeron 2 ventral margin evenly curved; epimeron 3 ventral margin straight.

Uropod 1 reaching to end of uropod 2 rami; inner ramus longer than outer ramus; outer ramus slender, about 6 times as long as broad.

Uropod 2 inner ramus longer than outer ramus.

Uropod 3 (Fig. 3E) peduncle much longer than broad (2.2 times as wide as long), 1.8 times as long as rami, 2 inner marginal robust setae, marginal slender setae present, with 7 distal peduncular robust setae; rami long, about twice as long as broad; outer ramus subequal in length to inner ramus, with 2 large recurved distal robust setae, and with 3 dorsomarginal robust setae, with lateral setal fringe; inner ramus with 4 distal robust setae, with 3 and 2 lateral robust setae on both inner and outer margins, respectively.

Telson (Fig. 3F) subtriangular, apically rounded, with small apical cusps; with 3 groups of lateral plumose setae, and a pair row of setae on submedial margins.

Depth zone. Littoral (1–2 m).

Distribution. Jeju, East Sea of Korea, South Sea of Korea.

Remarks. This species has been recognized as *A. tarasovi* in Korea since the species was described and illustrated by Shin et al. (2010). However, the type material of




Figure 3. *Ampithoe changbaensis* sp. nov., male holotype **A** gnathopod 1 **B** gnathopod 2 **C** pereopod 5 **D** pereopod 7 **E** uropod 3 **F** telson. Scale bars: 2 mm (**A–D**); 0.25 mm (**E, F**).

A. tarasovi described from Russia was examined here, and as a result, important morphological differences were detected between Korean material and the Russian type specimen. The characteristics differing between the Korean specimens and the type material are as follows. In male gnathopod 1, the basis is expanded anterodistally in the Korean specimens, while it is narrow and straight in the type material. The length

ratios of carpus and propodus of the Korean and Russian specimens are 1.3 and 1.8, respectively. The shape of gnathopod 1 carpus is more rectangular in the Russian material than in the Korean sample. The propodus of male gnathopod 2 is longer and more rectangular in the Russian material compared to the Korean sample. The apical margin of the telson is round in the Korean specimen, while it is subacute in the Russian type material. Based on the morphological differences mentioned above, the Korean material has been assigned to a new species, *A. changbaensis* sp. nov.

Ampithoe changbaensis sp. nov. is similar to *A. prolata* Hughes & Peart, 2013; however, it can be distinguished from this species by the following characteristics: (1) presence of marginal setae on merus, carpus and propodus of male gnathopod 1; (2) swollen basis of male gnathopod 1; (3) subrectangular and trapezoid shape of propodus of male gnathopod 2; and (4) truncated posterior margin of carpus of female gnathopod 1.

Ampithoe tarasovi Bulycheva, 1952

Figures 4–8

Ampithoe tarasovi Bulycheva, 1952: 246, fig. 38. Tzvetkova 1967: 190.

Type material. Male, collected by Tarasov from De-Kastri, Sea of Japan, 3 Aug. 1929.

Description. Based on holotype male, 14.3 mm (re-measured along the midbody line from the tip of the rostrum to the posterior end of urosomite 3), deposited at the Moscow Museum, Russia (no. 1/21349).

Head (Fig. 4B). Upper lip (Fig. 4C) with mid-lateral notch on margins.

Mandible (Fig. 4D, E) molar well developed, tritulating; accessory setal row with 9 robust setae; palp apically setose, 3-articulate; mandibular palp (Fig. 4F) article 1 shorter than article 2 (0.6 times article 2); article 2 shorter than article 3 (0.7 times article 3); article 3 long (3.3 times as long as wide), longer than article 1 (2.3 times article 1).

Lower lip (Fig. 4G) outer plates forming a medial excavation, lateral lobe much longer than medial lobe; mandibular lobe curved laterally, rounded apically.

Maxilla 1 (Fig. 5A) inner plate with 1 slender seta; palp well developed, with apical robust setae.

Maxilla 2 (Fig. 5B) inner plate narrower than outer plate, with oblique setal row.

Maxilliped (Figs 5C–E, 6A) outer plate with developed row of large robust setae along medial margin.

Pereon. Gnathopod 1 (Fig. 6B) sexually dimorphic, smaller than gnathopod 2, carpus and propodus with numerous plumose setae on both anterior and posterior margins; coxa subequal to coxa 2 in length, broader than deep, anterior margin straight, anter-oventral corner produced, rounded; basis longer than coxa, anterodistal lobe large and rounded; ischium anterior margin with large rounded lobe; merus posterodistal corner subquadrate; carpus about 2 times as long as broad, longer than merus, longer than propodus (1.8 times propodus), with posterodistal lobe slightly overlapping propodus, posterior margin straight; propodus broad, 1.4 times as wide as long, subovoid; palm acute, convex, defining corner rounded with 1 robust seta; dactylus subequal in length to palm.




Figure 4. *Ampithoe tarasovi* Bulycheva, male holotype **A** epimeral plates 1–3 **B** head **C** upper lip **D** left mandible **E** right mandible **F** palp of mandible **G** lower lip. Scale bars: 1 mm (**A**, **B**); 0.2 mm (**C**, **G**); 0.1 mm (**D–F**).

Gnathopod 2 (Figs 6C, 7A) sexually dimorphic; basis longer than coxa, with sparse slender setae, anterodistal lobe large and rounded, not reaching beyond ischium; ischium anterior margin with subquadrate lobe; carpus much shorter than propodus (0.3 times propodus), subtriangular; propodus narrow, 2.3 times as long as wide, subrectangular; palm transverse, with a sloped quadrate mid-medial hump and an apically




Figure 5. *Ampithoe tarasovi* Bulycheva, male holotype **A** maxilla 1 **B** maxilla 2 **C** inner lobe of maxilliped **D** outer lobe of maxilliped **E** maxilliped. Scale bars: 0.1 mm (**A–D**); 1 mm (**E**).

rounded defining tooth on posterodistal corner; dactylus slightly overreaching palm, curved, robust, apically blunt, without unguis.

Pereopod 3 (Fig. 8C) basis narrow; merus narrow; carpus about twice as long as broad.

Pereopod 4 (Fig. 8D) basis similar to pereopod 3.

Pereopod 5 (Fig. 8E) coxa simple and subrectangular. Pereopods 5–7 lost.




Figure 6. *Ampithoe tarasovi* Bulycheva, male holotype **A** palp of maxilliped **B** gnathopod 1 **C** gnathopod 2. Scale bars: 0.1 mm (**A**); 1 mm (**B**); 0.5 mm (**C**).

Pleon. Epimera 1–3 (Fig. 4A) with lateral ridges; ventral margin of epimera 2 and 3 straight, with distinct tooth on each posteroventral angle. Epimeron 1 subrounded posterodistally, with tooth on posteroventral angle; epimeron 2 subrounded posterodistally; epimeron 3 straight and sloped posterodistally.

Uropod 1 (Fig. 7B, C) reaching to end of uropod 2 rami; peduncle with 10 robust setae; inner ramus longer than outer ramus, with 5 marginal robust setae; outer ramus slender, about 6 times as long as broad, with 14 marginal robust setae.




Figure 7. *Ampithoe tarasovi* Bulycheva, male holotype **A** gnathopod 2 **B** lateral view of urosomite 3 **C** uropod 1 **D** uropod 2 **E** uropod 3 **F** telson. Scale bars: 1 mm (**A–C, E**); 0.2 mm (**D, F**).

Uropod 2 (Fig. 7D) peduncle with 8 robust setae; inner ramus longer than outer ramus, with 8 marginal robust setae; outer ramus 11 marginal robust setae.

Uropod 3 (Fig. 7E) peduncle much longer than broad (2.3 times width), 2.2 times as long as rami, with 3 marginal robust setae, marginal slender setae present, with 8 distal peduncular robust setae; rami long, about twice as long as broad; outer ramus shorter than inner ramus, with 2 large recurved distal robust setae and 1 dorsal robust setae, with lateral setal fringe; inner ramus with 6 distal robust setae, with 2 lateral robust setae both inner and outer margins, respectively.




Figure 8. *Ampithoe tarasovi* Bulycheva, female **A** gnathopod 1 **B** gnathopod 2 **C** pereopod 3 **D** pereopod 4 **E** coxa 5. Scale bars: 1 mm (**A–E**).

Telson (Fig. 7F) subtriangular, apically subacute, with small apical cusps, with 4 or 5 of lateral setae on both margins, and 2 pairs of lateral plumose setae, with 2 or 3 submedial setae on both margins.

Sexual dimorphic female, 15.3 mm.

Gnathopod 1 (Fig. 8A) subequal in size to gnathopod 2; coxa about as broad as long, anterior margin concave, anteroventral corner produced and subacute; basis sub-

Table 1. Distribution of the species of *Ampithoe* in coastal regions of Korea: EC, Eastern coast; WC, Western coast; SC, Southern coast; JC, Jeju coast (distribution data cited from Kim [2011] and Jung and Yoon [2014]).

Species	Coastal region			
	EC	WC	SC	JC
<i>A. akuolaka</i>	-	-	⊙	-
<i>A. brevipalma</i>	⊙	-	⊙	-
<i>A. lacertosa</i>	⊙	⊙	⊙	⊙
<i>A. ramondi</i>	⊙	-	⊙	⊙
<i>A. shimizuensis</i>	⊙	⊙	⊙	⊙
<i>A. valida</i>	⊙	⊙	⊙	⊙
<i>A. youngsanensis</i>	-	-	⊙	⊙
<i>A. changbaensis</i> sp. nov.	⊙	-	⊙	⊙

equal in length to coxa, with sparse slender setae, anterodistal lobe large and rounded; carpus subequal in length to propodus (1.1 times propodus); propodus narrow, 2 times as long as wide, subtriangular; palm acute, straight, defining corner subrounded with 1 robust seta; dactylus slightly overreaching palm, inner margin crenate.

Gnathopod 2 (Fig. 8B) basis shorter than coxa, with sparse slender setae, anterodistal lobe large and rounded, not reaching beyond ischium; ischium anterior margin without distinct lobe; carpus shorter than propodus (0.7 times propodus); propodus narrow, 1.6 times as long as wide, subrectangular; palm acute, defining corner subrounded with 1 robust seta; dactylus slightly overreaching palm, tapering evenly, apically acute, inner margin crenate.

Depth zone. Sublittoral (0–24 m).

Distribution. Peter the Great Bay, Sea of Japan.

Remarks. This species has the following characteristics: the apical and medial lobes of the outer lobes are separated in the lower lip; the carpus of male gnathopod 1 is about 1.8 times as long as the propodus; the palm of the male gnathopod 2 has a sloped quadrate hump and posterodistal tooth. Bulycheva (1952) noted that *A. tarasovi* is very abundant in macroalgae and reefs in Petra Velikogo Bay and in the northern Sea of Japan.

Here we provide a key to the Korean species of *Ampithoe* and distributional information in four coastal regions of Korea in Table 1.

- 1 In male, gnathopod 2, propodus subovoid or not large; palm acute or not.... 2
- In male, gnathopod 2, propodus large subrectangular; palm transverse or nearly so..... 5
- 2 Gnathopod 2, palm extremely acute, not defined in male.... *A. youngsanensis*
- Gnathopod 2, palm not acute in male..... 3
- 3 Gnathopod 2, palm concave, defined with angle in male *A. brevipalma*
- Gnathopod 2, palm incised, with distinct lobe in male 4
- 4 Gnathopod 2, propodus with pointed thumb-like lobe in male ...*A. akuolaka*
- Gnathopod 2, propodus with rounded thumb-like lobe in male*A. ramondi*

- 5 Epimeron 3 without tooth on posteroventral angle 6
 – Epimeron 3 with subacute tooth on posteroventral angle 7
 6 In male antenna 2, peduncular article 4 compressed and expanded
 *A. shimizuensis*
 – In male antenna 2, peduncular article 4 not expanded, ordinary *A. valida*
 7 In male, gnathopod 2, carpus and propodus with dense marginal setae; palm
 with quadrate hump; gnathopod 1, basis expanded anterodistally in male....
 *A. changbaensis* sp. nov.
 – In male, gnathopod 2, carpus and propodus without dense setae; palm with-
 out hump; gnathopod 1, basis not expanded anterodistally in male 7

Acknowledgements

We are thankful to the anonymous reviewers for valuable comments on the manuscript. We are grateful to Dr Jim Lowry (Australian Museum, Sydney, NSW, Australia) for permitting us to use his DELTA database of amphitoid species of the world. We would like to thank Natalia L. Demchenko for kindly translating the Russian label on the specimens of *A. tarasovi*.

This study was supported by the National Marine Biodiversity Institute Research Program (2021M00100).

References

- Audouin V (1826) Explication sommaire des planches de crustacés de l’Egypte et de la Syrie, publiées par Jules-Cesar Savigny, membre de l’Inst.; offrant un exposé des caractères naturels des genres avec la distinction des espèces. Description de l’Egypte; Histoire naturelle 1(4): 77–98.
- Barnard JL (1970) Sublittoral Gammaridea (Amphipoda) of the Hawaiian Islands. Smithsonian Contributions to Zoology 34: 1–286. <https://doi.org/10.5479/si.00810282.34>
- Bate CS (1858) On some new genera and species of Crustacea Amphipoda. Annals and Magazine of Natural History (Series 3) 1: 361–362. <https://doi.org/10.1080/00222935808696932>
- Boeck A (1871) Crustacea Amphipoda borealia et arctica. Forhandlingar i Videnskabs-Selskabet i Christiania 1870: 83–280. <https://doi.org/10.5962/bhl.title.2056>
- Bulycheva AI (1952) Novue vidy bokoplavov (Amphipoda, Gammaridea) iz Japonskogo Morja. Akademiia Nauk SSSR. Trudy Zoologicheskogo Instituta 12: 195–250.
- Coleman CO (2003) “Digital inking”: How to make perfect line drawings on computers. Organism, Diversity and Evolution, Electronic Supplement 14: 1–14. <https://doi.org/10.1078/1439-6092-00081>
- Coleman CO (2009) Drawing setae the digital way. Zoosystematics and Evolution 85(2): 305–310. <https://doi.org/10.1002/zoos.200900008>
- Dallwitz MJ (2005) Overview of the DELTA System. <http://delta-intkey.com>

- Horton T, Lowry J, De Broyer C, Bellan-Santini D, Coleman CO, Corbari L, Costello MJ, Daneliya M, Dauvin JC, Fišer C, Gasca R, Grabowski M, Guerra-García JM, Hendrycks E, Hughes L, Jaume D, Jazdzewski K, Kim YH, King R, Krapp-Schickel T, LeCroy S, Lörz AN, Mamos T, Senna AR, Serejo C, Sket B, Souza-Filho JF, Tandberg AH, Thomas JD, Thurston M, Vader W, Väinölä R, Vonk R, White K, Zeidler W (2021) World Amphipoda Database. *Ampithoe* Leach, 1814. World Register of Marine Species. <http://www.marine-species.org/aphia.php?p=taxdetails&cid=101459> [accessed on 2021-8-11]
- Hughes LE, Peart RA (2013) New species and new records of Ampithoidae (Peracarida: Amphipoda) from Australian waters. *Zootaxa* 3719(1): 1–102. <https://doi.org/10.11646/zootaxa.3719.1.1>
- Jung TW, Yoon SM (2014) A New Record of the ampithoid species, *Ampithoe akuolaka* (Crustacea: Amphipoda: Ampithoidae) from Korea. *Animal Systematics, Evolution and Diversity* 30(3): 215–224. <https://doi.org/10.5635/ASED.2014.30.3.215>
- Kim HS, Kim CB (1987) Marine gammaridean Amphipoda (Crustacea) of Cheju Island and its adjacent waters, Korea. *Korean Journal of Systematic Zoology* 3(1): 1–23.
- Kim HS, Kim CB (1988) Marine gammaridean Amphipoda (Crustacea) of the family Ampithoidae from Korea. *The Korean Journal of Systematic Zoology* 2: 107–134.
- Kim YH (2011) Invertebrate fauna of Korea. National Institute of Biological Resources, Ministry of Environment 21: 1–129.
- Leach WE (1814) Crustaceology. *The Edinburgh Encyclopaedia* 7: 383–429.
- Myers AA, Lowry JK (2003) A phylogeny and a new classification of the Corophiidea Leach, 1814 (Amphipoda). *Journal of Crustacean Biology* 23: 443–485. <https://doi.org/10.1163/20021975-99990353>
- Peart RA, Ah Yong ST (2016) Phylogenetic analysis of the family Ampithoidae Stebbing, 1899 (Crustacea: Amphipoda), with a synopsis of the genera. *Journal of Crustacean Biology* 36(4): 456–474. <https://doi.org/10.1163/1937240X-00002449>
- Shin MH, Hong JS, Kim W (2010) Redescriptions of two ampithoid amphipods, *Ampithoe lacertosa* and *A. tarasovi* (Crustacea: Amphipoda), from Korea. *The Korean Journal of Systematic Zoology* 26: 295–305. <https://doi.org/10.5635/KJSZ.2010.26.3.295>
- Shin MH, Coleman CO, Hong JS, Kim W (2015) A new species of *Peramphithoe* (Amphipoda: Ampithoidae) from South Korea, with morphological diagnoses of the world congeneric species. *Journal of Crustacean Biology* 35(2): 255–270. <https://doi.org/10.1163/1937240X-00002323>
- Smith SI (1873) Crustacea. In: Verrill AE (Ed.) Report upon the invertebrate animals of Vineyard Sound and the adjacent waters, with an account of the physical characters of the region. Report of the United States Commissioner of Fisheries 1: 545–580.
- Tzvetkova NL (1967) On the fauna and ecology of amphipods (Amphipoda, Gammaridea) of the Possjet Bay (the Sea of Japan). *Akademija Nauk SSSR, Zoologicheskii Institut, Issledovaniya Fauny Morei* 5: 160–195.