

Updated species checklist of fishes from Lake Dongting in Hunan Province, South China: Species diversity and conservation

Xiao Chen^{1,2}, Man Wang^{1,2}, E Zhang¹

1 *The Key Laboratory of Aquatic Biodiversity and Conservation, Institute of Hydrobiology, Chinese Academy of Sciences, Wuban, China* **2** *University of Chinese Academy of Sciences, Beijing, China*

Corresponding author: E Zhang (zhange@ihb.ac.cn)

Academic editor: Maria Elina Bichuette | Received 30 December 2021 | Accepted 20 May 2022 | Published 23 June 2022

<http://zoobank.org/A8AB836D-326C-4E1F-BBBD-13BF08ABE82B>

Citation: Chen X, Wang M, Zhang E (2022) Updated species checklist of fishes from Lake Dongting in Hunan Province, South China: Species diversity and conservation. ZooKeys 1108: 51–88. <https://doi.org/10.3897/zookeys.1108.79960>

Abstract

A lack of an updated checklist of freshwater fish species from Lake Dongting is a great hindrance to further biodiversity analysis. A seasonal survey of fishes in the lake was conducted from October 2017 to January 2019. Based on the data obtained during the field survey and coupled with known literature and the latest taxonomic development of relevant taxa, the species checklist of fishes from Lake Dongting was updated. A total of 130 species from 12 orders, 30 families and 76 genera has been documented, containing 126 native species and four alien species. Its fish fauna is dominated by the Xenocypridae that has the highest number of included species (30), followed by the Gobionidae (25) and Acheilognathidae (11). This checklist comprises 20 species undergoing nomenclatural changes and 11 new records, eight of which are native and three exotic. It excludes 20 species, which have been reported in error in historical works, due to synonyms, erroneous records, taxonomic changes and unconfirmed records. Unsourced in this survey were 34 species that are ecologically specialised: migratory, rheophilic, predatory, shellfish-dependent or pelagic-egg-spawning. While some of these species eluded capture likely due to the paucity of population, others may have been extirpated in Lake Dongting perhaps owing to human perturbations, such as river damming across affluents or the Chang-Jiang mainstem, sand dredging, overfishing or water pollution. The updated checklist lays a sound foundation for biodiversity conservation of fishes in Lake Dongting.

Keywords

Annotated list, biodiversity, ichthyofauna, taxonomy, threatened species

Introduction

Freshwater ecosystem and freshwater fish may well face one of the greatest threats in the world in the context of global biodiversity crisis (Dudgeon et al. 2006; Strayer and Dudgeon 2010). In comparison with other vertebrates, freshwater fishes are being more severely threatened by human interferences when the usage of water resources is strengthened (Reid et al. 2019; Barbarossa et al. 2021). The biodiversity conservation of freshwater fishes, nevertheless, has received disproportionate attention compared to terrestrial vertebrates (Tedesco et al. 2017). It is imperative and also in urgent need to protect the freshwater ecosystem and its biodiversity (Jackson et al. 2001; Liermann et al. 2012). Species inventories are beyond simply lists of names; they are actually representing an efficient method for obtaining updated information regarding species composition and distributions (Marta et al. 2019). This information provides valuable inputs of biodiversity monitoring which serves many conservational purposes, such as prioritising protection areas and directing conservation actions (Brooks et al. 2004). An update species inventory of a given area is of vital significance for biodiversity conservation.

The freshwater ecosystem of the Chang-Jiang (= Yangtze River; Jiang, Shui and He in Chinese mean river), the third largest river of the world and the largest river of China, supports rich biodiversity of aquatic organisms (Chen et al. 2020). The middle reaches of this river are regarded as one of the hotspots for freshwater fish diversity in Asia (Kottelat and Whitten 1996). Lake Dongting is one of two largest river-linked freshwater lakes in China, lying within the lower Chang-Jiang (= the mid-lower Chang-Jiang) basin which forms a freshwater ecoregion of the world for biodiversity conservation (Abell et al. 2008). This lake is an important portion of fluvio-lacustrine complex ecosystems of the mid-lower Chang-Jiang basin (Wang et al. 2019b), and also one of the priority areas for biodiversity conservation in China (Li et al. 2016a) and the Ramsar-listed floodplain wetlands (Dong et al. 2021), which serves as crucial habitats of migratory birds of East Asia-Australian flyway (Zou et al. 2019). Moreover, Lake Dongting, as a flood buffer zone, provides *Elaphurus davidianus* (Milu or Père David's deer) with seasonal sanctuaries (Yang et al. 2016). It also provides refuge for charismatic mammals like *Neophocaena asiaeorientalis* (Yangtze finless porpoise) (Zhang 2011; Huang et al. 2017) and the feeding grounds of large-sized flagship fishes, such as *Acipenser sinensis* (Chinises sturgeon) and *Psephurus gladius* (Chinese paddlefish) and economically-important potamodromous fishes like four major Chinese carps: *Aristichthys nobilis* (Bighead Carp), *Ctenopharyngodon idella* (Grass Carp), *Hypophthalmichthys molitrix* (Silver Carp) and *Mylopharyngodon piceus* (Black Carp) (Liu et al. 2010; Zhang et al. 2020a). Apparently, the lake plays a vital role in the conservation of aquatic biodiversity of the Chang-Jiang Basin.

The aquatic biodiversity of Lake Dongting is greatly imperilled by anthropogenic activities, like sand dredging, overfishing, alien species invasion, water pollution from industrial, agricultural and domestic sewage discharges and so forth (Dou and Jiang 2000; Fu et al. 2021; Jiang et al. 2022). It is also indirectly impacted by dam building across the Chang-Jiang mainstem and affluents of the lake owing to the continuity of the aquatic ecosystem (Wang et al. 2016; Liu and Wang 2018). Fishes, as top feed-

ers of the aquatic ecosystem and an important source of proteins in human food, are severely threatened by these factors (Zhao et al. 2019; Tregidgo et al. 2021). In the latest Red List assessment of Chinese freshwater fishes (Zhang and Cao 2021a), there are 12 imperilled species from Lake Dongting. *Psephurus gladius* (Martens, 1862) was recently declared to be extinct or functionally extinct (Zhang et al. 2020a). Such species as *Luciobrama macrocephalus* (Lacepède, 1803), *Ochetobius elongatus* (Kner, 1867) and *Tenuialosa reevesii* (Richardson, 1846) have been not seen in capture fisheries for decades or are likely extirpated (Wu et al. 2015; Zhang and Cao 2021b). The current status of freshwater fish diversity of Lake Dongting is, therefore, of particular conservation concern.

An updated checklist of fishes from Lake Dongting remains to be provided. The first checklist of freshwater fishes from this lake was given by Tang and Qian (1979), who recorded 114 fish species. A total of 117 species of the lake was later included in Anonymous' (1980) book entitled "Fish of Hunan Province". Subsequent species inventories of fishes from Lake Dongting primarily followed the book and three authoritative monographs of Chinese freshwater fishes authored by Chen (1998), Chu et al. (1999) and Yue (2000). Nevertheless, the species inventory of fishes from this lake needs to be regularly updated for biodiversity conservation, especially with lots of taxonomic revisions of freshwater fishes from the Chang-Jiang basin over the past decades. Therefore, three seasonal field sampling of fishes in Lake Dongting were conducted by us during 2017–2019. Coupled with the data collected in this survey, we aim to synthesise existing knowledge of freshwater fish diversity and systematics to provide an updated checklist of fish of the lake.

History of taxonomic research

The taxonomic history of fishes from Lake Dongting could be traced back to the mid-19th century. Père Heude, a French Jesuit catholic priest, made a collection of fish specimens at the lake from 1869 to 1884 (Luo 2005). Subsequently, Kreyenberg and Pappenheim (1908) reported 22 species from the Chang-Jiang and its tributaries, two of which were new species from Lake Dongting: *Coilia brachygnathus* Kreyenberg & Pappenheim, 1908 and *Culter oxycephaloides* Kreyenberg & Pappenheim, 1908. At the same time, Regan (1908) described three new Chinese species, two of which were *Glyptothorax sinensis* (Regan, 1908) and *Hemisalanx prognathus* Regan, 1908 from the lake. In 1921, Clifford Pope made a collection of fish specimens in Huping College, Yochow (= Yueyang City near East Lake Dongting) (Luo 2005). Nichols (1925a) proposed two new subspecies *Misgurnus anguillicaudatus tungting* Nichols, 1925 and *Misgurnus mohoity leopardus* Nichols, 1925 from Lake Dongting, both being regarded as invalid to date. Nichols (1925b) recorded three species of *Botia* Gray, 1831 from this lake, two of which were new to sciences, namely *B. citrauratea* Nichols, 1925 and *B. purpurea* Nichols, 1925. Both are now placed in *Leptobotia* Bleeker, 1870, but the

latter has been synonymised with *L. taeniops* (Sauvage, 1878). Simultaneously, Nichols (1925c-e) described five new species from Lake Dongting, viz. *Gobio longipinnis* (= *Rhinogobio ventralis* Sauvage & Dabry de Thiersant, 1874), *Gobius cliffordpopei* [= *Rhinogobius cliffordpopei* (Nichols, 1925)], *Hemiculterella engraulis* [= *Pseudolaubuca engraulis* (Nichols, 1925)], *Hemiculter clupeioides* [= *Hemiculter leucisculus* (Basilewsky, 1855)] and *Varicorhnus tungting* [= *Decorus tungting* (Nichols, 1925)]. Later, Nichols and his co-authors (1926, 1927) named a new subspecies *Sarcocheilichthys nigripinnis tungting* Nichols & Pope, 1927 and two new species, i.e. *Acheilognathus gracilis* Nichols, 1926 and *Pseudogobio tungtingensis* [= *Microphysogobio tungtingensis* (Nichols, 1926)] from this lake. Nichols (1928) recorded 71 nominal species from Lake Dongting in his provisional checklist of Chinese freshwater fishes. Subsequent taxonomic contributions to fishes of Lake Dongting were also made by many authors, such as Wu (1930), Tchang (1933) and Kimura (1934), who made a small collection of fish specimens in the lake. Chu (1931) recorded 74 fish species from Hunan Province, the majority of which were from Lake Dongting. Hora (1932) described a new species *Lepturichthys nicholsi* [= *Lepturichthys fimbriatus* (Günther, 1888)]. Nichols (1943), in his book entitled “The fresh-water fishes of China”, recognised 79 species or subspecies for fish specimens collected by Clifford Pope in East Dongting Lake in 1921.

More studies were focused on the species inventory of fishes from Lake Dongting following the establishment of P. R. China in 1949. Forty-three species of the lake were involved in Chu’s (1955) study on the distribution of fish species in the Yichang section of the Chang-Jiang. Liang and Liu (1959) compiled a species list of 69 fishes from Lake Dongting and its affluent (Xiang-Jiang). Later, Liang and Liu (1966), in their checklist of fishes from Hunan Province, recorded 119 species from Lake Dongting. Anonymous (1976) also reported 84 species from Lake Dongting in the book entitled “Fishes of the Chang-Jiang”. Tang and Qian (1979) were the first to provide a checklist of 114 fish species or subspecies from the lake. Anonymous (1980) included 117 species from Lake Dongting in the book entitled “Fishes of Hunan Province”. Although Dou and Jiang (2000) compiled a checklist of 104 species from the lake, this work was mainly based on their collections of fish specimens made during 1974–1975.

As from the 1990s, increasing research interests have centred on the fish diversity of Lake Dongting. Survey of fishery resources carried out by Liao et al. (2002), Liao et al. (2006) and Li (2006) into Lake Dongting from 1994 to 2005 found 117, 111 and 117 species, respectively. Ru and Liu (2013) identified 69 species in their surveys conducted into East and South Dongting Lake during 2004–2005. Li (2013) reported a total of 85 fish species, based on his field sampling from March to December in 2012. Eighty species were identified by Jiang et al. (2019) in their research on the spatio-temporal patterns of fish assemblages in Lake Dongting from 2012 to 2014. Sixty-two fish species were recorded by Qin et al. (2019) from the outlet channel of the lake from 2013 to 2015. Eighty-five and 66 fish species were sampled during 2002–2003 and 2012–2014 field surveys to monitor the changes of fish community structure at West Dongting Lake before and after the operation of Three Gorges Dam (Zhu et al. 2014). All these inventories were conducted, particularly in relation to environment impact assessment prepared for hydropower projects or fisheries investigations and the data

collection used for biodiversity analyses. Some surveys were focused on fish resources assessment; small-sized or less commercially valuable species were largely neglected. Others were not based on examination of collected specimens, but compiled through desk review or interview, containing little or no reliable information on fish diversity, although they claimed to have studied biodiversity. More importantly, these inventories required critical scrutiny from an ichthyological perspective as most of them, if not all, were not conducted by trained ichthyologists in the field; information on unrecognised species is impossibly captured through these surveys, therefore giving rise to a grossly underestimated biodiversity value and taxonomic impediment (Sluys 2013).

Material and method

Lake Dongting (28°44'N–29°35'N, 111°53'E–113°05'E) is located in the northern part of Hunan Province, connected to the middle Chang-Jiang mainstem (Wang and Dou 1998). This water-carrying floodplain lake receives not only runoff waters from four main affluents (Xiang-Jiang, Zi-Shui, Yuan-Jiang and Li-Shui), but flood water from the Chang-Jiang mainstem via three inlet channels (Songzi, Hudu and Ouchi Channel) and lake water then flows out into the mainstem of the river again via Chenglingji Channel (Dou and Jiang 2000). Generally, Lake Dongting is divided into three sub-lakes, i.e. East Dongting Lake, South Dongting Lake and West Dongting Lake (Zhao et al. 2005). The lake covers a surface area of 2625 km² at a water level of 33.5 m at Chenglingji Station (Dou and Jiang 2000).

Twenty sampling sites were selected in this study, based on habitat heterogeneity (Fig. 1; Suppl. material 1: Table S1). Field sampling was taken from October to November 2017, July to August 2018 and December 2018 to January 2019. Fish surveys were conducted in different types of habitats to ensure maximum representation of species diversity occurring in this area. Multiple sampling methods were thus applied. Three-layer gill nets were used for pelagic fish sampling, while trap nets were applied to catch demersal fishes. Additionally, fish specimens were collected from local fish markets and their sampling localities were restricted to Lake Dongting.

All collected specimens were identified to species level. The initial identification in the field principally followed Chen (1998), Chu et al. (1999) and Yue (2000). The caught specimens were fixed in 95% alcohol solution for molecular analysis, in general and DNA extraction, in particular or initially fixed in 10% formalin and then transferred to 70% ethanol for morphological examination and also for permanent collection. All specimens were deposited in the ichthyological collection of the Museum of Aquatic Organisms at the Institute of Hydrobiology, Chinese Academy of Sciences.

In addition to the data collected during our field sampling, known research works were referred. Reference was made to the following main historical records of fishes in the lake: Liang and Liu (1959, 1966), Tang and Qian (1979), Anonymous (1980), Li (2006), Ru (2008), Cao et al. (2012) and Li (2013).

Fish classifications are being transformed greatly as latest molecular phylogenies provide evidence in support for natural groups which were unanticipated by previous




Figure 1. Field sampling sites of Lake Dongting in this study.

studies (Betancur-R et al. 2017). For the order Cypriniformes, the dominant group of freshwater fishes worldwide, significant advances have been made in its familial-level classification; some new familial (Acheilognathidae, Gobionidae and Xenocyprididae) and subfamilial names (Acrossocheilinae and Spinibarbinae) have been proposed (Tan and Armbruster 2018). Except for Cypriniformes, the taxonomic revisions of other orders were referred from Van Der Laan et al. (2014). The species checklist of fishes in Lake Dongting was systematically arranged by order, family and subfamily in accordance with the latest developments made in the taxonomic ranks (Van Der Laan et al. 2014).

Table 1. Annotated checklist of the fish fauna from Lake Dongting. The species under each family or subfamily are sorted by alphabetical order. Notes are labelled with taxonomic alteration, synonymisation, misidentification and other meanings.

Valid species name	Previous studies	Note
Acipenseriformes		
Acipenseridae		
001 <i>Acipenser dabryanus</i> Duméril, 1869		○P
002 <i>Acipenser sinensis</i> Gray, 1835		○D
Polyodontidae		
003 <i>Psephurus gladius</i> (Martens, 1862)		○D
Anguilliformes		
Anguillidae		
004 <i>Anguilla japonica</i> Temminck & Schlegel, 1846		⊕D
Clupeiformes		
Clupeidae		
005 <i>Tenualosa reevesii</i> (Richardson, 1846)		○D
Engraulidae		
006 <i>Coilia brachygnathus</i> Kreyenberg & Pappenheim, 1908		⊕P
007 <i>Coilia nasus</i> Temminck & Schlegel, 1846		○D
Cypriniformes		
Catostomidae		
008 <i>Myxocyprinus asiaticus</i> (Bleeker, 1864)		⊕P
Botiidae		
009 <i>Leptobotia citraurata</i> (Nichols, 1925)		⊕P
	<i>Leptobotia elongata</i> (Bleeker, 1870)	⊖M
010 <i>Leptobotia rubrilabris</i> (Dabry de Thiersant, 1872)		○P
011 <i>Leptobotia taeniops</i> (Sauvage, 1878)		⊕P
012 <i>Parabotia banarescui</i> (Nalbant, 1965)		⊕P
013 <i>Parabotia fasciata</i> Dabry de Thiersant, 1872		⊕P
Cobitidae		
014 <i>Cobitis macrostigma</i> Dabry de Thiersant, 1872		⊕P
015 <i>Cobitis sinensis</i> Sauvage & Dabry de Thiersant, 1874		⊕P
016 <i>Misgurnus anguillicaudatus</i> (Cantor, 1842)		⊕P
017 <i>Paramisgurnus dabryanus</i> Dabry de Thiersant, 1872		⊕P
Balitoridae		
018 <i>Lepturichthys fimbriatus</i> (Günther, 1888)		○P
Cyprinidae		
Labeoninae		
019 <i>Cirrhinus cirrhosus</i> Bloch, 1795		+AP
020 <i>Cirrhinus molitorella</i> (Valenciennes, 1844)		⊕AP
021 <i>Decorus tungting</i> (Nichols, 1925)	<i>Bangana tungting</i> (Nichols, 1925)	○TP
022 <i>Pseudogyrinocheilus prochilus</i> (Sauvage & Dabry de Thiersant, 1874)		+P
Cyprininae		
023 <i>Carassius auratus</i> (Linnaeus, 1758)		⊕P
	<i>Cyprinus carpio</i> Linnaeus, 1758	⊖T
024 <i>Cyprinus rubrofasciatus</i> Lacepède, 1803		⊕P
	<i>Procypris nabaudi</i> (Tchang, 1930)	⊖M
Acrossocheilinae		
025 <i>Onychostoma rarum</i> (Lin, 1933)		○P
026 <i>Onychostoma simum</i> (Sauvage & Dabry de Thiersant, 1874)		○P
Spinibarbininae		
027 <i>Spinibarbus caldwelli</i> (Nichols, 1925)		○P

Valid species name	Previous studies	Note
	<i>Spinibarbus hollandi</i> Oshima, 1919	⊙T
	<i>Spinibarbus sinensis</i> Bleeker, 1871	⊙M
Xenocypridae		
028 <i>Aristichthys nobilis</i> (Richardson, 1845)		⊕P
029 <i>Chanodichthys dabryi</i> (Bleeker, 1871)	<i>Culter dabryi</i> Bleeker, 1871	⊕TP
030 <i>Chanodichthys erythropterus</i> (Basilewsky, 1855)	<i>Culter alburnus</i> Basilewsky, 1855	⊕TP
031 <i>Chanodichthys mongolicus</i> (Basilewsky, 1855)	<i>Culter mongolicus</i> Basilewsky, 1855	⊕TP
032 <i>Chanodichthys oxycephalus</i> (Bleeker, 1871)	<i>Culter oxycephalus</i> Bleeker, 1871	○TP
033 <i>Chanodichthys oxycephaloides</i> (Kreyenberg & Pappenheim, 1908)	<i>Culter oxycephaloides</i> Kreyenberg & Pappenheim, 1908	⊕TP
034 <i>Culter alburnus</i> (Basilewsky, 1855)	<i>Cultrichthys erythropterus</i> (Basilewsky, 1855)	⊕TP
035 <i>Ctenopharyngodon idella</i> (Valenciennes, 1844)		⊕P
036 <i>Distoechodon tumirostris</i> Peters, 1881		○P
037 <i>Elopichthys bambusa</i> (Richardson, 1845)		⊕P
038 <i>Hemiculter bleekeri</i> Warpachowski, 1888		⊕P
039 <i>Hemiculter leucisculus</i> (Basilewsky, 1855)		⊕P
040 <i>Hypophthalmichthys molitrix</i> (Valenciennes, 1844)		⊕P
041 <i>Luciobrama macrocephalus</i> (Lacepède, 1803)		○P
042 <i>Megalobrama amblycephala</i> Yih, 1955		⊕P
043 <i>Megalobrama mantschuricus</i> (Basilewsky, 1855)	<i>Megalobrama skolkovii</i> Dybowski, 1872	⊕SP
044 <i>Mylopharyngodon piceus</i> (Richardson, 1846)		⊕P
045 <i>Ochetobius elongatus</i> (Kner, 1867)		○P
046 <i>Opsariichthys bidens</i> Günther, 1873		⊕P
047 <i>Parabramis pekinensis</i> (Basilewsky, 1855)		⊕P
048 <i>Plagiognathops microlepis</i> (Bleeker, 1871)	<i>Xenocypris microlepis</i> Bleeker, 1871	○TP
049 <i>Pseudobrama simoni</i> (Bleeker, 1864)		⊕P
050 <i>Pseudolaubuca engraulis</i> (Nichols, 1925)		○P
051 <i>Pseudolaubuca sinensis</i> Bleeker, 1864		⊕P
052 <i>Sinibrama macrops</i> (Günther, 1868)		⊕P
	<i>Sinibrama wui</i> (Rendahl, 1933)	⊙S
053 <i>Squaliobarbus curriculus</i> (Richardson, 1846)		⊕P
054 <i>Toxabramis swinhonis</i> Günther, 1873		⊕P
055 <i>Xenocypris davidi</i> Bleeker, 1871		⊕P
056 <i>Xenocypris macrolepis</i> Bleeker, 1871	<i>Xenocypris argentea</i> Günther, 1868	⊕SP
057 <i>Zacco acanthogenys</i> (Boulenger, 1901)		⊕P
	<i>Zacco platypus</i> (Temminck & Schlegel, 1846)	⊙T
Acheilognathidae		
058 <i>Acheilognathus barbatulus</i> Günther, 1873		+P
059 <i>Acheilognathus barbatus</i> Nichols, 1926		○P
060 <i>Acheilognathus chankaensis</i> (Dybowski, 1872)		○P
061 <i>Acheilognathus gracilis</i> Nichols, 1926		⊕P
	<i>Acheilognathus imberbis</i> Günther, 1868	⊙U
062 <i>Acheilognathus hypselonotus</i> (Bleeker, 1871)		○P
063 <i>Acheilognathus macromandibularis</i> Doi, Arai & Liu, 1999		+P
064 <i>Acheilognathus macropterus</i> (Bleeker, 1871)		⊕P
065 <i>Acheilognathus polylepis</i> (Wu, 1964)		⊕P
066 <i>Acheilognathus tonkinensis</i> (Vaillant, 1892)		○P
	<i>Acheilognathus taenianalis</i> (Günther, 1873)	⊙S
067 <i>Rhodeus ocellatus</i> (Kner, 1866)		⊕P
068 <i>Rhodeus sinensis</i> Günther, 1868		⊕P
Gobionidae		
069 <i>Abbottina rivularis</i> (Basilewsky, 1855)		⊕P
070 <i>Coreius heterodon</i> (Bleeker, 1864)		⊕P

Valid species name	Previous studies	Note
	<i>Coreius guichenoti</i> (Sauvage & Dabry de Thiersant, 1874)	⊙M
071 <i>Gobiobotia filifer</i> (Garman, 1912)		⊕P
072 <i>Gobiobotia meridionalis</i> Chen & Cao, 1977	<i>Gobiobotia longibarba meridionalis</i> Chen & Cao, 1977	⊕TP
073 <i>Gobiobotia nicholsi</i> Bănărescu & Nalbant, 1966		○P
074 <i>Gobiobotia lii</i> Chen, Wang, Cao & Zhang, 2022		+P
	<i>Gobiobotia pappenbeimi</i> Kreyenberg, 1911	⊙M
	<i>Xenophysogobio boulengeri</i> (Tchang, 1929)	⊙M
075 <i>Hemibarbus labeo</i> (Pallas, 1776)		⊕P
076 <i>Hemibarbus maculatus</i> Bleeker, 1871		⊕P
077 <i>Microphysogobio tungtingensis</i> (Nichols, 1926)		⊕P
078 <i>Paracanthobrama guichenoti</i> Bleeker, 1864		⊕P
079 <i>Pseudogobio vaillanti</i> (Sauvage, 1878)		+P
080 <i>Pseudorasbora parva</i> (Temminck & Schlegel, 1846)		⊕P
081 <i>Rhinogobio cylindricus</i> Günther, 1888		○P
082 <i>Rhinogobio typus</i> Bleeker, 1871		⊕P
083 <i>Rhinogobio ventralis</i> Sauvage & Dabry de Thiersant, 1874		○P
	<i>Sarcocheilichthys kiangsiensis</i> Nichols, 1930	⊙T
084 <i>Sarcocheilichthys nigripinnis</i> (Günther, 1873)		⊕P
085 <i>Sarcocheilichthys tungtingensis</i> Nichols & Pope, 1927		○P
086 <i>Sarcocheilichthys sinensis</i> Bleeker, 1871		⊕P
087 <i>Saurogobio dabryi</i> Bleeker, 1871		⊕P
088 <i>Saurogobio dumerili</i> Bleeker, 1871		○P
089 <i>Saurogobio gymnocheilus</i> Lo, Yao & Chen, 1998		⊕P
090 <i>Saurogobio gracilicaudatus</i> Yao & Yang, 1977		+P
091 <i>Saurogobio lissilabris</i> Bănărescu & Nalbant, 1973		⊕P
092 <i>Saurogobio xiangiangensis</i> Tang, 1980		+P
093 <i>Squalidus argentatus</i> (Sauvage & Dabry de Thiersant, 1874)		⊕P
	<i>Squalidus nitens</i> (Günther, 1873)	⊙U
Siluriformes		
Bagridae		
094 <i>Hemibagrus macropterus</i> Bleeker, 1870	<i>Mystus macropterus</i> (Bleeker, 1870)	⊕P
095 <i>Tachysurus crassilabris</i> (Günther, 1864)	<i>Leiocassis crassilabris</i> Günther, 1864	⊕TP
096 <i>Tachysurus dumerili</i> (Bleeker, 1864)	<i>Leiocassis longirostris</i> Günther, 1864	⊕TP
097 <i>Tachysurus eupogon</i> (Boulenger, 1892)	<i>Pelteobagrus eupogon</i> (Boulenger, 1892)	⊕TP
098 <i>Tachysurus mica</i> (Gromov, 1970)	<i>Leiocassis argentivittatus</i> (Regan, 1905)	+TP
099 <i>Tachysurus nitidus</i> (Sauvage & Dabry de Thiersant, 1874)	<i>Pelteobagrus nitidus</i> (Sauvage & Dabry de Thiersant, 1874)	⊕TP
100 <i>Tachysurus sinensis</i> (Lacepède, 1803)	<i>Pelteobagrus fulvidraco</i> (Richardson, 1846)	⊕TP
	<i>Tachysurus tenuis</i> (Günther, 1873)	⊙M
101 <i>Tachysurus ussuriensis</i> (Dybowski, 1872)	<i>Pseudobagrus ussuriensis</i> (Dybowski, 1872)	○TP
102 <i>Tachysurus vachellii</i> (Richardson, 1846)	<i>Pelteobagrus vachellii</i> (Richardson, 1846)	⊕TP
103 <i>Tachysurus zhangfei</i> Shao, Cheng & Zhang, 2021	<i>Pseudobagrus albomarginatus</i> (Rendahl, 1928)	⊕TP
Amblycipitidae		
104 <i>Liobagrus aequilabris</i> Wright & Ng, 2008		⊕P
Sisoridae		
105 <i>Glyptothorax sinensis</i> (Regan, 1908)		⊕P
Siluridae		
106 <i>Silurus asotus</i> Linnaeus, 1758		⊕P
107 <i>Silurus meridionalis</i> Chen, 1977		⊕P
Ictaluridae		
108 <i>Ictalurus punctatus</i> (Rafinesque, 1818)		+AP

Valid species name	Previous studies	Note
Osmeriformes		
Salangidae		
109 <i>Hemisanalx prognathus</i> Regan, 1908		⊕F
	<i>Hemisanalx brachyrostralis</i> (Fang, 1934)	⊙S
110 <i>Neosalanx brevirostris</i> (Pellegrin, 1923)		○F
	<i>Neosalanx taihuensis</i> Chen, 1956	⊙S
111 <i>Neosalanx jordani</i> Wakiya & Takahashi, 1937		○F
	<i>Neosalanx oligodontis</i> Chen, 1956	⊙S
112 <i>Protosalanx hyalocranius</i> (Abbott, 1901)		○F
Gobiiformes		
Odontobutidae		
113 <i>Micropercops cinctus</i> (Dabry de Thiersant, 1872)	<i>Micropercops swinhonis</i> (Günther, 1873)	⊕SV
114 <i>Odontobutis sinensis</i> Wu, Chen & Chong, 2002		⊕V
Gobiidae		
115 <i>Mugilogobius myxodermus</i> (Herre, 1935)		○V
	<i>Rhinogobius brunneus</i> (Temminck & Schlegel, 1845)	⊙U
116 <i>Rhinogobius cliffordpopei</i> (Nichols, 1925)		○V
117 <i>Rhinogobius similis</i> Gill, 1859		⊕V
	<i>Rhinogobius giurinus</i> Gill, 1859	⊙S
Synbranchiformes		
Mastacembelidae		
118 <i>Sinobdella sinensis</i> (Bleeker, 1870)		⊕P
Synbranchidae		
119 <i>Monopterus albus</i> (Zuiew, 1793)		⊕P
Anabantiformes		
Osphronemidae		
120 <i>Macropodus opercularis</i> (Linnaeus, 1758)		⊕P
Channidae		
121 <i>Channa argus</i> (Cantor, 1842)		⊕P
122 <i>Channa asiatica</i> (Linnaeus, 1758)		⊕P
Beloniformes		
Hemiramphidae		
123 <i>Hyporhamphus intermedius</i> (Cantor, 1842)		⊕V
Adrianichthyidae		
124 <i>Oryzias latipes</i> (Temminck & Schlegel, 1846)		○F
Tetraodontiformes		
Tetraodontidae		
125 <i>Takifugu obscurus</i> (Abe, 1949)		○D
Centrarchiformes		
Centrarchidae		
126 <i>Micropterus salmoides</i> (Lacepède, 1802)		+AP
Sinipercidae		
127 <i>Siniperca chuatsi</i> (Basilewsky, 1855)		⊕P
128 <i>Siniperca kneri</i> Garman, 1912		⊕P
129 <i>Siniperca roulei</i> Wu, 1930	<i>Coreosiniperca roulei</i> (Wu, 1930)	⊕TP
130 <i>Siniperca scherzeri</i> Steindachner, 1892		⊕P

Note: ⊕ Historically recorded species caught in this study; ○ Historically recorded species uncollected in this study; ⊙ Historically recorded species excluded from the updated checklist; + Newly-recorded species; A-Alien or introduced species; S-Junior synonym species; M-Previously misidentified species; T-Taxonomically altered species; U-Unconfirmed species; P-Primary freshwater species; F-Secondary freshwater species; D-Diadromous species; V-Vicarious species.

Results

Analysis of the species checklist

A total of 130 fish species, identified from 12 orders, 30 families and 76 genera, have been documented from Lake Dongting (Table 1). Amongst these species, there are 126 native and four exotic species. Ninety-six species from 10 orders, 24 families and 61 genera collected during 2017–2019 fish survey are included.

For species richness, the order representing the greatest number of species were Cypriniformes (86 species, 66.15% of the total), followed by the Siluriformes (15, 11.54%), Centrarchiformes (5, 3.85%), Gobiiformes (5, 3.85%), Osmeriformes (4, 3.08%), Anabantiformes (3, 2.31%), Clupeiformes (3, 2.31%), Acipenseriformes (3, 2.31%), Synbranchiformes (2, 1.54%), Beloniformes (2, 1.54%), Tetrodontiformes (1, 0.77%) and Anguilliformes (1, 0.77%). The family Xenocyprididae has the highest number (30) of fish species, accounting for 23.08% of the total, followed by the Gobionidae and Acheilognathidae, with 25 and 11 species contributing to 19.23% and 8.46%, respectively. The subsequent families included the Bagridae (10, 7.69%), Cyprinidae (9, 6.92%) and so forth (Table 2).

Lake Dongting harboured 27 migratory fishes, six of which are diadromous and 21 potamodromous, and 103 sedentary fishes, accounting for 20.77% and 79.23% of the total freshwater fishes, respectively. There are 113 (86.92% of the total species) primary freshwater fishes (species spending the whole life in freshwater; Kottelat et al. (2012)), five (3.85%) secondary freshwater fishes (species related to marine families, but living in fresh or sometimes brackish water), six (4.62%) diadromous species (species migrating between fresh and brackish water, but staying in freshwater for part of their life), six (4.62%) vicarious species (species of otherwise largely marine families, but spending their whole life in freshwater, for example, some gobies species) (See Table 1).

The updated checklist of fishes from Lake Dongting includes 49 species endemic to China, 22 endemic to the Chang-Jiang and nine endemic to the mid-lower Chang-Jiang, respectively. This survey yielded 35 Chinese endemics (accounting for 71.43% of the total Chinese endemics from Lake Dongting), 13 endemic species of the Chang-Jiang (59.09% of the total endemic species of the river from the lake) and six endemic species of the mid-lower Chang-Jiang (66.67% of the total endemic species of these reaches from the lake), respectively.

Annotated species checklist

The updated checklist of fishes in Lake Dongting recognises a total of 130 species, based on the data collected in this survey and historical records. Amongst them, 93 native fish species were observed in this fish survey, including eight new records (See Table 1; note '+'). Thirty-four historically recorded species, unsampled in this field surveys, are contained in the updated checklist (Note 'o'). Other 20 historically re-

Table 2. The taxonomic composition of fish species in Lake Dongting.

Order	Family	Genus	Species
Acipenseriformes	Acipenseridae	1	2
	Polyodontidae	1	1
Anguilliformes	Anguillidae	1	1
Clupeiformes	Clupeidae	1	1
	Engraulidae	1	2
Cypriniformes	Catostomidae	1	1
	Botiidae	2	5
	Cobitidae	3	4
	Balitoridae	1	1
	Cyprinidae	7	9
	Xenocyprididae	22	30
	Acheilognathidae	2	11
	Gobionidae	12	25
Siluriformes	Bagridae	2	10
	Amblycipitidae	1	1
	Sisoridae	1	1
	Siluridae	1	2
	Ictaluridae	1	1
Osmeriformes	Salangidae	3	4
Gobiiformes	Odontobutidae	2	2
	Gobiidae	1	3
Synbranchiformes	Mastacembelidae	1	1
	Sybranchidae	1	1
Anabantiformes	Osphronemidae	1	1
	Channidae	1	2
Beloniformes	Hemiramphidae	1	1
	Adrianichthyidae	1	1
Tetraodontiformes	Tetraodontidae	1	1
Centrarchiformes	Centrarchidae	1	1
	Sinipercidae	1	4
12	30	76	130

corded species are excluded, including six being synonymised with other species (Note ‘S’), seven misidentified or having an erroneous record in the lake (Note ‘M’), four experiencing taxonomical changes (Note ‘T’) and three having unconfirmed records (Note ‘U’). A number of nomenclatural changes also occur for valid species included in the updated checklist (20 species). Taxonomic comments were appended to discuss its validity and occurrence where relevant.

Acipenseridae & Polyodontidae

The Acipenseridae has two representatives in the lake, namely *Acipenser sinensis* Gray, 1835 and *A. dabryanus* Duméril, 1869, while the Polyodontidae is presented only by a single species *Psephurus gladius*. All three large-sized sturgeons were not collected in Lake Dongting during this field survey. One juvenile individual (4340 mm SL, 566.0 g) of *A. sinensis* was collected from East Dongting Lake during the 2012–2013 field sur-

vey (unpublished data). The specimen is likely a captive-bred individual released into the wild. This conservation measure has been implemented in the upper Chang-Jiang Basin for nearly twenty years (Du et al. 2013). Nichols (1928) was the first to report on the existence of *A. dabryanus* in Lake Dongting. Liang and Liu (1959, 1966) included the sturgeon in their species inventories of the lake. This species has vanished in Lake Dongting since the Gezhouba Dam was constructed across the Chang-Jiang mainstem (Zhang and Cao 2021a). Likely, *A. dabryanus* became highly depleted as no individuals have been collected in the river as from 1995 (Zhang et al. 2017). The capture record on *P. gladius* showed a similar trend to that of *A. dabryanus* as no records of Chinese paddlefish have been reported since 1995 (Zhang et al. 2020a).

Anguillidae

Anguilla japonica Temminck & Schlegel, 1846, a delicious food fish of economic importance in China and even across the Globe, is the only representative of the family in Lake Dongting. Historically, the lake and its affluents were utilised by this catadromous fish as feeding grounds (Anonymous 1980), but it is hardly seen in fish capture presently (Liu et al. 2013). One small individual (295 mm SL, 33.9 g) was captured at Chenglingji Channel during this field survey. It might be an individual which escaped from reservoirs where cage culture was used to farm this fish, in terms of local fishermen.

Clupeidae & Tetraodontidae

The family Clupeidae and Tetraodontidae are each represented in Lake Dongting by a single species. The two diadromous fishes, *Tenuulosa reevesii* and *Takifugu obscurus* (Abe, 1949), are hardly seen in this lake so far. The last capture of *T. reevesii* (one individual) was at Jiangsu provincial section of the Chang-Jiang in 1998 (Liu et al. 2002). *Takifugu obscurus* is occasionally encountered in the lower Chang-Jiang Basin so far (Wang et al. 2016; Chen et al. 2020).

Engraulidae

This family has only two representatives in Lake Dongting: *Coilia brachygnathus* and *C. nasus* Temminck & Schlegel, 1846. So far, *C. brachygnathus* abounds in this lake where it is a delicious food fish of economic importance, but *C. nasus* is a rarely encountered fish. *Coilia nasus* is even regarded to have been extinct due to anthropogenic interferences for nearly two decades (Wang et al. 2016); however, this anadromous fish has recently been found to persist in Lake Dongting (Xuan et al. 2020).

Salangidae

This family has four representatives in Lake Dongting: *Hemisalanx prognathus*, *Neosalanx brevirostris* (Pellegrin, 1923), *N. jordani* Wakiya & Takahashi, 1937 and *Protosalanx hyalocranius* (Abbott, 1901). So far, the taxonomy of Chinese icefishes

still remains controversial (Fu et al. 2005; Zhang et al. 2007). Based on the latest taxonomic advances of this family, three formerly recorded species were removed from the updated checklist. *Neosalanx taihuensis* Chen, 1956 was treated as a synonym of *N. brevirostris* (Zhang et al. 2007). *Hemisanx brachyrostralis* (Fang, 1934) was synonymised with *H. prognathous* and so was *Neosalanx oligodontis* Chen, 1956 with *N. jordani* (Guo et al. 2011).

Catostomidae

This family has a single representative in China: *Myxocyprinus asiaticus* (Bleeker, 1864). *Myxocyprinus asiaticus* had long been considered as a migratory fish (Anonymous 1976). Nevertheless, Zhang and Zhao's (2001) examination on collection specimens found that all individuals caught from the mid-lower Chang-Jiang Basin were small-sized, but large-sized individuals came from the upper reaches of this river, so concluding that *M. asiaticus* may be not a migratory species. One specimen (382 mm SL and 675.1 g) of this species was captured at the estuary of the Xiang-Jiang into Lake Dongting during our field survey. It is probably a captive-bred individual released into the Xiang-Jiang at Hengyang section yearly, according to local fishermen.

Cyprinidae

The Cyprinidae, as traditionally delimited, contains species with one to three rows of pharyngeal teeth, barbels present or absent and Weberian apparatus (Chen 1998; Nelson et al. 2016). A recent re-classification of the Cypriniformes was provided by Tan and Armbruster (2018), based on Yang et al.'s (2015a) phylogenetic relationships of this order inferred from both mitochondrial and nuclear genes. The Cyprinidae *s. l.* splits into ten families, namely Acheilognathidae, Cyprinidae *s. str.*, Danionidae, Gobionidae, Leptobarbidae, Leuciscidae, Sundadanionidae, Tanichthyidae, Tincidae and Xenocyprididae. The Cyprinidae *s. str.* is further subdivided into eleven subfamilies, i.e. Acrossocheilinae, Barbinae, Cyprininae, Labeoninae, Poropuntiinae, Probarbinae, Schizopygopsinae, Schizothoracinae, Smiliogastrinae, Spinibarbinae and Torinae. Amongst them, four subfamilies have their representatives in this lake: Acrossocheilinae, Cyprininae, Labeoninae and Spinibarbinae.

Acrossocheilinae

This subfamily was newly erected to include species currently designated to *Folifer*, *Onychostoma* and *Acrossocheilus* (Yang et al. 2015a; Tan and Armbruster 2018), but its generic classification still needs in-depth study. According to historical records (Tang and Qian 1979; Li 2006), *Onychostoma* has two representatives in Lake Dongting: *Onychostoma simum* (Sauvage & Dabry de Thiersant, 1874) and *O. rarum* (Lin, 1933). The two rheophilic fishes were not caught in this survey, though.

Cyprininae

Previously, *Cyprinus carpio* Linnaeus, 1758 was extensively utilised as the available specific name for the common carp widespread in China. This species, however, is currently regarded as the endemic species of Europe (Kottelat and Freyhof 2007). The East Asian populations of the common carp represent a distinct species from *C. carpio*. The available specific name for it is *C. haematopterus* Temminck & Schlegel, 1846 (Zhou et al. 2003), a junior synonym of *C. rubrofasciatus* Lacepède, 1803 (Kottelat 2006, 2013). Specimens previously reported by Li (2006) as *Procypris rabaudi* (Tchang, 1930) from Lake Dongting are possibly misidentified. This species is a rheophilic fish usually found in headwaters of rivers, but not in the lentic environment (Zhang and Zhao 2016).

Labeoninae

This subfamily has four representatives in Lake Dongting: *Decorus tungting*, *Cirrhinus cirrhosus* Bloch, 1795, *C. molitorella* (Valenciennes, 1844) and *Pseudogyrinocheilus prochilus* (Sauvage & Dabry de Thiersant, 1874). The first species were firstly designated to *Sinilabeo* Rendahl, 1933 (Wu 1977; Tang et al. 2001) and later moved into *Bangana* Hamilton, 1822 (Zhang and Chen 2006). Recently, Zheng et al. (2019) assigned this species, along with *Bangana decora* (Peters, 1881) from the Zhu-Jiang Basin, *B. lemasoni* (Pellegrin & Chevey, 1936) from the Red River Basin and *B. rendahli* (Kimura, 1934) from the upper Chang-Jiang Basin, to their own genus named as *Decorus*. Both *Cirrhinus cirrhosus* and *Pseudogyrinocheilus prochilus* are two new records of this lake. The former was introduced into China as cultured fish from India during the 1990s (Wang and Zhang 2021); it, like *C. molitorella* in southern China, has widely been farmed as food fish for cultured Mandarin fish (Ye et al. 2016). Individuals of two *Cirrhinus* fishes, caught from Lake Dongting in this field survey, probably escaped from farming waters. The latter *P. prochilus* was collected at Chenglingji, the outlet channel from Lake Dongting into the Chang-Jiang mainstem. This means that the species has an extended distribution in this lake. The rheophilic fish is mainly found in the upper Chang-Jiang and Zhu-Jiang presently (Zhang 1994; Zheng et al. 2010). It was even recorded from the Li-Shui, Yuan-Jiang and Xiang-Jiang (Liang and Liu 1966; Anonymous 1980; Cao et al. 2012).

Spinibarbinae

The subfamily is represented in Lake Dongting by a single species: *Spinibarbus caldwelli* (Nichols, 1925). *Spinibarbus caldwelli* was previously recognised as *S. hollandi* Oshima, 1919 (Chu and Chen 1989; Yue 2000), a species widespread in Asian mainland (Tang et al. 2005). Indeed, *S. hollandi* is endemic to Taiwan Island of China (Tang et al. 2005). The available scientific name for Asian mainland specimens of this species is *S. caldwelli* (Tang et al. 2005). *Spinibarbus sinensis* Bleeker, 1871 is a species mainly found in the upper Chang-Jiang Basin (Chu and Chen 1989; Ding 1994; Yue

2000; Zhang and Zhao 2016). It was also reported from Lake Dongting by Tang and Qian (1979) and Li (2006). This identification, nevertheless, needs confirmation when specimens become available.

Xenocyprididae

The family is the dominant group of the ichthyofauna of Lake Dongting, with 30 species identified from 22 genera: *Aristichthys* Oshima, 1919 (one species), *Chanodichthys* Bleeker, 1860 (five), *Ctenopharyngodon* Steindachner, 1866 (one), *Culter* Basilewsky, 1855 (one), *Distoechodon* Peters, 1881 (one), *Elopichthys* Bleeker, 1860 (one), *Hemiculter* Bleeker, 1860 (two), *Hypophthalmichthys* Bleeker, 1860 (one), *Luciobrama* Bleeker, 1870 (one), *Megalobrama* Dybowski, 1872 (two), *Mylopharyngodon* Peters, 1881 (one), *Ochetobius* Günther, 1868 (one), *Opsariichthys* Bleeker, 1863 (one), *Parabramis* Bleeker, 1864 (one), *Plagiognathops* Berg, 1907 (one), *Pseudobrama* Bleeker, 1870 (one), *Pseudolaubuca* Bleeker, 1864 (two), *Sinibrama* Wu, 1939 (one), *Squaliobarbus* Günther, 1868 (one), *Toxabramis* Günther, 1873 (one), *Xenocypris* Günther, 1868 (two) and *Zacco* Jordan & Evermann, 1902 (one). The large majority of these species are widespread in the lowlands of south or east China.

Several previously-recorded species from Lake Dongting have synonymisations or taxonomic changes. *Xenocypris argentea* Günther, 1868 was synonymised with *X. macrolepis* Bleeker, 1871 (Kottelat 2013). *Xenocypris microlepis* Bleeker, 1871 had been referred to *Plagiognathops* Berg, 1907 (Kottelat 2013). *Zacco acanthogenys* (Boulenger, 1901) had long been synonymised with *Zacco platypus* (Temminck & Schlegel, 1846) until Wang (2019) and Zhu et al. (2020) revalidated it. The type locality of *Z. platypus* is in Japan (Liu et al. 2011), but *Z. acanthogenys* occurs in the mid-lower Chang-Jiang Basin. Specimens under the name of *Sinibrama wui* (Rendahl, 1933) from Lake Dongting are referred to as *S. macrops* (Günther, 1868), following Xie et al. (2003) and Zhang et al. (2004). Specimens, previously recognised as *Megalobrama skolkovii* Dybowski, 1872, from the lake are identified as *M. mantschuricus* (Basilewsky, 1855), following Vasil'eva and Makeeva (2003) and Bogutskaya et al. (2008).

The taxonomy of three genera *Chanodichthys* Bleeker, 1860, *Culter* Basilewsky, 1855 and *Cultrichthys* Smith, 1938 is hitherto in a chaotic status in Chinese literature. The type species of *Chanodichthys* is *Leptocephalus mongolicus* Basilewsky, 1855 [type locality: China: Mongolia (presently Inner Mongolia Province) and Manchuria (now northeast China)], that of *Culter* is *C. alburnus* Basilewsky, 1855 [type locality: China: rivers flowing into the Gulf of Tschili (today's Hebei Province)] and that of *Cultrichthys* is *C. brevicauda* Günther, 1868 (type locality: Taiwan, China). Bănărescu (1997) synonymised *Cultrichthys* with *Culter*. This synonymisation was subsequently accepted by some researchers (Bogutskaya and Naseka 2004; Bogutskaya et al. 2008). The type species of *Culter*, though, was misplaced in *Cultrichthys* in Chinese literature (Luo 1994; Luo and Yue 1996; Chen 1998). This misplacement can be traced back to Yi and Zhu (1959), who took it for granted that *Cultrichthys erythropterus* (Basilewsky, 1855), as indicated by the species name, is the available scientific name for the species

with pink pectoral, pelvic and anal fins. This character, along with a long keel extending along the mid-line of the chest and belly, is typical for *Culter alburnus* (Bogutskaya et al. 2008). *Cultrichthys erythropterus* (*sensu* Chen 1998) is, thus, the misidentification of *Culter alburnus* (*sensu* Chen 1998) and vice versa. *Culter*, as here delimited, includes two species: *C. alburnus* and *C. compressocorpus* Yih & Chu, 1959. All other species currently placed to *Culter* by Chinese authors should be referred to as *Chanodichthys*.

Gobionidae

Twenty-five species of gudgeons from Lake Dongting are placed in 12 genera, namely *Abbottina* Jordan & Fowler, 1903 (one species), *Coreius* Jordan & Starks, 1905 (one), *Gobiobotia* Kreyenberg, 1911 (four), *Hemibarbus* Bleeker, 1860 (two), *Microphysogobio* Mori, 1934 (one), *Paracanthobrama* Bleeker, 1864 (one), *Pseudogobio* Bleeker, 1860 (one), *Pseudorasbora* Bleeker, 1860 (one), *Rhinogobio* Bleeker, 1870 (three), *Sarcocheilichthys* Bleeker, 1860 (three), *Saurogobio* Bleeker, 1870 (six) and *Squalidus* Dybowski, 1872 (one). Most of these species are often seen in the mid-lower Chang-Jiang basin or even lowland areas of the southern China.

Sarcocheilichthys is represented in Lake Dongting by three species, namely *S. nigripinnis* (Günther, 1873), *S. tungtingensis* Nichols & Pope, 1927 and *S. sinensis* Bleeker, 1871. Our ongoing taxonomy of this genus demonstrates that *S. kiangsiensis* Nichols, 1930 occurs only in the lake Poyang system and that specimens, formerly identified as this species from Lake Dongting, belong to *S. tungtingensis* (An 2020). A critical revision of *Sarcocheilichthys* from China is underway; the species diversity of this genus has been highly underestimated so far. *Saurogobio* is presently represented in Lake Dongting by six species, namely *S. dabryi* Bleeker, 1871, *S. dumerili* Bleeker, 1871, *S. gracilicaudatus* Yao & Yang, 1977, *S. gymnocheilus* Lo, Yao & Chen, 1998, *S. lissilabris* Bănărescu & Nalbant, 1973 and *S. xiangjiangensis* Tang, 1980. Both *S. gracilicaudatus* and *S. xiangjiangensis* are new records and so is *P. vaillanti* (Sauvage, 1878). The species status of *S. lissilabris* was suspected by some researchers (Wu 1977; Chen 1998) or even it was synonymised with *S. gymnocheilus* (Dai et al. 2014). Tang et al. (2018) considered it as a valid species on the basis of molecular evidence and their examination on its type. Two historically documented species are removed from the updated species checklist: *Coreius guichenoti* (Sauvage & Dabry de Thiersant, 1874) and *Squalidus nitens* (Günther, 1873). The former is hitherto found only in the upper Chang-Jiang (Zhang et al. 2019; Liu et al. 2020a) and the latter, whose type locality is in Shanghai City (Günther 1873), has not been found in Lake Dongting for decades.

Four species of *Gobiobotia* were formerly reported from Lake Dongting: *G. filifer* (Garman, 1912), *G. meridionalis* Chen & Cao, 1977, *G. nicholsi* Bănărescu & Nalbant, 1966 and *G. pappenheimi* Kreyenberg, 1911. The first species is to date endemic to the Chang-Jiang Basin downstream of Yibin City. The second species had long been treated as a subspecies of *G. longibarba* Fang & Wang, 1931 until Chen (1998) regarded it as a full species. It is extensively known from the middle reaches of the Chang-Jiang and Zhu-Jiang Basins (Chen 1998; Tang et al. 2001; Zhang and Zhao 2016). The two species

were collected from this lake during this field survey. The third species was originally described from Lake Dongting (Bănărescu and Nalbant 1966), but later synonymised with *G. filifer* (Wu 1977). Our ongoing taxonomy of Chinese species of *Gobiobotia* shows that *G. nicholsi* is a valid species of the lake Dongting system, but it is so far known merely by its type specimens. Although Bănărescu and Nalbant (1966) reported on the occurrence of *G. pappenheimi* and *Xenophysogobio boulengeri* (Tchang, 1929) in Lake Dongting, no additional specimens have since been collected. Generally, *G. pappenheimi* (type locality: northern China: Tianjin City) is mainly found in the Hai-He and Huang-He (Wang 1984) and *X. boulengeri* (type locality: southwest China: Sichuan Province) occurred in the upper Chang-Jiang Basin (Zhang et al. 2019). Our photograph examination indicated that specimens, identified by Bănărescu and Nalbant (1966) as *G. pappenheimi* and *X. boulengeri* from Lake Dongting, are conspecific with *G. nicholsi*. Nevertheless, this identification still needs confirmation when topotypical specimens become available. Provisionally, these two species are here regarded to have an erroneous record in the Lake. Recently, a new species from Lake Dongting is here found, based on morphological and molecular evidence (Chen et al. 2022b). Therefore, the eight-barbel gudgeons have four representatives in the lake: *Gobiobotia filifer*, *G. lii*, *G. meridionalis* and *G. nicholsi*.

Acheilognathidae

The bitterlings have eleven representatives in Lake Dongting: *A. macropterus* (Bleeker, 1871), *A. barbatulus* Günther, 1873, *A. macromandibularis* Doi, Arai & Liu, 1999, *A. polylepis* (Wu, 1964), *A. gracilis*, *A. barbatus* Nichols, 1926, *A. chankaensis* (Dybowski, 1872), *A. tonkinensis* (Vaillant, 1892), *A. hypselonotus* (Bleeker, 1871), *Rhodeus ocellatus* (Kner, 1866) and *R. sinensis* Günther, 1868. The first one was formerly misidentified as *A. taenianalis* (Günther, 1873) (Tang and Qian 1979; Li 2006); however, *A. taenianalis* has been shown to be a junior synonym of *A. macropterus* (Kottelat 2013). The second and third bitterlings are new records for this lake (Doi et al. 1999; Li 2013). *Acheilognathus imberbis* Günther, 1868, previously documented from Lake Dongting (Ru 2008), is removed from the species checklist. Its type locality remains unclear, but it is reportedly present in the lower Chang-Jiang Basin so far (Li et al. 2016c; Zhang et al. 2020b). Moreover, this species has not been found in the Lake Dongting system over the past several decades. Although Yu et al. (2005) reported on its distribution in Xiang-Jiang, the identification still needs confirmation.

Botiidae

This family is so far represented in Lake Dongting by five species, three of which are from *Leptobotia* [*L. citrauratea* (Nichols, 1925), *L. rubrilabris* (Dabry de Thiersant, 1872) and *L. taeniops* (Sauvage, 1878)] and two from *Parabotia* [*P. fasciata* Dabry de Thiersant, 1872 and *P. banarescui* (Nalbant, 1965)]. Nichols (1925b) reported on the occurrence of *Botia rubrilabris* in the Lake Dongting system and described *B. purpurea* and *B. citrauratea* as two new species from the lake. The three sympatrically existing

congeneric species were later transferred to *Leptobotia* where *B. citrauratea* and *B. purpurea* were synonymised, respectively with *L. elongata* and *L. taeniops* (Bleeker, 1870) (Chen 1980; Kottelat 2004, 2012). Recently, *L. citrauratea* was resurrected from the synonym of *L. elongata*, based on examination of the type and morphological data (Bohlen and Šlechtová 2017). Guo and Zhang (2021) also affirmed that *L. citrauratea* survives in Lake Dongting (type locality). Only a single small-sized individual of *L. rubrilabris* was collected by Anonymous (1980) in this lake. Our field survey yielded no specimens of this species. Likely, it has been extirpated in Lake Dongting. The taxonomy of *Leptobotia* species from China needs a critical revision.

Bagridae

The taxonomy of the bagrid catfishes from China is notoriously poorly understood. This family is represented in Lake Dongting by two genera: *Hemibagrus* Bleeker, 1862 and *Tachysurus* Lacepède, 1803. Species previously referred to *Mystus* Scopoli, 1777 are misidentification of *Hemibagrus* in Chinese literature (Liu et al. 2013; Yuan et al. 2019; Yang et al. 2020). All species, formerly placed in *Pelteobagrus* Bleeker, 1864 and *Pseudobagrus* Bleeker, 1858, are currently referred to *Tachysurus* (Ng and Freyhof 2007; Kottelat 2013) and so are Chinese species formerly placed in *Leiocassis* Bleeker, 1857 (Cheng and Zhang 2012), which is in fact a genus endemic to Southeast Asia (Ng and Kottelat 2007).

The Bagridae is represented in Lake Dongting by 10 species, namely *Tachysurus crassilabris* (Günther, 1864), *T. dumerili* (Bleeker, 1864), *T. eupogon* (Boulenger, 1892), *T. mica* (Gromov, 1970), *T. nitidus* (Sauvage & Dabry de Thiersant, 1874), *T. sinensis* Lacepède, 1803, *T. ussuriensis* (Dybowski, 1872), *T. vachellii* (Richardson, 1846), *T. zhangfei* Shao, Cheng & Zhang, 2021 and *Hemibagrus macropterus* Bleeker, 1870. *Tachysurus dumerili* is a senior subjective synonym of *T. longirostris* Günther, 1864 (Kottelat 2013). Specimens of *T. mica* were formerly misidentified as the juveniles of other catfishes owing to their small size (Chu et al. 1999), but our ongoing taxonomy of Chinese *Tachysurus* indicates that it is a valid species. *Tachysurus sinensis* is a senior subjective synonym of *T. fulvidraco* (Richardson, 1846) (Ng and Kottelat 2007). Specimens, previously recognised as *T. albomarginatus* (Rendahl, 1928), from Lake Dongting represent an undescribed species, which was named as *T. zhangfei* (Shao et al. 2021). Possibly, Ru's (2012) specimens, under the name of *T. tenuis* (Günther, 1873), from Lake Dongting were misidentified as it is hitherto known merely from the type locality, Chongming Island, Shanghai City (Kottelat 2013; Cheng et al. 2021). This species is tentatively excluded from this updated species checklist.

Ictaluridae & Centrarchidae

The family Ictaluridae and Centrarchidae are each represented in Lake Dongting by a single species. *Ictalurus punctatus* (Rafinesque, 1818) and *Micropterus salmoides* (Lacepède, 1802), introduced as cultured fishes to China, are sporadically found in the lakes from southern China (Li et al. 2016d).

Sinipercidae

This family is so far represented in Lake Dongting by four species of the genus *Siniperca* Gill, 1862: *S. chuatsi* (Basilewsky, 1855), *S. knerii* Garman, 1912, *S. roulei* Wu, 1930 and *S. scherzeri* Steindachner, 1892 (Tang and Qian 1979; Li 2006; Li 2013). This third perch was previously assigned to *Coreosiniperca* Fang & Chong, 1932, but this genus has been shown to be invalid (Liu and Chen 1994).

Gobiidae

Five gobies of *Mugilogobius* Smitt, 1900 and *Rhinogobius* Gill, 1859 were previously recorded from Lake Dongting: *M. myxodermus* (Herre, 1935), *R. brunneus* (Temminck & Schlegel, 1845), *R. cliffordpopei*, *R. giurinus* Gill, 1859 and *R. similis* Gill, 1859 (Tang and Qian 1979; Li 2006). Specimens, under the name of *R. giurinus*, have been shown to be misidentification of *R. similis* (Suzuki et al. 2016; Suzuki et al. 2017). The current identification of *R. brunneus* from this lake remains suspicious. Its type locality is in Japan (Temminck and Schlegel 1845). Chinese specimens of this goby were referred to as different species (Wu and Zhong 2008). Nevertheless, whether the species exists in Chinese freshwaters remains unsolved yet. Temporarily, the goby is removed from this updated checklist. Thus, only three gobies are here recognised from Lake Dongting: *M. myxodermus*, *R. cliffordpopei* and *R. similis*.

Discussion

Species diversity

Lake Dongting, as the second-largest river-connected freshwater lake lying within the floodplain areas of the mid-lower Chang-Jiang Basin, supports diversified freshwater fish species. A total of 130 fish species is here reported from the Lake. This number accounts for ca. 31.48% of the total freshwater fishes of the Chang-Jiang Basin where 413 native species were recently documented (Zhang and Cao 2021a). According to the recently-published book entitled “The fish fauna of Hunan Province”, the Dongting Lake system harbours up to 218 freshwater fish species (Wu et al. 2021). The lake alone contributes to 59.63% of the total number of freshwater fishes from the system. In addition to serving as favourable habitats of the Yangtze finless porpoise (Huang et al. 2017) and the crucial stopover and breeding grounds of plentiful migrating birds (Fang et al. 2006; Zou et al. 2019), this lake is also used as sanctuaries or nursery grounds by numerous larvae of drifting-egg-spawning or potamodromous fishes like four major Chinese carps, as spawning grounds by some anadromous fishes like *Coilia nasus* and *Tenualosa reevesii* and as feeding grounds by some catadromous fishes like *Anguilla japonica* and *Takifugu obscurus* (Dou and Jiang 2000). Evidently, Lake Dongting is the key biodiversity area of this lake system or the Chang-Jiang Basin.

The total number of freshwater fish species of Lake Dongting given in this updated checklist is actually comparable to that of Lake Poyang, the first-largest river-connected floodplain lake of the mid-lower Chang-Jiang Basin, where a total of 136 fish species has been recorded so far (Zhang and Li 2007; Yang et al. 2015b; Fang et al. 2016). This number seems to be higher than that of Lake Dongting, but remains doubtful. From the latest species checklist of freshwater fishes from the Gan-Jiang—the largest river flowing into Lake Poyang, 36 historically recorded species were removed (Wang and Zhang 2021). Amongst them, at least ten species were contained in checklists of fish species of Lake Poyang by Zhang and Li (2007) and Fang et al. (2016); these ten species were also components of the ichthyofauna of Lake Poyang system compiled by Huang et al. (2013) and Hu et al. (2019). Both Lakes Dongting and Poyang support rich fish species diversity that is unmatched by any other lake in the mid-lower Chang-Jiang Basin, such as Lake Chao (54 fish species, Guo et al. 2007), Lake Tai (107, Zhu et al. 2007), Lake Hongze (88, Lin et al. 2013) or Lake Hong (84, unpublished data) and far higher than that of lakes located in the Yunnan-Guizhou Plateau (Yuan et al. 2010). This can be plausibly explained by uniqueness of these two large-sized floodplain subtropical lakes: the permanent lateral hydrological connection with the Chang-Jiang mainstem and co-existence of lentic and lotic environments. The assembly mechanism maintaining fish community within Lake Dongting has been addressed in Chen et al. (2022a).

The present study shows that fish species diversity of Lake Dongting remains insufficiently understood. The number of species, collected from the Lake in this survey, is lower compared with the frontrunners (Liang and Liu 1959, 1966; Tang and Qian 1979; Li 2006). Eight newly-recorded native species are added likely due to the multiple sampling methods used and three seasonal samplings during our survey from 2017 to 2019. More sampling efforts lead to the discovery of higher species richness (Hughes et al. 2021; Pompeu et al. 2021). Twenty historically-recorded species are excluded from the checklist, mainly due to the following reasons: (1) Species misidentification. This is the case for seven species which do not exist in the lake presently, namely *Coreius guichenoti*, *Gobiobotia pappenheimi*, *Leptobotia elongata*, *Procypris rabaudi*, *Pseudobagrus tenuis* (= *Tachysurus tenuis*), *Spinibarbus sinensis* and *Xenophysogobio boulengeri*; (2) Taxonomic alteration. Species, formerly identified as *Cyprinus carpio*, *Sarcocheilichthys kiangsiensis*, *Spinibarbus hollandi* and *Zacco platypus* from this lake or China, are now referred to as *Cyprinus rubrofasciatus*, *Sarcocheilichthys tungtingensis*, *Spinibarbus caldwelli* and *Zacco acanthogenys*, respectively; (3) Unconfirmed records. Whether *Acheilognathus imberbis*, *Rhinogobius brunneus* and *Squalidus nitens* occur in Lake Dongting remains controversial; (4) Synonymisation. The following six species are to date regarded as invalid: *Acheilognathus taenianalis*, *Hemisanx brachyrostralis*, *Neosalanx oligodontis*, *Neosalanx taihuensis*, *Rhinogobius giurinus*, and *Sinibrama wui*. It is apparent that problems with the current identification of some fish species in Lake Dongting still remains.

This checklist includes 20 species which experienced nomenclatural alterations, viz. *Bangana tungting* (= *Decorus tungting*), *Coreosiniperca roulei* (= *Siniperca roulei*), *Culter alburnus* (= *Chanodichthys erythropterus*), *Culter dabryi* (= *Chanodichthys dabryi*), *Culter mongolicus* (= *Chanodichthys mongolicus*), *Culter oxycephaloides*

(= *Chanodichthys oxycephaloides*), *Culter oxycephalus* (= *Chanodichthys oxycephalus*), *Cultrichthys erythropterus* (= *Culter alburnus*), *Gobiobotia longibarba meridionalis* (= *Gobiobotia meridionalis*), *Leiocassis argentivittatus* (= *Tachysurus mica*), *Leiocassis crassilabris* (= *Tachysurus crassilabris*), *Leiocassis longirostris* (= *Tachysurus dumerili*), *Mystus macropterus* (= *Hemibagrus macropterus*) *Pelteobagrus eupogon* (= *Tachysurus eupogon*), *Pelteobagrus fulvidraco* (= *Tachysurus sinensis*), *Pelteobagrus nitidus* (= *Tachysurus nitidus*), *Pelteobagrus vachellii* (= *Tachysurus vachellii*), *Pseudobagrus albomarginatus* (= *Tachysurus zhangfei*), *Pseudobagrus ussuriensis* (= *Tachysurus ussuriensis*), *Xenocypris microlepis* (= *Plagiognathops microlepis*). Two species, *Gobiobotia nicholsi* and *Sarcocheilichthys tungtingensis* are, for the time being, regarded as valid. Their taxonomic status needs to be confirmed when specimens from their type locality (today's East Dongting Lake) become available.

Biodiversity conservation

Amongst 130 freshwater fish species of Lake Dongting, 12 (9.23% of the total) are labelled as threatened species in Zhang and Cao's (2021a) assessment of the Red List of Chinese freshwater fishes, viz. *Acipenser sinensis* (CR), *A. dabryanus* (CR), *Anguilla japonica* (EN), *Decorus tungting* (EN), *Leptobotia rubrilabris* (VU), *Luciobrama macrocephalus* (CR), *Myxocyprinus asiaticus* (CR), *Ochetobius elongatus* (CR), *Onychostoma rarum* (VU), *Psephurus gladius* (CR), *Rhinogobio ventralis* (EN) and *Tenualosa reevesii* (CR) (see Table 3). Three species, *Psephurus gladius*, *Acipenser sinensis* and *A. dabryanus*, are also listed in the Appendices II of the Convention on International Trade in Endangered Species (CITES 2019). *Psephurus gladius* was declared to have been functionally extinct in the Chang-Jiang Basin, due to a permanent lack of reproduction or recruitment since 1993 (Zhang et al. 2020a). No wild individuals of *A. dabryanus* have been monitored since 1995 (Zhang et al. 2017). The critically endangered status of *A. sinensis* was mainly owing to a dramatic decline in population after 2000 (Zhang and Cao 2021a). Field surveys conducted from 2002 to 2009 found a trend of a drastic decrease in its juvenile population year by year (Wang et al. 2011; Wu et al. 2015). No spawning individuals were monitored from 2013 to 2015 into the Chang-Jiang mainstem downstream of the Gezhouba Dam (Wu et al. 2017; Zhang et al. 2020a), therefore indicating that the population of this freshwater megafauna species is extremely impacted by river damming (Zhang et al. 2017). Only one small individual of *A. sinensis* was collected from Lake Dongting during 2012 (unpublished data). This clearly means that the lake can be utilised as nursery or feeding grounds by the juveniles and, hence, plays an important role in the conservation of the sturgeon. Nevertheless, the young sturgeon is also likely the captive-bred juveniles released into the upper Chang-Jiang, given that restocking, one salvaging measure taken to conserve this fish, has been in place for several decades (Du et al. 2013). Two imperilled species, *Anguilla japonica* (EN) and *Myxocyprinus asiaticus* (CR) were collected in this field survey, indicating that both still persist here. More attention should be paid to the remaining threatened fish species unsampled in this survey. Whether they eluded capture or have been extirpated, their populations are in a continuous decline and salvaging actions should be adopted immediately.

Five species are also on the latest List of Key Protected Wild Animals in China, namely *Leptobotia rubrilabris*, *Luciobrama macrocephalus*, *Myxocyprinus asiaticus*, *Rhinogobio ventralis* and *Tenualosa reevesii* (Anonymous 2021). *Tenualosa reevesii* is of importance for capture fisheries, particularly in the mid-lower Chang-Jiang Basin. The population of this anadromous fish had been in remarkable decrease as from 1992 when Wan'an Dam was constructed across the Gan-Jiang, where its spawning grounds were located (Tang et al. 1993; Liu 2002). During the past twenty years, no individuals have been collected (Wang and Zhang 2021). The fish, like *Psephurus gladius*, has probably been extinct in the Chang-Jiang Basin (Zhang and Cao 2021a). *Myxocyprinus asiaticus* is rarely encountered in Lake Dongting due to a sharp decline in population resulting from anthropogenic disturbances (Fang et al. 2006). One small individual of 382 mm SL, which was caught during our field survey, is likely a captive-bred juvenile released into the wild to restock its population. *Luciobrama macrocephalus* used to be widely distributed in southern China, but this food fish of high value has become an occasionally-encountered species. The carnivorous fish has long been regarded as the target species to be eradicated as its juveniles prey on fries of other farmed fishes, thus having negative impacts on lake or reservoir fisheries. Deliberate removal of this apex predator was mainly responsible for its current endangerment status. *Rhinogobio ventralis* was initially described from Lake Dongting (Sauvage and Thiersant 1874), but the gudgeon has vanished since Liang and Liu's (1959, 1966) report on its existence in the lake. *Leptobotia rubrilabris*, originally described from the upper Chang-Jiang Basin, was also recorded from Lake Dongting (Nichols 1925b). The latest report on its survival in the lake was Anonymous (1980), who caught a single specimen of 80 mm SL. Field survey of fishes conducted from 2014 to 2019 into Lake Dongting yielded no specimens of this fish (Guo and Zhang 2021). Likely, it has already been extirpated in this system.

Besides three species (*Luciobrama macrocephalus*, *Myxocyprinus asiaticus* and *Tenualosa reevesii*), there are another 12 species currently included in Hunan provincial key protected wildlife list (The Forest Department of Hunan Province 2015): *Channa asiatica*, *Coilia nasus*, *Decorus tungting*, *Macropodus opercularis*, *Microphysogobio tungtingensis*, *Neosalanx brevirostris*, *Ochetobius elongatus*, *Onychostoma rarum*, *O. simum*, *Saurogobio xiangjiangensis*, *Siniperca roulei* and *Spinibarbus caldwelli* (Table 3). No specific conservation actions, however, have been in place for these species. It is worth pointing out that, except for the *Channa asiatica* and *Macropodus opercularis*, all these species seem to be of local economic importance in the mid-lower Chang-Jiang Basin or Lake Dongting system.

Lake Dongting harbours nine fish species endemic to the mid-lower Chang-Jiang Basin, namely *Acheilognathus hypselonotus*, *A. macromandibularis*, *A. macropterus*, *Coilia brachygnathus*, *Decorus tungting*, *Leptobotia citraurata*, *Megalobrama amblycephala*, *Microphysogobio tungtingensis* and *Saurogobio gracilicaudatus*. These species have a high risk of being imperilled by anthropogenic perturbation. More efforts should be dedicated to monitor their population size and trend. *Decorus tungting*, a popular food fish of local economic importance in Lake Dongting system before 1980s, is currently restricted only to some sections of the Yuan-Jiang and Zi-Shui, two affluents of Lake Dongting (Bian et al. 2011). Owing to a sharp decrease in population over the past 30 years, this

rheophilic species was assessed as Endangered (EN) in the latest assessment of Chinese freshwater fish Red List (Zhang and Cao 2021a). No doubt, salvaging actions should be taken to conserve this species. All these species, except *D. tungting*, were not included in this Red List. Nevertheless, two fishes were listed as Data Deficient (DD): *Coilia brachygnathus* and *Microphysogobio tungtingensis*. *Leptobotia citrauratea* can also be assessed in this category and was recently revalidated (Guo and Zhang 2021). These three species are possibly under the same threat as *D. tungting* and, thus, deserve special attention.

Table 3. Endemics of the mid-lower Chang-Jiang Basin and protected fish species in Lake Dongting.

Species	CITES	China	Hunan	IUCN	Endemics
<i>Psephurus gladius</i>	√	I		CR	
<i>Acipenser sinensis</i>	√	I		CR	
<i>Acipenser dabryanus</i>	√	I		CR	
<i>Coilia nasus</i>			√	LC	
<i>Coilia brachygnathus</i>				DD	√
<i>Tenualosa reevesii</i>		I	√	CR	
<i>Neosalanx brevirostris</i>			√	DD	
<i>Anguilla japonica</i>				EN	
<i>Myxocyprinus asiaticus</i>		II	√	CR	
<i>Onychostoma simum</i>			√	NT	
<i>Onychostoma rarum</i>			√	VU	
<i>Spinibarbus caldwelli</i>			√	LC	
<i>Luciobrama macrocephalus</i>		II	√	CR	
<i>Ochetobius elongatus</i>			√	CR	
<i>Decorus tungting</i>			√	EN	√
<i>Megalobrama amblycephala</i>				LC	√
<i>Acheilognathus macropterus</i>				LC	√
<i>Acheilognathus hypselonotus</i>				LC	√
<i>Acheilognathus macromandibularis</i>				LC	√
<i>Microphysogobio tungtingensis</i>			√	DD	√
<i>Saurogobio gracilicaudatus</i>				LC	√
<i>Saurogobio xiangjiangensis</i>			√	LC	
<i>Rhinogobio ventralis</i>		II		EN	
<i>Leptobotia citrauratea</i>				DD	√
<i>Leptobotia rubrilabris</i>		II		VU	
<i>Siniperca roulei</i>			√	NT	
<i>Channa asiatica</i>			√	LC	
<i>Macropodus opercularis</i>			√	NT	
Total	3	8	15		9

Thirty-four historically documented fish species were not collected from Lake Dongting during this field survey; their fate is of particular concern. These species fall within five categories. The first one is migrating species, like *Acipenser dabryanus*, *A. sinensis*, *Coilia nasus*, *Psephurus gladius*, *Tenualosa reevesii* and *Takifugu obscurus*. The main reasons for the extirpation of the first four species in the lake are mentioned above. Although the last two species eluded capture during this field survey, both were

reportedly collected in exceptional years (Ren et al. 2015; Wang et al. 2016; Chen et al. 2020). Since 1970s, more and more dams have been built across the affluents of Lake Dongting and also the Chang-Jiang mainstem (Wang et al. 2019a). The blockage of migration ways and the shrinkage of favourable habitats were the key factors leading to a sharp decrease in the population of the two diadromous fishes (Wang et al. 2016). Small population size makes it difficult for them to migrate for such a long distance from the estuary of Chang-Jiang into Lake Dongting, particularly when all fishes, being of economic importance in the river, were under high pressure from fishing during the past 20 years. The second category is such potamodromous or drifting-egg-spawning fishes as *Luciobrama macrocephalus*, *Ochetobius elongatus*, *Pseudolaubuca engraulis*, *Rhinogobio cylindricus*, *R. ventralis* and *Saurogobio dumerili*, which are susceptible to dam construction. River damming makes inundated reaches shift from lotic to lentic habitat, which have adverse impacts on the spawning of these species. The third category is rheophilic species such as *Decorus tungting*, *Leptobotia rubrilabris*, *Lepturichthys fimbriatus*, *Onychostoma rarum*, *O. simum*, *Spinibarbus caldwelli*, *Tachysurus ussuriensis* and *Zacco acanthogenys*. Their extirpation in Lake Dongting is mainly attributed to river damming in its affluents, which not only led to a remarkable decline in the population of these fishes, but also blocked their short migration into the lake. The fourth category is bitterlings, such as *Acheilognathus barbatus*, *A. chankaensis*, *A. hypselonotus* and *A. tonkinensis*, which depend on freshwater mussels for spawning. The absence of these bitterlings in Lake Dongting may be related to the decrease or disappearance of mussels caused by degrading water quality or sand extraction (Meng et al. 2018; Liu et al. 2020b; Wang and Zhang 2021). The fifth category includes some fishes of economic value, such as *Distoechodon tumirostris* Peters, 1881, *Neosalanx brevirostris*, *N. jordani*, *Plagiognathops microlepis* and *Protosalanx hyalocranius*. These fishes eluded capture mainly due to small population size led by overfishing and habitat loss or degradation. Overall, most of the unsampled fish species during this field survey have ecologically specialised preferences, for example, migratory, rheophilic, carnivorous, drifting-egg-producing or mussel-dependent. These fishes are susceptible to human disturbances and, thus, can act as biological indicators of aquatic ecosystem health. Their lack of samples clearly indicates that the freshwater ecosystem of Lake Dongting has been severely threatened by human perturbations including river damming, overfishing, habitat degradation and sanding dredging.

The Chang-Jiang basin is an area with over 400 million residents, highly impacted by anthropogenic interferences. It is also the most rapidly growing area of China's economic development. The loss of aquatic diversity and, thus, its ecological service function in this river is becoming a pressing challenge. It is urgently needed to take practical actions to conserve the freshwater ecosystem of the Chang-Jiang Basin. To this end, the Chinese government made a decision of implementing the conservation measure of 'ten-year fishing ban' in all natural water bodies of the mainstem and major tributaries of the Chang-Jiang since 2020 (Pan and Liu 2021). Whether it is an effective protection action for conserving the fish diversity of Lake Dongting is of much public concern. In this context, adequate information about the current status of fish

diversity, including species composition, distribution, population size and imperilled status, is an urgent requirement in the future to answer the question. This updated species checklist will be very useful for further biodiversity analysis and conservation of freshwater fishes from Chang-Jiang.

Acknowledgements

This work was granted by the special fund for Biodiversity Survey & Assessment Project for Biodiversity Conservation of Lake Dongting (2017HB2096001006) and National Science & Technology Fundamental Resources Investigation Program of China (2019FY101800). We are very grateful to Dr. Liang Cao, Chang-Ting An, Li-Jun Zhang, Zi-Tong Wang, Wei-Han Shao, Dong-Ming Guo (IHB), Dinh Tao Nguyen (CCNU), Unisa Conteh Kanu and Long-Hui Qiu (HZAU) for their help with field sampling. Our sincere thanks should go to Prof. Jianzhong Shen (HZAU) for his assistance in fieldworks. Special thanks should be given to Radford Arrindell (AMNH) for providing specimens photographs of *Gobiobotia nicholsi*, *G. pappenheimi*, *G. filifer* and *Xenophysogobio boulengeri*. We greatly appreciate all valuable comments of two Reviewers Jie Zhang and Fan Li.

References

- Abell R, Thieme ML, Revenga C, Bryer M, Kottelat M, Bogutskaya N, Coad B, Mandrak N, Balderas SC, Bussing W, Stiassny MLJ, Skelton P, Allen GR, Unmack P, Naseka A, Ng R, Sindorf N, Robertson J, Armijo E, Higgins JV, Heibel TJ, Wikramanayake E, Olson D, López HL, Reis RE, Lundberg JG, Sabaj Pérez MH, Petry P (2008) Freshwater ecoregions of the world: A new map of biogeographic units for freshwater biodiversity conservation. *Bioscience* 58(5): 403–414. <https://doi.org/10.1641/B580507>
- An C (2020) Integrative taxonomy of the gudgeon genus *Sarcocheilichthys* Bleeker, 1859 *sensu lato* (Cyprinidae: Gobioninae) in China. PhD thesis, Wuhan: Institute of Hydrobiology, Chinese Academy of Sciences.
- Anonymous (1976) Fishes of the Chang-Jiang. Science Press, Beijing, 286 pp.
- Anonymous (1980) Fish of Hunan Province. Hunan Science and Technology Press, Changsha, 231 pp.
- Anonymous (2021) List of Key Protected Wild Animals in China. National Forestry and Grassland Administration and Ministry of Agriculture and Rural Affairs, Beijing, 38 pp.
- Bănărescu P (1997) The status of some nominal genera of Eurasian Cyprinidae (Osteichthyes, Cypriniformes). *Revue Roumaine de Biologie Serie de Biologie Animale* 42: 19–30.
- Bănărescu P, Nalbant TT (1966) Notes on the genus *Gobiobotia* (Pisces, Cyprinidae) with description of three new species. *Annotationes Zoologicae et Botanicae* 27: 1–16.
- Barbarossa V, Bosmans J, Wanders N, King H, Bierkens MFP, Huijbregts MAJ, Schipper AM (2021) Threats of global warming to the world's freshwater fishes. *Nature Communications* 12(1): e1701. <https://doi.org/10.1038/s41467-021-21655-w>

- Betancur-R R, Wiley EO, Arratia G, Acero A, Bailly N, Miya M, Lecointre G, Orti G (2017) Phylogenetic classification of bony fishes. *BMC Evolutionary Biology* 17(1): e162. <https://doi.org/10.1186/s12862-017-0958-3>
- Bian W, Li C, Yu C, Liang Z, Zhang Z, Liu M, Yang D (2011) Biological characteristic and resource dynamic of *Sinilabeo decorus tungting*. *Journal of Hydroecology* 32: 67–73. <https://doi.org/10.3969/j.issn.1003-1278.2011.04.013>
- Bogutskaya NG, Naseka AM (2004) Catalogue of Agnathans and Fishes of Fresh and Brackish Waters of Russia with comments on nomenclature and taxonomy. Russian Academy of Sciences, KMK Scientific Press Ltd, Moscow, 389 pp.
- Bogutskaya NG, Naseka AM, Shedko SV, Vasil'eva ED, Chereshev IA (2008) The fishes of the Amur River: Updated check-list and zoogeography. *Ichthyological Exploration of Freshwaters* 19: 301–366. <https://doi.org/10.1093/icesjms/fsn132>
- Bohlen J, Šlechtová V (2017) *Leptobotia micra*, a new species of loach (Teleostei: Botiidae) from Guilin, southern China. *Zootaxa* 4250(1): 11. <https://doi.org/10.11646/zootaxa.4250.1.7>
- Brooks TM, Da Fonseca GAB, Rodrigues ASL (2004) Species, Data, and Conservation Planning. *Conservation Biology* 18(6): 1682–1688. <https://doi.org/10.1111/j.1523-1739.2004.00457.x>
- Cao Y, Liao F, Wu Y (2012) Aquatic fauna of Xiangjiang River. Hunan Science and Technology Publishing House, Changsha, 452 pp.
- Chen J (1980) A study on the classification of the Botoid fishes of China. *Zoological Research* 1: 3–26.
- Chen Y (1998) Fauna Sinica: Osteichthyes Cypriniformes II. Science Press, Beijing, 531 pp.
- Chen T, Wang Y, Gardner C, Wu F (2020) Threats and protection policies of the aquatic biodiversity in the Yangtze River. *Journal for Nature Conservation* 58: 125931. <https://doi.org/10.1016/j.jnc.2020.125931>
- Chen X, Li Z, Boda P, Fernandes I, Xie Z, Zhang E (2022a) Environmental filtering in the dry season and spatial structuring in the wet: different fish community assembly rules revealed in a large subtropical floodplain lake. *Environmental Science and Pollution Research*. [J] <https://doi.org/10.1007/s11356-022-20529-y>
- Chen X, Wang M, Cao L, Zhang E (2022b) *Gobiobotia lii*, a new species of gudgeon (Teleostei, Gobiionidae) from the middle Chang-Jiang Basin, central China, with notes on the validity of *G. nicholsi* Bănărescu & Nalbant, 1966. *Zoosystematics and Evolution* 98(1): 93–107. [J] <https://doi.org/10.3897/zse.98.80547>
- Cheng J, Zhang E (2012) A taxonomic research situation of the bargrid catfish genus *Pseudobagrus*. *Journal of Jinggangshan University* 033: 94–98. <https://doi.org/10.3969/j.issn.1674-8085.2012.02.024> [Natural Science]
- Cheng J, Shao W, López JA, Zhang E (2021) *Tachysurus lani*, a new catfish species (Teleostei: Bagridae) from the Pearl River basin, South China. *Ichthyological Exploration of Freshwaters*: 1–17. <https://doi.org/10.23788/IEF-1156>
- Chu Y (1931) Index Piscium Sinensium. Department of Biology, ST. John's University, Shanghai, 290 pp.
- Chu Y (1935) Comparative studies on the scales and on the pharyngeals and their teeth in Chinese Cyprinids, with particular reference to taxonomy and evolution. PhD thesis, Shanghai: St John's University. <https://doi.org/10.2307/1436747>

- Chu X (1955) On fishes of Ichang, with notes on their distribution in the Yangtze River. Shui Sheng Sheng Wu Hsueh Bao •••: 81–95. <http://ir.ihb.ac.cn/handle/152342/7396>
- Chu X, Chen Y (1989) The fishes of Yunnan, China (I). Science Press, Beijing, 387 pp.
- Chu X, Zheng B, Dai D (1999) Fauna Sinica: Osteichthyes Siluriformes. Science Press, Beijing, 243 pp.
- CITES (2019) Checklist of Convention on International Trade in Endangered Species. <https://checklist.cites.org/#/en/search/> [Accessed 14 September 2021]
- Dai Z, Zhang E, Jiang Z, Wang X (2014) Re-description of the gudgeon species *Saurogobio gracilicaudatus* Yao & Yang in Luo, Yue & Chen, 1977 (Teleostei: Cyprinidae) from the Chang-Jiang basin, South China, with a note on its generic classification. Zootaxa 3847(2): 283–291. <https://doi.org/10.11646/zootaxa.3847.2.8>
- Ding R (1994) The fishes of Sichuan, China. Sichuan Publishing House of Science and Technology, Chengdu, 661 pp.
- Doi A, Arai R, Liu H (1999) *Acheilognathus macromandibularis*, a new bitterling (Cyprinidae) from the lower Changjiang basin, China. Ichthyological Exploration of Freshwaters 10: 303–308.
- Dong R, Wang Y, Lu C, Lei G, Wen L (2021) The seasonality of macroinvertebrate β diversity along the gradient of hydrological connectivity in a dynamic river-floodplain system. Ecological Indicators 121: 107112. <https://doi.org/10.1016/j.ecolind.2020.107112>
- Dou H, Jiang J (2000) Dongting Lake. Chinese Scientific and Technology University Press, Hefei, 344 pp.
- Du H, Wang CY, Wei QW, Zhang H, Wu JM, Li L (2013) Distribution and movement of juvenile and sub-adult Chinese sturgeon (*Acipenser sinensis* Gray, 1835) in the Three Gorges Reservoir and the adjacent upstream free-flowing Yangtze River section: A re-introduction trial. Journal of Applied Ichthyology 29(6): 1383–1388. <https://doi.org/10.1111/jai.12343>
- Dudgeon D, Arthington AH, Gessner MO, Kawabata Z-I, Knowler DJ, Lévêque C, Naiman RJ, Prieur-Richard A-H, Soto D, Stiassny MLJ, Sullivan CA (2006) Freshwater biodiversity: Importance, threats, status and conservation challenges. Biological Reviews of the Cambridge Philosophical Society 81(02): 163–182. <https://doi.org/10.1017/S1464793105006950>
- Fang J, Wang Z, Zhao S, Li Y, Tang Z, Yu D, Ni L, Liu H, Xie P, Da L, Li Z, Zheng C (2006) Biodiversity changes in the lakes of the Central Yangtze. Frontiers in Ecology and the Environment 4(7): 369–377. [https://doi.org/10.1890/1540-9295\(2006\)004\[0369:BCITLO\]2.0.CO;2](https://doi.org/10.1890/1540-9295(2006)004[0369:BCITLO]2.0.CO;2)
- Fang C, Chen W, Zhou H, Zhang Y, Fu P, He G, Wu B, Wang S (2016) Fish resources in Poyang Lake and suggestions on their utilization. Jiangsu Agriculture and Technology 044: 233–242, 243. <http://doi.org/10.15889/j.issn.1002-1302.2016.09.067>
- Fricke R, Eschmeyer W, Van Der Laan R (2021) Eschmeyer's Catalog of fishes: Genera, Species, References. <http://researcharchive.calacademy.org/> [Accessed 14 September 2021]
- Fu C, Wu J, Chen J, Wu Q, Lei G (2003) Freshwater fish biodiversity in the Yangtze River basin of China: Patterns, threats and conservation. Biodiversity and Conservation 12(8): 1649–1685. <https://doi.org/10.1023/A:1023697714517>
- Fu C, Luo J, Wu J, López JA, Zhong Y, Lei G, Chen J (2005) Phylogenetic relationships of salangid fishes (Osmeridae, Salanginae) with comments on phylogenetic placement of the

- salangids based on mitochondrial DNA sequences. *Molecular Phylogenetics and Evolution* 35(1): 76–84. <https://doi.org/10.1016/j.ympev.2004.11.024>
- Fu H, Wang X, Ge D, Li W, Tan X, Yuan G, Jeppesen E (2021) Human activities uncouple the cascading effects of hydrological gradients on plant diversity and ecosystem functions in the Lake Dongting wetland. *Ecohydrology* n/a: e2359. <https://doi.org/10.1002/eco.2359>
- Garman S (1912) Pisces. In: Some Chinese vertebrates. Harvard College, Cambridge, 630 pp.
- Günther ACLG (1873) Report on a collection of fishes from China. *The Annals and magazine of natural history; zoology, botany, and geology* 12: 239–250. <https://doi.org/10.1080/00222937308680749>
- Guo D, Zhang E (2021) Re-description of the loach species *Leptobotia citrauratea* (Teleostei, Bortiidae), with the description of *L. brachycephala* from southern Zhejiang Province, China. *ZooKeys* 1017: 89–109. <https://doi.org/10.3897/zookeys.1017.57503>
- Guo L, Xie P, Ni L, Hu W, Li H (2007) The status of fishery resources of Lake Chaohu and its response to eutrophication. *Shui Sheng Sheng Wu Hsueh Bao* 31(5). <https://doi.org/10.3321/j.issn:1000-3207.2007.05.015>
- Guo L, Li J, Wang Z, Cuizhang F (2011) Phylogentic relationships of noodle-fishes (Osmeriformes: Salangidae) based on four mitochondrial genes. *Shui Sheng Sheng Wu Hsueh Bao* 35: 79–89. <https://doi.org/10.3724/SP.J.1035.2011.00449>
- Hora SL (1932) Classification, bionomics and evolution of homalopterid fishes. *Memoirs of the Indian Museum* v. 12(no. 2): 263–330. [Pls 10–12]
- Hu M, Wang C, Liu Y, Zhang X, Jian S (2019) Fish species composition, distribution and community structure in the lower reaches of Ganjiang River, Jiangxi, China. *Scientific Reports* 9(1): e10100. <https://doi.org/10.1038/s41598-019-46600-2>
- Huang L, Wu Z, Li J (2013) Fish fauna, biogeography and conservation of freshwater fish in Poyang Lake Basin, China. *Environmental Biology of Fishes* 96(10–11): 1229–1243. <https://doi.org/10.1007/s10641-011-9806-2>
- Huang S, Mei Z, Hao Y, Zheng J, Wang K, Wang D (2017) Saving the Yangtze finless porpoise: Time is rapidly running out. *Biological Conservation* 210: 40–46. <https://doi.org/10.1016/j.biocon.2016.05.021>
- Hughes RM, Herlihy AT, Peck DV (2021) Sampling efforts for estimating fish species richness in western USA river sites. *Limnologia* 87: 125859. <https://doi.org/10.1016/j.limno.2021.125859>
- Jackson DA, Peres-Neto PR, Olden JD (2001) What controls who is where in freshwater fish communities—the roles of biotic, abiotic, and spatial factors. *Canadian Journal of Fisheries and Aquatic Sciences* 58: 157–170. <https://doi.org/10.1139/cjfas-58-1-157>
- Jiang Z, Cao L, Zhang E (2019) Spatio-temporal variations of fish assemblages in the Dongting Lake. *Shui Sheng Sheng Wu Hsueh Bao (Supplement 43)*: 42–48. <https://doi.org/10.7541/2019.165>
- Jiang X, Wang J, Pan B, Li D, Wang Y, Liu X (2022) Assessment of heavy metal accumulation in freshwater fish of Dongting Lake, China: Effects of feeding habits, habitat preferences and body size. *Journal of Environmental Sciences* 112: 355–365. <https://doi.org/10.1016/j.jes.2021.05.004>

- Kimura S (1934) Description of the fishes collected from the Yangtze-kiang, China by the late Dr. K. Kishinouye and his party in 1927–1929. Journal of Shanghai Science Institute Section 3: 1.
- Kottelat M (2004) *Botia kubotai*, a new species of loach (Teleostei: Cobitidae) from the Ataran River basin (Myanmar), with comments on botiine nomenclature and diagnosis of a new genus. Zootaxa 401(1): 1–18. <https://doi.org/10.11646/zootaxa.401.1.1>
- Kottelat M (2006) Fishes of Mongolia: a check-list of the fishes known to occur in Mongolia with comments on systematics and nomenclature. Environment and Social Development, East Asia and Pacific Region, World Bank, Washington, D.C., 117 pp.
- Kottelat M (2012) Conspectus cobitidum: An inventory of the loaches of the world (Teleostei: Cypriniformes: Cobitoidei). The Raffles Bulletin of Zoology (Supplement 26): 1–199.
- Kottelat M (2013) The fishes of the inland waters of Southeast Asia: A catalogue and core bibliography of the fishes known to occur in freshwaters, mangroves and estuaries. The Raffles Bulletin of Zoology •••: 1–663. <https://doi.org/10.1186/1742-9994-10-72>
- Kottelat M, Freyhof J (2007) Handbook of European Freshwater Fishes. Cornol & Freyhof, Berlin, 646 pp.
- Kottelat M, Whitten A (1996) Freshwater biodiversity in Asia with special reference to fish. World Bank Technical Paper, 59 pp. <https://doi.org/10.1596/0-8213-3808-0>
- Kottelat M, Baird I, Kullander S, Ng HH, Parenti L, Rainboth W, Vidthayanon C (2012) The status and distribution of freshwater fishes of Indo-Burma. 35–65.
- Kreyenberg W, Pappenheim P (1908) Ein Beitrag zur Kenntnis der Fische der Jangtze und seiner Zuflüsse. Sitzungsberichte der Gesellschaft Naturforschender Freunde zu Berlin 1908: 95–109. <https://doi.org/10.5962/bhl.part.12852>
- Li C (2006) Study on the investigation of main economic fishery resource and the law of its variety in the Dongting Lake. Master thesis, Changsha: Hunan Agricultural University.
- Li J (2013) Ecological study on fish community and conservation strategies in Dongting Lake. Master thesis, Changsha: Central South University of Forestry & Technology.
- Li J, Jin Y, Wang W, Zhao Z, Wu X (2016a) Priority areas for land biodiversity conservation in China. Science Press, Beijing, 268 pp.
- Li Q, Liu Y, Zhou J, Gong Q, Li H, Lai J, Li L (2016b) The complete mitochondrial genome of *Gobiobotia filifer* (Teleostei, Cypriniformes: Cyprinidae). Mitochondrial DNA. Part A, DNA Mapping, Sequencing, and Analysis 27(5): 3325–3326. <https://doi.org/10.3109/19401736.2015.1018205>
- Li Q, Yan Y, Chu L, Zhu R, Gao J, Gao Y (2016c) Spatial and temporal patterns of stream fish assemblages within Taihu Basin. Hupo Kexue 28(6): 1371–1380. <https://doi.org/10.18307/2016.0623>
- Li S, Chen J, Wang X (2016d) Global distribution, entry routes, mechanisms and consequences of invasive freshwater fish. Shengwu Duoyangxing 24(6): 672–685. <https://doi.org/10.17520/biods.2015374>
- Liang Q, Liu S (1959) Fishes in Xiangjiang River and Dongting Lake. Journal of Hunan Normal University 3: 67–73. [Abstract] [Natural Science]
- Liang Q, Liu S (1966) Fishes in Hunan Province. Journal of Hunan Normal University 5: 85–111. [Natural Science]

- Liao F, He W, Huang X, Jing Q, He X (2002) Studies on present situation and change trend of Dongting Lake fishery resources and environment. *Shui Sheng Sheng Wu Hsueh Bao* 26: 5. <https://doi.org/10.3321/j.issn:1000-3207.2002.06.008>
- Liao F, He X, He W, Wang H, Xu D (2006) Status and protective regulation countermeasure in fishery resources and its environment of Dongting Lake. *Journal of Yueyang Vocational Technical College* 4: 6. <https://doi.org/10.3969/j.issn.1672-738X.2006.06.009>
- Liermann CR, Nilsson C, Robertson J, Ng RY (2012) Implications of dam obstruction for global freshwater fish diversity. *Bioscience* 62(6): 539–548. <https://doi.org/10.1525/bio.2012.62.6.5>
- Lin M, Zhang T, Ye S, Li W, Ren P, Yang Z, Liu J, Li Z (2013) Status of fish resources, historical variation and fishes management strategies in the Hongze Lake. *Shui Sheng Sheng Wu Hsueh Bao* 37: 1118–1127.
- Liu G (2002) Cause analysis and countermeasures of decline of fisheries resources about *Tenuulosa reevesii* in Xiajiang County. *Jiangxi Agricultural Science and Technology*: 40–41.
- Liu H, Chen Y (1994) Phylogeny of the Siniperine fishes with some taxonomic notes. *Zoological Research* 15 zk: 1–12.
- Liu X, Wang H (2018) Effects of loss of lateral hydrological connectivity on fish functional diversity. *Conservation Biology* 32(6): 1336–1345. <https://doi.org/10.1111/cobi.13142>
- Liu S, Chen D, Duan X, Qiu S, Wang L (2002) The resources status quo and protection strategies on Chinese shad. *Shui Sheng Sheng Wu Hsueh Bao* 26: 679–684.
- Liu MD, Chen DQ, Duan XB, Wang K, Liu SP (2010) Assessment of ecosystem health of upper and middle Yangtze River using fish-index of biotic integrity. *Changjiang Kexueyuan Yuanbao* 27: 1–6.
- Liu F, Wu J, Wang J (2011) Growth and reproductive characteristics of *Ancherythroculter kurematsui* Kimura. *Shui Sheng Sheng Wu Hsueh Bao* 35: 586–595.
- Liu L, Yang C, Yang P, Wang W, Zou W, Han Q (2013) Status and diversity of fish resources of Yuanshui River in Hunan Province, China. *Oceanologia et Limnologia Sinica* 44: 148–158.
- Liu H, Guo C, Qu X, Xiong F, Paukert CP, Chen Y, Su W (2020a) Fish diversity, endemism, threats, and conservation in the Jinsha River Basin (Upper Yangtze River), China. *North American Journal of Fisheries Management* 41(12): 1–18. <https://doi.org/10.1002/nafm.10441>
- Liu Z, Meng X, Li Z, Zhang J, Xu J, Yin S, Xie Z (2020b) Diversity assessment and protection strategies for the mollusk community in the southern Dongting Lake. *Shengwu Duoyangxing* 28(2): 155–165. <https://doi.org/10.17520/biods.2019287>
- Luo Y (1994) Some clarifications on the Cultrinae fishes of China. *Shui Sheng Sheng Wu Hsueh Bao* 18: 45–49. <http://ir.ihb.ac.cn/handle/152342/5108>
- Luo G (2005) History of western botanical and zoological studies in China. Shandong Education Press, Ji'nan, 434 pp.
- Luo Y, Yue P (1996) Preliminary studies on phylogeny of subfamily Cultrinae (Cypriniformes: Cyprinidae). *Shui Sheng Sheng Wu Hsueh Bao* 020: 182–185. <http://ir.ihb.ac.cn/handle/152342/4694>
- Marta S, Lacasella F, Romano A, Ficetola GF (2019) Cost-effective spatial sampling designs for field surveys of species distribution. *Biodiversity and Conservation* 28(11): 2891–2908. <https://doi.org/10.1007/s10531-019-01803-x>

- Meng X, Jiang X, Li Z, Wang J, Cooper KM, Xie Z (2018) Responses of macroinvertebrates and local environment to short-term commercial sand dredging practices in a flood-plain lake. *The Science of the Total Environment* 631–632: 1350–1359. <https://doi.org/10.1016/j.scitotenv.2018.03.086>
- Nelson J, Grande T, Wilson M (2016) *Fishes of the World*, 5th Edn. 750 pp. <https://doi.org/10.1002/9781119174844>
- Ng HH, Freyhof J (2007) *Pseudobagrus nubilosus*, a new species of catfish from central Vietnam (Teleostei: Bagridae), with notes on the validity of *Pelteobagrus* and *Pseudobagrus*. *Ichthyological Exploration of Freshwaters* 18: 9–16.
- Ng HH, Kottelat M (2007) The identity of *Tachysurus sinensis* La Cepède, 1803, with the designation of a neotype (Teleostei: Bagridae) and notes on the identity of *T. fulvidraco* (Richardson, 1845). *Electronic Journal of Ichthyology* 3: 35–54.
- Nichols JT (1925a) An analysis of Chinese loaches of the genus *Misgurnus*. *American Museum Novitates* 169: 1–7.
- Nichols JT (1925b) Some Chinese fresh-water fishes. I. Loaches of the genus *Botia* in the Yangtze Basin. II. A new Minnow-like Carp from Szechwan. III. The Chinese Sucker, *Myxocyprinus*. *American Museum Novitates* 177: 1–10.
- Nichols JT (1925c) Some Chinese fresh-water fishes. IV. Gudgeons of the genus *Coripareius*. V. Gudgeons related to the European *Gobio gobio*. VI. New gudgeons of the genera *Gnathopogon* and *Leucogobio*. *American Museum Novitates* 181: 1–8.
- Nichols JT (1925d) Some Chinese fresh-water fishes. VII. New carps of the genera *Varicorhinus* and *Xenocypris*. *American Museum Novitates* 182: 1–8.
- Nichols JT (1925e) Some Chinese fresh-water fishes. X. Subgenera of bagrin catfishes. XI. Certain apparently undescribed carps from Fukien. XII. A small goby from the central Yangtze. XIII. A new minnow referred to *Leucogobio*. XIV. Two apparently undescribed fishes. *American Museum Novitates* 185: 1–8.
- Nichols JT (1926) Some Chinese fresh-water fishes. XV. Two apparently undescribed catfishes from Fukien. XVI. Concerning gudgeons related to *Pseudogobio*, and two new species of it. XVII. Two new rhodeins. *American Museum Novitates* 214: 1–7.
- Nichols JT (1928) Chinese fresh-water fishes in the American Museum of Natural History's collections: A provisional check-list of the fresh-water fishes of China. *Bulletin of the American Museum of Natural History* 58: 1–62. [AMNH]
- Nichols JT (1943) *The fresh-water fishes of China*. The American Museum of Natural History, New York, 388 pp.
- Nichols JT, Pope CH (1927) The fishes of Hainan. *Bulletin of the American Museum of Natural History* 54: 321–398.
- Pan B, Liu X (2021) A review of water ecology problems and restoration in the Yangtze River Basin. *Changjiang Kexueyuan Yuanbao* 38(3): 1–8. <https://doi.org/10.11988/ckyyb.202007872021>
- Pompeu PS, de Carvalho DR, Leal CG, Leitão RP, Alves CBM, Braga DF, Castro MA, Junqueira NT, Hughes RM (2021) Sampling efforts for determining fish species richness in megadiverse tropical regions. *Environmental Biology of Fishes* 104(11): 1487–1499. <https://doi.org/10.1007/s10641-021-01184-7>

- Qin X, Gong Z, Liu H (2019) Lateral migration of fish between China's second largest freshwater lake (Dongting Lake) and the mainstem of the Yangtze River. *Environmental Biology of Fishes* 102(4): 527–539. <https://doi.org/10.1007/s10641-019-00851-0>
- Regan CT (1908) Descriptions of three new freshwater fishes from China. *Annals & Magazine of Natural History* 1(8): 120. <https://doi.org/10.1080/00222930808692364>
- Reid AJ, Carlson AK, Creed IF, Eliason EJ, Gell PA, Johnson PTJ, Kidd KA, MacCormack TJ, Olden JD, Ormerod SJ, Smol JP, Taylor WW, Tockner K, Vermaire JC, Dudgeon D, Cooke SJ (2019) Emerging threats and persistent conservation challenges for freshwater biodiversity. *Biological Reviews of the Cambridge Philosophical Society* 94(3): 849–873. <https://doi.org/10.1111/brv.12480>
- Ren P, He H, Song Y, Cheng F, Xie S (2015) The spatial pattern of larval fish assemblages in the lower reach of the Yangtze River: Potential influences of river–lake connectivity and tidal intrusion. *Hydrobiologia* 766: 365–379. <https://doi.org/10.1007/s10750-015-2471-2>
- Ru H (2008) Spatio-temporal patterns and river-lake migration rhythms of fish assemblage in Dongting Lake, a large Yangtze-connected lake. Master thesis, Lanzhou: Northwest Normal University.
- Ru H (2012) Life history processes of river-lake migratory fishes in the area of the Dongting Lake, a large river-connected lake in the Yangtze floodplain. PhD thesis, Wuhan: Institute of Hydrobiology, Chinese Academy of Sciences.
- Ru H, Liu X (2013) River-lake migration of fishes in the Dongting Lake area of the Yangtze floodplain. *Journal of Applied Ichthyology* 29(3): 594–601. <https://doi.org/10.1111/jai.12116>
- Sauvage HE, Thiersant Dd (1874) Notes sur les poissons des eaux douces de Chine. In: Audouin JV, Bouvier EL, Grassé P-P, Milne-Edwards H, Milne-Edwards A, Perrier E (Eds) *Annales des sciences naturelles (Zoologie et Paléontologie)*. Crochard, 162–180.
- Shao W, Cheng J, Zhang E (2021) Eight in One: Hidden Diversity of the Bagrid Catfish *Tachysurus albomarginatus* s.l. (Rendhal, 1928) Widespread in Lowlands of South China. *Frontiers in Genetics* 12: 713793. <https://doi.org/10.3389/fgene.2021.713793>
- Sluys R (2013) The unappreciated, fundamentally analytical nature of taxonomy and the implications for the inventory of biodiversity. *Biodiversity and Conservation* 22(4): 1095–1105. <https://doi.org/10.1007/s10531-013-0472-x>
- Strayer DL, Dudgeon D (2010) Freshwater biodiversity conservation: Recent progress and future challenges. *Journal of the North American Benthological Society* 29(1): 344–358. <https://doi.org/10.1899/08-171.1>
- Suzuki T, Shibukawa K, Senou H, Chen IS (2016) Redescription of *Rhinogobius similis* Gill 1859 (Gobiidae: Gobionellinae), the type species of the genus *Rhinogobius* Gill 1859, with designation of the neotype. *Ichthyological Research* 63(2): 227–238. <https://doi.org/10.1007/s10228-015-0494-3>
- Suzuki T, Shibukawa K, Aizawa M (2017) *Rhinogobius mizunoi*, a new species of freshwater goby (Teleostei: Gobiidae) from Japan. *Kanagawa Kenritsu Hakubutsukan Kenkyu Hokoku, Shizen Kagaku* 46: 79–95. https://doi.org/10.32225/bkpmnh.2017.46_79
- Tan M, Armbruster JW (2018) Phylogenetic classification of extant genera of fishes of the order Cypriniformes (Teleostei: Ostariophysii). *Zootaxa* 4476(1): 34. <https://doi.org/10.11646/zootaxa.4476.1.4>

- Tang J, Qian M (1979) Fish Fauna in Dongting Lake. *Freshwater Fisheries*: 24–32.
- Tang W, Liu H, Ma J, Xiao R (1993) Effects of Jiangxi Wan'an dam on *Tenualosa reevesii* breeding in Ganjiang and its countermeasures. *Water Fisheries* 65: 18–19. <http://ir.ihb.ac.cn/handle/152342/5254>
- Tang W, Chen Y, Wu H (2001) Fish species diversity of Wulin Mountains region and its zoogeographic analyses. *Journal of Shanghai Fisheries University*.
- Tang Q, Liu H, Yang X, Nakajima T (2005) Molecular and morphological data suggest that *Spinibarbus caldwelli* (Nichols) (Teleostei: Cyprinidae) is a valid species. *Ichthyological Research* 52(1): 77–82. <https://doi.org/10.1007/s10228-004-0259-x>
- Tang Q, Li X, Yu D, Zhu YR, Ding BQ, Liu H, Danley PD (2018) *Saurogobio punctatus* sp. nov., a new cyprinid gudgeon (Teleostei: Cypriniformes) from the Yangtze River, based on both morphological and molecular data. *Journal of Fish Biology* 92(2): 347–364. <https://doi.org/10.1111/jfb.13498>
- Tchang TL (1933) The study of Chinese cyprinoid fishes, part 1. *Zoologia Sinica B* 2: 1–247.
- Tedesco PA, Beauchard O, Bigorne R, Blanchet S, Buisson L, Conti L, Cornu J-F, Dias MS, Grenouillet G, Hugueny B, Jezequel C, Leprieur F, Brosse S, Oberdorff T (2017) A global database on freshwater fish species occurrence in drainage basins. *Scientific Data* 4(1): e170141. <https://doi.org/10.1038/sdata.2017.141>
- Temminck CJ, Schlegel H (1845) Pisces. Siebold, P F de (ed): *Fauna Japonica, sive descriptio animalium, quae in itinere per Japoniam suscepto annis 1823–1830 collegit, notis, observationibus et adumbrationibus illustravit Ph Fr de Siebold Lugduni Batavorum* [Leiden] (A Arnz et soc) Parts 7–9: 113–172[, Pls 1–143 + A].
- The Forest Department of Hunan Province (2015) Hunan provincial key protected wildlife list. <http://lyj.hunan.gov.cn/> [Accessed 11 November 2021]
- Tregidgo D, Parry L, Barlow J, Pompeu PS (2021) Urban market amplifies strong species selectivity in Amazonian artisanal fisheries. *Neotropical Ichthyology* 19(3): e200097. <https://doi.org/10.1590/1982-0224-2021-0097>
- Van Der Laan R, Eschmeyer WN, Fricke R (2014) Family-group names of recent fishes. *Zootaxa* 3882(1): 1–230. <https://doi.org/10.11646/zootaxa.3882.1.1>
- Vasil'eva ED, Makeeva AP (2003) Taxonomic status of the black Amur bream and some remarks on problems of taxonomy of the genera *Megalobrama* and *Sinibrama* (Cyprinidae, Cultrinae). *Journal of Ichthyology* 43: 607–623.
- Wang H (1984) *Fishes in Beijing*. Beijing Press, 121 pp.
- Wang X (2019) Population genetic structure and spatial ecological process of two fish species in the Chishui River. PhD thesis, Wuhan: University of Chinese Academy of Sciences.
- Wang Z (2021) Updated species checklist and species diversity of fishes in the Gan-Jiang Basin of Jiangxi Province, South China. Master thesis, Wuhan: Institute of Hydrobiology, Chinese Academy of Sciences.
- Wang S, Dou H (1998) *Chinese Lakes*. Science Press, Beijing, 598 pp.
- Wang Z, Zhang E (2021) An updated species checklist of freshwater fishes from the Gan-Jiang. *Biodiversity Science* 29(9): 1256–1264. <https://doi.org/10.17520/biods.2021119>
- Wang JH, Wei QW, Zou YC (2011) Conservation strategies for the Chinese sturgeon, *Acipenser sinensis*: An overview on 30 years of practices and future needs. *Journal of Applied Ichthyology* 27(2): 176–180. <https://doi.org/10.1111/j.1439-0426.2011.01716.x>

- Wang H, Liu X, Wang H (2016) The Yangtze River Floodplain: Threats and Rehabilitation. American Fisheries Society Symposium, 263–291.
- Wang H, Liu X, Wang H (2019a) The Yangtze river-floodplain ecosystem: Multiple threats and holistic conservation. *Shui Sheng Sheng Wu Hsueh Bao* 43(S1): 157–182. <https://doi.org/10.7541/2019.178>
- Wang X, Liu F, Yu D, Liu H (2019b) Mitochondrial divergence suggests unexpected high species diversity in the opsariichthine fishes (Teleostei: Cyprinidae) and the revalidation of *Opsariichthys macrolepis*. *Ecology and Evolution* 9(5): 2664–2677. <https://doi.org/10.1002/ece3.4933>
- Wang D, Gao L, Tian H, Dong W, Duan X, Liu S, Chen D (2020) Population genetics and sympatric divergence of the freshwater gudgeon, *Gobiobotia filifer*, in the Yangtze River inferred from mitochondrial DNA. *Ecology and Evolution* 10(1): 50–58. <https://doi.org/10.1002/ece3.5746>
- Wu H (1930) Description de poissons nouveaux de Chine. *Bulletin du Muséum National d'Histoire Naturelle (Série 2)* 2: 255–259.
- Wu H (1977) Fish of Chinese Cyprinidae (II). Shanghai Scientific & Technical Publishers, Shanghai, 298 pp.
- Wu H, Zhong J (2008) Fauna Sinica, Osteichthyes Perciformes (V) Gobioidi [M], Science Press, Beijing, 568–635.
- Wu X, Chen Y, Chen X, Chen J (1981) The phylogeny of subfamilies and the relationship between phylogeny of subfamilies of *Cyprinus carpio*. *Science China* •••: 115–122.
- Wu JM, Wang CY, Zhang H, Du H, Liu ZG, Shen L, Wei QW, Rosenthal H (2015) Drastic decline in spawning activity of Chinese sturgeon *Acipenser sinensis* Gray 1835 in the remaining spawning ground of the Yangtze River since the construction of hydrodams. *Journal of Applied Ichthyology* 31(5): 839–842. <https://doi.org/10.1111/jai.12882>
- Wu J, Wang C, Zhang S, Zhang H, Du H, Liu Z, Wei Q (2017) From continuous to occasional: Small-scale natural reproduction of Chinese sturgeon occurred in the Gezhouba spawning ground, Yichang, China. *Journal of Fishery Sciences of China* 24(3): 425–431. <https://doi.org/10.3724/SPJ.1118.2017.17095>
- Wu Y, Li H, Liao F, Yang X, Xie Z (2021) The fish fauna of Hunan Province. Science Press, Beijing, 488 pp.
- Xie Z, Xie C, Zhang E (2003) Morphological variations among the Chinese species of *Sinibrama* (Pisces: Teleostei: Cyprinidae), with comments on their species validities. *Zoological Research* 24: 321–330. <https://doi.org/10.3321/j.issn:0254-5853.2003.05.001>
- Xuan Z, Jiang T, Liu H, Qiu C, Chen X, Yang J (2020) Are there still anadromous the estuarine tapetail anchovies *Colia nasus* in Dongting Lake? *Shui Sheng Sheng Wu Hsueh Bao* 44: 838–843. <https://doi.org/10.7541/2020.100>
- Yang J, Xiao W, Kuang X, Wei Z, Liu R (2000) Studies on the distribution, population size and the active regularity of *Lipotes vexillifer* and *Neophocaena phocaenoides* in Dongting Lake and Boyang Lake. *Changjiang Liuyu Ziyuan Yu Huanjing* 9: 444–450.
- Yang L, Sado T, Vincent Hirt M, Pasco-Viel E, Arunachalam M, Li J, Wang X, Freyhof J, Saitoh K, Simons AM, Miya M, He S, Mayden RL (2015a) Phylogeny and polyploidy: Resolving the classification of cyprinine fishes (Teleostei: Cypriniformes). *Molecular Phylogenetics and Evolution* 85: 97–116. <https://doi.org/10.1016/j.ympev.2015.01.014>

- Yang S, Li M, Zhu Q, Wang M, Liu H (2015b) Spatial and temporal variations of fish assemblages in Poyanghu Lake. *Changjiang Liuyu Ziyuan Yu Huanjing* 24: 54–64. <https://doi.org/10.11870/cjlyzyyhj201501008>
- Yang D, Song Y, Ma J, Li P, Zhang H, Price MRS, Li C, Jiang Z (2016) Stepping-stones and dispersal flow: Establishment of a meta-population of Milu (*Elaphurus davidianus*) through natural re-wilding. *Scientific Reports* 6(1): e27297. <https://doi.org/10.1038/srep27297>
- Yang T, Yu D, Gao X, Liu H (2020) Mechanism of fish community assembly in middle reaches of the Yangtze River. *Shui Sheng Sheng Wu Hsueh Bao* 44(5): 1045–1054. <https://doi.org/10.7541/2020.121>
- Ye J, Wu J, Yang X, Cong N, Dong T, Wang D (2016) Evaluation of suitability of *Squaliobarbus curriculus* and *Cirrhinus mrigala* as food fish for Mandarin fish farming. *Shanghai Haiyang Daxue Xuebao* ••: 569–574. <https://doi.org/10.12024/jso.2015100158>
- Yi B, Zhu Z (1959) Review of the genera *Culter* and *Erythroculter* of China. *Shui Sheng Sheng Wu Hsueh Bao* 3(002): 170–196. <http://ir.ihb.ac.cn/handle/152342/7266>
- Yu X, Luo T, Zhou H (2005) Large-scale patterns in species diversity of fishes in the Yangtze River Basin. *Shengwu Duoyangxing* 13(6): 473. <https://doi.org/10.1360/biodiv.050121>
- Yuan G, Ru H, Liu X (2010) Fish diversity and fishery resources in lakes of Yunnan Plateau during 2007–2008. *Hupo Kexue* 22: 837–841.
- Yuan X, Yang X, Ge H, Li H, Deng D (2019) Temporal distribution of fish community structure in Dongting Lake estuary. *Agricultural Sciences* 10(03): 294–301. <https://doi.org/10.4236/as.2019.103025>
- Yue P (2000) *Fauna Sinica: Osteichthyes Cypriniformes III*. Science Press, Beijing, 674 pp.
- Zhang E (1994) Phylogenetic relationship of the endemic Chinese cyprinid fish *Pseudogyrinocheilus prochilus*. *Zoological Research* 15: 26–35.
- Zhang X (2011) Population ecology of Yangtze finless porpoise in Dongting Lake and the adjacent waters. PhD thesis, Wuhan: Institute of Hydrobiology, Chinese Academy of Sciences.
- Zhang E, Cao W (2021a) China's Red List of Biodiversity-Fish (I). Science Press, Beijing, 377 pp.
- Zhang E, Cao W (2021b) China's Red List of Biodiversity-Fish (II). Science Press, Beijing, 525 pp.
- Zhang E, Chen Y (2006) Revised diagnosis of the genus *Bangana* Hamilton, 1822 (Pisces: Cyprinidae), with taxonomic and nomenclatural notes on the Chinese species. *Zootaxa* ••: 41–54. <http://biostor.org/reference/16771>
- Zhang Q, Hu G (2020) Utilization of species checklist data in revealing the spatial distribution of fish diversity. *Journal of Fish Biology* 97(3): 817–826. <https://doi.org/10.1111/jfb.14437>
- Zhang T, Li Z (2007) Fish resources and fishery utilization of Lake Poyang. *Hupo Kexue* 19(4): 434–444. <https://doi.org/10.18307/2007.0412>
- Zhang C, Zhao Y (2001) Migration of the Chinese sucker (*Myxocyprinus asiaticus*) in Yangtze River Basin with a discussion on the potential effects of the dams on fish. *Dong Wu Xue Bao* 47: 518–521. <https://doi.org/10.3969/j.issn.1674-5507.2001.05.007>
- Zhang C, Zhao Y (2016) Species Diversity and Distribution of Inland Fishes in China. Science Press, Beijing, 296 pp.

- Zhang E, Xie Z, Xie C (2004) Morphological variation between *Sinibrama macrops* and *S. wui*, with notes on their validities. *Shui Sheng Sheng Wu Hsueh Bao* •••: 511–518. <https://doi.org/10.3321/j.issn:1000-3207.2004.05.009>
- Zhang J, Li M, Xu M, Takita T, Wei F (2007) Molecular phylogeny of icefish Salangidae based on complete mtDNA cytochrome b sequences, with comments on estuarine fish evolution. *Biological Journal of the Linnean Society. Linnean Society of London* 91(2): 325–340. <https://doi.org/10.1111/j.1095-8312.2007.00785.x>
- Zhang H, Li JY, Wu JM, Wang CY, Du H, Wei QW, Kang M (2017) Ecological effects of the first dam on Yangtze main stream and future conservation recommendations: A review of the past 60 years. *Applied Ecology and Environmental Research* 15(4): 2081–2097. https://doi.org/10.15666/aecer/1504_20812097
- Zhang C, Yang J, Zhao Y, Pan X (2019) Fishes in the Jinsha Jiang River Basin, the upper reaches of the Yangtze River, China. Science Press, Beijing, 607 pp.
- Zhang H, Jaric I, Roberts DL, He Y, Du H, Wu J, Wang C, Wei Q (2020a) Extinction of one of the world's largest freshwater fishes: Lessons for conserving the endangered Yangtze fauna. *The Science of the Total Environment* 710: 136242. <https://doi.org/10.1016/j.scitotenv.2019.136242>
- Zhang X, Yang T, Luo X, Yuan C, Liu H (2020b) Fish phylogenetic community structure in the Poyang Lake and its tributary the Xiushui River in summer. *Shui Sheng Sheng Wu Hsueh Bao*. <https://doi.org/10.7541/2020.151>
- Zhao S, Fang J, Miao S, Gu B, Tao S, Peng C, Tang Z (2005) The 7-decade degradation of a large freshwater lake in central Yangtze River, China. *Environmental Science & Technology* 39: 431–436. <https://doi.org/10.1021/es0490875>
- Zhao K, García Molinos J, Zhang H, Zhang M, Xu J (2019) Contemporary changes in structural dynamics and socioeconomic drivers of inland fishery in China. *The Science of the Total Environment* 648: 1527–1535. <https://doi.org/10.1016/j.scitotenv.2018.08.196>
- Zheng L, Chen X, Yang J (2010) A new species of genus *Pseudogyrinocheilus* (Teleostei: Cyprinidae) from Guangxi, China. *Environmental Biology of Fishes* 87(2): 93–97. <https://doi.org/10.1007/s10641-009-9555-7>
- Zheng L, Chen X, Yang J (2019) Molecular phylogeny and systematic revision of *Bangana sensu lato* (Teleostei, Cyprinidae). *Journal of Zoological Systematics and Evolutionary Research* 57(4): 884–891. <https://doi.org/10.1111/jzs.12294>
- Zhou J, Wu Q, Ye Y, Tong J (2003) Genetic divergence between *Cyprinus carpio carpio* and *Cyprinus carpio haematopterus* as assessed by Mitochondrial DNA analysis, with emphasis on origin of European Domestic Carp. *Genetica* 119(1): 93–97. <https://doi.org/10.1023/A:1024421001015>
- Zhu S (1995) Synopsis of freshwater fishes of China. Jiangsu Science and Technology Publishing House, Nanjing, 562 pp.
- Zhu S, Liu Z, Gu X (2007) Changes of the fish fauna and fish yield analysis in Lake Taihu. *Hupo Kexue* 19(6): 664–669. <https://doi.org/10.18307/2007.0607>
- Zhu Y, Lv C, Hu H, Wang Z, Jia Y, He M, Huang X, Lei G (2014) Changes in fish community structure in West Dongting Lake after the operation of the Three Gorges Dam. *Hupo Kexue* 000: 844–852. <https://doi.org/10.18307/2014.0605>

- Zhu L, Yu D, Liu H (2020) *Zacco sinensis* sp. nov. (Cypriniformes: Cyprinidae), a new fish species from Northern China. *Sichuan Journal of Zoology* 039: 168–176. <https://doi.org/10.11984/j.issn.1000-7083.20190353>
- Zou Y, Zhang P, Zhang S, Chen X, Li F, Deng Z, Yang S, Zhang H, Li F, Xie Y (2019) Crucial sites and environmental variables for wintering migratory waterbird population distributions in the natural wetlands in East Dongting Lake, China. *The Science of the Total Environment* 655: 147–157. <https://doi.org/10.1016/j.scitotenv.2018.11.185>

Supplementary material I

Table S1

Authors: Xiao Chen

Data type: Table (docx. file)

Explanation note: Table S1. Geographical coordinates of 20 sampling sites in Lake Dongting.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.1108.79960.suppl1>