

Two new species and one new record of *Parakiefferiella* Thienemann, 1936 from China (Diptera, Chironomidae)

Wenbin Liu¹, Bingjiao Sun¹, Chuncai Yan², Chao Song¹, Xinhua Wang¹

1 College of Life Science, Nankai University, Tianjin 300071, China **2** Tianjin Key Laboratory of Animal and Plant Resistance, Tianjin Normal University, Tianjin, 300387, China

Corresponding author: Xinhua Wang (xhwang@nankai.edu.cn)

Academic editor: F. L. da Silva | Received 2 April 2015 | Accepted 2 November 2015 | Published 14 December 2015

<http://zoobank.org/FEC231B1-B08E-4008-92C7-2D319E551C42>

Citation: Liu W, Sun B, Yan C, Song C, Wang X (2015) Two new species and one new record of *Parakiefferiella* Thienemann, 1936 from China (Diptera, Chironomidae). ZooKeys 545: 139–148. doi: 10.3897/zookeys.545.6000

Abstract

Two new species of the genus *Parakiefferiella* Thienemann, 1936 (*P. fasciata* Liu & Wang, **sp. n.** and *P. liupanensis* Liu & Wang, **sp. n.**) are described as adult males from China. *Parakiefferiella tamatriangulata* Sasa is recorded from China for the first time. A key to the known adult males of *Parakiefferiella* from China is presented. The generic diagnosis of male *Parakiefferiella* given by Cranston et al. (1989) is emended.

Keywords

Chironomidae, *Parakiefferiella*, new species, key, taxonomy

Introduction

The genus *Parakiefferiella* was erected by Thienemann in 1936, with *P. coronata* (Edwards, 1929) as type species. According to Cranston et al. (1989), it can be separated from other orthoclad genera by the following combination of characters: eyes bare, without dorsomedial extension; acrostichals absent but scutum with median tuft of microtrichia, sometimes on hump; wing membrane without setae, with fine punctation; squama bare; transverse sternapodeme gently curved, with developed oral projections. Virga strongly developed. Gonostylus with more or less well pronounced curve

or bend and slight or absent crista dorsalis. Larval stages of the genus are found in both running and standing waters. Adults fly during spring and summer, rarely in large numbers.

The genus is probably quite speciose. Tuiskunen (1986) described four new species and redescribed eight species from Fennoscandian region. Moubayed (1991, 1994) and recently Moubayed-Breil and Langton (2004) described three additional species: *P. pyrenaica* Moubayed, 1991 (France); *P. wuelkeri* Moubayed, 1994 (western Europe and north Africa); *P. normandiana* Moubayed-Breil & Langton, 2004 (France, Germany and England). According to the catalog of Japanese Orthoclaadiinae (Yamamoto 2004), ten valid species were recorded in Japan. Nine species were reported in the Russian Far East by Makarchenko and Makarchenko (2010). To date, 44 species of the genus have been recorded worldwide, of which 33 are Palaearctic, seven are Nearctic, three are Oriental, four are Neotropical and four are Afrotropical (Ashe and O'Connor 2012; Ree, Nam and Jeong 2012).

So far, three species of the genus (*P. bathophila* (Kieffer, 1912), *P. coronata* (Edwards, 1929) and *P. tipuliformis* (Tokunaga, 1940)) were recorded in China (Wang 2000). The species *P. tipuliformis* (Tokunaga, 1940) from Taiwan province previously placed in *Spaniotoma* (*Smittia*) by Tokunaga (1940) was transferred to *Parakiefferiella* by Sasa and Kikuchi (1995). The species diagnostic characters fit the genus *Parakiefferiella*. Moreover, *P. coronata* (Edwards, 1929) was recorded just as larval stage (Wang 2000).

Based on specimens from China, two new species are described in this paper, and a key to the species of *Parakiefferiella* in China is provided.

Materials and methods

Morphology and terminology follow Sæther (1980). The material examined was slide-mounted following the procedures outlined by Sæther (1969). Measurements are given as the range followed by the mean. Color is described as observed in specimen preserved in alcohol. The specimens examined in this study are deposited in the College of Life Sciences, Nankai University, China (BDN).

Taxonomy

Parakiefferiella Thienemann, 1936

Emended diagnosis. Based on the material examined and references, the generic diagnosis of male *Parakiefferiella* (Cranston et al. 1989) must be emended as follows: antenna with 12–13 or occasionally with five (*P. gynocera* (Edwards, 1937)) or ten (*P. fasciata* Liu & Wang, sp. n. and *P. liupanensis* Liu & Wang, sp. n.) flagellomeres; wing anal lobe weakly to moderately developed (*P. bathophila* (Kieffer, 1912)); R and R₁ with few setae, occasionally all veins bare (*P. fasciata* Liu & Wang, sp. n. and *P.*

liupanensis Liu & Wang, sp. n.); anal point with 2–8 basal setae, sometimes bare (*P. fasciata* Liu & Wang, sp. n.).

***Parakiefferiella fasciata* Liu & Wang, sp. n.**

<http://zoobank.org/4FC0DE0C-3F8D-4746-938E-DBD5E5188AB6>

Figs 1–7

Type material. Holotype: ♂ (BDN. No.1165), China, Shandong Province, Yantai City, Kunyu Mountain, 37°30'10"N, 121°23'40"E, 24.viii.1987, sweeping net, Wang XH. The specimens were sealed with Canada balsam on slides. Paratypes: 3 ♂♂, data as holotype.

Diagnosis. The adult male can be distinguished from known species of the genus by the following combination of characters: anal point lacking keel, very broad at base, rounded apically, without basal setae; virga consisting of four spines; antenna with ten flagellomeres, AR 0.47–0.50; tergites III and VIII dark brown, others tergites pale yellow; HR 1.98–2.06; HV 2.65–3.00.

Description. Male imago (n = 4). Total length 1.04–1.09, 1.07 mm. Wing length 0.71–0.72, 0.71 mm. Total length/wing length 1.45–1.56, 1.47. Wing length/length of profemur 3.24–3.29, 3.26.

Coloration. Head brown. Thorax brown with dark spot. Tergites (Fig. 4) III and VIII dark brown, others tergites pale yellow.


Head. Antenna (Fig. 2) with ten flagellomeres. AR 0.47–0.50, 0.49. Ultimate flagellomere 137–144, 140 µm long. Inner vertical 1. Clypeus with four setae. Tentorium 55–72, 66 µm long, 5–7, 6 µm wide. Palpomere lengths (in µm): 10–12, 11; 14–19, 16; 26–29, 28; 43–48, 47; 88–96, 92. L: 5th/3rd: 3.26–3.55, 3.43

Wing (Fig. 1). Anal lobe reduced. VR 1.12–1.21, 1.17. Costal extension 40 µm long. Brachiolum with one seta. All veins bare. Squama bare.

Thorax (Fig. 3). Anteprenotal lobes developed. Anteprenotum with one lateral seta. Dorsocentrals 2–4, 3; one prealars. Scutellum with 2–6, 5 setae. Mesonotum with a tuft of hairs.

Legs. Spur of fore tibia 24–25, 24 µm long; spurs of mid tibia 9–12, 11 µm and 12–14, 13 µm long; spurs of hind tibia 9–10, 9 µm and 20–22, 21 µm long. Hind tibial comb with 9–12, 11 spines, 19–38, 30 µm long. Width at apex of fore tibia 17–19, 19 mm, of mid tibia 19–20, 19 mm, of hind tibia 21–24, 23 mm. Lengths (in µm) and proportions of legs as in Table 1.

Hypopygium (Figs 5–7). Anal point triangular and hyaline, very broad at base, with rounded apex and without basal seta. Laterosernite IX with 2–4, 3 setae. Phallapodeme 20–24, 23 µm long. Transverse sternapodeme 48–54, 53 µm long. Virga consisting of four spines, 20–35, 25 µm long. Tergites III and VIII bearing colored bands. Gonocoxite 72–74, 73 µm long, outer margins concave medially. Superior volsella absent; inferior volsella rounded and very large, occupying nearly half of the total length of gonocoxite, lacking dorsal setae and densely covered with microtrichia (Fig. 6); ventral



Figures 1–7. *Parakiefferiella fasciata* Liu & Wang, sp. n., male. **1** wing **2** antenna **3** thorax **4** abdomen tergites coloration **5** hypopygium (dorsal view) **6** inferior; **7** hypopygium (ventral view).

margin strongly swollen medially. Gonostylus 36–38, 37 μm long, anterior margin nearly straight, posterior margin slightly curved in dorsal view. Megaseta 5–7, 7 μm long. Crista dorsalis reduced, slightly visible in lateral view. HR 1.98–2.06, 2.01. HV 2.65–3.00, 2.75.

Table 1. Lengths (in μm) and proportions of legs of *Parakiefferiella fasciata* Liu & Wang, sp. n. (n = 4)

	P ₁	P ₂	P ₃
fe	216–220, 219	288–300, 293	240–258, 256
ti	247–264, 254	240–249, 243	276–282, 278
ta ₁	120–130, 121	108–120, 112	124–139, 131
ta ₂	57–67, 59	57–67, 62	60–68, 66
ta ₃	48–52, 49	43–48, 47	67–76, 71
ta ₄	29–31, 30	29–31, 30	40–43, 41
ta ₅	31–36, 34	30–33, 32	33–38, 34
LR	0.45–0.52, 0.48	0.41–0.44, 0.43	0.45–0.51, 0.48
BV	3.33–3.53, 3.43	3.61–3.87, 3.73	3.07–3.10, 3.09
SV	3.86–4.00, 3.98	4.89–5.00, 4.90	3.88–4.13, 4.01
BR	2.50–2.75, 2.71	3.08–3.25, 3.12	3.50–3.75, 3.54

Etymology. The specific name is from Latin *fasciata*, referring to different colored bands in its tergites.

Remarks. The new described species is distinguished from all other *Parakiefferiella* species by the following unusual combination of characters: antenna with ten flagellomeres; tergites III and VIII with colored bands; anal point lacking both dorsal setae and keel; absence of dorsal setae on inferior volsella, which is bearing only microtrichia.

While the new species close related to *P. coronata* (Edwards, 1929; Makarchenko and Makarchenko 2010). However, the new species differs from the latter on the basis of following characters: (1) the antenna of the new species with ten flagellomeres, whereas *P. coronata* has 13 flagellomeres; (2) the superior volsella of *P. coronata* is large, whereas in the new species it is absent; (3) the new species has abdomen pale yellow with tergites III–VIII dark brown, whereas *P. coronata* has abdomen dark with tergites IV, VI–VII shiny.

Female and immature stages unknown.


***Parakiefferiella liupanensis* Liu & Wang, sp. n.**

<http://zoobank.org/816E951D-6283-4643-9640-C9EDC141A97E>

Figs 8–13

Type material. Holotype: ♂ (BDN. No.1168), China, Ningxia Hui Autonomous Region, Liupan Mountain, Erlong river, 35°38'40"N, 106°31'40"E, 6.viii.1987, sweeping method, Wang XH. Paratypes: 4 ♂♂, as holotype.

Diagnosis. The adult male can be distinguished from known species of the genus by the following combination of characters: anal point obtuse triangular with pointed and hyaline apex, lacking keel and bearing four setae placed laterally (two on each side); antenna with ten flagellomeres; all of the veins bare; inferior volsella square with



Figures 8–13. *Parakiefferiella liupanensis* Liu & Wang, sp. n., male. **8** wing **9** antenna **10** thorax **11** hypopygium (dorsal view) **12** inferior **13** hypopygium (ventral view).

rounded inner margin, not projected, occupying about half of the gonocoxite length, lacking dorsal setae and covered with microtrichia; ventral margin swollen medially; AR 0.33–0.37; HR 2.20–2.28; HV 2.36–2.43.

Description. Male imago (n = 5). Total length 1.20–1.25, 1.23 mm. Wing length 0.78–0.85, 0.81 mm. Total length/wing length 1.45–1.47, 1.46. Wing length/length of profemur 3.38–3.58, 3.45.

Coloration of preserved specimens. Head and tergites brown. Thorax brown with dark spot.

Head. Antenna (Fig. 9) with ten flagellomeres. AR 0.33–0.37. Ultimate flagellomere 108–115, 112 μm long. Temporal seta one, including one outer vertical. Clypeus with 3–4 setae. Tentorium 62–72, 69 μm long, 7–9, 8 μm wide. Palpomere lengths (in μm): 12–14, 13; 24–26, 25; 28–34, 31; 28–36, 32; 52–54, 53. L: 5th/3rd: 1.61–1.83, 1.81.

Wing (Fig. 8). Anal lobe moderate reduced. VR 1.02–1.18, 1.14. Costal extension 40 μm long. Brachiolum with one seta. All of the veins bare. Squama bare.

Thorax (Fig. 10). Anteprenotal lobes developed. Anteprenotum with one lateral seta. Dorsocentrals 5–7, 6; acrostichals absent; two prealars. Scutellum with four setae. Mesonotum with a tuft of hairs.

Legs. Spur of fore tibia 12–21, 18 μm long; spurs of mid tibia 9–12, 11 μm and 12–14, 13 μm long; spurs of hind tibia 9–10, 9 μm and 20–22, 21 μm long. Hind tibial comb with 9–12, 10 spines, 19–38, 23 μm long. Width at apex of fore tibia 19–22, 20 mm, of mid tibia 12–22, 17 mm, of hind tibia 18–22, 21 mm. Lengths (in μm) and proportions of legs as in Table 2.

Hypopygium (Figs 11–13). Anal point obtuse triangular with pointed and hyaline apex, lacking keel and bearing 4 setae placed laterally (2 on each side). Laterosernite IX with 2–4, 3 setae. Phallapodeme 19–21, 20 μm long. Transverse sternapodeme, straight 44–48, 46 μm long. Virga consisting of two spines, 20–25, 23 μm long. Gonocoxite 79–82, 80 μm long. Superior volsella absent; inferior volsella square, occupying about half of the gonocoxite length, lacking dorsal setae and covered with microtrichia (Fig. 12); ventral margin swollen medially. Gonostylus 36–38, 37 μm long, narrowed apically, distinctly bent and curved inward in dorsal view; crista dorsal

Table 2. Lengths (in μm) and proportions of legs of *Parakiefferiella liupanensis* Liu & Wang, sp. n. (n = 5)

	P ₁	P ₂	P ₃
fe	236–242, 239	264–272, 268	239–248, 241
ti	288–302, 298	216–241, 232	252–272, 265
ta ₁	141–153, 145	108–120, 109	132–137, 136
ta ₂	98–116, 111	60–64, 63	72–80, 74
ta ₃	72–76, 74	50–54, 52	74–79, 76
ta ₄	36–40, 38	26–31, 29	36–48, 43
ta ₅	31–36, 34	30–33, 31	33–38, 36
LR	0.49–0.53, 0.51	0.45–0.50, 0.49	0.48–0.52, 0.50
BV	2.62–2.72, 2.65	3.61–3.66, 3.63	2.85–2.89, 2.87
SV	3.50–3.66, 3.61	4.69–4.77, 4.71	3.73–3.84, 3.79
BR	1.25–1.75, 1.50	2.50–2.75, 2.60	3.40–3.75, 3.60

absent. Megaseta 5–7, 7 μm long. Without crista dorsalis. HR 2.20–2.28, 2.23. HV 2.36–2.43, 2.41.

Etymology. The specific name refers to the Liupan Mountain where the species was collected.

Remarks. The new species can easily be separated from other related *Parakiefferiella* species by the following characters: antenna with 10 segments; anal point square, lacking keel and bearing four setae laterally; inferior volsella without dorsal setae and covered with microtrichia.

P. liupanensis Liu & Wang, sp. n. is closely related to *P. bathophila* (Kiffer, 1912). However, the new species differs from the latter on the basis of the following characters: (1) the antenna of the new species with ten flagellomeres, whereas *P. bathophila* possesses 13 flagellomeres; (2) the hypopygium of *P. bathophila* with triangular superior volsella, whereas in the new species it is absent; (3) the anal point of the new species obtuse triangular, not tapered apically, whereas *P. bathophila* gradually tapered to apex; (4) the shape of inferior volsella in the new species square, without dorsal setae, whereas *P. bathophila* triangular, with dorsal setae.

Female and immature stages unknown.

Parakiefferiella tamatriangulata Sasa, 1981

Parakiefferiella tamatriangulata Sasa, 1981: 94; Sasa and Arakawa 1994: 99.

Specimens examined. 1 ♂, Liaoning Province, Dandong City, Fengcheng County, 40°51'00"N, 124°07'00"E, 25.iv. 1992, Sweeping net, Wang JC.

Diagnostic characters. Total length 2.06–2.29 mm. Wing length 1.02–1.37 mm. AR 0.38–0.53. Anal lobe reduced. Laterosernite IX with five setae. Phallapodeme 36–38 μm long. Transverse sternapodeme 74–81 μm long. Anal point small, triangular and with pointed apex, bare and hyaline excepting the basal portion with microtrichia. Inferior volsella semicircular, bearing dorsal setae. Gonostylus 6 μm long, narrowed apically, distinctly bent and curved inward in dorsal view. Megaseta 10 μm long. HR 2.18. HV 1.56.

Remarks. The Chinese specimens mainly agree with the original description by Sasa (1981), but they have higher AR (0.53) than the specimens from Japan (0.38–0.48).

Distribution. China (Palearctic China: Liaoning Province), Japan.

Key to known adult males of the genus *Parakiefferiella* in China

- | | | |
|---|---|---------------------------------------|
| 1 | Tergites banded | 2 |
| – | Tergites unicolor..... | 3 |
| 2 | Tergites III and VIII dark brown, others pale yellow; all veins bare..... | |
| | | <i>P. fasciata</i> Liu & Wang, sp. n. |

- All tergites with a narrow dark band on caudal margin; veins very delicate and colorless *P. tipuliformis* (Tokunaga)
- 3 Antenna with 13 flagellomeres; inferior volsella triangular or semicircular, bearing dorsal setae 4
- Antenna with 10 flagellomeres; inferior volsella square lacking dorsal setae...
..... *P. liupanensis* Liu & Wang, sp. n.
- 4 Anal lobe developed; inferior volsella large, roughly triangular.....
..... *P. bathophila* (Kieffer)
- Anal lobe reduced; inferior volsella semicircular..... *P. tamatriangulata* Sasa

Acknowledgements

Financial support from National Natural Science Foundation of China (NSFC, grant No. 31272284; 31301908; 31460572; J1210005), Fauna of China (FY120100) and Natural Science Foundation of Tianjin (14JCQNJC14600) are acknowledged with thanks. We are grateful to Bingchun Ji and Yufen Li who made parts of slide preparations.

References

- Ashe P, O'Connor JP (2012) A World Catalogue of Chironomidae (Diptera). Part 2B. Orthocladiinae. Irish Biogeographical Society & National Museum of Ireland, Dublin, 457–466.
- Cranston PS, Oliver DR, Sæther OA (1989) The adult males of Orthocladiinae (Diptera: Chironomidae) of the Holarctic region - Keys and diagnoses. In: Wiederholm T (Ed.) Chironomidae of the Holarctic region. Keys and diagnoses. Part 3. Adult males. Entomological Scandinavica, Supplement 34: 219–220.
- Edwards FW (1929) British non-biting midges (Diptera: Chironomidae). Transactions of the Royal Entomological Society of London 77: 279–439. doi: 10.1111/j.1365-2311.1929.tb00692.x
- Edwards FW (1937) Chironomidae (Diptera) collected by Prof. Aug. Thienemann in Swedish Lappland. Annals and Magazine of Natural History 20(10): 140–148. doi: 10.1080/00222933708655324
- Kieffer JJ (1912) Quelques nouveaux Tendipedides (Diptera) obtenus declosion (1re Note). Bulletin de la Société Entomologique de France 31: 86–88.
- Makarchenko EA, Makarchenko MA (2010) A review of *Parakiefferiella* Thienemann (Diptera, Chironomidae, Orthocladiinae) from the Russian Far East. Evraziatskii entomologicheskii Zhurnal 9(3): 397–410.
- Moubayed J (1991) Male, female and pupal exuviae of *Parakiefferiella pyrenaica* sp. n. (Diptera, Chironomidae). Nouvelle Revue d'Entomologie (Nouvelle Série) 8(1): 71–75.
- Moubayed J (1994) *Parakiefferiella wuelkeri* n. sp. (Diptera: Chironomidae) from western Europe and North Africa. British Journal of Entomology and Natural History 7: 7–10.

- Moubayed-Breil J, Langton PH (2004) Description of *Parakiefferiella normandiana* sp. n. from France (Diptera: Chironomidae). *Ephemera* 3(1): 9–16.
- Ree HI, Jeong KY (2010) Fauna of non-biting midges (Diptera, Chironomidae) from Soyang River in Chuncheson-si, Gangwon-do, Korea. *Korean Journal of Systematic Zoology* 26(2): 123–226. doi: 10.5635/ASED.2012.28.1.002
- Ree HI, Nam SH, Jeong KY (2012) Two new and four unrecorded species of Chironomidae (Diptera) in Korea. *Animal systematics, Evolution and Diversity* 28(1): 2–11.
- Sæther OA (1969) Some Nearctic Podonominae, Diamesinae and Orthocladiinae (Diptera: Chironomidae). *Bulletin of the Fishers Research Board of Canada* 170: 1–154.
- Sæther OA (1980) Glossary of Chironomid morphology terminology (Diptera: Chironomidae). *Entomologica Scandinavica, Supplement* 14: 1–51.
- Sasa M (1981) Studies on Chironomid midges of the Tama River. Part 4. Chironomidae recorded at a winter survey. *Research Report from the National Institute for Environmental Studies*. 29: 79–148.
- Sasa M, Arakawa R (1994) Seasonal changes of the chironomid species emerging from Lake Furudo. *Research Report Toyama Prefectural Environmental Pollution Research Center*. Part 3: 88–109.
- Sasa M, Kikuchi M (1995) Chironomidae (Diptera) of Japan. University of Tokyo Press, 333 pp.
- Thienemann A (1936) Alpine Chironomiden. *Ergebnisse von Untersuchungen in der Gegend von Garmisch-Partenkirchen, Oberbayern*. *Archiv für Hydrobiologie* 30: 167–262.
- Tokunaga M (1940) Chironomidae from Japan. XII. New or little known Ceratopogonidae and Chironomidae. *Philippine Journal Science* 72(3): 255–317.
- Tuiskunen J (1986) The Fennoscandian species of *Parakiefferiella* Thienemann (Diptera, Chironomidae, Orthocladiinae). *Annales Zoologici Fennici* 23: 175–196.
- Wang XH (2000) A revised checklist of Chironomidae from China (Diptera). In: Hoffrichter O (Ed.) *Late 20th Century Research on Chironomidae: An Anthology from the 13th International Symposium on Chironomidae*. Shaker Verlag, Aachen, 639.
- Yamamoto M (2004) A catalog of Japanese Orthocladiinae (Diptera: Chironomidae). *Makunagi* 21: 1–121.