

Complete mitochondrial genome of the freshwater fish *Onychostoma lepturum* (Teleostei, Cyprinidae): genome characterization and phylogenetic analysis

I-Chen Wang^{1*}, Hung-Du Lin^{2*}, Chih-Ming Liang¹, Chi-Chun Huang³,
Rong-Da Wang⁴, Jin-Quan Yang⁴, Wei-Kuang Wang¹

1 Department of Environmental Engineering and Science, Feng Chia University, Taichung 407, Taiwan **2** The Affiliated School of National Tainan First Senior High School, Tainan 701, Taiwan **3** Taiwan Endemic Species Research Institute, Nantou 552, Taiwan **4** Shanghai Universities Key Laboratory of Marine Animal Taxonomy and Evolution, Shanghai Ocean University, Shanghai 201306, China

Corresponding author: Wei-Kuang Wang (wkwang@fcu.edu.tw)

Academic editor: M.E. Bichuette | Received 13 August 2020 | Accepted 25 November 2020 | Published 18 December 2020

<http://zoobank.org/92DDFB54-0FC4-404D-BF2C-7C0B373E2992>

Citation: Wang I-C, Lin H-D, Liang C-M, Huang C-C, Wang R-D, Yang J-Q, Wang W-K (2020) Complete mitochondrial genome of the freshwater fish *Onychostoma lepturum* (Teleostei, Cyprinidae): genome characterization and phylogenetic analysis. ZooKeys 1005: 57–72. <https://doi.org/10.3897/zookeys.1005.57592>

Abstract

The cyprinid genus *Onychostoma* Günther, 1896 consists of 24 valid species distributed in Southeast Asia, including Taiwan, Hainan, mainland China and the Indochina region. In the present study, we determined the complete mitochondrial genome of *O. lepturum*, which is 16,598 bp in length, containing 13 protein-coding genes, two rRNA genes, 22 tRNA genes and a typical control region (D-loop). To verify the molecular phylogeny of the subfamily Acrossocheilinae, we provide new insights to better understand the taxonomic status of *Acrossocheilus*, *Onychostoma* and *Folifer brevifilis*. The phylogenetic trees presented three major clades based on the 13 protein-coding genes from 28 Acrossocheilinae species. Clades I and II represent the *Onychostoma* and *Acrossocheilus* groups, respectively. Species of *Acrossocheilus*, *Onychostoma* and *F. brevifilis* are included in Clade III, which is considered as an ancestral group. This work provides genomic variation information and improves our understanding of the Acrossocheilinae mitogenome, which will be most valuable in providing new insights for phylogenetic analysis and population genetics research.

Keywords

Cyprinid, mitogenome, *Onychostoma lepturum*, phylogeny, population genetics, Southeast Asia

* These authors contributed equally to this work.

Introduction

The cyprinid genus *Onychostoma* Günther, 1896 consists of 24 valid species distributed in Southeast Asia, including Taiwan, Hainan, mainland China and the Indochina region (Song et al. 2018; Froese 2019). *Onychostoma* is composed of mountain stream cyprinids that are characterized primarily by the possession of a sharp, cornified sheath cutting edge in the lower jaw and no fleshy lower lip (Shan et al. 2000). Among these species, four species are only distributed in the Indochina region, including Laos Vietnam and Thailand, eleven species are endemic to mainland China, one species is endemic to Taiwan island, and the remaining seven species are shared with mainland China and the Indochina region (Froese 2019). *Onychostoma lepturum* (Boulenger, 1900) is distributed in Laos and Vietnam and can also be found in the Yuanjiang River in mainland China and Hainan Island (Shan et al. 2000; Kottelat 2001).

The Cyprinidae family has the most species of any freshwater fish family. The family encompasses 11 subfamilies, with the genus *Onychostoma* belonging to the Acrossocheilinae subfamily (Yang et al. 2015). The taxonomic placement of the Asian genus *Onychostoma* has undergone many changes. *Onychostoma* was synonymized with the African genus *Varicorhinus* (e.g., Taki 1975), but recent studies demonstrate that *Onychostoma* and *Varicorhinus* fall in two distinct major clades in terms of chromosome numbers and molecular markers (Yang et al. 2015). According to previous studies, *Acrossocheilus* and *Onychostoma* were reported to be paraphyletic or polyphyletic (Yang et al. 2015; Wang et al. 2016; Zheng et al. 2016). Among these were Acrossocheilini, *Acrossocheilus*, *Onychostoma* and *Folifer brevifilis* (Peters, 1881), which formed a clade in the molecular analysis (Yang et al. 2015; Wang et al. 2016; Zheng et al. 2016). Three genera (*Acrossocheilus*, *Onychostoma* and *Folifer*) have been previously placed in different taxa together with many other cyprinines (Chen et al. 1984; Yue 2000). However, the phylogenetic relationships of these three genera, based on different molecular markers, has been controversial.

Folifer brevifilis is closely related to *O. simum* (Sauvage & Dabry de Thiersant, 1874) based on the mitochondrial and nuclear markers (Yang et al. 2015; Wang et al. 2016; Zheng et al. 2016). According to mitochondrial genomes, *O. simum* and *O. gerlachi* (Peters, 1881) were in sister groups (Zhang et al. 2018). The genus *Acrossocheilus* represents three separate lineages: the barred species (e.g., *A. beijiangensis* (Wu & Lin, 1977) and *A. iridescens* (Nichols & Pope, 1927)), non-barred species (e.g., *A. yunnanensis* (Regan, 1904)) and *A. monticola* (Günther, 1888) based on mitochondrial DNA sequences (Zheng et al. 2016). However, Hou et al. (2020) reassigned species of the *Acrossocheilus* cluster into two separate clades: Clade I (an ancestral clade), including *A. monticola* and *A. yunnanensis*, and Clade II, which was further divided into two sublineages (subclades A and B) based on available whole mitochondrial genomes. Subclade B clustered with *A. longipinnis* (Wu, 1939), *A. iridescens* and *A. barbodon* (Nichols & Pope, 1927), and subclade A included other *Acrossocheilus* species (e.g., *A. parallens* (Nichols, 1931), *A. hemispinus* (Nichols, 1925), *A. jishouensis* (Zhao, Chen & Li, 1997) (Hou et al. 2020). Yang et al. (2015) and Zheng et al. (2016) proposed that members of *Onychostoma* be divided into three groups based on morphological and molecular data. Zhang et al. (2018) also showed that the

eight species of *Onychostoma* cluster into three separate lineages based on the whole mitochondrial genome sequence. However, similar research in Zhai et al. (2020) only identified two lineages among nine species of *Onychostoma* with the same molecular markers. Interestingly, Zhang et al. (2018) proposed that *O. rarum* (Lin, 1933) was the sister group of *O. alticorpus* (Oshima, 1920), but recently another study showed that *O. rarum* seems to be more closely related to *O. barbatulum* (Pellegrin, 1908) and *O. barbatum* (Lin, 1931) (Zhai et al. 2020). In the previous studies, the classification of the three groups was inconsistent (Yang et al. 2015; Zheng et al. 2016; Zhang et al. 2018). For example, *O. lepturum* (Boulenger, 1900) clustered together with *O. meridionale* (Kottelat, 1998) (Yang et al. 2015). In addition, Wang et al. (2016) and Zheng et al. (2016) proposed that *O. lepturum* was nested with *O. gerlachi*.

Recent population analyses suggested that the nucleotide diversity of cyprinids on Hainan Island was lower than that of cyprinids in mainland China (Zhou et al. 2017). A decline in freshwater fish resources has been observed due to the effects of overfishing, water pollution and environmental deterioration in China (Kang et al. 2013, 2014). The phylogenetic relationships among species in the genera *Onychostoma* and *Acrossocheilus* have been studied based on morphological data and on nuclear and mitochondrial genes (Wang et al. 2007; Xin 2008; Wang et al. 2016; Zheng et al. 2016). However, the current understanding among members of the genera *Onychostoma* and *Acrossocheilus* and their internal phylogenetic relationships remains confusing. The complete mitochondrial genome has alternative molecular markers for phylogenetic analysis capable of providing much more robust phylogenetic reconstructions than smaller portions of the mtDNA (Huang et al. 2017; Hou et al. 2020). Mitogenomes are thought to be reliable markers for reconstructing phylogenies in recent taxonomic and phylogenetic studies of cyprinids (Huang et al. 2017; Chung et al. 2020).

Previous studies suggest an inclusive phylogenetic clade including species from *Acrossocheilus*, *Onychostoma*, and *Folifer brevifilis* based on molecular markers (Yang et al. 2015; Zheng et al. 2016). Although a previous study has characterized the complete mitochondrial genome of *O. lepturum* (Zhai et al. 2020), the genome annotation, comparative analysis and the phylogenetic relationships of *Onychostoma* remain poorly understood due to the limited genomic data used. Our approach better informs the conservation of this species; thus, we determined the complete mitochondrial genome of *O. lepturum* based on next-generation sequencing data and assessed its phylogenetic relationships with another 11 available mitogenomes in the genus *Onychostoma* and 16 available mitogenomes in the genus *Acrossocheilus* and *F. brevifilis*, with an available mitogenome in *Spinibarbus hollandi* (Oshima, 1919) used as an outgroup.

Materials and methods

Sample and DNA extraction

The sample of *Onychostoma lepturum* was caught from the Lingshui River in Baoting County of Hainan in China (18°42'07"N, 109°40'44"E). Samples were collected from

the field sites with seines, fatally anesthetized with MS-222 (Sigma, St. Louis, MO) and fixed and stored in 100% ethanol. All specimens are lodged in the laboratory of Jin-Quan Yang, Shanghai Ocean University, Key Laboratory of Exploration and Utilization of Aquatic Genetic Resources. All animal experiments were carried out in accordance with the guidelines and with approval of the Animal Research and Ethics Committee of Shanghai Ocean University (permissions, SHOU-DW-2018-021). Total genomic DNA was extracted from muscle tissue using the Genomic DNA Purification Kit (Genra Systems, Valencia, CA) in the laboratory.

Sequencing and genome annotation and analysis

The complete mitogenome of *O. lepturum* was obtained from high-throughput sequencing of whole-genome DNA with a HiSeqX Ten platform (Illumina, San Diego, CA) with a paired-end 150 bp approach. Next-generation sequencing (NGS) was used to perform low-coverage whole-genome sequencing to obtain the complete mitogenome according to a previous protocol (Chiu et al. 2018). By using commercial software (Geneious V9, Auckland, New Zealand), approximately 1.1% of raw reads (34,001 out of 29,140,518) were assembled de novo to produce a single, circular form of the complete mitogenome with an average coverage of 245 X. Compared to the corresponding complete mtDNA sequence of genus *Onychostoma*, 13 protein-coding genes and ribosomal RNA (rRNA) genes were identified using Clustal X 1.83 (<http://www.clustal.org/>). Codon usage, nucleotide substitution and base composition were determined using MEGA X (Kumar et al. 2018), and the skewing of the nucleotide composition was measured in terms of AT- and GC-skews according to the following formulas: AT-skew = $(A - T)/(A + T)$ and GC-skew = $(G - C)/(G + C)$ (Perna and Kocher 1995).

Phylogenetic analysis

Phylogenetic analyses using a total of 11 mitogenomes of *Onychostoma* species, 16 mitogenomes of *Acrossocheilus* species and one mitogenome of *F. brevifilis* were performed based on Neighbor-joining (NJ), Maximum-likelihood (ML), and Bayesian (BI) methods, with *Spinibarbus hollandi* as the outgroup. Twenty-nine mitogenomes were downloaded from NCBI and were aligned using MEGA X (alignment with CLUSTALW) with default settings (Kumar et al. 2018). The best model GTR + G (General Time Reversible model with Gamma distributed rates among sites) was chosen based on the Akaike information criterion (AIC) using the smart model selection algorithm (Lefort et al. 2017), and the ML trees were constructed using PhyML 3.0 software (Guindon et al. 2010). The statistical confidences were assessed through the bootstrap test inferred from 1000 replicates. A NJ tree was constructed based on the Kimura 2-parameter model with 1000 bootstrap replicates using MEGA X (Kumar et al. 2018). The Bayesian inference (BI) tree was conducted using the GTR+G model strategy with MrBayes 3.2.6 (Ronquist and Huelsenbeck 2003), and two independent Markov Chains Monte Carlo (MCMC) chains were run for 5×10^7 generations; the first 50,000 trees before stationarity were discarded as burn-in, and the remaining trees were used to construct

the majority-rule consensus trees. Effective sample size (ESS) values, as computed by plotting the log likelihood scores against the generation times using the program Tracer 1.7 (Rambaut et al. 2018), were above 200 for the convergence of MCMC runs.

Results

Mitochondrial genomic structure and composition

In the present study, the complete mitochondrial genome sequence of *O. lepturum* derived by NGS was found to be 16,598 bp and was deposited in GenBank (accession MT258556). The mitogenome contained 37 typical mitochondrial genes with 13 typical vertebrate protein-coding genes, 2 ribosomal RNA (rRNA) genes, 22 tRNAs, and a control region (D-Loop) (Fig. 1). Most of the *O. lepturum* mitochondrial genes were encoded on the H-strand, although the ND6 and eight tRNA genes (tRNA^{Gln}, tRNA^{Ala}, tRNA^{Asn}, tRNA^{Cys}, tRNA^{Tyr}, tRNA^{Ser}, tRNA^{Glu} and tRNA^{Pro}) were encoded on the L-strand (Fig. 1). The total length was found to be similar to that of the other *Onychostoma* sequences compared, differing from them by between 1 and 9 bp. The mitogenome base composition is 31.3% A, 16.2% G, 24.0% T, and 28.6% C, with a slight AT bias (55.3%). Eleven of thirteen protein-coding genes in *O. lepturum* started with a typical ATG codon, except for the COI and ATP6 genes, which were GTG. Seven protein-coding genes ended with the termination codon TAA (ND1, CO1, ATP8, ATP6, ND4L, ND5, and ND6), while the remaining six genes terminated with a single base T (Table 1). The 22 tRNA genes ranged in size from 67 to 76 bp, and the length of tRNA^{Cys} gene (67 bp) was the shortest, whereas the longest were the tRNA^{Leu} and tRNA^{Lys} genes (76 bp). The noncoding control region (D-loop) is located between tRNA^{Phe} and tRNA^{Pro} and is 937 bp in length (Table 1). The genes in the *O. lepturum* mitogenome were closely arranged with overlapping and interval phenomena. There is a total of 22 bp in overlaps between six gene junctions, and each single overlap ranged in size from 1 to 7 bp, with the longest overlapping region (7 bp) located between ATP8/ATP6 and ND4L/ND4, 4 bp of overlapping regions between ND5 and ND6 and fewer than 2 bp at the remaining three positions (Table 1). However, there are 11 intergenic spacer regions ranging in size from 1 to 35 bp (67 bp in total), and the largest spacer (35 bp) is located between tRNA^{Asn} and tRNA^{Cys} (Table 1).

We assessed the amino acid (AA) codon usage by calculating the relative synonymous codon usage (RSCU) values in 13 PCGs, which are shown in Fig. 2a. A total of 3803 codons were encoded by 13 PCGs, and the most frequently used codons were CUA (4.7%), ACA (3.4%) and ACC (3.1%). In the PCGs of the *O. lepturum* mitogenome, the AA components and their codon usage reveal that one codon family (Trp) represents more than 100 codons per thousand codons (CDpT), three codon families (Cys, Met and Ser2) between 50 CDpT and 100 CDpT, and the other twenty codon families less than 50 CDpT (Fig. 2b).

Comparative analysis of nucleotide base composition showed that the composition of *O. lepturum* is identical to that of the other *Onychostoma* fishes, and most of the




Figure 1. Gene map of the mitochondrial genome of *Onychostoma lepturum*. Two rRNA genes (in red); 13 coding genes (in green); 22 tRNA genes and control region (D-loop) (in yellow). (Color figure online).

genes within these *Onychostoma* species maintain a consistent position and direction (Table 2). The analysis of our complete dataset nucleotide frequencies confirmed a bias for A, showing averages of 31.3% of A, 16.2% G, 24.0% T, and 28.6% C, with a slight AT bias (55.3%), which is similar to the patterns found in most fish mitogenomes. The PCGs have a slightly higher AT content (54.7%) than ribosomal RNA genes (53.6%) (Table 2). Furthermore, the A+T content at codon site 2 of PCGs (58.7%) was slightly higher than that at site 3 (58.5%), while the A+T content at the first site was the lowest (47.1%). The A+T-rich region was the main noncoding region (control region) of mitogenomes, has the highest A+T content (66.5%), which was significantly higher than other genes of the mitogenome and was typical of animal mitochondrial genomes (Zhang and Hewitt 1997; Zhou et al. 2017). The AT and GC skews of *O. lepturum* are 0.132 and -0.277, respectively (Table 2). The GC-skew of all genes was negative and revealed a similar pattern of base composition behavior for the *O. lepturum* mitog-

Table 1. Organization of the mitochondrial genome of *Onychostoma lepturum*.

| Locus | Position | | | Codon | | | | |
|---------------------|----------|-------|----------|-------|------|------------|------------------------------------|---------------------|
| | start | stop | size(bp) | start | stop | anti-codon | intergenic nucleotide ⁺ | strand ⁺ |
| tRNA ^{Phe} | 1 | 69 | 69 | | | GAA | 0 | H |
| 12s rRNA | 70 | 1030 | 961 | | | | 0 | H |
| tRNA ^{Val} | 1031 | 1102 | 72 | | | TAC | 0 | H |
| 16s rRNA | 1103 | 2783 | 1681 | | | | 0 | H |
| tRNA ^{Leu} | 2784 | 2859 | 76 | | | TAA | 0 | H |
| <i>ND1</i> | 2860 | 3834 | 975 | ATG | TAA | | 0 | H |
| tRNA ^{Leu} | 3840 | 3911 | 72 | | | GAT | 4 | H |
| tRNA ^{Gln} | 3910 | 3980 | 71 | | | TTG | -2 | L |
| tRNA ^{Met} | 3983 | 4051 | 69 | | | CAT | 2 | H |
| <i>ND2</i> | 4052 | 5096 | 1045 | ATG | T- | | 0 | H |
| tRNA ^{Tyr} | 5097 | 5168 | 72 | | | TCA | 0 | H |
| tRNA ^{Ala} | 5171 | 5239 | 69 | | | TGC | 2 | L |
| tRNA ^{Asn} | 5241 | 5313 | 73 | | | GTT | 1 | L |
| tRNA ^{Cys} | 5349 | 5415 | 67 | | | GCA | 35 | L |
| tRNA ^{Tyr} | 5415 | 5486 | 72 | | | GTA | -1 | L |
| <i>COI</i> | 5488 | 7038 | 1551 | GTG | TAA | | 1 | H |
| tRNA ^{Ser} | 7039 | 7109 | 71 | | | TGA | 0 | L |
| tRNA ^{Asp} | 7113 | 7184 | 72 | | | GTC | 3 | H |
| <i>COII</i> | 7197 | 7887 | 691 | ATG | T- | | 12 | H |
| tRNA ^{Lys} | 7888 | 7963 | 76 | | | TTT | 0 | H |
| <i>ATP8</i> | 7965 | 8129 | 165 | ATG | TAG | | 1 | H |
| <i>ATP6</i> | 8123 | 8805 | 683 | GTG | TA- | | -7 | H |
| <i>COIII</i> | 8806 | 9590 | 785 | ATG | TAA | | 0 | H |
| tRNA ^{Gly} | 9591 | 9662 | 72 | | | TCC | 0 | H |
| <i>ND3</i> | 9663 | 10011 | 349 | ATG | T- | | 0 | H |
| tRNA ^{Arg} | 10012 | 10081 | 70 | | | TCG | 0 | H |
| <i>ND4L</i> | 10082 | 10378 | 297 | ATG | TAA | | 0 | H |
| <i>ND4</i> | 10372 | 11752 | 1381 | ATG | T- | | -7 | H |
| tRNA ^{His} | 11753 | 11821 | 69 | | | GTG | 0 | H |
| tRNA ^{Ser} | 11822 | 11890 | 69 | | | GCT | 0 | H |
| tRNA ^{Leu} | 11892 | 11964 | 73 | | | TAG | 1 | H |
| <i>ND5</i> | 11965 | 13788 | 1824 | ATG | TAA | | 0 | H |
| <i>ND6</i> | 13785 | 14306 | 522 | ATG | TAG | | -4 | L |
| tRNA ^{Glu} | 14307 | 14375 | 69 | | | TTC | 0 | L |
| <i>Cytb</i> | 14381 | 15521 | 1141 | ATG | T- | | 5 | H |
| tRNA ^{Thr} | 15522 | 15594 | 73 | | | TGT | 0 | H |
| tRNA ^{Pro} | 15594 | 15663 | 70 | | | TGG | -1 | L |
| D-loop | 15664 | 16598 | 935 | | | | 0 | H |

Table 2. Nucleotide composition of the *Onychostoma lepturum* mitochondrial genome.

| | Length (bp) | T% | C% | A% | G% | A+T% | AT-skew% | GC-skew% |
|--------------------------------|-------------|------|------|------|------|------|----------|----------|
| Genome | 16598 | 24 | 28.6 | 31.3 | 16.2 | 55.3 | 0.132 | 0.277 |
| PCGs | 11409 | 25.7 | 29.4 | 29 | 15.8 | 54.7 | 0.06 | -0.3 |
| 1 st codon position | 3807 | 20.3 | 27 | 26.8 | 25.8 | 47.1 | 0.138 | -0.023 |
| 2 nd codon position | 3802 | 40.1 | 27.7 | 18.6 | 13.6 | 58.7 | -0.366 | -0.341 |
| 3 rd codon position | 3800 | 16.8 | 33.6 | 41.7 | 8 | 58.5 | 0.423 | -0.615 |
| rRNA | 2642 | 19.1 | 25.8 | 34.5 | 20.6 | 53.6 | 0.287 | -0.112 |
| tRNA | 1566 | 26.6 | 21.5 | 28.2 | 23.7 | 54.8 | 0.029 | 0.049 |
| D-loop | 935 | 32.5 | 20.3 | 34 | 13.2 | 66.5 | 0.022 | -0.212 |

enome, except for tRNA which was positive (0.049) (Table 2). Strand asymmetry in nucleotide composition is usually reflected by AT and GC skews, which is a remarkable feature of animal mitochondrial genomes (Wei et al. 2010). The mitogenome contents of 11 *Onychostoma* species were calculated and showed an A+T bias, ranging




Figure 2. Comparison of codon usage in mitochondrial genomes of *Onychostoma lepturum* **a** Relative synonymous codon usage (RSCU) in the *Onychostoma lepturum* mitogenome. Codon families are provided on the X-axis, and the RSCU values, on the Y-axis **b** Codon distribution in the *Onychostoma lepturum* mitogenome. CDspT, codons per thousand codons. Codon families are provided on the X-axis.

from 55.3% (*O. lepturum*) to 56.6% (*O. barbatulum*) (Suppl. material 1: Table S1). The analysis of *Onychostoma* mitochondria populations showed distinct skew patterns, in which AT was positive and GC was negative (Suppl. material 1: Table S1).

The 22 tRNA genes in the *O. lepturum* mitogenome are interspersed between rRNA and protein-coding genes, with sizes ranging from 67 to 76 bp; tRNA^{Cys} was the shortest (67 bp), while tRNA^{Leu} and tRNA^{Lys} were the longest (76 bp). The arrangement of 8 L-strand encoded and 14 H-strand-encoded tRNA genes is similar to the distributions observed in other *Onychostoma* species. Two rRNA genes were identified on the L-strand in *O. lepturum*, which is similar to the other *Onychostoma* species, with a total length of




Figure 3. Phylogenetic trees derived from Maximum-Likelihood (ML) and Neighbor Joining (NJ) approaches based on whole mitochondrial genomes. The numbers on the nodes are the bootstrap values of ML/NJ. The number after the species name is the GenBank Accession Number.

2642 bp. The 16S rRNA is located between $tRNA^{Val}$ and $tRNA^{Leu}$, with a length of 1681 bp, whereas the 12S rRNA is located between $tRNA^{Phe}$ and $tRNA^{Val}$, with a length of 961 bp. Regarding the two rRNA genes, the GC-skew is slightly negative (-0.112), but the AT-skew is strongly positive (0.287). The total A+T content of the rRNA genes (53.1%) is lower than those of the total tRNA genes (55.0%) and the total PCG genes (55.6%).

Phylogenetic analyses

To further investigate the phylogenetic position of *O. lepturum* within the genera *Acrossocheilus* and *Onychostoma*, the concatenated set of nucleotide sequences of available whole mitochondrial genomes from 10 *Onychostoma* species, 16 *Acrossocheilus* species and *F. brevifilis* were derived for phylogenetic reconstruction, along with *S. hollandi* as an outgroup. The ML, NJ and BI analyses showed the same topology, representing the three main lineages (Clades I, II and III). The phylogenetic tree revealed that Clade I is the *Onychostoma* group, which can be separated into two subclades (subclades I-A and I-B) with strong support (Fig. 3). The phylogenetic po-

sition of *O. lepturum* is closer to that of *O. meridionale* than to that of *O. simum* and *O. gerlachi* in subclade I-A. Subclade I-B clusters with five species of *Onychostoma*, including *O. barbatulum*, *O. barbatum*, *O. fangi*, *O. lini* and *O. macrolepis*. Clade II is the *Acrossocheilus* group and includes 14 species, which is in accordance with traditional classifications based on morphology and previous phylogenetic studies based on whole mitochondrial genomes (Yuan et al. 2015; Hou et al. 2020). Species of *Acrossocheilus*, *Onychostoma* and *F. brevifilis* are included in Clade III, which is considered as an ancestral group, including *A. monticola*, *A. yunnanensis*, *O. rarum*, *O. alticorpus* and *F. brevifilis*.

Discussion

Mitochondrial genome of *O. lepturum*

The total length of the *Onychostoma* mitogenomes ranged between 16,590 (*O. rarum*) and 16,601 bp (*O. simum* and *O. gerlachi*), while that of the *O. lepturum* is a typical closed circular DNA molecule with a length of 16,598 bp (Fig. 1; GenBank No. MK296412), making it similar to other *Onychostoma* sequences, which differ by between 3 and 8 bp (16,590 bp for *O. rarum* and 16,601 bp for *O. simum*, *O. gerlachi*) (Suppl. material 1: Table S1). We suggest that the variability observed in closely related mitogenome length can be caused by variations in tandem repeat elements within the control region (D-Loop); differences in the lengths of intergenic regions are also a likely explanation for the gene overlaps (Fig. 1). The nucleotide composition of the *O. lepturum* mitogenome is highly biased toward A+T (55.3%), which is similar to the values in other *Onychostoma* species and displays strand skewness consistent with asymmetrical mutation pressure (Bielawski and Gold 2002) (Suppl. material 1: Table S1). Of the 22 identified tRNAs, the AT and GC skews were both positive in the *O. lepturum* mitogenome, which is common in *Onychostoma* mitogenomes. Comparative mitogenomic structure, organization and gene arrangement analyses of all *Onychostoma* mitogenomes are conserved without any structural rearrangement. Among *Onychostoma* fishes that also exhibited highly similar nucleotide compositions and codon usage patterns, a slight difference was also observed in some species. Ten of the thirteen total PCGs used ATG as the initiation codon (ND1, ND2, ND4, ND4L, ND5, ND6, COII, COIII, ATP8, and Cyt *b*), whereas the COI gene started with the GTG codon in all *Onychostoma* species. Most protein-coding genes started with a traditional ATG codon except for COI in *Onychostoma* mitogenomes, which is consistent with previous reports for other fish mitogenomes (Satoh et al. 2016). In the ATP6 gene, only *O. lepturum* was found with GTG as the initiation codon, and the other *Onychostoma* species used the typical ATG codon. In contrast, only ND3 held diverse start codons in *Onychostoma* mitogenomes. Five *Onychostoma* species were initiated by GTG, including *O. alticorpus*, *O. rarum*, *O. simum*, *O. gerlachi* and *O. meridionale*. The other *Onychostoma* species was initiated by ATG (Table 3). In addition, six PCGs

(ND1, COI, ATP8, ND4L, ND5 and ND6) harbored the typical termination codons (TAA and TAG) in the *Onychostoma* mitochondrial genome, and the remaining PCGs were terminated with incomplete codons (T- and TA-). The incomplete termination codon T in the *Onychostoma* mitochondrial genome was a common termination codon (ND2, ND3, ND4, COII and Cyt *b*), except for *O. alticorpus*, which was terminated by TAA in the ND3 gene (Table 3); however, the COI, ND4L and ND5 genes used TAA, the ATP8 gene used TAG, while ND1 and ND6 used TAG or TAA (Table 3). ND1 and ND6 are terminated with the codon TAA for most *Onychostoma* species, which is different from the termination codon (TAG) observed for ND1 in *O. barbatulum*, *O. simum* and *O. gerlachi* and ND6 in *O. lepturum*, *O. barbatulum*, *O. alticorpus*, *O. simum*, *O. gerlachi* and *O. meridionale* (Table 3). We suggest that ND1 and ND6 appear to have evolved relatively rapidly in these *Onychostoma* species, and a similar observation has also been reported in a previous study of *Acrossocheilus* mitogenomes (Hou et al. 2020).

Molecular phylogeny of *Onychostoma*

Taki (1975) suggested that two groups were subgenera of the genus *Onychostoma*: *Onychostoma* and *Gymnostomus*. These two groups are divided based on their possession of osseous simple dorsal rays; the subgenus *Gymnostomus* includes those having non-osseous rays, while the subgenus *Onychostoma* includes those with osseous simple dorsal rays (Taki 1975). There are five species (*O. lepturum*, *O. barbatulum*, *O. barbatum*, *O. macrolepis* and *O. alticorpus*) belonging to the subgenus *Gymnostomus*, whereas the remaining species belongs to the subgenus *Onychostoma*. According to the mouth width, mouth-opening shape, and post-labial groove length, previous studies divided members of *Onychostoma* into three groups (Chen 1989; Xin et al. 2009). Our results showed that the *Onychostoma* with osseous simple dorsal rays (*O. simum*, *O. rara*, *O. lini*, *O. meridionale*, *O. gerlachi* and *O. fangi*) could not be successfully clustered together (Fig. 3). In addition, molecular evidence revealed that Clade I, including subclades I-A (*O. lepturum*, *O. meridionale*, *O. simum* and *O. gerlachi*) and subclade I-B (*O. barbatulum*, *O. barbatum*, *O. fangi*, *O. lini* and *O. macrolepis*), comprise a stable monophyletic group.

Comparison was made between the phylogenetic trees constructed by Yang et al. (2015), Wang et al. (2016) and Zheng et al. (2016), whereby input sequences such as mtDNA and nuclear sequences were used for phylogeny. One of the similarities between all studies is that Clade I in the *Onychostoma* group comprises a stable monophyletic group distinct from *Acrossocheilus* and *F. brevifilis*. Following our study, *O. simum* was consistently grouped closely with *O. gerlachi* in the *Onychostoma* group (Clade I-A), as in the previous study based on the mitogenome data (Zhang et al. 2018). However, one distinct difference among these above mentioned studies is that in the trees constructed by Yang et al. (2015), Wang et al. (2016) and Zheng et al. (2016) based on nuclear and mitochondrial genes, *O. simum* was found to be more closely related to *O. alticorpus* and *F. brevifilis*. *Onychostoma simum* was more closely related

Table 3. Composition and skewness in mitogenomes of 12 *Omychostoma* species.

| Species | ND1 | ND2 | COI | COII | ATP8 | ATP6 | COIII | ND3 | ND4L | ND4 | ND5 | ND6 | Cytb | GenBank |
|-----------------------|---------|--------|---------|--------|---------|---------|---------|---------|---------|--------|---------|---------|--------|------------|
| <i>O. lepturum</i> | ATG/TAA | ATG/T- | GTG/TAA | ATG/T- | ATG/TAG | GTG/TA- | ATG/TA- | ATG/T- | ATG/TAA | ATG/T- | ATG/TAA | ATG/TAG | ATG/T- | MT258556 |
| <i>O. macrolepis</i> | ATG/TAA | ATG/T- | GTG/TAA | ATG/T- | ATG/TAG | ATG/TA- | ATG/TA- | ATG/T- | ATG/TAA | ATG/T- | ATG/TAA | ATG/TAA | ATG/T- | KF999680.1 |
| <i>O. lini</i> | ATG/TAA | ATG/T- | GTG/TAA | ATG/T- | ATG/TAG | ATG/TA- | ATG/TA- | ATG/T- | ATG/TAA | ATG/T- | ATG/TAA | ATG/TAA | ATG/T- | JQ343982.1 |
| <i>O. barbatum</i> | ATG/TAA | ATG/T- | GTG/TAA | ATG/T- | ATG/TAG | ATG/TA- | ATG/TA- | ATG/T- | ATG/TAA | ATG/T- | ATG/TAA | ATG/TAA | ATG/T- | JX646870.1 |
| <i>O. fangi</i> | ATG/TAA | ATG/T- | GTG/TAA | ATG/T- | ATG/TAG | ATG/TA- | ATG/TA- | ATG/T- | ATG/TAA | ATG/T- | ATG/TAA | ATG/TAA | ATG/T- | AP011193.1 |
| <i>O. barbatulum</i> | ATG/TAG | ATG/T- | GTG/TAA | ATG/T- | ATG/TAG | ATG/TA- | ATG/TA- | ATG/T- | ATG/TAA | ATG/T- | ATG/TAA | ATG/TAG | ATG/T- | KC896762.1 |
| <i>O. alticorpus</i> | ATG/TAA | ATG/T- | GTG/TAA | ATG/T- | ATG/TAG | ATG/TA- | ATG/TA- | GTG/TAA | ATG/TAA | ATG/T- | ATG/TAA | ATG/TAG | ATG/T- | KC791686.1 |
| <i>O. narum</i> | ATG/TAA | ATG/T- | GTG/TAA | ATG/T- | ATG/TAG | ATG/TA- | ATG/TA- | GTG/T- | ATG/TAA | ATG/T- | ATG/TAA | ATG/TAA | ATG/T- | KF626377.1 |
| <i>O. sinum</i> | ATG/TAG | ATG/T- | GTG/TAA | ATG/T- | ATG/TAG | ATG/TA- | ATG/TA- | GTG/T- | ATG/TAA | ATG/T- | ATG/TAA | ATG/TAG | ATG/T- | KF021233.1 |
| <i>O. gerlachi</i> | ATG/TAG | ATG/T- | GTG/TAA | ATG/T- | ATG/TAG | ATG/TA- | ATG/TA- | GTG/T- | ATG/TAA | ATG/T- | ATG/TAA | ATG/TAG | ATG/T- | KP244449.1 |
| <i>O. meridionale</i> | ATG/TAA | ATG/T- | GTG/TAA | ATG/T- | ATG/TAG | ATG/TA- | ATG/TA- | GTG/T- | ATG/TAA | ATG/T- | ATG/TAA | ATG/TAG | ATG/T- | AP011351.1 |
| <i>O. brevisbarba</i> | ATG/TAA | ATG/T- | GTG/TAA | ATG/T- | ATG/TAG | ATG/TA- | ATG/TA- | ATG/T- | ATG/TAA | ATG/T- | ATG/TAA | ATG/TAG | ATG/T- | MG523272.1 |

to *O. gerlachi* and was distributed in the Lancang Jiang and Red River basins (Song et al. 2018), as described in a previous study based on the biogeography of *Onychostoma* (Zhang et al. 2018). The resulting relationships are not consistent with earlier conclusions based on morphological characteristics (Taki 1975; Xin et al. 2009) or on nuclear and mitochondrial genes (Wang et al. 2007; Yang et al. 2016; Zheng et al. 2016). However, the single-gene evolutionary tree of the COI gene is inconsistent with that of combined complete mitochondrial genome, suggesting that *O. simum* (KF021233.1) were incorrectly identified. According to NCBI data and Wang et al. (2013), specimens KF021233.1 did not have an associated collection site (and we suggest that specimens KF021233.1 were *O. gerlachi*). Our results show that *O. lepturum* was found to be more closely related to *O. meridionale* than *O. gerlachi*. *Onychostoma lepturum* and *O. meridionale* occur in Laos and Vietnam, while *O. lepturum* is distributed in the Yuanjiang River in mainland China and Hainan Island. According to the previous biogeographic studies, the Gulf of Tonkin once formed part of the coastal plain of mainland China and the River in Vietnam and Hainan Island drained into the Gulf of Tonkin during Pleistocene glaciations (Zhang et al. 2020).

Recent studies have revealed similar scenarios in the genetic patterns of *Garra orientalis* (Yang et al. 2016), *Aphyocypris normalis* (Huang et al. 2019) and *Opsariichthys hainanensis* (Zhang et al. 2020). Xin et al. (2009) proposed that *O. lini* and *O. barbata* were found to be related closer to *O. macrolepis* based on morphological characters. This result suggested that the phylogenetic relationship of subclade I-B are concordant with the molecular phylogenetic and morphological analyses. In addition, Clade II is the *Acrossocheilus* group, which includes 14 species, comprising a stable monophyletic group distinct from the *Onychostoma* group in the ML, NJ and BI trees (Fig. 3). The present study strongly supported that some species of *Acrossocheilus*, *Onychostoma* and *F. brevifilis* belong to the monophyletic group (Clade III), including *A. monticola*, *A. yunnanensis*, *O. rarum*, *O. alticorpus* and *F. brevifilis*. Interestingly, our results corroborate the previous finding that *O. rarum* is the sister group of *O. alticorpus* (Hoang et al. 2015; Song et al. 2018; Zhang et al. 2018), but Zhai et al. (2020) suggested that *O. rarum* appeared to be more closely related with *O. barbatulum* and *O. barbatum*. We suggested that the specimens of *O. rarum* (NC022869.1) were also misidentified. Moreover, all of these members were previously confirmed as a monophyletic group (Yuan et al. 2015; Hou et al. 2020). Overall, the complete mtDNA sequence of *O. lepturum* provides useful genetic data for addressing further questions in the systematics and evolutionary history of *Onychostoma*, for understanding its molecular diversity and for genetic conservation applications.

Acknowledgements

This work was supported by grants from the Ministry of Science and Technology, Taiwan (MOST 108-2221-E-035-057 and MOST 109-2221-E-035-034). There are no conflicts of interest in this study.

Reference

- Bielawski JP, Gold JR (2002) Mutation patterns of mitochondrial H-and L-strand DNA in closely related cyprinid fishes. *Genetics* 161: 1589–1597.
- Boulenger GA (1900) On the reptiles, batrachians, and fishes collected by the late Mr. John Whitehead in the interior of Hainan. In *Proceedings of the Zoological Society of London* 189: 956–962.
- Chen XL, Yue PQ, Lin RD (1984) Major groups within the family Cyprinidae and their phylogenetic relationships. *Acta zootaxonomica sinica* 9: 424–440.
- Chen Y (1989) Anatomy and phylogeny of the cyprinid fish genus *Onychostoma* Gunther, 1896. *Bulletin of British Museum Nature History (Zoology)* 55: 109–121.
- Chiu YW, Chang CW, Lin HD, Shen KN (2018) The complete mitogenome of the winged argonaut *Argonauta hians* and its phylogenetic relationships in Octopoda. *Conservation Genetics Resources* 10: 359–362. <https://doi.org/10.1007/s12686-017-0824-z>
- Chung HH, Kamar CKA, Lim LWK, Liao Y, Lam TT, Chong YL (2020) Sequencing and Characterisation of Complete Mitogenome DNA for *Rasbora hobelmani* (Cyprinidae) with Phylogenetic Consideration. *Journal of Ichthyology* 60: 90–98. <https://doi.org/10.1134/S0032945220010014>
- Froese RE (2019) FishBase. World Wide Web electronic publication. <http://www.fishbase.org>
- Guindon S, Dufayard J-F, Lefort V, Anisimova M, Hordijk W, Gascuel O (2010) New Algorithms and Methods to Estimate Maximum-Likelihood Phylogenies: Assessing the Performance of PhyML 3.0. *Systematic Biology* 59: 307–321. <https://doi.org/10.1093/sysbio/syq010>
- Hoang HD, Pha, HM, Tran NT (2015) Two new species of shovel-jaw carp *Onychostoma* (Teleostei: Cyprinidae) from southern Vietnam. *Zootaxa*, 3962: 123–138. <https://doi.org/10.11646/zootaxa.3962.1.6>
- Hou XJ, Lin HD, Tang WQ, Liu D, Han CC, Yang JQ (2020) Complete mitochondrial genome of the freshwater fish *Acrossocheilus longipinnis* (Teleostei: Cyprinidae): genome characterization and phylogenetic analysis. *Biologia* 75: 1871–1880. <https://doi.org/10.2478/s11756-020-00440-y>
- Huang SP, Wang FY, Wang TY (2017) Molecular phylogeny of the *Opsariichthys* group (Teleostei: Cypriniformes) based on complete mitochondrial genomes. *Zoological studies* 56: e40.
- Huang XX, Hsu KC, Kang B, Kuo PH, Tsai WH, Liang CM, Lin HD, Wang WK (2019) Population structure of *Aphyocypris normalis*: phylogeography and systematics. *ZooKeys* 872: 77–90. <https://doi.org/10.3897/zookeys.872.33105>
- Kang B, Deng J, Huang X, Chen L, Feng Y (2013) Explaining freshwater fish biogeography: history versus environment versus species personality. *Reviews in Fish Biology and Fisheries* 23: 523–536. <https://doi.org/10.1007/s11160-013-9314-x>
- Kang B, Deng J, Wu Y, Chen L, Zhang J, Qiu H, Lu Y, He D (2014) Mapping China's freshwater fishes: diversity and biogeography. *Fish and Fisheries* 15: 209–230. <https://doi.org/10.1111/faf.12011>
- Kottelat M (2001) Freshwater fishes of Northern Vietnam: a preliminary check-list of the fishes known or expected to occur in northern Vietnam with comments on systematics and nomenclature. Environment and Social Development Unit, East Asia and Pacific Region, World Bank, Washington, 184 pp.

- Kumar S, Stecher G, Li M, Knyaz C, Tamura K (2018) MEGA X: Molecular Evolutionary Genetics Analysis across Computing Platforms. *Molecular Biology and Evolution* 35: 1547–1549. <https://doi.org/10.1093/molbev/msy096>
- Lefort V, Longueville J-E, Gascuel O (2017) SMS: Smart Model Selection in PhyML. *Molecular Biology and Evolution* 34: 2422–2424. <https://doi.org/10.1093/molbev/msx149>
- Perna NT, Kocher TD (1995) Patterns of nucleotide composition at fourfold degenerate sites of animal mitochondrial genomes. *Journal of Molecular Evolution* 41: 353–358. <https://doi.org/10.1007/BF01215182>
- Rambaut A, Drummond AJ, Xie D, Baele G, Suchard MA (2018) Posterior Summarization in Bayesian Phylogenetics Using Tracer 1.7. *Systematic Biology* 67: 901–904. <https://doi.org/10.1093/sysbio/syy032>
- Ronquist F, Huelsenbeck JP (2003) MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19: 1572–1574. <https://doi.org/10.1093/bioinformatics/btg180>
- Satoh TP, Miya M, Mabuchi K, Nishida M (2016) Structure and variation of the mitochondrial genome of fishes. *BMC Genomics* 17: e719. <https://doi.org/10.1186/s12864-016-3054-y>
- Shan X, Lin R, Yue P, Chu X (2000) *Onychostoma*. In: Yue P (Ed.) *Fauna Sinica, Osteichthyes*. Science Press, Beijing, Cypriniformes III, 126–147.
- Song X-L, Cao L, Zhang E (2018) *Onychostoma brevibarba*, a new cyprinine fish (Pisces: Teleostei) from the middle Chang Jiang basin in Hunan Province, South China. *Zootaxa* 4410: 147–163. <https://doi.org/10.11646/zootaxa.4410.1.8>
- Taki Y (1975) Cyprinid fishes of the genera *Onychostoma* and *Scaphiodonichthys* from Upper Laos, with remarks on the dispersal of the genera and their allies. *Japanese Journal of Ichthyology* 22: 143–150.
- Wang JP, Cheng CY, Ho CW, Ueng YT (2015) Complete mitochondrial DNA genome of *Onychostoma sima* (Cypriniformes: Cyprinidae). *Mitochondrial DNA* 26: 135–136. <https://doi.org/10.3109/19401736.2013.815173>
- Wang X, Li J, He S (2007) Molecular evidence for the monophyly of East Asian groups of Cyprinidae (Teleostei: Cypriniformes) derived from the nuclear recombination activating gene 2 sequences. *Molecular Phylogenetics and Evolution* 42: 157–170. <https://doi.org/10.1016/j.ympev.2006.06.014>
- Wang X, Gan X, Li J, Chen Y, He S (2016) Cyprininae phylogeny revealed independent origins of the Tibetan Plateau endemic polyploid cyprinids and their diversifications related to the Neogene uplift of the plateau. *Science China Life Sciences* 59: 1149–1165. <https://doi.org/10.1007/s11427-016-0007-7>
- Wei SJ, Shi M, Chen XX, Sharkey MJ, Achterberg C van, Ye GY, He JH (2010) New views on strand asymmetry in insect mitochondrial genomes. *PLoS ONE* 5(9): e12708. <https://doi.org/10.1371/journal.pone.0012708>
- Xin Q (2008) Taxonomic revision of species and phylogenetic analysis of interspecific relationships within the cyprinid genus *Onychostoma sensu lato* Günther, 1896. Dissertation. The Graduate School of Chinese Academy of Sciences.
- Xin Q, Zhang E, Cao WX (2009) *Onychostoma virgatum*, a new species of cyprinid fish (Pisces: Teleostei) from southern Anhui Province, South China. *Ichthyological Exploration of Freshwaters* 20: e255.

- Yang JQ, Hsu KC, Liu ZZ, Su LW, Kuo PH, Tang WQ, Zhou ZC, Liu D, Bao BL, Lin HD (2016) The population history of *Garra orientalis* (Teleostei: Cyprinidae) using mitochondrial DNA and microsatellite data with approximate Bayesian computation. *BMC Evolutionary Biology* 16: e.73. <https://doi.org/10.1186/s12862-016-0645-9>
- Yang L, Sado T, Vincent Hirt M, Pasco-Viel E, Arunachalam M, Li J, Wang X, Freyhof J, Saitoh K, Simons AM, Miya M, He S, Mayden RL (2015) Phylogeny and polyploidy: Resolving the classification of cyprinine fishes <https://doi.org/10.1016/j.ympev.2015.01.014> (Teleostei: Cypriniformes). *Molecular Phylogenetics and Evolution* 85: 97–116.
- Yuan LY, Liu XX, Zhang E (2015) Mitochondrial phylogeny of Chinese barred species of the cyprinid genus *Acrossocheilus Oshima*, 1919 (Teleostei: Cypriniformes) and its taxonomic implications. *Zootaxa* 4059: 151–168. <https://doi.org/10.11646/zootaxa.4059.1.8>
- Yue P (2000) *Fauna Sinica. Osteichthyes. Cypriniformes III*. Science Press. Beijing, 661 pp.
- Zhai D, Xie Z, Wang Y, Yu J, Chen Y, Xia M, Liu H, Xiong F (2020) Complete mitochondrial genome of *Onychostoma leptura* and phylogenetic analysis of *Onychostoma*. *Mitochondrial DNA Part B* 5: 2297–2298. <https://doi.org/10.1080/23802359.2020.1772692>
- Zhang C, Cheng Q, Geng H, Lin AH, Wang HY (2018) Molecular phylogeny of *Onychostoma* (Cyprinidae) based on mitochondrial genomes. *Acta Hydrobiologica Sinica* 42: 512–516.
- Zhang DX, Hewitt GM (1997) Insect mitochondrial control region: A review of its structure, evolution and usefulness in evolutionary studies. *Biochemical Systematics and Ecology* 25: 99–120. [https://doi.org/10.1016/S0305-1978\(96\)00042-7](https://doi.org/10.1016/S0305-1978(96)00042-7)
- Zhang WJ, Wang JJ, Li C, Chen JQ, Li W, Jiang SY, Hsu KC, Zhao M, Lin HD, Zhao J (2020) Spatial genetic structure of *Opsariichthys hainanensis* in South China. *Mitochondrial DNA Part A* 31: 98–107. <https://doi.org/10.1080/24701394.2020.1741564>
- Zheng LP, Yang JX, Chen XY (2016) Molecular phylogeny and systematics of the Barbinae (Teleostei: Cyprinidae) in China inferred from mitochondrial DNA sequences. *Biochemical Systematics and Ecology* 68: 250–259. <https://doi.org/10.1016/j.bse.2016.07.012>
- Zhou TQ, Lin HD, Hsu KC, Kuo PH, Wang WK, Tang WQ, Liu D, Yang JQ (2017) Spatial genetic structure of the cyprinid fish *Onychostoma lepturum* on Hainan Island. *Mitochondrial DNA Part A* 28: 901–908. <https://doi.org/10.1080/24701394.2016.1209193>

Supplementary material I

Table S1

Authors: I-Chen Wang, Hung-Du Lin, Chih-Ming Liang, Chi-Chun Huang, Rong-Da Wang, Jin-Quan Yang, Wei-Kuang Wang

Data type: DNA

Explanation note: Composition and skewness in the genus *Onychostoma* mitogenomes.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.1005.57592.suppl1>